



Near minimally normed spline quasi-interpolants on uniform partitions

Domingo Barrera-Rosillo, Maria José Ibañez-Pérez, Paul Sablonnière, Driss Sbibih

► To cite this version:

Domingo Barrera-Rosillo, Maria José Ibañez-Pérez, Paul Sablonnière, Driss Sbibih. Near minimally normed spline quasi-interpolants on uniform partitions. *Journal of Computational and Applied Mathematics*, 2005, 181 (1), pp.211-233. 10.1016/j.cam.2004.11.031 . hal-00001430v2

HAL Id: hal-00001430

<https://hal.science/hal-00001430v2>

Submitted on 8 Nov 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Near minimally normed spline quasi-interpolants on uniform partitions

D. Barrera^a M. J. Ibáñez^a P. Sablonnière^b D. Sbibih^c

^a*Departamento de Matemática Aplicada, Universidad de Granada,
Campus universitario de Fuentenueva s/n, 18071 Granada, Spain*

^b*INSA de Rennes, 20 Avenue des Buttes de Coësmes, CS 14315,
35043 Rennes Cedex, France*

^c*Département de Mathématiques et Informatique, Faculté des Sciences,
Université Mohammed 1er, 60000 Oujda, Morocco*

Abstract

Spline quasi-interpolants are local approximating operators for functions or discrete data. We consider the construction of discrete and integral spline quasi-interpolants on uniform partitions of the real line having small infinite norms. We call them near minimally normed quasi-interpolants: they are exact on polynomial spaces and minimize a simple upper bound of their infinite norms. We give precise results for cubic and quintic quasi-interpolants. Also the quasi-interpolation error is considered, as well as the advantage that these quasi-interpolants present when approximating functions with isolated discontinuities.

1 Introduction

Usually, the construction of spline approximants requires the solution of linear systems. Spline quasi-interpolants (abbr. QIs) are local approximants avoiding this problem, so they are very convenient in practice. For a given nondecreasing biinfinite sequence $\mathbf{t} = (t_i)_{i \in \mathbb{Z}}$ such that $|t_i| \rightarrow +\infty$ as $i \rightarrow \pm\infty$, and $t_i < t_{i+k}$ for all i , let $N_{i,k}$ be the i th B-spline of order $k \in \mathbb{N}$, and $S_{k,\mathbf{t}}$ the linear space spanned by these B-splines (see e.g. [27]). In [4,5], quasi-interpolants $Qf = \sum_{i \in \mathbb{Z}} \lambda_i(f) N_{i,k}$ were constructed, where the linear form λ_i uses both functional and derivative values (see [6, chapter XII], [13, chapter 5], and [27, chapter 6], as well as [22,26] for the uniform case). In [17], another B-spline quasi-interpolation method is defined with λ_i involving functional values in a divided difference scheme. A discrete QI (abbr. dQI) is obtained when $\lambda_i(f)$ is a finite linear combination of functional values. This problem has been considered in [10] (see also [22–24]). When $\lambda_i(f)$ is the inner product of f

with a linear combination of B-splines, Q is an integral QI (abbr. iQI). For details on this topic, see for example [12,14,20,21,25] (and references therein).

All these QIs are exact on the space $\mathbb{P}_{k-1}$ of polynomials of degree at most $k-1$ (or a subspace of $\mathbb{P}_{k-1}$). In [16], one can find a general construction of univariate spline QIs reproducing the spline space $S_{k,t}$.

Once the QI is constructed, a standard argument (see e.g. [13, p. 144]) shows that, if $\mathcal{S}$ is the reproduced space, then $\|f - Qf\|_\infty \leq (1 + \|Q\|_\infty) \text{dist}(f, \mathcal{S})$. This leads us to the construction of QIs with minimal infinite norm (see [7, p. 73], [22] in the box-spline setting). This problem has been considered in the discrete case in [1,15].

Here, we are interested in spline QIs Q on uniform partitions of the real line such that (a) $\lambda_i(f) := \lambda(f(\cdot + i))$ is either a linear combination of values of f at points lying in a neighbourhood of the support of the i th B-spline, or the inner product of f with a linear combination of B-splines; (b) Q reproduces the polynomials in the spline space; and (c) a simple upper bound of the infinite norm of Q is minimized.

The paper is organized as follows. In Section 2, we establish the exactness conditions on polynomials of the discrete or integral QIs. In Section 3, we define a minimization problem whose solution will be called near minimally normed discrete or integral quasi-interpolant. In Sections 4 and 5, we describe some near minimally normed cubic and quintic discrete and integral QIs respectively. In Section 6, we establish some error bounds for cubic discrete and integral QIs. In Section 7, we show that these QIs diminish the overshoot when applying them to the Heaviside function, so seem suitable for the approximation of functions with isolated discontinuities.

Only the even order B-splines are considered here, because the results for the others are similar.

2 Discrete and integral QIs on uniform partitions

Consider the sequence $\mathbf{t} = \mathbb{Z}$ of integer knots. Let $M := M_{2n}$ be the B-spline of even order $2n$, $n \geq 2$, with support $[-n, n]$ (see e.g. [26,27]). We deal with discrete and integral spline quasi-interpolation operators

$$Qf = \sum_{i \in \mathbb{Z}} \lambda_i(f) M_i,$$

where $M_i := M(\cdot - i)$, and the linear form λ_i has one of the two following forms

- (i) $\lambda_i(f) = \sum_{j=-m}^m \gamma_j f(i+j)$ when Q is a dQI, and
- (ii) $\lambda_i(f) = \sum_{j=-m}^m \gamma_j \langle f, M_{i+j} \rangle$ when Q is an iQI, with $\langle f, g \rangle := \int_{\mathbb{R}} f g$,

for $m \geq n$ and $\gamma_j \in \mathbb{R}$, $-m \leq j \leq m$.

Defining the fundamental function

$$L := L_{2n,m} = \sum_{j=-m}^m \gamma_j M_j, \quad (1)$$

then Qf can be written in the form

$$Qf := Q_{2n,m}f = \sum_{i \in \mathbb{Z}} \chi_i(f) L(\cdot - i), \quad (2)$$

where $\chi_i(f)$ is equal to $f(i)$ or $\langle f, M_i \rangle$. It is clear that L is symmetric with respect the origin if and only if $\gamma_{-j} = \gamma_j$, $j = 1, 2, \dots, m$. In the sequel we only consider *symmetric* dQIs

$$Qf = \sum_{i \in \mathbb{Z}} \left(\gamma_0 f(i) + \sum_{j=1}^m \gamma_j (f(i+j) + f(i-j)) \right) M_i, \quad (3)$$

and *symmetric* iQIs

$$Qf = \sum_{i \in \mathbb{Z}} \left(\gamma_0 \langle f, M_i \rangle + \sum_{j=1}^m \gamma_j (\langle f, M_{i-j} \rangle + \langle f, M_{i+j} \rangle) \right) M_i. \quad (4)$$

Usually, the coefficients γ_j are determined in such a way that the operator Q be exact on the space $\mathbb{P}_{2n-1}$ of polynomials of degree at most $2n-1$. In order to determine the linear constraints that are equivalent to the exactness of Q on $\mathbb{P}_{2n-1}$, we need the expressions of the monomials $e_k(x) := x^k$, $k = 0, 1, \dots, 2n-1$, as linear combinations of the integer translates of the B-spline M (see [8, Th. 6.2.1, p. 464]).

Proposition 1 *For $k = 0, 1, \dots, n-1$, there hold*

$$e_{2k} = \sum_{i \in \mathbb{Z}} \left\{ i^{2k} + \sum_{l=1}^k \frac{(2k)!}{(2k-2l)!} \beta(2l, 2n) i^{2k-2l} \right\} M_i \quad (5)$$

$$e_{2k+1} = \sum_{i \in \mathbb{Z}} \left\{ i^{2k+1} + \sum_{l=1}^k \frac{(2k+1)!}{(2k+1-2l)!} \beta(2l, 2n) i^{2k+1-2l} \right\} M_i \quad (6)$$

where the numbers $\beta(2l, 2n)$ are provided by the expansion

$$\left(\frac{x/2}{\sin(x/2)} \right)^{2n} = \sum_{l=0}^{\infty} (-1)^l \beta(2l, 2n) x^{2l}. \quad (7)$$

Remark 2 In [26], the coefficients in (7) are denoted by $\gamma_{2l}^{(2n)}$, and the fact that $(-1)^l 2^l (2l)! \gamma_{2l}^{(2n)}$ is a polynomial at n of degree l is indicated. On the other hand, in [8, Prop. 6.2.1, p. 464], it was proved that these coefficients are related to the central factorial numbers $t(r, s)$ of the first kind (see [8, 9]). More specifically, for $0 \leq l \leq n-1$ we have (cf. [8, p. 469])

$$\beta(2l, 2n) = \frac{(2n-1-2l)!}{(2n-1)!} t(2n, 2n-2l).$$

Tables 1 and 2 show some values of $t(2n, 2l)$ and $\beta(2l, 2n)$ respectively.

Table 1

Some values of the central factorial numbers of the first kind $t(2n, 2l)$

	n									
l	0	1	2	3	4	5	6	7	8	9
0	1	0	0	0	0	0	0	0	0	0
1	0	1	-1	4	-36	576	-14400	518400	-25401600	1625702400
2	0	0	1	-5	49	-820	21076	-773136	38402064	-2483133696
3	0	0	0	1	-14	273	-7645	296296	-15291640	1017067024
4	0	0	0	0	1	-30	1023	-44473	2475473	-173721912
5	0	0	0	0	0	1	-55	3003	-191620	14739153
6	0	0	0	0	0	0	1	-91	7462	-669188
7	0	0	0	0	0	0	0	1	-140	16422
8	0	0	0	0	0	0	0	0	1	-204
9	0	0	0	0	0	0	0	0	0	1

In order to express the exactness conditions of Q on $\mathbb{P}_{2n-1}$ in terms of the parameters γ_j , we define

$$\begin{aligned} \Gamma_0 &= \gamma_0 + 2 \sum_{j=1}^m \gamma_j, \\ \Gamma_{2l} &= 2 \sum_{j=1}^m j^{2l} \gamma_j, \quad 1 \leq l \leq n-1. \end{aligned}$$

Lemma 3 For each $d \leq n-1$, the d QI Q given by (3) reproduces the monomials e_{2k} , $0 \leq k \leq d$, if and only if

$$\Gamma_{2l} = (2l)! \beta(2l, 2n), \quad l = 0, 1, \dots, d. \quad (8)$$

Table 2

Values of $\beta(2l, 2n)$ for $2 \leq n \leq 9$.

n	l								
	0	1	2	3	4	5	6	7	8
2	1	$-\frac{1}{6}$							
3	1	$-\frac{1}{4}$	$\frac{1}{30}$						
4	1	$-\frac{1}{3}$	$\frac{7}{120}$	$-\frac{1}{140}$					
5	1	$-\frac{5}{12}$	$\frac{13}{144}$	$-\frac{41}{3024}$	$\frac{1}{630}$				
6	1	$-\frac{1}{2}$	$\frac{31}{240}$	$-\frac{139}{6048}$	$\frac{479}{151200}$	$-\frac{1}{2772}$			
7	1	$-\frac{7}{12}$	$\frac{7}{40}$	$-\frac{311}{8640}$	$\frac{37}{6480}$	$-\frac{59}{79200}$	$\frac{1}{12012}$		
8	1	$-\frac{2}{3}$	$\frac{41}{180}$	$-\frac{67}{1260}$	$\frac{2473}{259200}$	$-\frac{4201}{2993760}$	$\frac{266681}{1513512000}$	$-\frac{1}{51480}$	
9	1	$-\frac{3}{4}$	$\frac{23}{80}$	$-\frac{757}{10080}$	$\frac{2021}{134400}$	$-\frac{4679}{1900800}$	$\frac{3739217}{10897286400}$	$-\frac{63397}{1513512000}$	$\frac{1}{218790}$

PROOF. We have

$$\begin{aligned}
Qe_{2k} &= \sum_{i \in \mathbb{Z}} \left\{ \gamma_0 i^{2k} + \sum_{j=1}^m \gamma_j \left((i+j)^{2k} + (i-j)^{2k} \right) \right\} M_i \\
&= \sum_{i \in \mathbb{Z}} \left\{ \Gamma_0 i^{2k} + \sum_{l=1}^k \binom{2k}{2l} \Gamma_{2l} i^{2k-2l} \right\} M_i.
\end{aligned}$$

According to (5) of Proposition 1, Q reproduces $e_0, e_2, \dots$, and e_{2d} if and only if

$$\binom{2k}{2l} \Gamma_{2l} = \frac{(2k)!}{(2k-2l)!} \beta(2l, 2n), \quad l = 0, 1, \dots, d,$$

and the claim follows. $\square$

The exactness relations for the iQI Q given by (4) are more involved. They need the moments of the B-spline M , denoted by

$$\mu_l(2n) := \int_{\mathbb{R}} e_l(x) M(x) dx, \quad l \geq 0.$$

In particular, $\mu_0(2n) = 1$, and $\mu_{2l+1}(2n) = 0$ for $l \geq 0$.

Lemma 4 *For each $d \leq n-1$, the iQI Q given by (4) reproduces the monomials e_{2k} , $0 \leq k \leq d$, if and only if*

$$\sum_{r=0}^k \binom{2k}{2r} \Gamma_{2r} \mu_{2k-2r}(2n) = (2k)! \beta(2k, 2n), \quad k = 0, 1, \dots, d. \quad (9)$$

PROOF. Let $d = 0$. Since

$$\lambda_i(e_0) = \gamma_0 \langle e_0, M_i \rangle + \sum_{j=1}^m \gamma_j \langle e_0, M_{i-j} + M_{i+j} \rangle = \Gamma_0,$$

Q reproduces e_0 if and only if $\Gamma_0 = 1$. Let us assume that Q reproduces the monomials $e_0, e_2, \dots$, and e_{2d} if and only if (9) holds. We will prove that $Qe_{2k} = e_{2k}$ for $k = 0, 1, \dots, d+1$ if and only if

$$\sum_{r=0}^k \binom{2k}{2r} \Gamma_{2r} \mu_{2k-2r}(2n) = (2k)! \beta(2k, 2n), \quad k = 0, 1, \dots, d+1.$$

Let us assume that $Qe_{2k} = e_{2k}$, $k = 0, 1, \dots, d+1$. By hypothesis, (9) holds, and moreover $Qe_{2d+2} = e_{2d+2}$. We have

$$Qe_{2d+2} = \sum_{i \in \mathbb{Z}} q_i(e_{2d+2}) M_i,$$

where

$$\lambda_i(e_{2d+2}) = \gamma_0 \langle e_{2d+2}, M_i \rangle + \sum_{j=1}^m \gamma_j \langle e_{2d+2}, M_{i-j} + M_{i+j} \rangle.$$

Since

$$\langle e_{2d+2}, M_i \rangle = \int_{\mathbb{R}} x^{2d+2} M_i dx = \int_{\mathbb{R}} (x+i)^{2d+2} M(x) dx$$

and

$$\begin{aligned} \langle e_{2d+2}, M_{i-j} + M_{i+j} \rangle &= \int_{\mathbb{R}} x^{2d+2} (M_{i-j}(x) + M_{i+j}(x)) dx \\ &= \int_{\mathbb{R}} \left\{ (x+i+j)^{2d+2} + (x+i-j)^{2d+2} \right\} M(x) dx \\ &= \sum_{l=0}^{d+1} \binom{2d+2}{2l} 2j^{2l} \gamma_j \int_{\mathbb{R}} (x+i)^{2d+2-2l} M(x) dx, \end{aligned}$$

we deduce that

$$\lambda_i(e_{2d+2}) = \Gamma_0 \int_{\mathbb{R}} (x+i)^{2d+2} M(x) dx + \sum_{l=1}^{d+1} \binom{2d+2}{2l} \Gamma_{2l} \int_{\mathbb{R}} (x+i)^{2d+2-2l} M(x) dx.$$

By using the moments of M , we get

$$\int_{\mathbb{R}} (x+i)^{2d+2} M(x) dx = \sum_{r=0}^{d+1} \binom{2d+2}{2r} \mu_{2r}(2n) i^{2d+2-2r},$$

and

$$\int_{\mathbb{R}} (x+i)^{2d+2-2l} M(x) dx = \sum_{s=0}^{d+1-l} \binom{2d+2-2l}{2s} \mu_{2s}(2n) i^{2d+2-2l-2s}.$$

As $\Gamma_0 = 1$, we obtain

$$\begin{aligned}
\lambda_i(e_{2d+2}) &= i^{2d+2} + \sum_{r=1}^{d+1} \binom{2d+2}{2r} \mu_{2r}(2n) i^{2d+2-2r} \\
&\quad + \sum_{l=1}^{d+1} \binom{2d+2}{2l} \Gamma_{2l} \sum_{s=0}^{d+1-l} \binom{2d+2-2l}{2s} \mu_{2s}(2n) i^{2d+2-2l-2s} \\
&= i^{2d+2} + \sum_{r=1}^{d+1} \binom{2d+2}{2r} \mu_{2r}(2n) i^{2d+2-2r} \\
&\quad + \sum_{r=0}^d \left\{ \sum_{s=0}^r \binom{2d-2s}{2r-2s} \binom{2d+2}{2s+2} \mu_{2r-2s}(2n) \Gamma_{2s+2} \right\} i^{2d-2r} \\
&= i^{2d+2} + \sum_{r=0}^d \left\{ \binom{2d+2}{2r} \mu_{2r+2}(2n) + \right. \\
&\quad \left. \sum_{s=0}^r \binom{2d-2s}{2r-2s} \binom{2d+2}{2s+2} \mu_{2r-2s}(2n) \Gamma_{2s+2} \right\} i^{2d-2r}
\end{aligned}$$

Taking into account that

$$\binom{2d-2s}{2r-2s} \binom{2d+2}{2s+2} = \binom{2d+2}{2r+2} \binom{2r+2}{2s+2},$$

we have

$$\lambda_i(e_{2d+2}) = i^{2d+2} + \sum_{r=0}^d \binom{2d+2}{2r+2} T_r i^{2d-2r}$$

with

$$T_r = \mu_{2r+2}(2n) + \sum_{s=0}^r \binom{2r+2}{2s+2} \mu_{2r-2s}(2n) \Gamma_{2s+2}.$$

By hypothesis,

$$\begin{aligned}
\lambda_i(e_{2d+2}) &= i^{2d+2} + \sum_{r=0}^{d-1} \binom{2d+2}{2r+2} (2r+2)! \beta(2r+2, 2n) i^{2d-2r} \\
&\quad + \left\{ \mu_{2d+2}(2n) + \sum_{s=0}^d \binom{2d+2}{2s+2} \mu_{2d-2s}(2n) \Gamma_{2s+2} \right\} \\
&= i^{2d+2} + \sum_{r=1}^d \frac{(2d+2)!}{(2d+2-2r)!} \beta(2r, 2n) i^{2d+2-2r} \\
&\quad + \left\{ \mu_{2d+2}(2n) + \sum_{s=0}^d \binom{2d+2}{2s+2} \mu_{2d-2s}(2n) \Gamma_{2s+2} \right\}
\end{aligned}$$

From (5), we find that

$$e_{2d+2} = \sum_{i \in \mathbb{Z}} \left\{ i^{2d+2} + \sum_{l=1}^{d+1} \frac{(2d+2)!}{(2d+2-2l)!} \beta(2l, 2n) i^{2d+2-2l} \right\} M_i.$$

Hence, $Qe_{2d+2} = e_{2d+2}$ implies that

$$\mu_{2d+2}(2n) + \sum_{s=0}^d \binom{2d+2}{2s+2} \mu_{2d-2s}(2n) \Gamma_{2s+2} = (2d+2)! \beta(2d+2, 2n).$$

This is the equality obtained by taking $k = d+1$ in (9).

The proof of the converse implication is similar. $\square$

As a consequence of the symmetry properties, the discrete and integral QIs also reproduce the monomials of odd degree.

Lemma 5 *Let Q be the dQI given by (3) or the iQI given by (4). Then, for each $d < n$, the fact that $Qe_{2k} = e_{2k}$, $k = 0, 1, \dots, d$, implies that $Qe_{2d+1} = e_{2d+1}$.*

PROOF. Let Q be the dQI given by (3). Then,

$$\begin{aligned} Qe_{2d+1} &= \sum_{i \in \mathbb{Z}} \left\{ \gamma_0 i^{2d+1} + \sum_{j=1}^m \gamma_j \left((i+j)^{2d+1} + (i-j)^{2d+1} \right) \right\} M_i \\ &= \sum_{i \in \mathbb{Z}} \left\{ \Gamma_0 i^{2d+1} + \sum_{l=1}^d \binom{2d+1}{2l} \Gamma_{2l} i^{2d+1-2l} \right\} M_i \end{aligned}$$

According to (8), we have

$$Qe_{2d+1} = \sum_{i \in \mathbb{Z}} \left\{ i^{2d+1} + \sum_{l=1}^d \frac{(2d+1)!}{(2d+1-2l)!} \beta(2l, 2n) i^{2d+1-2l} \right\} M_i.$$

Finally, by (6), $Qe_{2d+1} = e_{2d+1}$.

Now, let Q be the iQI given by (4). As in the proof of Lemma 4,

$$Qe_{2d+1} = \sum_{i \in \mathbb{Z}} \left\{ i^{2d+1} + \sum_{r=0}^{d-1} \binom{2d+1}{2r+2} T_r i^{2d-2r-1} \right\} M_i.$$

Using (9), we get

$$\begin{aligned} Qe_{2d+1} &= \sum_{i \in \mathbb{Z}} \left\{ i^{2d+1} + \sum_{r=0}^{d-1} \binom{2d+1}{2r+2} (2r+2)! \beta(2r+2, 2n) i^{2d-2r-1} \right\} M_i \\ &= \sum_{i \in \mathbb{Z}} \left\{ i^{2d+1} + \sum_{r=1}^d \frac{(2d+1)!}{(2d+1-2r)!} \beta(2r, 2n) i^{2d+1-2r} \right\} M_i \end{aligned}$$

and $Qe_{2d+1} = e_{2d+1}$ follows from (6). $\square$

Remark 6 *The moments of a B-spline can be expressed in terms of central factorial numbers $T(p, q)$ of the second kind (cf. [8, p. 423]). For the even order B-spline, this relationship is*

$$\mu_{2l}(2n) = \frac{T(2l+2n, 2n)}{\binom{2l+2n}{2n}}.$$

Tables 3 and 4 show some values of $T(2n, 2l)$ and $\mu_{2l}(2n)$, respectively.

Let $\Gamma := (\Gamma_0, \Gamma_2, \dots, \Gamma_{2n-2})^T$, and

$$\beta := (1, 2!\beta(2, 2n), \dots, (2n-2)!\beta(2n-2, 2n))^T.$$

Let $A := (a_{ij})_{1 \leq i, j \leq n}$ be the lower triangular matrix given by

$$a_{r+s, s} = \binom{2r+2s-2}{2s-2} \mu_{2r}(2n), \quad r = 0, 1, \dots, n-1, \quad s = 1, 2, \dots, n-r.$$

It is clear that $a_{ii} = 1$, $i = 1, 2, \dots, n$. Its inverse $B = (b_{i,j})_{1 \leq i, j \leq n}$ is also a lower triangular matrix, and $b_{ii} = 1$.

Proposition 7 (1) *The dQI Q given by (3) is exact on $\mathbb{P}_{2n-1}$ if and only if*

$$\gamma_0 + 2 \sum_{j=1}^m \gamma_j = 1 \quad \text{and} \quad 2 \sum_{j=1}^m j^{2l} \gamma_j = (2l)! \beta(2l, 2n), \quad 1 \leq l \leq n-1 \quad (10)$$

(2) *The iQI Q given by (4) is exact on $\mathbb{P}_{2n-1}$ if and only if*

$$\begin{aligned} \gamma_0 + 2 \sum_{j=1}^m \gamma_j &= 1, \\ 2 \sum_{j=1}^m j^{2l} \gamma_j &= \sum_{r=1}^{l+1} (2r-2)! b_{l+1, r} \beta(2r-2, 2n), \quad 1 \leq l \leq n-1 \end{aligned} \quad (11)$$

PROOF. Equations (10) follow from (8) of Lemma 3 with $d = n-1$. According to equations (9) given in Lemma 4, the iQI (4) is exact on $\mathbb{P}_{2n-1}$ if and only if $A\Gamma = \beta$. Equivalently, $\Gamma = B\beta$, and (11) follows. $\square$

Table 3

Some values of the central factorial numbers of the second kind $T(2n, 2l)$.

	n										
l	0	1	2	3	4	5	6	7	8	9	10
0	1	0	0	0	0	0	0	0	0	0	0
1	0	1	1	1	1	1	1	1	1	1	1
2	0	0	1	5	21	85	341	1365	5461	21845	87381
3	0	0	0	1	14	147	1408	13013	118482	1071799	9668036
4	0	0	0	0	1	30	627	11440	196053	3255330	53157079
5	0	0	0	0	0	1	55	2002	61490	1733303	46587905
6	0	0	0	0	0	0	1	91	5278	251498	10787231
7	0	0	0	0	0	0	0	1	140	12138	846260
8	0	0	0	0	0	0	0	0	1	204	25194
9	0	0	0	0	0	0	0	0	0	1	285
10	0	0	0	0	0	0	0	0	0	0	1

Table 4

Values of $\mu(2l, 2n)$ for $2 \leq n \leq 9$.

	l								
n	0	1	2	3	4	5	6	7	8
2	1	$\frac{1}{3}$	$\frac{3}{10}$	$\frac{17}{42}$	$\frac{31}{45}$	$\frac{15}{11}$	$\frac{54612}{1820}$	$\frac{257}{36}$	$\frac{1533}{85}$
3	1	$\frac{1}{2}$	$\frac{7}{10}$	$\frac{32}{21}$	$\frac{13}{3}$	$\frac{651}{44}$	$\frac{63047}{1092}$	$\frac{7483}{30}$	
4	1	$\frac{2}{3}$	$\frac{19}{15}$	$\frac{80}{21}$	$\frac{457}{30}$	$\frac{2455}{33}$	$\frac{164573}{390}$		
5	1	$\frac{5}{6}$	2	$\frac{215}{28}$	$\frac{713}{18}$	$\frac{11095}{44}$			
6	1	1	$\frac{29}{10}$	$\frac{569}{49}$	$\frac{2569}{30}$				
7	1	$\frac{7}{6}$	$\frac{119}{30}$	$\frac{131}{6}$					
8	1	$\frac{4}{3}$	$\frac{26}{5}$						
9	1	$\frac{3}{2}$							

Equations (10) and (11) show that the exactness on $\mathbb{P}_{2n-1}$ of the dQIs or iQIs is expressed by linear systems having a common fixed matrix while the right hand side depends on the type of the QI considered. Therefore, we study jointly the construction of near minimally normed dQIs and iQIs.

3 Near minimally normed discrete and integral QIs

Once the exactness conditions have been established for a symmetric dQI or iQI Q , we consider a simple upper bound of its infinite norm.

Let $\gamma := (\gamma_0, \gamma_1, \dots, \gamma_m)^T$, and $\nu(\gamma) := \nu_{2n,m}(\gamma) = |\gamma_0| + 2 \sum_{j=1}^m |\gamma_j|$. Let f be a function such that $\|f\|_\infty \leq 1$. From (2), we have

$$|Qf| \leq \sum_{i \in \mathbb{Z}} |\chi_i(f)| |L(\cdot - i)|.$$

As, $|\chi_i(f)| \leq 1$, we deduce that

$$|Qf| \leq \Lambda,$$

where Λ is the Lebesgue function

$$\Lambda := \Lambda_{2n,m} = \sum_{i \in \mathbb{Z}} |L(\cdot - i)|.$$

Hence, from (1), we conclude that

$$\|Q\|_\infty \leq \nu(\gamma).$$

We propose the construction of discrete and integral QIs that minimize the bound $\nu(\gamma)$ under the linear constraints consisting of reproducing all monomials in $\mathbb{P}_{2n-1}$. In general, it is difficult to minimize the infinite norm of Q , which is equal to the Chebyshev norm of the Lebesgue function.

Let Q be the QI given by (3) (resp. (4)). Then, let us define

$$V := V_{2n,m} = \left\{ \gamma \in \mathbb{R}^{m+1} : \gamma \text{ satisfies (10) (resp. (11))} \right\}. \quad (12)$$

Problem 8 *Solve* $\text{Min } \{\nu(\gamma), \gamma \in V\}$.

Definition 9 *Let γ be a solution of Problem 8. Then the corresponding discrete (resp. integral) QI given by (3) (resp. (4)) is said to be a near minimally normed (abbr. NMN) dQI (resp. iQI) of order $2n$ relative to m and exact on $\mathbb{P}_{2n-1}$.*

Solving system (10) (resp. (11)) in $\gamma_0, \gamma_{m-n+2}, \dots, \gamma_m$ by Cramer's rule yields

$$\gamma_i = \gamma_i^* - \sum_{j=1}^{m-n+1} c_{i,j} \gamma_j, \quad i \in \{0, m-n+2, \dots, n\} \text{ and } c_{i,j} \in \mathbb{R}, \quad (13)$$

where $(\gamma_0^*, \gamma_{m-n+2}^*, \dots, \gamma_m^*)^T \in \mathbb{R}^n$ is the unique solution of (10) (resp. (11)) when $\gamma_1 = \gamma_2 = \dots = \gamma_{m-n+1} = 0$.

Let $\gamma^* = (\gamma_0^*, 0, \dots, 0, 2\gamma_{m-n+2}^*, \dots, 2\gamma_m^*)^T \in \mathbb{R}^{m+1}$. From (13),

$$\begin{aligned} \nu(\gamma) &= \left| \gamma_0^* - \sum_{j=1}^{m-n+1} c_{0,j} \gamma_j \right| + \sum_{i=1}^{m-n+1} |2\gamma_i| + 2 \sum_{k=n-m+2}^m \left| \gamma_i^* - \sum_{j=1}^{m-n+1} c_{i,j} \gamma_j \right| \\ &= \|\gamma^* - G\tilde{\gamma}\|_1 \end{aligned}$$

with $\tilde{\gamma} = (\gamma_1, \gamma_2, \dots, \gamma_{m-n+1})^T$, and

$$G = \begin{pmatrix} c_{0,1} & c_{0,2} & \cdots & c_{0,m-n+1} \\ 2 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 2 \\ 2c_{m-n+2,1} & 2c_{m-n+2,2} & \cdots & 2c_{m-n+2,m-n+1} \\ \vdots & \vdots & \ddots & \vdots \\ 2c_{m,1} & 2c_{m,2} & \cdots & 2c_{m,m-n+1} \end{pmatrix}. \quad (14)$$

Proposition 10 *γ is a solution of Problem 8 if and only if $G\tilde{\gamma}$ is a best linear l_1 -approximation of γ^* in $\{Gc : c \in \mathbb{R}^{m-n+1}\}$. Therefore, Problem 8 has at least one solution.*

The existence of NMN dQIs and iQIs is guaranteed, but not the uniqueness. A solution can be calculated by using the simplex method, because the minimization problem is equivalent to a linear programming one. Other methods can be used to determine a solution of the minimization problem (cf. [18]).

4 Some examples of near minimally normed discrete QIs

In this section, we give some examples of the discrete QIs studied in the preceding sections. Concretely, we will consider the cubic and quintic cases.

4.1 Cubic discrete QIs

Let us consider the cubic case $n = 2$. Relations (5) and (6) give

$$e_0 = \sum_{i \in \mathbb{Z}} M_i, \quad e_1 = \sum_{i \in \mathbb{Z}} i M_i, \quad e_2 = \sum_{i \in \mathbb{Z}} \left(i^2 - \frac{1}{3} \right) M_i, \quad \text{and} \quad e_3 = \sum_{i \in \mathbb{Z}} (i^3 - i) M_i.$$

Then, for $m \geq 2$, the dQI $Q_{4,m}$ is exact on $\mathbb{P}_3$ if and only if

$$\gamma_0 + 2 \sum_{j=1}^m \gamma_j = 1 \quad \text{and} \quad \sum_{j=1}^m j^2 \gamma_j = -\frac{1}{6}.$$

γ is a solution of this linear system if and only if

$$\gamma_0 = 1 + \frac{1}{3m^2} - \sum_{j=1}^{m-1} 2 \left(1 - \frac{j^2}{m^2}\right) \gamma_j \quad \text{and} \quad \gamma_m = -\frac{1}{6m^2} - \sum_{j=1}^{m-1} \frac{j^2}{m^2} \gamma_j.$$

Proposition 11 *Let $m \geq 1$ and $\tilde{Q}_{4,m}$ be the cubic dQI given by*

$$\tilde{Q}_{4,m} f = \sum_{i \in \mathbb{Z}} \left\{ \left(1 + \frac{1}{3m^2}\right) f(i) - \frac{1}{6m^2} (f(i+m) + f(i-m)) \right\} M_i. \quad (15)$$

Then

- (1) For $m = 1$, $\tilde{Q}_{4,1}$ is the unique dQI exact on $\mathbb{P}_3$ among all the cubic dQIs given by (3).
- (2) For each $m \geq 2$, $\tilde{Q}_{4,m}$ is a NMN cubic dQI exact on $\mathbb{P}_3$.
- (3) When $m \rightarrow +\infty$, $(\tilde{Q}_{4,m})_{m \geq 1}$ converges in the infinite norm to the Schoenberg's operator $S_4 : C(\mathbb{R}) \rightarrow S_{4,\mathbb{Z}}$ defined by $S_4 f = \sum_{i \in \mathbb{Z}} f(i) M_i$.
- (4) The following equalities hold: $\|\tilde{Q}_{4,1}\|_\infty = \frac{11}{9} \simeq 1.2222$, $\|\tilde{Q}_{4,2}\|_\infty = \frac{41}{36} \simeq 1.1389$, and

$$\|\tilde{Q}_{4,m}\|_\infty = 1 + \frac{2}{3m^2} \text{ for all } m \geq 3.$$

PROOF. (1) For $m = 1$, the dQI

$$Qf = \sum_{i \in \mathbb{Z}} (\gamma_0 f(i) + \gamma_1 (f(i+1) + f(i-1))) M_i$$

is exact on $\mathbb{P}_3$ if and only if $\gamma_0 = \frac{4}{3}$ and $\gamma_1 = -\frac{1}{6}$. Thus, $\tilde{Q}_{4,1}$ is obtained.

(2) We will prove that $\left(1 + \frac{1}{3m^2}, 0, \dots, 0, -\frac{1}{6m^2}\right)^T \in \mathbb{R}^{m+1}$ is a solution of Problem 8 (its associated cubic dQI is $\tilde{Q}_{4,m}$). Consider the dQI given by (3). The objective function $\nu_{4,m}(\gamma) = |\gamma_0| + 2 \sum_{i=1}^m |\gamma_i|$ coincides on the affine subspace $V_{4,m}$ with the polyhedral convex function $\|g(\tilde{\gamma})\|_1$, where $g(\tilde{\gamma}) :=$

$\gamma^* - G\tilde{\gamma}$, $\gamma^* = \left(1 + \frac{1}{3m^2}, 0, \dots, 0, -\frac{1}{3m^2}\right)^T$, and the matrix G given by (14) is

$$G = 2 \begin{pmatrix} 1 - \frac{1}{m^2} & 1 - \frac{2^2}{m^2} & \dots & 1 - \frac{(m-1)^2}{m^2} \\ 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \\ \frac{1}{m^2} & \frac{2^2}{m^2} & \dots & \frac{(m-1)^2}{m^2} \end{pmatrix}. \quad (16)$$

Hence, $\left(1 + \frac{1}{3m^2}, 0, \dots, 0, -\frac{1}{3m^2}\right)^T$ is a solution of Problem 8 if and only if $0 \in \mathbb{R}^{m-1}$ minimizes $\|g(\tilde{\gamma})\|_1$. Let $v = (v_0, v_1, \dots, v_m)^T$ be the vector of components $v_0 = 1$, $v_m = -1$ and, for $j = 1, \dots, m-1$,

$$v_j = 2 \frac{j^2}{m^2} - 1.$$

We have $g(0) = \gamma^*$. It is clear that $|v_j| \leq 1$, $0 \leq j \leq m-1$, $\text{sgn } v_0 = \text{sgn } \gamma_0^*$, $\text{sgn } v_m = \text{sgn } \gamma_m^*$, and $v^T G = 0$. Therefore (cf. [28, Th. 1.7, p. 16]), g attains its minimum at $0 \in \mathbb{R}^{m-1}$, and this completes the proof of (2).

(3) Let $f \in C(\mathbb{R})$ such that $\|f\|_\infty \leq 1$. We have

$$\begin{aligned} |\tilde{Q}_{4,m}f - S_4f| &\leq \sum_{i \in \mathbb{Z}} \left| -\frac{1}{6m^2}f(i-m) + \frac{1}{3m^2}f(i) - \frac{1}{6m^2}f(i+m) \right| M_i \\ &\leq \sum_{i \in \mathbb{Z}} \left(\frac{2}{3m^2} \|f\|_\infty \right) M_i \\ &\leq \frac{2}{3m^2}. \end{aligned}$$

Hence, $\|\tilde{Q}_{4,m} - S_4\| \rightarrow 0$ as $m \rightarrow +\infty$, and $(\tilde{Q}_{4,m})_{m \geq 1}$ converges to S_4 .

(4) For each $m \geq 4$, the supports of M_{-m} , M , and M_m have pairwise disjoint interiors. Therefore, the Lebesgue function $\tilde{\Lambda}_{4,m}$ corresponding to $\tilde{Q}_{4,m}$ can be written as

$$\tilde{\Lambda}_{4,m} = \sum_{i \in \mathbb{Z}} \left(\frac{1}{6m^2} M_{i+m} + \frac{3m^2+1}{3m^2} M_i + \frac{1}{6m^2} M_{i-m} \right),$$

i.e., $\tilde{\Lambda}_{4,m}$ is a constant function equal to $1 + \frac{2}{3m^2}$.

Let $m = 1$. By using the Bernstein-Bézier representation (see e.g. [11, chapter 1]), one can prove that

$$\|\tilde{Q}_{4,1}\|_{\infty} = \tilde{\Lambda}_{4,1}\left(\frac{1}{2}\right).$$

Thus, we obtain

$$\|\tilde{Q}_{4,1}\|_{\infty} = 2 \left(\left| \tilde{L}_{4,1}\left(\frac{1}{2}\right) \right| + \left| \tilde{L}_{4,1}\left(\frac{3}{2}\right) \right| + \left| \tilde{L}_{4,1}\left(\frac{5}{2}\right) \right| \right).$$

Since M is a symmetric function supported on $[-2, 2]$, $M\left(\frac{1}{2}\right) = \frac{13}{48}$, and $M\left(\frac{3}{2}\right) = \frac{1}{48}$, we get

$$\tilde{L}_{4,1}\left(\frac{1}{2}\right) = \frac{5}{9}, \quad \tilde{L}_{4,1}\left(\frac{3}{2}\right) = -\frac{5}{96}, \quad \text{and} \quad \tilde{L}_{4,1}\left(\frac{5}{2}\right) = -\frac{1}{288}.$$

The claim follows.

The values of $\|\tilde{Q}_{4,2}\|_{\infty}$ and $\|\tilde{Q}_{4,3}\|_{\infty}$ can be computed in a similar way. $\square$

Remark 12 *A detailed study of $\|\gamma^* - G\tilde{\gamma}\|_1$ shows that this function attains its absolute minimum at $0 \in \mathbb{R}^{m-1}$. Then, for $m \geq 2$ fixed, $\tilde{Q}_{4,m}$ is the unique NMN dQI exact on $\mathbb{P}_3$.*

The sequence $(\|\tilde{Q}_{4,m}\|_{\infty})_{m \geq 1}$ is strictly decreasing, and the new cubic dQIs are also better than the classical $\tilde{Q}_{4,1}$ with respect to the infinite norm.

4.2 Quintic discrete QIs

The quintic case ($n = 3$) can be studied in a similar way. Now, we have

$$\begin{aligned} e_0 &= \sum_{i \in \mathbb{Z}} M_i, \quad e_1 = \sum_{i \in \mathbb{Z}} i M_i, \quad e_2 = \sum_{i \in \mathbb{Z}} \left(i^2 - \frac{1}{2}\right) M_i, \\ e_3 &= \sum_{i \in \mathbb{Z}} \left(i^3 - \frac{3}{2}i\right) M_i, \quad e_4 = \sum_{i \in \mathbb{Z}} \left(i^4 - 3i^2 + \frac{4}{5}i\right) M_i, \quad e_5 = \sum_{i \in \mathbb{Z}} \left(i^5 - 5i^3 + 4i\right) M_i. \end{aligned}$$

For each $m \geq 3$, the dQI

$$Q_{6,m}f = \sum_{i \in \mathbb{Z}} \left(\gamma_0 f(i) + \sum_{j=1}^m \gamma_j (f(i+j) + f(i-j)) \right) M_i$$

is exact on $\mathbb{P}_5$ if and only if

$$a_0 + 2 \sum_{j=1}^m \gamma_j = 1, \quad \sum_{j=1}^m j^2 \gamma_j = -\frac{1}{4} \quad \text{and} \quad \sum_{j=1}^m j^4 \gamma_j = \frac{2}{5}.$$

The general solution of these equations satisfies

$$\gamma_{m-1} = \gamma_{m-1}^* - \sum_{j=1}^{m-2} \frac{j^2 (m^2 - j^2)}{(m-1)^2 (2m-1)} \gamma_j, \quad \gamma_m = \gamma_m^* - \sum_{j=1}^{m-2} \frac{j^2 (j^2 - (m-1)^2)}{m^2 (2m-1)} \gamma_j,$$

and

$$\gamma_0 = \gamma_0^* - 2 \sum_{j=1}^{m-2} \left(1 - \frac{j^2}{(m-1)^2 m^2} \right) \gamma_j,$$

where

$$\gamma_0^* = 1 + \frac{8 + 5(m^2 + (m-1)^2)}{10(m-1)^2 m^2}, \quad \gamma_{m-1}^* = -\frac{5m^2 + 8}{20(m-1)^2 (2m-1)},$$

and

$$\gamma_m^* = \frac{8 + 5(m-1)^2}{20m^2 (2m-1)}.$$

So, minimizing the objective function

$$\nu_{6,m}(\gamma) = |\gamma_0| + 2 \sum_{j=1}^m |\gamma_j|$$

subject to the exactness conditions is equivalent to the unconstrained minimization of

$$\begin{aligned} g(\tilde{\gamma}) = & \left| \gamma_0^* - 2 \sum_{j=1}^{m-2} \left(1 - \frac{j^2}{(m-1)^2 m^2} \right) \gamma_j \right| + 2 \sum_{j=1}^{m-2} |\gamma_j| \\ & + 2 \left| \gamma_{m-1}^* - \sum_{j=1}^{m-2} \frac{j^2 (m^2 - j^2)}{(m-1)^2 (2m-1)} \gamma_j \right| + 2 \left| \gamma_m^* - \sum_{j=1}^{m-2} \frac{j^2 (j^2 - (m-1)^2)}{m^2 (2m-1)} \gamma_j \right| \end{aligned}$$

The following result can be proved as in the cubic case.

Proposition 13 *Let $m \geq 2$ and $\tilde{Q}_{6,m} f = \sum_{i \in \mathbb{Z}} \tilde{\lambda}_i(f) M_i$ be the quintic dQI given by*

$$\tilde{\lambda}_i(f) = \gamma_0^* f(i) + \gamma_{m-1}^* (f(i+m-1) + f(i-m+1)) + \gamma_m^* (f(i+m) + f(i-m)).$$

Then

- (1) *For $m = 2$, $\tilde{Q}_{6,2}$ is the unique dQI exact on $\mathbb{P}_5$ among all the quintic dQIs given by (3).*
- (2) *For each $m \geq 3$, $\tilde{Q}_{6,m}$ is the unique NMN quintic dQI exact on $\mathbb{P}_5$.*
- (3) *When $m \rightarrow +\infty$, $(\tilde{Q}_{6,m})_{m \geq 1}$ converges in the infinite norm to the Schoenberg's operator $S_6 : C(\mathbb{R}) \rightarrow S_{6,\mathbb{Z}}$ defined by $S_6 f = \sum_{i \in \mathbb{Z}} f(i) M_i$.*

(4) The following equalities hold: $\|\tilde{Q}_{6,2}\|_\infty = \frac{37183}{28800} \simeq 1.2911$, $\|\tilde{Q}_{6,3}\|_\infty = \frac{61}{48} \simeq 1.2708$, $\|\tilde{Q}_{6,4}\|_\infty = \frac{23152727}{19353600} \simeq 1.1963$, $\|\tilde{Q}_{6,5}\|_\infty = \frac{853}{720} \simeq 1.1847$, and

$$\|Q_{6,m}\|_\infty = 1 + \frac{8 + 5m^2}{5(m-1)^2(2m-1)} \quad \text{for all } m \geq 6.$$

Once again, for each $m \geq 3$ the corresponding NMN quintic dQI is better than the classical one w.r.t. the infinite norm.

5 Some examples of near minimally normed integral QIs

In this section, we briefly describe some results on NMN cubic and quintic iQIs.

5.1 Cubic integral QIs

In the cubic case, equations (11) become

$$\gamma_0 + 2 \sum_{j=1}^m \gamma_j = 1 \quad \text{and} \quad \sum_{j=1}^m j^2 \gamma_j = -\frac{1}{3}.$$

A NMN cubic iQI relative to m and exact on $\mathbb{P}_3$ corresponds to a $\gamma = (\gamma_0, \gamma_1, \dots, \gamma_m)^T$ such that $\tilde{\gamma} = (\gamma_1, \dots, \gamma_{m-1})^T$ minimizes $g(\tilde{\gamma}) = \|\gamma^* - G\tilde{\gamma}\|_1$, where the matrix G is given by (16), and $\gamma^* = \left(1 + \frac{2}{3m^2}, 0, \dots, 0, -\frac{2}{3m^2}\right)^T$. A similar proof as that of Proposition 11 shows that g achieves its minimum at $0 \in \mathbb{R}^{m-1}$. Hence, we have the following result.

Proposition 14 *Let $m \geq 1$ and $\tilde{Q}_{4,m}$ be the cubic iQI given by*

$$\tilde{Q}_{4,m}f = \sum_{i \in \mathbb{Z}} \left\{ \left(1 + \frac{2}{3m^2}\right) \langle f, M_i \rangle - \frac{1}{3m^2} \langle f, M_{i-m} + M_{i+m} \rangle \right\} M_i. \quad (17)$$

Then

- (1) *For $m = 1$, $\tilde{Q}_{4,1}$ is the unique iQI exact on $\mathbb{P}_3$ among all the cubic iQIs given by (4).*
- (2) *For each $m \geq 2$, $\tilde{Q}_{4,m}$ is the unique NMN cubic iQI relative to m and exact on $\mathbb{P}_3$.*
- (3) *When $m \rightarrow +\infty$, $(\tilde{Q}_{4,m})_{m \geq 1}$ converges in the infinite norm to the Schoen-*

berg's operator defined by

$$S_4 f = \sum_{i \in \mathbb{Z}} \langle f, M_i \rangle M_i.$$

(4) The following equalities hold: $\|\tilde{Q}_{4,1}\|_\infty = \frac{55}{36} \simeq 1.5278$, $\|\tilde{Q}_{4,2}\|_\infty = \frac{23}{18} \simeq 1.2778$, $\|\tilde{Q}_{4,3}\|_\infty = \frac{31}{27} \simeq 1.1481$, and

$$\|\tilde{Q}_{4,m}\|_\infty = 1 + \frac{4}{3m^2} \text{ for all } m \geq 4.$$

5.2 Quintic integral QIs

The quintic case is qualitatively different. The iQI is exact on $\mathbb{P}_5$ if and only if

$$\Gamma_0 = 1, \quad \mu_2(6) + \Gamma_2 = 2!\beta(2,6), \quad \text{and} \quad \mu_4(6) + 6\mu_2(6)\Gamma_2 + \Gamma_4 = 4!\beta(4,6),$$

that is,

$$\gamma_0 + 2 \sum_{j=1}^m \gamma_j = 1, \quad \sum_{j=1}^m j^2 \gamma_j = -\frac{1}{2} \quad \text{and} \quad \sum_{j=1}^m j^4 \gamma_j = \frac{31}{20}.$$

Proposition 15 (1) When $m = 2$, the unique iQI exact on $\mathbb{P}_5$ among all the quintic iQIs given by (4) is $\tilde{Q}_{6,2}f = \sum_{i \in \mathbb{Z}} \langle f, C_{i,2} \rangle M_i$, with

$$C_{i,2} = \frac{111}{35}M_i - \frac{71}{60}(M_{i-1} + M_{i+1}) + \frac{41}{240}(M_{i-2} + M_{i+2}).$$

(2) When $m = 3$ or 4, there is a unique NMN iQI $\tilde{Q}_{6,3}f = \sum_{i \in \mathbb{Z}} \langle f, C_{i,m} \rangle M_i$ exact on $\mathbb{P}_5$, given by

$$\begin{aligned} C_{i,3} &= \frac{531}{360}M_i - \frac{121}{400}(M_{i-2} + M_{i+2}) + \frac{71}{900}(M_{i-3} + M_{i+3}), \\ C_{i,4} &= \frac{1721}{1440}M_i - \frac{191}{1260}(M_{i-3} + M_{i+3}) + \frac{121}{2240}(M_{i-4} + M_{i+4}). \end{aligned}$$

(3) When $m = 5$, for each $\alpha \in \left[-\frac{281}{2880}, 0\right]$ the iQI given by

$$\tilde{Q}_{6,5}f = \sum_{i \in \mathbb{Z}} \langle f, C_{i,5} \rangle M_{i,6},$$

with

$$\begin{aligned} C_{i,5} &= \frac{4441 - 2240\alpha}{4000}M_i - \frac{281 + 2880\alpha}{2880}(M_{i-4} + M_{i+4}) \\ &\quad + \frac{191 + 1260\alpha}{4500}(M_{i-5} + M_{i+5}) \end{aligned}$$

is a NMN quintic iQI .

(4) When $m = 6$, there is a unique NMN quintic iQI , given by

$$\tilde{Q}_{6,6}f = \sum_{i \in \mathbb{Z}} \langle f, C_{i,6} \rangle M_i,$$

with

$$C_{i,6} = \frac{49534}{47385} M_i - \frac{14479}{568620} (M_{i-3} + M_{i+3}) + \frac{317}{113724} (M_{i-6} + M_{i+6}).$$

Observe that $\left(\frac{49534}{47385}, 0, 0, -\frac{14479}{568620}, 0, 0, \frac{317}{113724}\right)^T$ is the unique solution of the integral Problem 8 when $n = 3$ and $m = 6$.

Remark 16 As in the cubic case, it is easy to check the following results.

$$\begin{aligned} \|\tilde{Q}_{6,2}\|_{\infty} &\simeq 2.0313. \\ \|\tilde{Q}_{6,3}\|_{\infty} &= \frac{719}{432} \simeq 1.6643, \text{ and } \|\tilde{Q}_{6,4}\|_{\infty} = \frac{27520099}{19353600} \simeq 1.4220. \\ \|\tilde{Q}_{6,5}\|_{\infty} &= \frac{1721}{1440} \simeq 1.1951 \text{ for } \alpha = -\frac{281}{2880}, \text{ and } \|\tilde{Q}_{6,5}\|_{\infty} = \frac{88469567}{69120000} \simeq 1.2799 \\ &\text{for } \alpha = 0. \\ \|\tilde{Q}_{6,6}\|_{\infty} &= \frac{70529153}{57396000} \simeq 1.2288. \end{aligned}$$

6 Some error bounds for cubic discrete and integral QIs

In this section, we give some bounds for the error $f - \tilde{Q}_{4,m}f$ associated with the cubic QIs given by (15) and (17). The exactness of $\tilde{Q}_{4,m}$ on $\mathbb{P}_3 \subset S_{4,\mathbb{Z}}$ yields

$$\|f - \tilde{Q}_{4,m}f\|_{\infty} \leq (1 + \|\tilde{Q}_{4,m}\|_{\infty}) \text{dist}(f, S_{4,\mathbb{Z}}).$$

We observe that the class of the approximated function f does not take place in the construction of the NMN QI. Then, as we will show below, a better bound is obtained if we take into account the regularity of f .

Proposition 17 Let $m \geq 1$ and $\xi \in [0, 1]$. For any $f \in C^4(\mathbb{R})$, the quasi-interpolation error for the dQI given by (15) (resp. the iQI given by (17)) satisfies

$$|f(\xi) - \tilde{Q}_{4,m}f(\xi)| \leq \tilde{C}_m \|f^{(4)}\|_{\infty, [0,1]}, \quad (18)$$

with $\tilde{C}_1 = \frac{37}{1728} \simeq 0.0214$, and $\tilde{C}_m = \frac{96m^3 - 105m^2 + 120m - 38}{3456m^2}$ for $m \geq 2$ (resp. $\tilde{C}_1 = \frac{135937}{2903040} \simeq 0.0468$, $\tilde{C}_2 = \frac{151213}{1935360} \simeq 0.0781$, $\tilde{C}_3 = \frac{70031}{580608} \simeq 0.1206$, and $\tilde{C}_m = \frac{80640m^3 - 118281m^2 + 181440m - 81374}{1451520m^2}$ for $m \geq 4$).

PROOF. Let $\tilde{Q}_{4,m}$ be the dQI given by (15). Let $\mathcal{L}$ be the linear form defined as $\mathcal{L}f = f(\xi) - \tilde{Q}_{4,m}f(\xi)$. The Peano's Theorem (see e.g. [19, chapter 4]) gives

$$f(\xi) - \tilde{Q}_{4,m}f(\xi) = \int_0^1 k_{4,m}(\xi, t) f^{(4)}(t) dt,$$

where

$$3!k_{4,m}(\xi, t) = \mathcal{L}(\cdot - t)_+^3 = (\xi - t)_+^3 - Q_{4,m}(\cdot - t)_+^3(\xi).$$

Since $(\xi - t)_+^3 = 0$ for $\xi < t$, in this case it follows easily that

$$k_{4,m}(\xi, t) = \sum_{i=-1}^2 \alpha_i(t) M_4(\xi - i),$$

with

$$\begin{aligned} \alpha_{-1}(t) &= \begin{cases} 0, & \text{if } m = 1, \\ \frac{1}{36}m^2(-1 + m - t)^2, & \text{if } m \geq 2, \end{cases} \\ \alpha_0(t) &= \frac{1}{36m^2}(m - t)^3, \\ \alpha_1(t) &= \frac{1}{36m^2}(-2(3m^2 + 1)(1 - t)^3 + (1 + m - t)^3), \\ \alpha_2(t) &= \begin{cases} \frac{1}{36}((1 - t^3) - 8(2 - t)^3 + (3 - t)^3), & \text{if } m = 1, \\ \frac{1}{36m^2}(-2(3m^2 + 1)(2 - t)^3 + (2 + m - t)^3), & \text{if } m \geq 2. \end{cases} \end{aligned}$$

For $-1 \leq i \leq 2$, $M_4(\xi - i)$ is a cubic polynomial on $[0, 1]$. Let $b_i^3(t) = \binom{3}{i}(1 - t)^3 t^{3-i}$, $0 \leq i \leq 3$, be the i th Bernstein polynomial. It is well known (see e.g. [11, p. 13]) that

$$\begin{aligned} M_4(\xi + 1) &= \frac{1}{6}b_0^3(\xi), \\ M_4(\xi) &= \frac{1}{6}(4b_0^3(\xi) + 4b_1^3(\xi) + 2b_2^3(\xi) + b_3^3(\xi)), \\ M_4(\xi - 1) &= \frac{1}{6}(b_0^3(\xi) + 2b_1^3(\xi) + 4b_2^3(\xi) + 4b_3^3(\xi)), \\ M_4(\xi - 2) &= \frac{1}{6}b_3^3(\xi). \end{aligned}$$

Thus, we get

$$k_{4,m}(\xi, t) = \sum_{i=0}^3 \omega_i(t) b_i^3(\xi), \quad (19)$$

where

$$\omega = W\mathbf{b}, \quad (20)$$

with $\omega = (\omega_1, \omega_2, \omega_3, \omega_3)^T$, $\mathbf{b} = (b_0^3, b_1^3, b_2^3, b_3^3)^T$, and $W = (W_{ij})_{1 \leq i, j \leq 4}$ is given by

$$W = \frac{1}{216} \begin{pmatrix} 4 & 4 & 2 & 1 \\ 4 & 6 & 4 & 2 \\ 2 & 16 & 8 & 4 \\ -35 & 2 & 4 & 4 \end{pmatrix}$$

if $m = 1$, and

$$\begin{aligned} W_{11} &= W_{12} = W_{21} = W_{34} = W_{43} = W_{44} = \frac{1}{108m^2} (-1 + 3m - 3m^2 + 3m^3), \\ W_{13} &= W_{2,4} = W_{3,1} = W_{4,2} = \frac{1}{108m^2} (-2 + 6m - 6m^2 + 3m^3), \\ W_{2,2} &= W_{3,3} = \frac{1}{108m} (1 + 3m^2), \\ W_{2,3} &= W_{3,2} = \frac{1}{108m} (2 - 3m + 3m^2), \\ W_{1,4} &= \frac{1}{108m^2} (-2 + 4m - 3m^2 + m^3), \\ W_{4,1} &= \frac{1}{36m^2} (-2 + 4m - 3m^2 + m^3) \end{aligned}$$

for $m \geq 2$.

All the coefficients in the matrix W are positive for $m \geq 9$. Hence, in these cases the kernel $k_{4,m}$ is positive in $[0, 1] \times [0, 1]$. For the other values of m , the element $W_{4,1}$ is negative. Therefore, we will analyze carefully the kernel in the triangle T_1 of vertices $A_1 = (0, 0)$, $A_2 = (1, 1)$, and $A_3 = (0, 1)$.

For $m = 1$, let (u, v, w) be the barycentric coordinates of (ξ, t) with respect to T_1 , i.e.,

$$uA_1 + vA_2 + wA_3 = (\xi, t), \quad u + v + w = 1.$$

The kernel $k_{4,1}$ being a polynomial in (ξ, t) of total degree 6, it has a unique Bézier representation (see e.g. [11, chapter 5])

$$k_{4,1}(\xi, t) = \sum_{|l|=6} a_l b_l^6(u, v, w),$$

where $b_l^6(u, v, w) = \frac{6!}{l_1!l_2!l_3!} u^{l_1} v^{l_2} w^{l_3}$, $0 \leq l_1, l_2, l_3$ and $|l| := l_1 + l_2 + l_3 = 6$, are the Bernstein polynomials of degree 6, and the Bernstein–Bézier coefficients

a_l are as follows:

$$\begin{array}{rcl}
a_{0,0,6} & a_{0,1,5} & a_{0,2,4} & a_{0,3,3} & a_{0,4,2} & a_{0,5,1} & a_{0,6,0} & 20 & 30 & 44 & 59 & 72 & 80 & 80 \\
a_{1,0,5} & a_{1,1,4} & a_{1,2,3} & a_{1,3,2} & a_{1,4,1} & a_{1,5,0} & & 30 & 46 & 69 & 89 & 96 & 80 \\
a_{2,0,4} & a_{2,1,3} & a_{2,2,2} & a_{2,3,1} & a_{2,4,0} & & & 44 & 69 & 106 & 129 & 112 \\
a_{3,0,3} & a_{3,1,2} & a_{3,2,1} & a_{3,3,0} & & & & = \frac{1}{2592} & 59 & 89 & 129 & 146 \\
a_{4,0,2} & a_{4,1,1} & a_{4,2,0} & & & & & 72 & 96 & 112 \\
a_{5,0,1} & a_{5,1,0} & & & & & & 80 & 80 \\
a_{6,0,0} & & & & & & & 80
\end{array}$$

We then get the positivity of $k_{4,1}(\xi, t)$ on T_1 . The same conclusion can be drawn for $k_{4,m}$, $2 \leq m \leq 8$.

Now, let us determine the Peano kernel when $\xi \geq t$. Since $(\xi - t)_+^3 = (\xi - t)^3$, and $\tilde{Q}_{4,m}$ is exact on $\mathbb{P}_3$, we have

$$k_{4,m}(\xi, t) = \tilde{Q}_{4,m}((\cdot - t)^3 - (\cdot - t)_+^3)(\xi) = -\tilde{Q}_{4,m}(t - \cdot)_+^3(\xi).$$

After some calculations, we find that

$$k_{4,m}(\xi, t) = \sum_{i=0}^3 \omega_{3-i}(1-t) b_i^3(\xi). \quad (21)$$

Therefore, $k_{4,m}$ is also positive on the triangle T_2 of vertices $(0, 0)$, $(1, 0)$, and $(1, 1)$.

As $k_{4,m}$ does not changes sign, there exists $\tau \in [0, 1]$ such that

$$f(\xi) - \tilde{Q}_{4,m}f(\xi) = f^{(4)}(\tau) K_{4,m}(\xi),$$

where

$$K_{4,m}(\xi) := \int_0^1 k_{4,m}(t) dt.$$

Let $X = \xi(1 - \xi)$. From (19), (20), and (21), we obtain for $m = 1$

$$K_{4,1}(\xi) = \frac{1}{864} (36X^2 + 21X + 11),$$

and for $m \geq 2$

$$\begin{aligned}
K_{4,m}(\xi) = \frac{1}{1728m^2} & \left(X^2 + 12(7 - 12m + 9m^2)X \right. \\
& \left. + 4(12m^3 - 21m^2 + 24m - 10) \right)
\end{aligned}$$

For all $m \geq 1$, the function $K_{4,m}$ attains its maximum value at $\xi = \frac{1}{2}$, and $\tilde{C}_m = K_{4,m}\left(\frac{1}{2}\right)$.

The proof for the integral QI given by (17) runs as before. $\square$

From (18), the quasi-interpolation error for the scaled QI $\tilde{Q}_{m,4}^h$, $h > 0$, associated with $\tilde{Q}_{m,4}$ satisfies

$$\|f - \tilde{Q}_{4,m}^h f\|_{\infty} \leq \tilde{C}_m h^4 \|f^{(4)}\|_{\infty}. \quad (22)$$

7 Application to functions with discontinuities

When the classical scaled dQI $\tilde{Q}_{4,1}^h$ is used to approximate the Heaviside function H defined as

$$H(x) = \begin{cases} 0, & \text{if } x < 0, \\ 1, & \text{if } x \geq 0, \end{cases}$$

the QI $\tilde{Q}_{4,1}^h H$ oscillates near the discontinuity point with an overshoot independent of f and equal to $\frac{11}{1800} (6 + \sqrt{11}) \simeq 5.7\%$. Since the scaled Schoenberg's dQI $S_4^h H$ does not oscillate near zero and the sequence $(\tilde{Q}_{4,m}^h)_{m \geq 1}$ converges to S_4 in the infinite norm when $m \rightarrow +\infty$, it seems natural to use a NMN cubic dQI in order to obtain another QI $\tilde{Q}_{4,m}^h H$ giving a smaller overshoot. For $m = 2$ (resp. $m = 3$) this equals $\frac{762+13\sqrt{26}}{22500} \simeq 3.7\%$ (resp. 1.85%). However, the constant $\tilde{C}_m$ in (22) given by (18) increases with m . Thus it is better to choose a low value of m . Table 5 shows the higher decrease of the upper bound $\tilde{\nu}_{4,m}$ and the infinite norm of $\tilde{Q}_{4,m}$ occurs for $m = 2$ and 3. Figure 1 shows how the overshoot is damped and shifted when the dQIs $\tilde{Q}_{4,2}$ and $\tilde{Q}_{4,3}$ are used.

Table 5

Values of $\tilde{\nu}_{4,m}$, $\|\tilde{Q}_{4,m}\|_{\infty}$, and $\tilde{C}_m$ for some cubic NMN QIs $\tilde{Q}_{4,m}$.

m	dQI			iQI		
	$\tilde{\nu}_{4,m}$	$\ \tilde{Q}_{4,m}\ _{\infty}$	$\tilde{C}_m$	$\tilde{\nu}_{4,m}$	$\ \tilde{Q}_{4,m}\ _{\infty}$	$\tilde{C}_m$
1	1.6667	1.2222	0.0214	2.3333	1.5278	0.0468
2	1.1667	1.1389	0.0398	1.3333	1.2778	0.0781
3	1.0741	1.0741	0.0633	1.1481	1.1481	0.1206
4	1.0417	1.0417	0.0887	1.0833	1.0833	0.1685

For a function with isolated discontinuities, when $h \rightarrow 0$ also happens the oscillatory behaviour near each discontinuity point. $\tilde{Q}_{4,2}$ and $\tilde{Q}_{4,3}$ behaves




Fig. 1. The Heaviside function and the QIs $\tilde{Q}_{4,1}H$, $\tilde{Q}_{4,2}H$, and $\tilde{Q}_{4,3}H$.

better than $\tilde{Q}_{4,1}$ in a neighbourhood of each discontinuity point. Far from them (22) holds. For cubic iQI a similar result holds.

8 Conclusion

In this paper, we have constructed discrete and integral spline QIs on uniform partitions of the real line with optimal approximation orders and small norms by minimizing a simple upper bound of the true norm. They can be used to approximate functions with isolated discontinuities. Although this method gives good results w.r.t. the infinite norm, the constant appearing in the standard estimation of the quasi-interpolation error is too much crude, because the bound is independent of the class of the function to be approximated. It will be interesting to construct QIs in order to obtain a better bound for enough smooth functions.

Future works would be include a similar treatment for nonuniform partitions in the univariate case (cf. [2]), as well as the construction of discrete and integral QIs based on B-splines on the uniform three or four-directional meshes of the plane, in the bivariate case (cf. [3]).

References

- [1] D. Barrera, M. J. Ibáñez and P. Sablonnière, Near-best discrete quasi-interpolants on uniform and nonuniform partitions, in: A. Cohen, J.-L. Merrien, and L. L. Schumaker (Eds.), *Curve and Surface Fitting: Saint-Malo 2002*, Nashboro Press, Brentwood, 2003, pp. 31–40.

- [2] D. Barrera, M. J. Ibáñez, P. Sablonnière and D. Sbibi, Near-best spline discrete quasi-interpolants on non-uniform partitions of $\mathbb{R}$, submitted.
- [3] D. Barrera, M. J. Ibáñez, P. Sablonnière and D. Sbibi, Near-best quasi-interpolants associated with H-splines on a three-direction mesh, submitted.
- [4] C. de Boor, On uniform approximation by splines, *J. Approx. Theory* 1 (1968) 219–235.
- [5] C. de Boor and G. F. Fix, Spline approximation by quasiinterpolants, *J. Approx. Theory* 8 (1973) 19–54.
- [6] C. de Boor, *A practical guide to splines*, Springer-Verlag, New York, 1978.
- [7] C. de Boor, K. Höllig and S. Riemenschneider, *Box splines*, Springer-Verlag, New York, 1993.
- [8] P. L. Butzer, M. Schmidt, E. L. Stark and L. Vogt, Central factorial numbers; their main properties and some applications, *Numer. Funct. Anal. and Optimiz.* 10, 5&6 (1989) 419–488.
- [9] P. L. Butzer and M. Schmidt, Central factorial numbers and their role in finite difference calculus and approximation, in: *Colloquia Mathematica Societatis János Bolyai* 58, Approximation Theory, Kecskémet, Hungary 1990, pp. 127–150.
- [10] G. Chen, C. K. Chui and M. J. Lai, Construction of real-time spline quasi-interpolation schemes, *Approx. Theory Appl.* 4 (1988) 61–75.
- [11] C. K. Chui, *Multivariate Splines*, SIAM, Philadelphia, 1988.
- [12] Z. Ciesielski, Local spline approximation and nonparametric density estimation, in: *Construction Theory of Functions'87*, Bulgarian Academy of Sciences, Sofia, 1988, pp. 79–84.
- [13] R. A. DeVore and G. G. Lorentz, *Constructive approximation*, Springer-Verlag, Berlin, 1993.
- [14] T. N. T. Goodman and A. Sharma, A modified Bernstein-Schoenberg operator, in: *Construction Theory of Functions'87*, Bulgarian Academy of Sciences, Sofia, 1988, pp. 168–173.
- [15] M. J. Ibáñez Pérez, *Quasi-interpolantes spline discretos de norma casi mínima. Teoría y aplicaciones*, Ph.D., Departamento de Matemática Aplicada, Universidad de Granada, Granada (Spain), 2003.
- [16] B.-G. Lee, T. Lyche and M. Mørken, Some examples of quasi-interpolants constructed from local spline projectors, in: T. Lyche, L. L. Schumaker (Eds.), *Math Methods for Curves and Surfaces: Oslo 2000*, Vanderbilt University Press, Nashville, 2000, pp. 243–252.
- [17] T. Lyche and L. L. Schumaker, Local spline approximation methods, *J. Approx. Theory* 15 (1975) 294–325.

- [18] M. R. Osborne, *Simplicial algorithms for minimizing polyhedral functions*, Cambridge University Press, Cambridge, 2001.
- [19] G. M. Phillips, *Interpolation and approximation by polynomials*, Springer-Verlag, New-York, 2003.
- [20] P. Sablonnière, Positive spline operators and orthogonal splines, *J. Approx. Theory* 52 (1988) 28–42.
- [21] P. Sablonnière and D. Sbibi, Spline integral operators exact on polynomials, *Approx. Theory Appl.* Vol. 10, No. 3 (1994) 56–73.
- [22] P. Sablonnière, Quasi-interpolantes splines sobre particiones uniformes, Meeting in Approximation Theory, Úbeda (Spain), July 2000. Prépublication IRMAR 00-38, 2000.
- [23] P. Sablonnière, On some multivariate quadratic spline quasi-interpolants on bounded domains, in: K. Jetter, M. Reiner, J. Stöckler (Eds.), *Modern developmets in multivariate approximation*, ISNM Vol. 145, Birkhäuser-Verlag, Basel, 2003, pp. 263–278.
- [24] P. Sablonnière, Quadratic spline quasi-interpolants on bounded domains of $\mathbb{R}^d$, $d = 1, 2, 3$, Spline and radial functions, *Rend. Sem. Univ. Pol. Torino*, Vol. 61 (2003) 61-78.
- [25] D. Sbibi, *Approximations des fonctions d’une ou deux variables par des opérateurs splines intégraux*, Ph.D., Université de Rennes, Rennes (France), 1987.
- [26] I. J. Schoenberg, *Cardinal Spline Interpolation*, SIAM, Philadelphia, 1973.
- [27] L. L. Schumaker, *Spline functions. Basic theory*, John Wiley & Sons, New York, 1981.
- [28] G. A. Watson, *Approximation Theory and Numerical Methods*, Wiley, Chichester, 1980.