



Synthesis of new bioisosteric hemiasterlin analogues with extremely high cytotoxicity

Chinh Pham The, Tuyet Anh Dang Thi, Thi Phuong Hoang, Quoc Anh Ngo,
Duy Tien Doan, Thu Ha Nguyen Thi, Tham Pham Thi, Thu Ha Vu Thi,
Mickael Jean, Pierre van de Weghe, et al.

► To cite this version:

Chinh Pham The, Tuyet Anh Dang Thi, Thi Phuong Hoang, Quoc Anh Ngo, Duy Tien Doan, et al.. Synthesis of new bioisosteric hemiasterlin analogues with extremely high cytotoxicity. *Bioorganic and Medicinal Chemistry Letters*, 2014, 24 (22), pp.5216-5218. 10.1016/j.bmcl.2014.09.065 . hal-01080910

HAL Id: hal-01080910

<https://univ-rennes.hal.science/hal-01080910>

Submitted on 7 Nov 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Synthesis of New Simplified Hemiasterlin Derivatives with α,β -Unsaturated Carbonyl Moiety

Chinh Pham The,^a Tuyet Anh Dang Thi,^a Thi Phuong Hoang,^a Quoc Anh Ngo,^a Duy Tien Doan,^a Thu Ha Nguyen Thi,^a Tham Pham Thi,^a Thu Ha Vu Thi,^a M. Jean,^b P. van de Weghe^{*b} and Tuyen Nguyen Van^{*a}

^aInstitute of Chemistry - Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam.

^bEquipe Produits Naturels, Synthèses et Chimie Médicinale (PNSCM), UMR CNRS 6226 – Institut des Sciences Chimiques de Rennes, Université de Rennes 1, UFR Sciences Pharmaceutiques et Biologique, 2, Avenue du Prof L. Bernard, F-35043 Rennes Cedex, France.

ABSTRACT

In this article, we report a convenient and efficient method for the synthesis of new simplified derivatives of hemiasterlin in which the α,α -dimethylbenzylic moiety A is replaced by α,β -unsaturated aryl groups as Michael acceptor. Most of these derivatives have a strong cytotoxic activity on three human tumor cell lines (KB, Hep-G₂ and MCF₇). Analogs **17b** and **17f** showed a high cytotoxicity against KB and Hep-G₂ cancer cell lines comparable to paclitaxel and ellipticine.

Hemiasterlins belong to a family of naturally occurring tripeptides from marine sponges.¹ The important derivatives of hemiasterlins are hemiasterlin A, hemiasterlin B, and hemiasterlin C, which were isolated from marine sponge *Auletta* and *Cymbastella* (Fig. 1) and exhibited potent cytotoxicity in vitro against murine leukemia P388 and human breast, ovarian, colon, and lung cancer cell lines.^{2,3} Hemiasterlins suppress microtubule depolymerization presumably by binding to the vinca-alkaloid binding-domain of tubulin and leading to mitotic arrest and cell death.^{4a} The synthetic analog HTI-286 (**2**) displayed especially potent cytotoxicity against paclitaxel (TaxolTM) resistant tumor cell lines in vitro and in vivo and is currently in clinical trials.^{4b}





Figure 1. Structures of hemiasterlin derivatives and HTI analogs.

There are several reports on the synthesis of new derivatives of hemiasterlin in which the indole aromatic ring in the moiety A of the original molecule was replaced by aryl functional groups.^{5,6a} However, asymmetric synthesis of the stereospecific amine group and especially the *gem*-dimethyl moiety were proved to be highly problematic. To overcome this difficulty, several studies explored modifications of segment A in which the *gem*-dimethyl moiety has been eliminated. Some of these derivatives showed promising cytotoxic activity.⁶


In recent years, previous works have shown that the presence of a α,β -unsaturated carbonyl system in peptide derivatives can improve the biological activity compared to the initial compound.^{7,8} Explanations may include either a limitation of free energy of the peptide by addition of a constrained system or by an electrophilic character of the compound leading to the possibility to form covalent bond with the protein target. Based on the idea that hemiasterlin derivatives containing α,β -unsaturated carbonyl systems could induce a remarkable cytotoxicity, we decided to synthesize new hemiasterlin derivatives in which the α,α -dimethylbenzylic group (fragment A) is replaced by a α,β -unsaturated carbonyl group.

New hemiasterlin derivatives were synthesized via classical peptide coupling reactions between two fragments **7a-h** and **14a,b**. A general procedure for the synthesis of compound **7** is outlined in Scheme 1. Compounds **7** were prepared from glycine in three steps.⁹ The synthesis started by acetylation of glycine with acetic anhydride in water at room temperature following by condensation with aryl aldehydes using sodium acetate in the presence of acetic anhydride at 90 °C for 12 h which afforded azalactones **6a-h** in 72-83% yields.⁹ Finally, azalactones **6a-h** were hydrolyzed in aqueous sodium hydroxide, followed by treatment with hydrochloric acid (12 N) at 100 °C for 4 h to give compounds **7a-h** in 65-88% yields.




Scheme 1. Reagents and conditions (i) 2.0 equiv Ac_2O , H_2O , rt, 20 h. (ii) 0.75 equiv ArCHO , 1.0 equiv AcONa , Ac_2O , 90 °C, 12 h. (iii) NaOH (1 N) then HCl (12 N), 100 °C, 4 h.

Compounds **14a,b** were obtained from Boc-L-valine in 6 steps as depicted in Scheme 2.⁴ In the first step, Boc-L-valines **8a,b** were converted to Weinreb amide **9a,b** in good yields (86 and 81%) by treatment with a mixture of *N,O*-dimethylhydroxylamine hydrochloride, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and hydroxybenzotriazole (HOBT) in the presence of N-ethylisopropylamine (*i*-PrNHEt) at room temperature for 12 h.¹⁰ Reduction of **9a,b** using LiAlH_4 in THF was carried out at room temperature for 1 h to give aldehydes **10a,b** in 78% and 65% yields, respectively.¹¹ Afterward, Wittig reaction of aldehydes **10a,b** with ethyl 2-(triphenylphosphoranylidene)propanoate was carried out at reflux in CH_2Cl_2 for 6 h to afford the alkenoates **11a** and **11b** in 86% and 88% yields, respectively.^{6,12} Then, removal of the Boc group using trifluoroacetic acid in CH_2Cl_2 at room temperature for 1 h led to **12a,b** in high yields (90 and 95%). The expected compounds **14a,b** are finally obtained after coupling **12a,b** and Boc-L-leucine in the presence of EDC/HOBt in DMF at room temperature for 12 h following by Boc deprotection in classical reaction conditions.



Scheme 2. Reagents and conditions (i) 1.1 equiv NH(Me)OMe, 1.1 equiv EDC, 1.1 equiv HOBt, 2.0 equiv *i*-PrNHEt, DMF, rt, 12 h. (ii) 4.0 equiv LAH, THF, rt, 1 h. (iii) Ph₃P=C(CH₃)CO₂E_t, CH₂Cl₂, reflux, 6 h. (iv) TFA, CH₂Cl₂, rt, 1 h. (v) 1.0 equiv Boc-L-Leucine, 1.1 equiv EDC, 1.1 equiv HOBt, 2.0 equiv *i*-PrNHEt, DMF, rt, 12 h. (vi) TFA, CH₂Cl₂, rt, 1 h.

The final compounds **16a-h** were then prepared in two steps after peptide coupling reactions of **14a** with amides **7a-h** following by the saponification of ester using a 1 N lithium hydroxide solution (Scheme 3-a). Similarly, the hemiasterlin derivatives **18a-g** were obtained from **14b** (Scheme 3-b).



Scheme 3. Reagents and conditions (i) 1.0 equiv **12a-h**, 1.1 equiv EDC, 1.1 equiv HOBt, 2.0 equiv *i*-PrNHEt, DMF, rt, 12 h. (ii) 10 equiv LiOH, MeOH : H₂O (2:1), rt, 10 h.

All compounds **15a-h**, **16a-h**, **17a-g** and **18a-g** were evaluated in vitro for their cytotoxic activity against four human tumor cell lines (KB, Hep-G₂, LU and MCF₇) and the results were summarized in Table 1. Eleven hemiasterlin derivatives showed strong activity against the KB, Hep-G₂ cell line with IC₅₀ value below 100 nM. Analogs **16b**, **17b** and **18f** exhibited potent cytotoxicity against the KB cell line with IC₅₀ = 8.2, 3.5 and 3.7 nM, respectively. Meanwhile analogs **16b**, **2217c** and **17f** displayed potent cytotoxicity against the Hep-G₂ cell line with IC₅₀ value 17.8, 16.3 and 3.7 nM, respectively. Derivatives **15a**, **15e** and **17a** presented a cytotoxic activity against MCF₇ cell line with IC₅₀ value 42.3, 60.0 and 67.9 nM. Concerning the last cell line, the LU cell line, the hemiasterlin analogs showed weak activities with IC₅₀ values above

269 nM. It is noteworthy to mention that two derivatives **17a** and **17f** present a cytotoxic activity against two cancer cell lines (KB, Hep-G₂) comparable with those of ellipticine and paclitaxel

Table 1. Cytotoxicity evaluation.

Entry	compound	IC ₅₀ (nM)			
		KB	Hep-G ₂	LU	MCF ₇
1	15a	30.8	39.7	>269	42.3
2	15b	160.5	214.2	>269	>269
3	15c	86.2	67.3	>269	>269
4	15d	107.4	57.6	>269	203
5	15e	77.4	47.7	228.4	62
6	15f	>269	>269	>269	>269
7	15g	>269	>269	>269	>269
8	15h	>269	>269	>269	>269
9	6a	63	33.4	>269	>269
10	6b	8.2	17.8	>269	>269
11	6c	155.4	>269	>269	>269
12	6d	>269	>269	>269	>269
13	6e	>269	>269	>269	>269
14	6f	>269	>269	>269	>269
15	6g	>269	>269	>269	>269
16	6h	>269	>269	>269	>269
17	17a	23.0	23.1	>269	67.9
18	17b	3.5	31.4	231.0	178.4
19	17c	24.1	16.3	158	112.8
20	17d	202	15.0	>269	>269
21	17e	>269	>269	>269	>269
22	17f	3.7	3.7	>269	215
23	17g	49.0	>269	>269	>269
24	18a	69.7	13.0	>269	>269
25	18b	188.0	>269	>269	>269
26	18c	>269	149.4	>269	>269
27	18d	234	63.2	>269	>269
28	18e	>269	>269	>269	>269
29	18f	15.8	63.8	>269	>269
30	18g	>269	>269	>269	>269
31	Ellipticine	1.26	1.26	1.82	2.15
32	Paclitaxel	3.9	0.19	-	-

In conclusion, a concise synthetic approach for new modified hemiasterlin derivatives was achieved in which the α,α -dimethylbenzylic group and amino NHMe moiety were replaced respectively by a α,β -unsaturated aryl and an amide NH-Ac group avoiding the synthesis of the chiral fragment A. Most of these derivatives possess strong cytotoxic activity on three human tumor cell lines (KB, Hep-G₂ and MCF₇) and two analogs, **17b** and **17f**, showed a comparable cytotoxicity activity to paclitaxel and ellipticine against KB and Hep-G₂ cancer cell lines. Based on previously reported work,^{6b} we can envisage that our hemiasterlin derivatives act as tubulin

polymerization inhibitors. From our best compounds, more detailed biological studies will be undertaken and will be reported in due course.

Acknowledgments

This work was financially supported in part by scientific research and technological development project (code: DT.NCCB-DHUD.2011-G/07).

Supplementary Material

Supplementary data associated with this article can be found, in the online version, at doi:

References and notes

1. Talpir, R.; Benayahu, Y.; Kashman, Y.; Pannell, L.; Schleyer, M. *Tetrahedron Lett.*, **1994**, 35, 4453.
2. Gamble, W. R.; Durso, N. A.; Fuller, R. W.; Westergaard, C. K.; Johnson, T. R.; Sackett, D. L.; Hamel, E.; Cardellina II, J. H.; Boyd, M. R. *Bioorg. Med. Chem.*, **1999**, 7, 1611.
3. Coleman, J. E.; Silva, E. D.; Kong, F.; Andersen, R. J.; Allen, T. M. *Tetrahedron*, **1995**, 51, 10653.
4. (a) Anderson, H. J.; Coleman, J. E.; Andersen, R. J.; Roberge, M. *Cancer Chemother. Pharmacol.*, **1997**, 39, 223. (b) Loganzo, F.; Discafani, C. M.; Annable, T.; Beyer, C.; Musto, S.; Hari, M.; Tan, X.; Hardy, C.; Hernandez, R.; Baxter, M.; Singanallure, T.; Khafizova, G.; Poruchynsky, M. S.; Fojo, T.; Nieman, J. A.; Ayral-Kaloustian, S.; Zask, A.; Andersen, R. J.; Greenberger, L. M. *Cancer Res.*, **2003**, 63, 1838-1845.
5. (a) Andersen, R.; Piers, E.; Nieman, J.; Coleman, J.; Roberge, M. WO9932509. (b) Yamashita, A.; Norton, E. B.; Kaplan, J. A.; Niu, C.; Loganzo, F.; Hernand, R.; Beyer, C. F.; Annable, T.; Musto, S.; Discafani, C.; Zask, A.; Ayral-Kaloustian, S. *Bioorg. Med. Chem. Lett.*, **2004**, 14, 5317. (c) Greenberger, L. M.; Loganzo, F. Jr; Discafani-Marro, C.; Zask, A.; Ayral-Kaloustian, S. WO2004026293. (d) Mokale, S. N.; Lokwani, D.; Shinde, D. *Bioorg. Med. Chem.*, **2012**, 20, 3119. (d) Mali, S. M.; Bandyopadhyay, A.; Jadhav, S. V.; Kumar, M. G.; Gopi, H. N. *Org. Biomol. Chem.*, **2011**, 9, 6566.
6. (a) Niew, J. A.; Coleman, J. E.; Wallace, D. J. ; Piers, E.; Lim, L. Y.; Roberge, M.; Andersen, R. J. *J. Nat. Prod.*, **2003**, 6, 183. (b) Simoni, D.; Lee, R. M.; Durrant, D.; Chi, N. W.; Baruchello, R.; Rondanin, R.; Rullo, C.; Marchetti, P. *Bioorg. Med. Chem. Lett.*, **2010**, 20, 3431. (c) Murray, W. V.; Sun, S.; Turchi, I. J.; Brown, F. K.; Gauthier, A. D. *J. Org. Chem.*, **1999**, 64, 5930. (d) Grison, C.; Geneve, S.; Halbin, E.; Coutrot, P. *Tetrahedron*, **2001**, 57, 4903.
7. Yamazaki, Y.; Tanaka, K.; Nicholson, B.; Deyanat-Yazdi, G.; Pott, B.; Yoshida, T.; Oda, A.; Kitagawa, T.; Orikasa, S.; Kiso, Y.; Yasui, H.; Akamatsu, M.; Chinen, T.; Usui, T.; Shinozaki, Y.; Yakushiji, F.; Miller, B. R.; Neuteboom, S.; Palladino, M.; Kanoth, K.; Lloyd, G. K.; Hayashi, Y. *J. Med. Chem.*, **2012**, 55, 1056.
8. Pijus, K. M.; Zhiyong, R.; Xiaomin, C.; Chiyi, X.; McMurray, J. S. *J. Med. Chem.*, **2009**, 52, 6126.
9. (a) Herbst, R. M. ; Shemin, D. *Org. Synth.*, **1939**, 19, 1. (b) Thomas, C.; Jodie, B.; Flavio, C. *Tetrahedron Lett.*, **2010**, 51, 625. (c) Mesaik, M. A.; Rahat, S.; Khan, K. M.; Zia-Ullah; Choudhary, M. I.; Murad, S.; Ismail, Z.; Atta-ur-Rahman; Ahmad, A. *Bioorg. Med. Chem.*, **2004**, 12, 2049. (d) RajanBabu, T. V.; Ayers, T. A.; Halliday, G. A.; You, K. K.; Calabrese, J. C. *J. Org. Chem.*, **1997**, 62, 6012. (e) Cutolo, M.;

- Fiandanese, V.; Naso, F.; Sciacovelli, O. *Tetrahedron Lett.*, **1983**, 24, 4603. (f) Baell, J. B.; Garrett, T. P. J.; Hogarth, P. M.; Matthews, B. R.; McCarthy, T. D.; Pietersz, G. A. WO2000015214. (g) Sawant, S. D.; Barvrkar, A. A.; Chabukswar, A. R.; Sarak, S. D. *Journal of Applicable Chemistry*, **2013**, 2, 372. (h) Blanco-Lomas, M.; Funes-Ardoiz, I.; Campos, P. J.; Sampedro, D. *Eur. J. Org. Chem.*, **2013**, 6611. (i) Funes-Ardoiz, I.; Blanco-Lomas, M.; Campos, P. J.; Sampedro, D. *Tetrahedron*, **2013**, 69, 9766.
10. (a) Taillier, C.; Bellotta, V.; Meyer, C.; Cossy, J. *Org. Lett.*, **2004**, 6, 2145. (b) Nahm, S.; Weinreb, S. M. *Tetrahedron Lett.*, **1981**, 39, 3815. (c) Tyrrell, E.; Brawn, P.; Carew, M.; Greenwood, I. *Tetrahedron Lett.*, **2011**, 52, 369. (d) Kim, J. G.; Jang, D. O. *Bull. Korean Chem. Soc.*, **2010**, 31, 171.
11. (a) Angelastro, M. R.; Mehdi, S.; Burkhardt, J. P.; Peet, N. P.; Bey, P. *J. Med. Chem.*, **1990**, 33, 11. (b) Burke, A. J.; Davies, S. G.; Garner, C. A.; McCarthy, T. D.; Roberts, P. M.; Smith, A. D.; Rodriguez-Solla, H.; Vicker, R. *J. Org. Biomol. Chem.*, **2004**, 2, 1387.
12. Murphy, J. A.; Commeueuc, A. G. J.; Snaddon, T. N.; McGuire, T. M.; Khan, T. A.; Hisler, K.; Dewis, M. L.; Carling, M. *Org. Lett.*, **2005**, 7, 1427.