

**Maternal occupation during pregnancy, birth weight and length of gestation:
combined analysis of 13 European birth cohorts**

Abstract

Objectives: We assess whether maternal employment during pregnancy, overall and in selected occupational sectors, is associated with birth weight, small for gestational age (SGA), term low birth weight (LBW), length of gestation, and preterm delivery in a population-based birth cohort design.

Methods: We used data from more than 200 000 mother-child pairs enrolled in 13 European birth cohorts. Employed *vs* non-employed women were compared. Among employees, we defined groups of occupations representing the main sectors of employment in women where potential reproductive hazards are considered to be present. The comparison group included all other employed women not included in the occupational sector being assessed. We performed meta-analyses of cohort-specific estimates and explored heterogeneity.

Results: Employees had a lower risk of preterm delivery than non-employees (OR_{adj} 0.86, 95%CI 0.81-0.91). Working in most of the occupational sectors studied was not associated with adverse birth outcomes. Nurse employment was associated with lower risk SGA infants (OR_{adj} 0.91, 95%CI 0.84-0.99) whereas food industry workers had an increased risk of preterm delivery (OR_{adj} 1.50, 95%CI 1.12-2.02). There was little evidence for heterogeneity between cohorts.

Conclusions: This study suggests that, overall, employment during pregnancy is associated with a reduction in the risk of preterm birth and that work in certain occupations may affect pregnancy outcomes. This study of exploratory nature provides an important platform on which to base further prospective studies focused on the

potential consequences of maternal occupational exposures during pregnancy on child development.

Keywords: occupation; birth weight; length of gestation; meta-analysis; Europe; cohort studies

Introduction

Today, women constitute almost half of the workforce in many European countries and most women work during their reproductive years (1). Studies examining the relationship between employment during pregnancy and reproductive health have yielded contradictory results (2-8). In most European countries, employment during pregnancy selects a population with specific demographic and social characteristics, and better access to prenatal health care (3). At the same time, work during pregnancy may entail strenuous or hazardous work conditions. A number of chemical and physical agents present in the workplace or related to work organization have been suspected to interfere with the normal reproductive process (9). For some chemicals such as anaesthetic gases, toxic metals, and specific solvents, the evidence is quite compelling (10) whereas for most agents and factors there is large uncertainty and if any association exists it seems of modest magnitude (11). Occupational sectors, for instance health sector or cleaning work, have the characteristics of employing a large number of women and cumulating several potential hazards according to distinct exposure patterns.

Most of the evidence comes from occupational cohort studies using retrospective data, case-control studies, and registry-based studies (12,13). The population-based birth cohort design has rarely been used in this field (2,14-16), mainly because of lack of power when studying rare exposures or occupations and outcomes such as congenital malformations. However, this study design has important strengths such as prospective data collection during pregnancy and potential for controlling relevant confounders. In the present study we used data from 13 European birth cohorts including more than 200 000 mother-child pairs to examine whether maternal employment status during pregnancy was associated with birth weight, small for gestational age (SGA), term low birth weight

(LBW), length of gestation, and preterm delivery. Among employed pregnant women, we examined these pregnancy outcomes among 10 major occupational sectors where exposures potentially hazardous to reproduction are considered to be present. Heterogeneity in effects of all the above mentioned associations across cohorts was explored. This study is of exploratory nature and is the first step towards carrying out further analyses on specific occupational exposures and their potential health effects in children.

Methods

Study populations and data collection

As part of the European Union funded Environmental Health Risks in European Birth Cohorts (ENRIECO) project, an inventory was made of environmental exposure data, including maternal occupation, and health data in existing European birth cohort studies (17). Among the 37 cohorts included in the inventory, 15 matched our inclusion criteria of 1) having information on the maternal occupations in the period starting one month before pregnancy until birth and 2) having coded this information into occupational codes. All these cohorts were invited to participate in this study and 13 cohorts accepted. Of the two eligible cohorts that did not participate, 1 declined participation due to a lack of resources and 1 did not reply to the invitation.

A data transfer agreement document was signed by each cohort and datasets were transferred to the Centre for Research in Environmental Epidemiology with personal identifiers removed using a Secure File Transfer Protocol. Informed consent was obtained from all study participants as part of the original studies and in accordance with each study's institutional review board. The population sample was restricted to live-born

singleton infants, with known values for birth weight or length of gestation and available occupational history (N=222 317) (Supplemental material Table 1). A live birth was defined as an infant showing signs of life at a length of gestation of at least 22 completed weeks or weighting 500 g or more.

Occupational history

Information whether the mother worked during the period starting one month before conception until birth was obtained by self-reports or from questionnaires carried out by trained interviewers at different time points during pregnancy in each participating cohort. In five cohorts, occupational history covered the whole pregnancy period, including one month before pregnancy, whereas for others this information covered one or two trimesters or birth (Table 1). Pregnant women classified as non-employed included those that reported not being employed during the period starting one month before conception until birth but also housewives and students when specified. Self-employed women, for example those working in agriculture, were identified and included as employed.

Table 1 Description of the 13 European birth cohorts: occupational history information and occupational coding

Cohort	Setting location	Time period of enrolment	Occupational history information		Occupational coding	
			Time of collection during pregnancy	Period of pregnancy covered	Original code	Job title considered for this analysis
ABCD	The Netherlands (Amsterdam)	2003-2004	1 st trimester	1 st trimester	SBC-92 ^b	Job during the 1 st trimester of pregnancy
BAMSE	Sweden (Stockholm)	1994-1996	Birth	Birth	NYK-96 ^c	Job at birth
DNBC	Denmark (Denmark)	1996-2002	12 th weeks	1 month before and 1 st trimester	DISCO-88 ^d	Job at 15 th pregnancy weeks
Generation R	The Netherlands (Rotterdam)	2001-2006	30 th pregnancy weeks	All trimesters until 30 th weeks	ISCO-88	Job at 30 th pregnancy weeks
Generation XXI	Portugal (Porto)	2005-2006	Birth	All trimesters	ISCO-88	Job at birth
INMA_Granada ^a	Spain (Granada)	2000-2002	Birth	Birth	CNO-94 ^e	Job at birth
INMA_new ^a	Spain (Asturias, Gipuzkoa, Sabadell, Valencia)	2004-2008	12 th and 32 th weeks	1 month before and all trimesters until 32 th weeks	CNO-94 ^e	Only the longest occupation reported during pregnancy was codified

Cohort	Setting location	Time period of enrolment	Occupational history information		Occupational coding	
			Time of collection during pregnancy	Period of pregnancy covered	Original code	Job title considered for this analysis
KANC	Lithuania (Kaunas)	2007-2009	3 rd trimester	1 month before and 1 st and 3 rd trimesters	ISCO-88	Job during the 3 rd trimester of pregnancy
MoBa	Norway	1999-2008	17 th pregnancy weeks	17 th pregnancy weeks	ISCO-88	Job at 17 th pregnancy weeks
NINFEA	Italy	2005+	Once from the beginning of the pregnancy to maternity leave	Variable during pregnancy	ISCO-88	Job during the 1 st trimester of pregnancy
PÉLAGIE	France (Brittany)	2002-2006	1 st trimester	1 month before and 1 st trimester	ISCO-68	Job during the 1 st trimester of pregnancy
REPRO_PL	Poland (Lodz, Wroclaw, Lask, Kielce, Katowice, Legnica, Lublin, Szczecin,	2007-2011	8-12 th , 20-24 th , and 30-34 th weeks	1 month before and all trimesters until 30-34 th	ISCO-88	Job during the 1 st trimester of pregnancy
RHEA	Greece (Heraklion, Crete)	2007-2008	1 st and 3 rd trimesters	1 month before and all trimesters	ISCO-88	Job during the 1 st trimester of pregnancy

^aThe INMA cohort was represented by two cohorts (INMA_new and INMA_Granada) because the occupational information was collected at different times during pregnancy using different questionnaires and because INMA_Granada only included boys.

^bStandard Classification of Occupations of 1992: <http://www.cbs.nl/nl-NL/menu/methoden/classificaties/overzicht/sbc/1992/default.htm>.

^cNordic Standard Classification of Occupations of 1996: http://www.scb.se/Grupp/Hitta_statistik/Forsta_Statistik/Klassifikationer/_Dokument/oversattningsnyckel.pdf.

^dStatistics Denmark's Classification of Occupations of 1988. DISCO-88 is a slightly modified version of ISCO-88 and no translation was required.

^eNational Classification of Occupations of 1994: http://www.ine.es/clasifi/cno94_ciuo88.htm.

Classification of Occupations of 1988 (ISCO-88) (18). In cohorts where another classification of occupations was used we translated the codes using the conversion matrices available online into ISCO-88 (Table 1). In cohorts where occupational history was collected more than once during pregnancy, only the longest or the occupation reported at the first interview was used (Table 1). In cohorts where the mother reported being employed in more than one job at the same time, the first job reported was used in the analysis. Some wrong ISCO-88 codes were replaced by missing codes in job title information.

First, we studied the impact of employment status during pregnancy on birth outcomes comparing employed *vs* non-employed women. Secondly, considering that an employee can be exposed to a wide range of environmental exposures at workplace, we defined groups of occupations representing the main sectors of employment in women where potential reproductive hazards are considered to be present (10,11,19). Occupational exposures present in these sectors include a variety of chemical, biological and physical factors, together with factors related to work organization. We thus defined 10 broad occupational sectors, based on job titles, each sector representing a distinct occupational exposure pattern to chemical, physical and/or biological agents (Supplemental material Table 2): (i) Health: infectious agents, pharmaceutical products, ionizing radiation; (ii) Day-care: cleaning agents, mild infectious agents; (iii) Industry, commerce and home cleaning: cleaning agents; (iv) Agriculture and gardening: pesticides; (v) Electricity/electronic/optic workers: solvents, paints, glues, inks, welding, brazing; (vi) Laboratory work and chemical industry: laboratory and chemical products; (vii) Food industry: food additives; (viii) Printing and painting: solvents, paints, glues, inks; (ix)

Hairdressers: cosmetics; and (x) Construction workers: solvents, paints. This classification was based on 4-digit ISCO-88 codes except in the Generation XXI cohort where only 3-digit codes were available. A detailed list of the ISCO-88 codes included in each occupational sector can be found in Supplemental material Table 2. Some occupations were classified in more than one occupational sector (e.g. “Painters and related workers” (ISCO-88: 7141) were classified in the Printing and Painting and in the Construction workers sectors). Within these occupational sectors we further considered certain subsectors or job titles of interest: (i) nurses; (ii) personal care and related workers; (iii) housekeeping and restaurant services; and (iv) domestic and related helpers (Supplemental material Table 2). We selected these subsectors because they accounted for a large proportion of the participants’ occupation in the sector and because previous reports focused on them (20,21).

Birth weight and length of gestation

Birth weight (in grams) and length of gestation at delivery (in days) were obtained from clinical records (Supplemental material Table 3a and 3b). The last menstrual period (LMP)-based length of gestation estimation was taken if it was consistent by ≤ 7 days with the ultrasound (US)-based estimation. When both estimations differed by > 7 days, or the LMP measurement was unavailable, the US-based estimation was taken. If both measurements (LMP and US) were unavailable, the length of gestation based on maternal report measurement was taken. For the construction of the SGA variable, we first performed a conservative clean-up excluding the most extreme combinations of birth weight and length of gestation ($n=37$) (22). A SGA binary variable was then defined for each cohort as the 10% with the lowest birth weight stratified by length of gestation, child’s sex, maternal age, and parity. Preterm delivery was defined as birth before 37

weeks of gestation. Term LBW was defined as low birth weights ($<2500\text{g}$) at term (≥ 37 weeks) compared to normal birth weights ($\geq 2500\text{g}$) at term.

Covariates

Information on tobacco smoking, country of birth, health, education, and other covariates was obtained by self-reported questionnaires, by antenatal health care visits, or by medical/national birth registries (Supplemental material Table 3a and 3b). Potential confounding variables were defined as similarly as possible among the participating cohorts (Supplemental material Table 4a and 4b). For maternal education (Supplemental material Table 5) and maternal country of origin, cohort-specific categories were standardized to create common variables.

Statistical analysis

To assess the association between being employed during pregnancy and birth weight and length of gestation we used linear or logistic regression models for continuous and binary birth outcomes, respectively; non-employed women were used as the reference group. The same models were used to assess the association between working during pregnancy in a given occupational sector and birth outcomes; in this comparison the reference group for each specific occupational sector included all pregnant women working during pregnancy except in the occupational sector being assessed (thus including women working in the other 9 sectors). Therefore, the size and characteristics of the reference group differed in the analysis of each occupational sector.

We first developed separate regression models for each cohort. Models included subjects with available information on the exposure, outcome and potential confounders; subjects

with missing covariates were excluded from analysis. Models in each cohort were adjusted for maternal age (continuous in years), parity (0, 1, ≥ 2), maternal education (low, medium, high), maternal smoking during pregnancy (non-smoking, ≤ 10 cigarettes/day, >10 cigarettes/day), maternal BMI (<18.5 kg/m², 18.5-24.9 kg/m², 25-29.9 kg/m², ≥ 30 kg/m²), marital status (living with the father, living alone), and child's sex (boy, girl) (Supplemental material Table 4a and 4b). We also adjusted for maternal country of origin (European, non-European) only in cohorts where this information was available and heterogeneous (i.e. European *and* non-European): ABCD, BAMSE, Generation R, INMA_new, NINFEA, and PELAGIE (Supplemental material Table 4a and 4b). Length of gestation was additionally included in the models assessing term LBW. Length of gestation and the square of length of gestation were included in the models assessing birth weight. Models assessing SGA were not adjusted for length of gestation, maternal age, parity, and child's sex because the SGA definition was based on these variables.

Meta-analyses were used to estimate the overall summary effects of each association. To test the consistency of effect estimates across studies we used the Cochran's Q -test (23). If the Q -test was statistically significant ($p < 0.05$), indicating heterogeneity across populations, we used random effects analyses including cohort as random effect and all other covariates as fixed effects. Since the Q -test has low statistical power in meta-analysis with few number of studies we also used the I^2 statistic considering an $I^2 \geq 25\%$ indicating moderate heterogeneity (24). Therefore, when the Q -test was not statistically significant but the $I^2 \geq 25\%$ we also used random effects analyses. In case of heterogeneity, we used meta-regression to assess whether differences across cohorts were due to the timing during pregnancy when occupational history was collected (whole pregnancy period, 1st trimester, 2nd trimester, 3rd trimester, birth), the geographical region (Southern

cohorts: GENERATION XXI, INMA_Granada, INMA_new, NINFEA, PELAGIE, and RHEA vs Northern cohorts: ABCD, BAMSE, DNBC, GENERATION R, KANC, MoBa, and REPRO_PL), or the period of enrolment (before or after 2003).

Various sensitivity analyses were performed to assess the robustness of our results. Firstly, to test the influence of the largest cohorts, we repeated the analyses excluding DNBC or MoBa; we also excluded the non-European women in cohorts where this information was available. Secondly, to test whether a high proportion of elected caesarean procedures in a given cohort might explain a reduction in birth weight or length of gestation, we excluded those women with elective caesareans from the models. Thirdly, to test whether maternal diabetes and hypertension during pregnancy could affect the studied associations, we included these variables in the models. Fourthly, potential effect modification by maternal smoking during pregnancy was explored in stratified analyses. Finally, we repeated the models of occupational sectors under study using another group as a reference, defined as working women considered not to be occupationally exposed to reprotoxic agents and with similar socioeconomic status (i.e. managers, computer workers, teachers, administrative). Hence, the size of this “unexposed” reference group was the same in all models (n=64 179). This “unexposed” group is potentially more selective, particularly in terms of maternal age, education, smoking, and BMI but not in terms of pregnancy outcomes (Supplemental material Tables 6 and 7), than the reference groups for the main analysis and therefore was only used for sensitivity analysis. All statistical analyses were conducted with Stata 12.0 statistical software (Stata Corporation, College Station, Texas).

Results

A total of 222 317 mothers had known values for birth weight or length of gestation and their occupational history data was available. Of these, 177 983 (80%) declared being employed during pregnancy (Tables 2a and 2b and Supplemental material Table 1). Among employed pregnant women, 134 312 (76%) had their occupation coded; in MoBa, only 43% of all employees had maternal occupations already coded at the time of the analysis (Tables 2a and 2b). Employed pregnant women without coded job title information were excluded from the analysis on occupational sectors.

Overall, employed women were older, more likely to be non-smokers, with higher educational levels and lower BMIs, and longer length of gestation, compared to women non-employed during pregnancy (Supplemental material Tables 6 and 7). The ten occupational sectors under study accounted for 52% of the employed pregnant women with the highest proportion of women working in the health, day-care, and industry, commerce and home cleaning sectors (Tables 2a and 2b). The distribution of these three occupational sectors varied between cohorts: health and day-care were more prevalent in the Northern cohorts whereas industry, commerce and home cleaning were more prevalent in the Southern cohorts (Tables 2a and 2b).

Table 2a Distribution of employed women during pregnancy, employees with ISCO-88 codes, and the 10 occupational sectors and subsectors in the total population and in each cohort (Number and %)

	Total		ABCD		BAMSE		DNBC		Generation R		Generation XXI		INMA_Gra	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Employed women^a	177 983	80.2	5741	73.7	3536	91.1	70 757	81.6	5328	82.7	6070	77.2	233	46.8
Employees with ISCO-88^b	134312	75.5	5365	93.5	3536	100	70 015	98.9	5207	97.7	5994	98.7	220	94.4
Occupational sector/subsector^c	70133	52.2	1694	31.6	1509	42.7	37 500	53.6	1792	34.4	3467	57.8	745	33.6
Health	26522	19.7	577	10.8	506	14.3	14642	20.9	621	11.9	664	11.1	16	7.3
Nurses	11052	8.2	184	3.4	165	4.7	5344	7.6	302	5.8	135	2.3	5	2.3
Day-care	27215	20.2	360	6.7	567	16.0	16294	23.3	597	11.5	815	13.6	8	3.6
Personal care and related workers	12070	9.0	90	1.7	442	12.5	7572	10.8	296	5.7	379	6.3	4	1.8
Industry, commerce and home cleaning	13053	9.7	455	8.5	271	7.7	5073	7.2	346	6.6	2227	37.2	40	18.2
Housekeeping and restaurant services	3605	2.7	98	1.8	78	2.2	1797	2.6	116	2.2	641	10.7	10	4.6
Domestic and related helpers	2635	2.0	94	1.8	63	1.8	1235	1.8	17	0.3	386	6.4	4	1.8
Agriculture and gardening	1458	1.1	8	0.1	17	0.5	870	1.2	40	0.8	99	1.7	1	0.5
Electricity/electronic/optic workers	2193	1.6	28	0.5	103	2.9	1489	2.1	27	0.5	240	4.0	0	0.0
Laboratory work and chemical industry	4258	3.2	52	1.0	99	2.8	2839	4.1	46	0.9	161	2.7	0	0.0
Food industry	968	0.7	12	0.2	17	0.5	607	0.9	5	0.1	47	0.8	1	0.5

Printing and painting	2144	1.6	182	3.4	72	2.0	1147	1.6	101	1.9	116	1.9	1	0.5
Hairdressers	1890	1.4	58	1.1	57	1.6	791	1.1	46	0.9	289	4.8	9	4.1
Construction workers	693	0.5	1	0.0	10	0.3	601	0.9	2	0.0	1	0.0	1	0.5
'Unexposed' occupational group	64179	47.8	3671	68.4	2027	57.3	32 515	46.4	3415	65.5	2527	42.2	146	66.4
(sensitivity analysis)														

^aPercentage of the total women with occupational history available (N=222 317) (Supplemental material Table 1).

^bPercentage of the total employed women (N=177 983) (Supplemental material Table 1).

^cPercentage of the total employees with ISCO-88 codes (N=134 312) (Supplemental material Table 1). Because some participant's occupations were classified in more than one sector, the total number of pregnant women is greater than 70 133.

Table 2b Distribution of employed women during pregnancy, employees with ISCO-88 codes, and the 10 occupational sectors and subsectors in each cohort (Number and %)

	INMA_new		KANC		MoBa		NINFEA		PELAGIE		REPRO_PL		RHEA	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Employed women^a	2122	85.3	3744	88.0	72 963	77.7	2626	91.7	2957	89.0	998	84.9	908	81.7
Employees with ISCO-88^b	2122	100	3538	94.5	31 019	42.5	2504	95.4	2918	98.7	996	99.8	878	96.7
Occupational sector/subsector^c	1088	51.3	1220	34.5	18970	61.2	543	21.7	1495	51.2	358	35.9	423	48.2
Health	172	8.1	311	8.8	8144	26.3	263	10.5	480	16.4	67	6.7	59	6.7
Nurses	65	3.1	32	0.9	4488	14.5	83	3.3	196	6.7	19	1.9	34	3.9
Day-care	134	6.3	58	1.6	7835	25.3	79	3.2	384	13.2	42	4.2	42	4.8
Personal care and related workers	75	3.5	14	0.4	2984	9.6	24	1.0	169	5.8	2	0.2	19	2.2
Industry, commerce and home cleaning	607	28.6	606	17.1	2617	8.4	94	3.8	411	14.1	163	16.4	143	16.3
Housekeeping and restaurant services	111	5.2	118	3.3	517	1.7	20	0.8	32	1.1	17	1.7	50	5.7
Domestic and related helpers	191	9.0	61	1.7	369	1.2	8	0.3	158	5.4	8	0.8	41	4.7
Agriculture and gardening	15	0.7	11	0.3	217	0.7	7	0.3	58	2.0	12	1.2	103	11.7
Electricity/electronic/optic workers	25	1.2	19	0.5	155	0.5	13	0.5	68	2.3	22	2.2	4	0.5
Laboratory work and chemical industry	29	1.4	58	1.6	784	2.5	62	2.5	101	3.5	16	1.6	11	1.3
Food industry	37	1.7	47	1.3	145	0.5	2	0.1	31	1.1	4	0.4	13	1.5

Printing and painting	56	2.6	58	1.6	292	0.9	35	1.4	29	1.0	6	0.6	19	2.2
Hairdressers	58	2.7	44	1.2	410	1.3	13	0.5	54	1.9	21	2.1	40	4.6
Construction workers	26	1.2	11	0.3	29	0.1	0	0.0	8	0.3	1	0.1	2	0.2
‘Unexposed’ occupational group (sensitivity analysis)	1034	48.7	2318	65.5	12 049	38.8	1961	78.3	1423	48.8	638	64.1	455	51.8

^aPercentage of the total women with occupational history available (N=222 317) (Supplemental material Table 1).

^bPercentage of the total employed women (N=177 983) (Supplemental material Table 1).

^cPercentage of the total employees with ISCO-88 codes (N=134 312) (Supplemental material Table 1). Because some participant’s occupations were classified in more than one sector, the total number of pregnant women is greater than 70 133.

Employed women during pregnancy had an average lower child's birth weight (β_{adj} -9 95% CI -14 -3), a higher risk of SGA (OR_{adj} 1.09 95% CI 1.02-1.13) and a lower risk of preterm delivery than non-employed women (OR_{adj} 0.86 95% CI 0.81-0.91) (Tables 3a and b, Figure 1 and Supplemental material Figure 1). Working in most of the ten occupational sectors was not associated with the birth outcomes studied compared to not working in the occupational sector being assessed (Tables 3a and 3b). Nurse employment was associated with an average increase in child's birth weight of 11 g (95% CI 2 21) and a lower risk of the child being SGA (OR_{adj} 0.91 95% CI 0.84-0.99) (Table 3a and Supplemental material Figure 2). Personal care and domestic related work was associated with a decrease in child's birth weight (β_{adj} -21 95% CI -42 0 and β_{adj} -20 g 95% CI -40 -0, respectively), although the former association was nearly statistically significant (Table 3a). Working in the food industry was associated with an average decrease in length of gestation of 1 day (95%CI -1.9 to -0.2) and an increase in the risk of preterm delivery (OR_{adj} 1.50 95% CI 1.12-2.02) (Table 3b and Supplemental material Figure 3). Working in construction was associated with an average reduction in child's birth weight of 39 g (95% CI -78 -1) and with an increased length of gestation of 1.2 days (95% CI 0.1-2.2) (Tables 3a and 3b). No other associations were observed between any other selected occupational sector and birth outcomes (Tables 3a and 3b).

Table 3a Combined adjusted^a associations between employed women, occupational sectors/subsectors and birth weight, small for gestational age, and term low birth weight. Effect estimates (β or OR – 95%CI), N (Numbers) and Cases are presented for the total population

	Birth weight (g)			Small for gestational age			Term low birth weight		
	N	β	95% CI	Cases	OR	95% CI	Cases	OR	95% CI
Employed women^b	136 090	-8.5	-13.9, -3.1	13146	1.09	1.02, 1.13*	1499	0.92	0.71, 1.20*
Occupational sector/subsector^c									
Health	20334	2.8	-3.9, 9.6	1821	0.93	0.88, 0.98	185	1.04	0.89, 1.23
Nurses	8626	11.4	1.5, 21.4	730	0.91	0.84, 0.99	65	1.01	0.77, 1.32
Day-care	20385	5.9	-0.9, 12.7	1906	0.95	0.90, 1.00	188	1.01	0.85, 1.19
Personal care and related workers	8845	-20.9	-42.0, 0.3*	892	1.07	0.92, 1.24*	91	0.98	0.78, 1.23
Industry, commerce and home cleaning	9906	-1.3	-10.8, 8.2	1005	0.98	0.90, 1.05	179	1.09	0.90, 1.31
Housekeeping and restaurant services	2677	-2.4	-19.1, 14.2	293	1.06	0.94, 1.21	56	1.23	0.91, 1.66
Domestic and related helpers	1877	-20.0	-39.8, -0.2	210	1.08	0.93, 1.25	29	1.38	0.75, 2.55*
Agriculture and gardening	1087	10.3	-15.5, 36.1	104	1.05	0.85, 1.29	16	1.66	0.97, 2.85
Electricity/electronic/optic workers	1756	-5.0	-25.3, 15.4	159	1.00	0.85, 1.18	21	1.17	0.74, 1.85
Laboratory work and chemical industry	3316	2.8	-12.3, 17.9	322	1.00	0.77, 1.30*	29	1.01	0.69, 1.48
Food industry	749	5.8	-25.3, 36.9	74	1.01	0.79, 1.29	11	1.36	0.72, 2.57
Printing and painting	1583	-0.4	-21.8, 21.0	147	0.97	0.81, 1.15	23	1.43	0.92, 2.22
Hairdressers	1208	8.2	-16.4, 32.8	122	1.03	0.85, 1.24	16	1.17	0.41, 3.35*

Construction workers	509	-39.3	-77.6, -1.0	64	1.22	0.93, 1.60	7	1.56	0.71, 3.41
----------------------	-----	-------	-------------	----	------	------------	---	------	------------

N: Number of pregnant women with the information about the outcome of interest; Cases: number of pregnancy women with the outcome of interest.

^aBirth weight model adjusted for maternal age, parity, maternal education, maternal smoking during pregnancy, maternal BMI, marital status, infant's sex, length of gestation, length of gestation², and maternal country of origin in ABCD, BAMSE, Generation R, INMA_new, NINFEA, and PELAGIE; SGA model adjusted for maternal education, maternal smoking during pregnancy, maternal BMI, marital status, and maternal country of origin in ABCD, BAMSE, Generation R, INMA_new, NINFEA, and PELAGIE; Term low birth weight model adjusted for maternal age, parity, maternal education, maternal smoking during pregnancy, maternal BMI, marital status, infant's sex, length of gestation, and maternal country of origin in ABCD, BAMSE, GENERATION R, INMA_new, NINFEA, and PELAGIE.

^bReference group: non-employed women during pregnancy (N=44 334)

^cReference group: all other pregnant women working during pregnancy and with an ISCO-88 code but not included in the occupational sector being assessed.

*Random effects model (Q -test $p < 0.05$ and/or $I^2 \geq 25\%$)

Table 3b Combined adjusted^a associations between employed women, occupational sectors/subsectors and length of gestation and preterm delivery. Effect estimates (β or OR – 95%CI), N (Numbers) and Cases are presented for the total population

	Length of gestation (days)			Preterm delivery		
	N	β	95% CI	Cases	OR	95% CI
Employed women^b	136 452	0.4	-0.0, 0.9*	6143	0.86	0.81, 0.91
Occupational sector/subsector^c						
Health	20373	-0.1	-0.3, 0.1	889	1.03	0.96, 1.12
Nurses	8657	0.1	-0.5, 0.6*	370	1.08	0.97, 1.21
Day-care	20539	0.1	-0.1, 0.3	905	1.04	0.88, 1.22*
Personal care and related workers	8942	-0.1	-0.4, 0.2	414	1.02	0.83, 1.25*
Industry, commerce and home cleaning	9956	0.2	-0.1, 0.4	539	1.07	0.96, 1.18
Housekeeping and restaurant services	2699	0.1	-0.3, 0.6	150	1.08	0.90, 1.28
Domestic and related helpers	1907	-0.1	-0.6, 0.5	108	1.19	0.97, 1.46
Agriculture and gardening	1099	0.4	-0.3, 1.1	57	1.04	0.78, 1.38
Electricity/electronic/optic workers	1777	0.3	-0.7, 1.3	81	1.05	0.84, 1.32
Laboratory work and chemical industry	3340	-0.1	-0.5, 0.4	143	0.93	0.78, 1.11
Food industry	758	-1.0	-1.9, -0.2	50	1.50	1.12, 2.02
Printing and painting	1597	-0.0	-0.6, 0.6	78	1.07	0.85, 1.35

Hairdressers	1477	-0.3	-1.0, 0.4	75	1.19	0.92, 1.54
Construction workers	513	1.2	0.1, 2.2	15	1.01	0.33, 3.13*

N: Number of pregnant women with the information about the outcome of interest; Cases: number of pregnancy women with the outcome of interest.

^aModels adjusted for maternal age, parity, maternal education, maternal smoking during pregnancy, maternal BMI, marital status, infant's sex, and maternal country of origin in ABCD, BAMSE, Generation R, INMA_new, NINFEA, and PELAGIE.


^bReference group: non-employed women during pregnancy (N=44 334)

^cReference group: all other pregnant women working during pregnancy and with an ISCO-88 code but not included in the occupational sector being assessed.

*Random effects model (Q -test $p < 0.05$ and/or $I^2 \geq 25\%$)

There was little evidence for heterogeneity across cohorts in most of the associations observed. Heterogeneity was only observed in the associations between workers and SGA and between personal care sector and birth weight, but none of the variables tested were explanatory factors of such heterogeneity. After excluding one or the other of the two largest cohorts all the associations observed with birth weight or length of gestation disappeared. The association between construction workers and birth weight was mainly driven by the DNBC cohort, since this cohort represented more than 80% of all construction workers (Tables 2a and 2b). The associations observed with SGA or preterm delivery did not change significantly. These associations did not also change after excluding the non-Europeans or elective caesareans (Figure 2). Diabetes and hypertension during pregnancy did not change any of the aforementioned associations (data not shown). Moreover, there was no indication of modification by smoking (Figure 2). The models of occupational sectors did not change substantially when the “unexposed” reference group was used (Figure 2 and Supplemental material Tables 8a and 8b).


Figure 1 Combined adjusted^a associations between employed and non-employed women and preterm delivery. Effect estimates (OR - 95%CI) and numbers (N) are presented across cohorts and for the total population (Overall) and represent the number of employed women with preterm deliveries



Footnote:

^aAdjusted for maternal age, parity, maternal education, maternal smoking during pregnancy, maternal BMI, marital status, infant's sex, and maternal country of origin in ABCD, BAMSE, Generation R, INMA_new, NINFEA, and PELAGIE.

Figure 2 Sensitivity analyses of the combined adjusted^a associations between nurses and small for gestational age and food industry workers and preterm delivery. Effect estimates (OR - 95%CI) and numbers (N) are presented for the total population



Footnotes:

Abbreviations: ref.: reference.

Base model in each association was similar to the one restricted to subjects with information on type of delivery.

^aSmall for gestational age model adjusted for maternal education, maternal smoking during pregnancy, maternal BMI, marital status, and maternal country of origin in ABCD, BAMSE, Generation R, INMA_new, NINFEA, and PELAGIE. Preterm delivery model adjusted for maternal age, parity, maternal education, maternal smoking during pregnancy, maternal BMI, marital status, infant's sex, and maternal country of origin in ABCD, BAMSE, Generation R, INMA_new, NINFEA, and PELAGIE.

Discussion

This large study including more than 200 000 pregnant women of 13 birth cohorts in Europe suggests that employment during pregnancy, as compared to non-employment, is associated with a lower risk of preterm delivery. Working during pregnancy in most occupational sectors is not associated with birth weight or length of gestation compared to not working in the occupational sector being assessed. However, some associations were observed: working as a nurse was associated with favourable birth outcomes whereas food industry employment was associated with less favourable pregnancy outcomes. These findings were consistent across cohorts and across the various birth outcomes studied.

Being healthy is related to access and permanence in the labour market as well as to pregnancy outcome. Results of studies assessing the effects of maternal occupation and pregnancy outcomes are inconsistent, with studies showing no indications that prenatal employment status affects birth outcomes (4-8) and others showing an increased risk of adverse birth outcomes among women working more than 40 hours/week (4,7,26). Employed women in our population, compared to non-employed, had a lower risk of preterm delivery, an increased risk of SGA babies and reduced child's birth weight. However, this reduction in birth weight was only observed in the MoBa cohort and among babies born at term (data not shown). The higher risk of SGA babies was not observed in all cohorts and we could not identify the potential factors of this heterogeneity. In contrast, the association observed with preterm delivery was consistent across cohorts and after performing a series of sensitivity analysis. Studies in Poland and Portugal (3,6) found an increased risk of preterm delivery among non-employed women, particularly among women seeking employment. Our 'non-employed' group was a mixture of women with various statuses, including unemployed, women holding maternity or permanent leave, housewives, and students, and in most of cohorts we were not able to separate these groups. It is worth mentioning, that the main group who presented a lower risk of preterm deliveries among employed women was the non-exposed occupational group (i.e. managers, computer workers, teachers, administrative). This group had a higher social economical status compared with non-workers and also compared with the majority of the 10 occupational groups selected for our analysis. Therefore, the specific association with preterm delivery could indicate that social factors that differ between employed and non-employed are acting and have not been accounted for in the adjustment. This

association could be explained by the so-called ‘healthy worker bias’ which means that the observed association might be the result of residual confounding (25).

Among employed women, working as a nurse in our study was associated with favourable birth outcomes. These results contradict most of the previous studies where unfavourable effects among nurses were observed, often in relation with specific exposures (15,20,27-29); only one study conducted in Canada showed a lower risk of low birth weight infants among nurses (30). Nurses may engage healthier behaviours during pregnancy than other employees because they are close to preventive messages and have better pregnancy follow-up (31). We also found that working in the food industry was related with an increase in preterm delivery and reduced length of gestation. An elevated risk of adverse birth outcomes in these occupations has been described previously (8,32,33). Our findings of low birth weight babies and longer length of gestation among women working in construction agree with some epidemiological studies (12,13,34), but in our study these associations could only be observed in the largest cohorts, especially in DNBC. We also observed that personal care and domestic related work, more prevalent in the Southern cohorts, was associated with lower child’s birth weight but these associations were not consistent across cohorts and were no longer statistically significant after changing the reference group or after excluding the largest cohort.

Important strengths of this study are its large sample size and the prospective design of the majority of cohorts (in BAMSE, Generation XXI and INMA_Granada data was collected at birth), which makes it less vulnerable to recall bias. We also included a large set of potential confounding factors, usually collected directly from the mother, that have been associated with employment status, occupational exposures, and reproductive

outcomes (11). Registry-based studies are usually country-specific and sometimes results cannot be extrapolated to other countries. The present study includes data from 13 countries with a relatively good coverage across Europe and with different periods of inclusion, reflecting exposure to job conditions in Europe in the past 1-2 decades. We used occupational sectors, based on job titles, as markers of occupational exposure because information on job tasks and specific hazard exposures in the work setting was not available. Further, most specific job titles did not include sufficient numbers of women. The use of broad occupational sectors may have introduced misclassification, as the level/prevalence of exposure to specific hazards may greatly vary between occupations. This may have hampered the detection of associations even in the case of true effects although this was partly avoided when studying occupational subsectors with sufficient numbers and known high levels and prevalence of exposure to specific hazards. Substantial differences in the timing during pregnancy when occupational history was collected were encountered. In cohorts where occupational history only covered one or two trimesters or birth, we assumed that the employment situation remained stable during the whole pregnancy. We considered it unlikely that this assumption affected our findings, since we did not find evidence that this factor predicted heterogeneity across cohorts. Moreover, in cohorts where the mother reported being employed in more than one job at the same time or during pregnancy, the first job reported was used in the analysis. This assumes that the second job reported will be less hazardous for fetal growth which seems reasonable since change in occupation during pregnancy will usually go towards less hazardous exposure. LMP and US measurements are considered the best approaches to estimate length of gestation although they can be subject to recall, misclassification or measurement errors (35). Eleven of our cohorts had information on LMP and/or US for most of the mothers; BAMSE had maternal-based approach as the

only measurement and ABCD did not have information on the exact method of estimation.

Baseline participation proportion varied considerably among the different cohorts (from 43 to 92%). This baseline selection may limit the representativeness to the source populations in some of the studies, especially for the prevalence of the different occupations. However, it is not likely to affect internal validity, i.e. the associational estimate between occupational exposures and reproductive outcomes (36), which are the main aim of the current analysis. We need to consider that women not included in our analysis (N=12 897) were more likely to be younger, smokers and less educated and had less favourable birth outcomes as compared to women included (N=222 317) (data not shown). This has led to an under-representation of pregnant women of the most disadvantaged groups and who are at higher risk of pregnancy complications. We have defined occupational sectors based on potential exposure to chemical, biological and physical factors but other risk factors related to work organization such as stress, physical load or shift work could also be present and may have had an influence on preterm deliveries in particular. Maternal education was the only indicator of maternal socio-economic status available and since occupation is strongly related with social position, residual confounding by socio-economic status is likely. The comparison group in our main analysis also included women potentially exposed to environmental hazards at work and this may have lead to attenuate associations. Although we have found few associations of modest magnitude, using another reference group in a sensitivity analysis has strengthened our findings. Also, we conducted other sensitivity analyses and evaluated the robustness of results across cohorts and across the various birth outcomes. Multiple comparisons may have given rise to some chance findings in our analyses.

Despite the large sample size of the present study, this was still not sufficient to produce stable estimates of associations in occupational strata with few cases of SGA, preterm deliveries, and term LBW.

This collaborative project represents a first step towards carrying out further analyses such as the application of Job Exposure Matrices, which would allow us to assign specific occupational exposures of interest, to a whole range of job titles in population based studies (37,38).

Conclusions

The results of this collaborative effort of European birth cohorts suggest that employment during pregnancy is associated with a reduced risk of preterm birth and that work in certain occupations may be associated with changes in child's birth weight and length of gestation. This study provides a good picture of the overall impact on the course of pregnancy in important occupational sectors of the women's workforce in Europe. It provides also a useful platform on which to base further prospective studies of more specific maternal occupational exposures and child health and development after birth.

Acknowledgments

The authors would particularly like to thank all the cohort participants for their collaboration.

Reference List

1. European Agency for Safety and Health at Work. Women and Health at Work. http://osha.europa.eu/en/priority_groups/gender/index_html (accessed 21th July 2013).
2. Farrow A, Shea KM, Little RE. Birthweight of term infants and maternal occupation in a prospective cohort of pregnant women. The ALSPAC Study Team. *Occup Environ Med*. 1998;55(1):18-23.
3. Hanke W, Saurel-Cubizolles MJ, Sobala W, Kalinka J. Employment status of pregnant women in central Poland and the risk of preterm delivery and small-for-gestational-age infants. *Eur J Public Health*. 2001;11(1):23-28.
4. Jansen PW, Tiemeier H, Verhulst FC, Burdorf A, Jaddoe VW, Hofman A. Employment status and the risk of pregnancy complications: the Generation R Study. *Occup Environ Med*. 2010.
5. Kozhimannil KB, Attanasio LB, McGovern PM, Gjerdingen DK, Johnson PJ. Reevaluating the relationship between prenatal employment and birth outcomes: a policy-relevant application of propensity score matching. *Womens Health Issues*. 2013;23(2):e77-e85.
6. Rodrigues T, Barros H. Maternal unemployment: an indicator of spontaneous preterm delivery risk. *Eur J Epidemiol*. 2008;23(10):689-693.
7. Saurel-Cubizolles MJ, Zeitlin J, Lelong N, Papiernik E, Di Renzo GC, Breart G. Employment, working conditions, and preterm birth: results from the Europop case-control survey. *J Epidemiol Community Health*. 2004;58(5):395-401.
8. Savitz DA, Olshan AF, Gallagher K. Maternal occupation and pregnancy outcome. *Epidemiology*. 1996;7(3):269-274.

9. Burdorf A, Figa-Talamanca I, Jensen TK, Thulstrup AM. Effects of occupational exposure on the reproductive system: core evidence and practical implications. *Occup Med (Lond)*. 2006;56(8):516-520.
10. Figa-Talamanca I. Occupational risk factors and reproductive health of women. *Occup Med (Lond)*. 2006;56(8):521-531.
11. Palmer KT, Bonzini M, Harris EC, Linaker C, Bonde JP. Work activities and risk of prematurity, low birth weight and pre-eclampsia: an updated review with meta-analysis. *Occup Environ Med*. 2013;70(4):213-222.
12. Ahmed P, Jaakkola JJ. Maternal occupation and adverse pregnancy outcomes: a Finnish population-based study. *Occup Med (Lond)*. 2007;57(6):417-423.
13. Li X, Sundquist J, Sundquist K. Parental occupation and risk of small-for-gestational-age births: a nationwide epidemiological study in Sweden. *Hum Reprod*. 2010;25(4):1044-1050.
14. Ronda E, Moen BE, Garcia AM, Sanchez-Paya J, Baste V. Pregnancy outcomes in female hairdressers. *Int Arch Occup Environ Health*. 2010;83(8):945-951.
15. Suarez-Varela MM, Kaerlev L, Zhu JL, Bonde JP, Nohr EA, Llopi-Gonzalez A et al. Hospital work and pregnancy outcomes: a study in the Danish National Birth Cohort. *Int J Occup Environ Health*. 2009;15(4):402-409.
16. Tuntiseranee P, Olsen J, Chongsuvivatwong V, Limbutara S. Socioeconomic and work related determinants of pregnancy outcome in southern Thailand. *J Epidemiol Community Health*. 1999;53(10):624-629.
17. Vrijheid M, Casas M, Bergstrom A, Carmichael A, Cordier S, Eggesbo M et al. European birth cohorts for environmental health research. *Environ Health Perspect*. 2012;120(1):29-37.

18. ILO (International Labor Organisation). International Standard Classification of Occupations 1988.
<http://www.ilo.org/public/english/bureau/stat/isco/isco88/index.htm>. 1990.
19. Garcia AM, Gonzalez-Galarzo MC, Kauppinen T, Delclos GL, Benavides FG. A job-exposure matrix for research and surveillance of occupational health and safety in Spanish workers: MatEmESp. *Am J Ind Med*. 2013;56(10):1226-1238.
20. Quansah R, Jaakkola JJ. Occupational exposures and adverse pregnancy outcomes among nurses: a systematic review and meta-analysis. *J Womens Health (Larchmt)*. 2010;19(10):1851-1862.
21. Riipinen A, Sallmen M, Taskinen H, Koskinen A, Lindbohm ML. Pregnancy outcomes among daycare employees in Finland. *Scand J Work Environ Health*. 2010;36(3):222-230.
22. Alexander GR, Himes JH, Kaufman RB, Mor J, Kogan M. A United States national reference for fetal growth. *Obstet Gynecol*. 1996;87(2):163-168.
23. Cochran WG. The combinatin of estimates from different experiments. *Biometrics*. 1954;10(1):101-129.
24. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21(11):1539-1558.
25. Hernan MA, Hernandez-Diaz S, Robins JM. A structural approach to selection bias. *Epidemiology*. 2004;15(5):615-625.
26. Vrijkotte TG, van der Wal MF, van Eijdsden M, Bonsel GJ. First-trimester working conditions and birthweight: a prospective cohort study. *Am J Public Health*. 2009;99(8):1409-1416.

27. Fransman W, Roeleveld N, Peelen S, de Kort W, Kromhout H, Heederik D. Nurses with dermal exposure to antineoplastic drugs: reproductive outcomes. *Epidemiology*. 2007;18(1):112-119.
28. Lawson CC, Whelan EA, Hibert EN, Grajewski B, Spiegelman D, Rich-Edwards JW. Occupational factors and risk of preterm birth in nurses. *Am J Obstet Gynecol*. 2009;200(1):51-58.
29. Simcox AA, Jaakkola JJ. Does work as a nurse increase the risk of adverse pregnancy outcomes? *J Occup Environ Med*. 2008;50(5):590-592.
30. Arbour LT, Beking K, Le ND, Ratner PA, Spinelli JJ, Teschke K et al. Rates of congenital anomalies and other adverse birth outcomes in an offspring cohort of registered nurses from British Columbia, Canada. *Can J Public Health*. 2010;101(3):230-234.
31. Connolly MA, Gulanick M, Keough V, Holm K. Health practices of critical care nurses: are these nurses good role models for patients? *Am J Crit Care*. 1997;6(4):261-266.
32. Li X, Sundquist J, Kane K, Jin Q, Sundquist K. Parental occupation and preterm births: a nationwide epidemiological study in Sweden. *Paediatr Perinat Epidemiol*. 2010;24(6):555-563.
33. Meyer JD, Nichols GH, Warren N, Reisine S. Maternal occupation and risk for low birth weight delivery: assessment using state birth registry data. *J Occup Environ Med*. 2008;50(3):306-315.
34. Ronda E, Hernandez-Mora A, Garcia AM, Regidor E. [Maternal occupation, pregnancy length and low birth weight]. *Gac Sanit*. 2009;23(3):179-185.

35. Slama R, Ballester F, Casas M, Cordier S, Eggesbo M, Iñiguez C et al. Epidemiologic tools to study the influence of environmental factors on fecundity and pregnancy-related outcomes. *Epidemiol Rev.* 2014;36(1):148-164.
36. Rothman KJ, Gallacher JE, Hatch EE. Why representativeness should be avoided. *Int J Epidemiol.* 2013;42(4):1012-1014.
37. Bowman JD, Touchstone JA, Yost MG. A population-based job exposure matrix for power-frequency magnetic fields. *J Occup Environ Hyg.* 2007;4(9):715-728.
38. Brouwers MM, van Tongeren M, Hirst AA, Bretveld RW, Roeleveld N. Occupational exposure to potential endocrine disruptors: further development of a job exposure matrix. *Occup Environ Med.* 2009;66(9):607-614.

Contributions

Maribel Casas, Sylvaine Cordier, Martine Vrijheid, Mark Nieuwenhuijsen, and Joelle Fevotte designed the study with the contribution and approval of all the other co-authors. Joelle Fevotte and Sylvaine Cordier did the initial definition of the occupational groups. Each cohort provided the corresponding dataset to CREAL, participated in the interpretation of the data, and did a critical revision of the manuscript (ABCD: Tanja GM Vrijkotte, Manon van Eijsden; BAMSE: Eva Hallner, Inger Kull; DNBC: Anne-Marie Nybo Andersen, Pernille Stemann Larsen; Generation R: Alex Burdorf, Claudia Snijder; Generation XXI: Henrique Barros, Ana Cristina dos Santos; INMA_Granada: Mariana F Fernandez; INMA_new: Ana M García, Adonina Tardón, Loreto Santa Marina; KANC: Gediminas Balcius, Regina Gražulevičienė; MoBa: Merete Eggesbø, Karl-Christian Nordby; NINFEA : Lorenzo Richiardi, Daniela Zugna; PÉLAGIE: Sylvaine Cordier, Christine Monfort; REPRO_PL: Wojciech Hanke, Kinga Polanska; RHEA: Vaso Melaki, Evridiki Patelarou). Maribel Casas and David Martínez harmonized the datasets and

performed all the analyses. Nathalie Costet constructed the small for gestational age variable for each cohort and also contributed to the statistical analyses. Jens Peter Bonde, Manolis Kogevinas, Mark Nieuwenhuijsen and Joelle Fevotte contributed as occupational experts in the manuscript. Maribel Casas interpreted the data and wrote the first draft of the manuscript. Sylvaine Cordier and Martine Vrijheid directed the progress of the study, interpreted the data, and did a critical revision of all versions of the manuscript. All authors reviewed the manuscript for important intellectual content, and carefully read and approved this final version.

Competing interests

Nothing to declare

Funding

This work has received funding from the European Union's Seventh Framework Programme under grant agreement CHICOS n° 241604 and ENRIECO n° 226285.

Funding per cohort:

ABCD (Amsterdam Born Children and their Development study): The Netherlands Organization for Health Research and Development (ZonMw; grant number: 2100.0076)

BAMSE (The Stockholm Children Allergy and Environmental Prospective Birth Cohort Study): Swedish Heart-Lung Foundation, the Stockholm County Council, the Swedish Research Council, the Swedish Research Council for Health, Working life and Welfare, and the MedALL.

DNBC (Danish National Birth Cohort): The Danish National Research Foundation has established the Danish Epidemiology of Science Centre that initiated and created the DNBC. The cohort is furthermore a result of a major grant from this foundation.

Additional support for the DNBC is obtained from the Pharmacy Foundation, the Egmont Foundation, the March of Dimes Birth Defect Foundation, the Agustinus Foundation, and the Health Foundation.

Generation R: The first phase of the Generation R Study was made possible by financial support from the Erasmus Medical Center Rotterdam, the Erasmus University Rotterdam and the Netherlands Organization for Health Research and Development (ZonMw). The present study was conducted with support from the European Commission Seventh Framework Programme (FP7) for Research and Technology Development for the CONTAMED Project, EU grant agreement no. 212502.

Generation XXI: The Generation XXI cohort was funded by “Fundação para a Ciência e Tecnologia”, “Fundação Calouste Gulbenkian”, “Programa Operacional de Saúde – Saúde XXI” and “Administração Regional de Saúde do Norte”.

INMA_Granada (Environment and Childhood): This study was funded by grants from Instituto de Salud Carlos III (Red INMA G03/176, CB06/02/0041), Spanish Ministry of Health (FIS-07/0252), the EU Commission (QLK4-1999-01422, QLK4-2002-00603 and CONTAMED FP7-ENV-212502), and Consejería de Salud de la Junta de Andalucía (grant number 183/07).

INMA_new (Environment and Childhood): This study was funded by grants from UE (FP7-ENV-2011 cod 282957 and HEALTH.2010.2.4.5-1), Instituto de Salud Carlos III (Red INMA G03/176, CB06/02/0041, FIS-FEDER 03/1615, 04/1509, 04/1112, 04/1931, 05/1079, 05/1052, 06/1213, 07/0314, 09/02647, 11/01007, 11/02591, CP11/00178, FIS-PI06/0867, and FIS-PS09/00090), Conselleria de Sanitat Generalitat Valenciana, Spanish Ministry of Health (FIS-PI041436, FIS- PI081151, FIS-PI042018, FIS-PI09/02311), Generalitat de Catalunya-CIRIT 1999SGR 00241, Obre Social Cajastur, Universidad de

Oviedo, Department of Health of the Basque Government (2005111093 and 2009111069), the Provincial Government of Gipuzkoa (DFG06/004 and DFG08/001).

KANC (Kaunas cohort): This study was supported by grant FP6-036224 from the European Commission.

MoBa (The Norwegian Mother and Child Cohort Study): The study is supported by The Norwegian Ministry of Health, NIH/NIEHS (grant no. N01-ES-85433), NIH/NINDS (grant no. 1 U01 NS 047537) and Norwegian Research Council/FUGE (grant no. 151918/S10).

NINFEA (Birth and Infancy: Effects of the Environment): Compagnia San Paolo Foundation and Piedmont Region

PELAGIE (Endocrine disruptors: longitudinal study on pregnancy abnormalities, infertility, and childhood): The PELAGIE cohort is funded by the National Institute of Health and Medical Research (INSERM), the French Ministry of Health, the French Ministry of Labor, ANSES, ANR and InVS.

REPRO_PL (Polish Mother and Child Cohort): REPRO_PL cohort is supported in part by funds from National Centre for Research and Development, Poland (grant no. PBZ-MEiN-/8/2/2006; contract no. K140/P01/2007/1.3.1.1) and grant PNRF-218-AI-1/07 from Norway through the Norwegian Financial Mechanism within the Polish-Norwegian Research Fund.

RHEA (Mother Child Cohort in Crete): This work was partly supported by the EU Integrated Project NewGeneris, 6th Framework Programme, (Contract no. FOOD-CT-2005-016320) and by the EU funded project HiWATE, 6th Framework Programme (Contract no Food-CT-2006-036224).