



HAL
open science

Vegetation maps based on remote sensing are informative predictors of habitat selection of grassland birds across a wetness gradient

Aurélien G. Besnard, Aurélie Davranche, Sébastien Maugenest, Jan-Bernard Bouzillé, Alain Vian, Jean Secondi

► To cite this version:

Aurélien G. Besnard, Aurélie Davranche, Sébastien Maugenest, Jan-Bernard Bouzillé, Alain Vian, et al. Vegetation maps based on remote sensing are informative predictors of habitat selection of grassland birds across a wetness gradient. *Ecological Indicators*, 2015, 58, pp.47-54. 10.1016/j.ecolind.2015.05.033 . hal-01161554

HAL Id: hal-01161554

<https://univ-rennes.hal.science/hal-01161554>

Submitted on 10 Jun 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Vegetation maps based on remote sensing are informative**
2 **predictors of habitat selection of grassland birds across a wetness**
3 **gradient**

4 Besnard, A.G. ^{1*}, Davranche A. ², Maugenest S. ¹, Bouzillé J.B. ³, Vian A. ⁴, Secondi, J. ¹

5 **Running title:** Vegetation mapping highlights habitat selection

6 **E-mail addresses:**

7 aurelien.besnard@univ-angers.fr

8 aurelie.davranche@univ-angers.fr

9 sebastien.maugenest@univ-angers.fr

10 jan-bernard.bouzille@univ-rennes1.fr

11 alain.vian@univ-angers.fr

12 jean.secondi@univ-angers.fr

13 * *corresponding author*

14 ¹ LUNAM University, University of Angers, GECCO (Group Ecology and Conservation
15 of Vertebrates), 49045 Angers, France; Telephone number: 33 241 735 030

16 ² LUNAM University, University of Angers, LETG - LEESA, 2 Boulevard Lavoisier,
17 49045 Angers, France

18 ³UMR Ecobio Equipe Dynamique des communautés Campus de Beaulieu, 35042 Rennes,
19 France

20 ⁴IRHS, 2 Boulevard Lavoisier - 49045 Angers, France

21

22 Keywords: floodplain, vegetation mapping, SPOT 5, phytosociology, agri-environmental scheme,

23 Whinchat, Corncrake, Yellow Wagtail, Corn Bunting, Reed Bunting.

24 **Abstract**

25 Vegetation is a major environmental factor influencing habitat selection in bird species. High
26 resolution mapping of vegetation cover is essential to model the distribution of populations and
27 improve the management of breeding habitats. However, the task is challenging for grassland birds
28 because microhabitat variations relevant at the territory scale cannot be measured continuously over
29 large areas to delineate areas of higher suitability. Remote sensing may help to circumvent this
30 problem. We addressed this issue by using SPOT 5 imagery and phytosociological data. We mapped
31 grassland vegetation in a floodplain using two methods. We (i) mapped the continuous Ellenberg
32 index of moisture and (ii) identified 5 vegetation classes distributed across the wetness gradient.
33 These two methods produced consistent output maps, but they also provided complementary
34 results. Ellenberg index is a valuable proxy for soil moisture while the class approach provided more
35 information about vegetation structure, and possibly trophic resources. In spite of the apparent
36 uniformity of meadows, our data show that birds do not settle randomly along the moisture and
37 vegetation gradients. Overall birds tend to avoid the driest vegetation classes, i.e. the highest
38 grounds. Thus, vegetation maps based on remote sensing could be valuable tools to study habitat
39 selection and niche partition in grassland bird communities. It is also a valuable tool for conservation
40 and habitat management.

41

42 **Highlights**

- 43 • We mapped grassland vegetation in a floodplain using two methods
- 44 • The Ellenberg index proves to be a valuable proxy for soil moisture
- 45 • Vegetation classes provided more information about vegetation structure

- 46 • Grassland birds do not settle randomly along the vegetation gradient

- 47 • Vegetation maps based on remote sensing are useful tools to study habitat selection

49 **1. Introduction**

50 Delimiting zones of protection is a major issue of conservation programs (McNeely, 1994;
51 Moilanen et al., 2009). In order to identify core protection areas and optimize management, policy
52 makers need robust background information like precise ecological requirements for target species.
53 Grasslands present an interesting case in that respect because they often appear to human eyes as
54 large expanses of uniform vegetation. For this reason, designing efficient protection areas in
55 grasslands may seem challenging but this objective needs to be met. Due to anthropogenic changes,
56 grassland birds are threatened in several part of the World (Azpiroz et al., 2012; Brennan and
57 Kuvlesky, 2005; Tryjanowski et al., 2011; Tucker et al., 1994). Agri-Environmental Schemes (AES) were
58 implemented in the 1990's within the European Union to subsidize grassland management
59 compatible with breeding but many species have continued to decline (Kleijn et al., 2006). Precise
60 knowledge of habitat selection for target species is therefore crucial to design new and more
61 ecologically oriented AES measures in areas where conservation objectives partly failed.

62 Although their breeding habitat may seem homogeneous, grassland species do not settle at
63 random in meadows. Spatial variations in density are frequently observed. For instance, many
64 species are area-sensitive and avoid small fragments of habitats (Besnard and Secondi, 2014; Davis
65 and Brittingham, 2004; Helzer and Jelinski, 1999). Even in larger patches, they tend to avoid
66 landscape features like hedges because of higher predation risk (Morris and Gilroy, 2008). Vegetation
67 itself offers various level of suitability for nesting. Vegetation structure is a major feature that
68 influences the settlement of grassland birds (Fisher and Davis, 2010; Jacobs et al., 2012). Plant
69 community largely determines arthropod assemblages (Schaffers et al., 2008) and therefore the
70 quantity and quality of available trophic resources (Britschgi et al., 2006). Grassland birds tend to

71 prefer areas with higher densities of flowers during the breeding season (Fischer et al., 2012). This
72 preference may reflect higher prey availability to feed the young (Oppermann, 1990). Vegetation
73 cover also determines predation risk (Ejsmond, 2008), particularly the ability of birds to hide their
74 nests under the canopy (Whittingam and Evans, 2004). Therefore, grassland birds tend to select nest
75 site with taller swards and denser vegetation (Davis, 2005). In addition, some species need
76 herbaceous perches for foraging and territorial defence (Fischer et al., 2012; Oppermann, 1990).

77 Soil moisture is a major factor of grassland ecology (Price, 2002; Suzuki et al., 2006). It is
78 considered as the main driver of vegetation patterns in most regions (Moeslund et al., 2013). Flood
79 is a recurrent phenomenon that generates a wetness gradient and structures vegetation
80 communities (Martinez and Letoan, 2007). However, wetness is also influenced by local
81 environmental conditions like altitude, microtopography, or soil. Fortunately, vegetation has the
82 property to integrate all the components of wetness (Goward et al., 1991). Mapping vegetation in
83 relation to its affinity for moisture is expected to provide informative predictors to analyse habitat
84 selection in grassland birds and improve the management of these habitats. Satellite remote sensing
85 techniques are promising tools in this regard. They provide vegetation data with a spatial resolution
86 high enough to analyse habitat selection in birds across ecological gradients. Furthermore, satellite
87 images often cover geographical ranges large enough to delineate areas with different levels of
88 conservation priority (Guo, 2004; Poulin et al., 2010).

89 We assessed two methodological approaches to describe the variation of grassland vegetation
90 across the wetness gradient in floodplains. We used the Ellenberg moisture index that attributes a
91 value to each vegetal species, corresponding to its affinity for soil moisture (Ellenberg et al., 1992).
92 We computed a continuous wetness gradient based on the mean Ellenberg index of the local
93 vegetation community to map the wetness gradient across a floodplain. In addition, we tested a

94 discrete method to map vegetation classes as defined by the phytosociological approach (Tichý,
95 2002). A vegetation class is “a system of vegetal organisms with a floristic composition that is
96 statistically repetitive” (Biondi, 2011). Each class may offer a specific level of suitability for birds
97 depending on its physical properties and the various resources it provides. In grasslands, these
98 vegetation classes may therefore be an informative proxy to describe habitat requirements of bird
99 species. This approach may be easier to apply for managers since habitats are determined and their
100 conservation prioritized according to vegetation associations, for instance in the European Union
101 (Davies et al., 2004). Finally, AES are implemented in our study area to protect grassland and birds.
102 We tested if AES types were related to vegetation community, i.e. the hydrological functioning of the
103 floodplain, and if the spatial distribution of AES matched habitat selection of birds as described by
104 vegetation.

105

106 **2. Methods**


107 *2.1 Study area*

108 The study area covers the floodplains of the Loire River and its main tributaries around Angers
109 city in France (47.48, -0.56) (Fig. 1). Hydrological flow is relatively undisturbed by anthropogenic
110 developments in contrast to other floodplains of similar size in Western Europe (Hesselink et al.,
111 2003). Due to frequent floods, extensively managed grasslands still represent the main land cover
112 type. Agricultural practices consist in mowing meadows once a year in June-July and in allowing low
113 intensity grazing by cattle during vegetation regrowth. Considering the shallow slope of the ground,
114 plant community mainly depends on soil wetness and submersibility. Owing to their management,
115 these grasslands still host several patrimonial bird species, like the Corncrake (*Crex crex*), known to
116 be highly sensitive to intensive agriculture (Green et al., 1997a). Many grassland bird species
117 collapsed in the second part of 20th century, mainly due to the intensification of practices (more
118 efficient machinery, fertilisation, advanced mowing schedule). Agri-environmental schemes (AES)
119 that were implemented locally mainly consisted in delaying hay mowing. The proportion of grassland
120 under AES is quite high in the study area (45% in 2011). Several levels of AES measures are present in
121 the study area, differing mainly by the earliest mowing date. For clarity, we aggregated parcels in two
122 sets: those mown in June (3 540 ha) and those mown in July (1 571 ha). Mowing before July causes
123 high mortality in broods (Broyer, 2007; Green et al., 1997b). Therefore, we considered that meadows
124 mown in June were not suitable for the sustainable breeding of grassland birds.

125

126

Figure 1: Map of grassland distribution in the study area. The inset shows the location in Europe.



128

129

130 *2.2 Vegetation relevés and phytosociological classification*

131 We conducted phytosociological relevés on 107 plots to describe vegetation community across
132 the whole study area. Survey was carried out from May 16th to June 10th 2011 to ensure the reliable
133 identification of plant species. A plot was defined by the standard 16m² quadrat method (Chytrý and
134 Otýpková, 2003). Quadrats were selected in a larger zone of homogeneous vegetation to limit the
135 environmental effects of other land cover bordering the pixel of the relevé during the remote sensing

136 process. We sampled as many contrasting situations along the flooding gradient as possible basing
137 on our knowledge of the study area. Sampling plots were positioned using a Differential Global
138 Positioning System (DGPS), through a Trimble[®] Juno[®] SB. The estimated georeferencing error after
139 post-processing was 6 m maximum. However it was lower than 4m for 93% of plots. All species in the
140 quadrat were identified using either a local specific flora (Corillon, 1981) or a vegetative flora
141 (Eggenberg and Möhl, 2013) when necessary. Area covered by each species was evaluated using a
142 Braun-Blanquet coefficient (Braun-Blanquet, 1964). We then classified relevés with JUICE software
143 (Tichý, 2002), using the TWINSpan method (Hill, 1979), including standard relevés from a previous
144 study conducted in the same area (Foucault, 1984) as referentials for the classification. This method
145 classifies relevés according to their similarity in order to identify phytosociological taxa. We identified
146 five taxa along the wetness gradient in the area: 1. Phalaridion arundinaceae, Kopecky 1961; 2.
147 Gratiolo officinalis-Oenanthetum fistulosae inferior, De Foucault, 1984 nom. Ined; 3. Gratiolo
148 officinalis-Oenanthetum fistulosae superior, De Foucault 1984 nom. Ined; 4. Senecio aquatici-
149 Oenanthetum mediae, Bournerias et al. 1978; 5. Arrhenaterion elatioris, Hoch, 1926. The wettest
150 and the driest have been considered at the alliance level owing to the diversity of associations
151 encountered in the field for these two taxa. The three intermediate classes (2-4) have been specified
152 to the association level because their observed compositions matched the original descriptions.
153 However, the Gratiolo officinalis-Oenanthetum fistulosae was split in two due to the wide area it
154 covered and named in accordance with the type relevés described by De Foucault (1984). We tested
155 the significance of those partitions with a non-parametric multivariate analysis of variance
156 (Anderson, 2001) using *adonis* from the *vegan* R-package (Oksanen, 2013).

157

158 *2.3 Wetness Ellenberg index*

159 In order to quantify the wetness gradient, we used the wetness Ellenberg index. Ellenberg
160 indexes determine the affinity of each species for environmental parameters like light, temperature,
161 or nitrogen in soil (Ellenberg et al., 1992). We used here moisture affinity scores which are available
162 for each species (Hill et al., 1999). A value of 1 is given to a specialist of the driest habitats and a
163 value of 12 for a specialist of the wettest habitats. The indicator value of Ellenberg scores has been
164 evaluated by a field study that concluded that they were very good descriptors of the local
165 environment (Schaffers and Sýkora, 2000). For each plot, we averaged Ellenberg values of all species,
166 regardless of their cover, to obtain a single value. We considered this value as a proxy of soil wetness
167 at the plot scale. We computed average Ellenberg scores weighted by species cover and unweighted.
168 The unweighted method gave more realistic outputs. This is because cover depends mainly on the
169 sociability, for example the cover of grasses species is often high whereas sparse species like
170 *Fritillaria meleagris* can have a more informative value. In addition, weighted average penalize rare
171 species that often display narrow niches and are informative indicators.

172

173 *2.4 Mapping*

174 A SPOT 5[®] satellite image 2.5m colour, correction level 2B (allocation accuracy less than 10m),
175 was acquired on May 5th 2011. These scenes are provided with three bands: B1 (green: 0.50 to 0.59
176 μm), B2 (red: 0.61 to 0.68 μm), B3 (near-infrared NIR: 0.79 to 0.89 μm). A mask of all the meadows
177 within the floodplains was applied on this scene (60 km long x 60 km wide) to delineate the whole
178 study area using in ARCGIS version 10 (Environmental Systems Research Institute, Meudon France).
179 All analyses were conducted with this mask. Conditions of image acquisition were optimal with a low
180 incidence angle (4.77°) and cloud cover less than 10%. Spring 2011 was relatively dry so all the study
181 area was emerged but a small 41-ha zone that is kept flooded to provide spawning sites for fishes. A

182 vector layer of vegetation plots (quadrats) was created using their GPS coordinates. Circular buffers
 183 (diameter 4 m) centred on each point were drawn to get a vector layer of polygons of the vegetation
 184 plots. Then, we used the Spatial Analyst module to extract the mean radiance values of every
 185 sampling plot for each spectral band. We then calculated for each plot multispectral indices (Table 1)
 186 adapted for SPOT bands (see the method from Davranche *et al.* (2010 and 2013)).

187

188 **Table 1** Multispectral indices used in this study.

ID index	Multispectral index	References
1	Differential Vegetation Index (DVI)	Richardson & Everitt, 1992
2	Index of free water (IFW)	Adell & Puech, 2003
3	Normalized Difference Vegetation Index (NDVI)	Rouse <i>et al.</i> , 1973
4	Normalized difference water index (NDWI)	Mc Feeters, 1996
5	Optimized Soil Adjusted Vegetation Index (OSAVI)	Rondeaux <i>et al.</i> , 1996
6	B3	Spot image
7	B2	Spot image
8	B2/B1	This study
9	Soil Adjusted Vegetation Index (SAVI)	Huete, 1988
10	Brightness index (BI)	Kauth and Thomas, 1976
11	Simple ratio (SR)	Pearson & Miller, 1972
12	B1	Spot image
13	B1-B2	This study
14	$(B1-B2)/(B1+B2)$	This study
15	$B1^2/B2$	This study
16	Vegetation indice (VI)	Lillesand & Kiefer, 1987
17	Water index (WI)	Davranche et al, 2013
18	Water impoundment index (WII)	Caillaud <i>et al.</i> , 1987

189

190

191 Classification of the 5 vegetation classes was performed using a decision tree with a dichotomous
192 partitioning (Breiman et al., 1984). It was performed for each class using a binary coding: presence or
193 absence. We also tested a multiple-class classification, where we attempted to identify all classes in
194 the same analysis, but this method gave poor results. The Rpart package (Recursive PARTitioning,
195 Therneau and Atkinson, 1997) in the R software version 2.15.2 software (Rdevelopment core team,
196 2012) was used. This method is based on the cost complexity parameter (cp) for pruning. As
197 described in Davranche et al. (2010) for unbalanced samples, we used the cross-validation procedure
198 called CV-1SE (Esposito et al., 1999) for pruning with 10 subsets and iterative runs of the algorithm
199 (Breiman et al., 1984) to select the cp and the prior parameter. Cross-validation is well suited to small
200 samples (Breiman et al., 1984), so it can be recommended when no additional independent sample
201 is available.

202 The distribution of wetness Ellenberg index was checked for normality. We used linear models to
203 test whether index values obtained from field relevés could be predicted from remote sensing
204 variables). Models with all combinations of predictors were computed with the 'lm' function from
205 the R-package 'stats'. Model selection was carried out using the 'dredge' function of the R-package
206 'MuMIn' (Barton, 2013). The best model was selected according to the lowest AICc value (Burnham
207 and Anderson, 2002) (Appendix 1 in Supplementary material). As advised by Symonds and Moussali
208 (2011), we assessed the goodness-of-fit of this best AICc model, by calculating the coefficient of
209 determination (R^2) and the normalized root-mean-square error (NRMSE) between the predicted and
210 observed values.

211

212 *2.5 Bird census*

213 The study area host several breeding grassland species, some of which have been declining for
214 several decades in Western Europe (Tucker et al., 1994). We sampled the 4 dominant passerine
215 species in this small avian community: the Whinchat *Saxicola rubetra*, the Yellow Wagtail *Motacilla*
216 *flava*, the Corn Bunting *Emberiza calandra* and the Reed Bunting *Emberiza schoeniclus* (Noël, 2003).
217 We scanned every hay meadows of the study area during the 2011 breeding season to locate birds.
218 We used binoculars (10x42) and a spotting scope (20x-60x, 66mm) to detect and identify species.
219 Fieldwork was performed from April 13th to June 17th 2011 in good weather conditions (no
220 precipitation and low wind) to optimize bird detection and avoid the recording of migrating birds.
221 The time of day was not considered to maximize the area surveyed during the breeding season.
222 However, the detection rate of these species remains relatively high throughout the day, even in the
223 late afternoon when males rest on the top of vegetation. All occurrences were located on a map
224 (IGN© 1:25 000), and reported on a GIS software (ArcGIS© 10) by the same observer. We also used
225 the Corncrake database from LPO France (French official BirdLife partner). These data were collected
226 in 2011 by volunteer ornithologists who located singing males at night. Like for passerines, Corncrake
227 data were compiled in a GIS dataset. Overall, we observed 451 Whinchats, 151 Yellow Wagtails, 114
228 Corn Buntings, 174 Reed Buntings, and we used 479 occurrences of Corncrake across the study area.

229

230 2.6 Statistical analyses

231 In order to compare Ellenberg index values on bird locations with background values, we defined
232 1 501 background plots distributed across a 250m-grid. We created a buffer zone of 100m around
233 each bird location and background plot. We extracted the average wetness index for all pixels
234 included in the buffer area. For each bird species, we then compared mean wetness values on
235 location buffers and background plots using a Wilcoxon test based on bootstrap (1000 iterations of

236 background values with $n \geq 114$ corresponding to the sample size of each species) with the R software
237 (version 3.0.2).

238 Similarly, we extracted the mean area covered by each vegetation class within 100m-buffers
239 around each bird location. Knowing that each class was calculated independently, a pixel can be
240 assigned to several classes. We compared values for each class to the area covered by the same class
241 in the whole study area, and tested the differences between the two variables using a Wilcoxon test.
242 We measured selection or avoidance by birds for each class by calculating the difference in the
243 distributions of vegetation classes between bird and random location using background plots
244 described above ($n=1\ 501$).

245 In order to compare the two methods (Ellenberg wetness index and vegetation classes), we used
246 44 transects of 500m long surrounded with a 100m-buffer (i.e. 13.1 ha). These transects were
247 distributed across the floodplain in open meadows selected for their suitable landscape for grassland
248 birds. These were previously used with success to investigate the interest of the topographic
249 wetness index as a predictor of grassland bird distribution (Besnard et al., 2013). For the two
250 vegetation maps, we extracted mean values (Ellenberg index and vegetation class) on these transects
251 and we fitted a linear model between these two datasets. We also used Chi-squared tests to test
252 whether vegetation classes and bird distributions were randomly distributed across mowing dates.
253 All tests were carried out using R software (version 3.0.2). Finally, we could determine the pattern of
254 flooding susceptibility in 35 plots spread across the study area during a spring flood (April-June) in
255 2013. We sampled these plots twice a week before, during and after the flood, and derived a flood
256 index which corresponds to the number of times the plots were flooded. To determine whether
257 Ellenberg index was related to submersibility, we fitted a linear model between flood index and
258 mean Ellenberg index on plots.

259

260 3. Results

261 Using remote sensing method we could map the wetness gradient based on Ellenberg moisture
262 index across the hay meadows of the study area (Fig. 2; Best model : $df = 7$, $\text{LogLik} = 110.97$, $\text{AICc} =$
263 237.08 , $\text{weight} = 0,00048$, $R^2 = 0.41$, $\text{NRMSE} = 19\%$; $Y = 114,10 + 0,19 * \text{DVI} + 49,73 * \text{NDWI} + 1,65 * \text{B2} +$
264 $164,96 * \text{B2} / \text{B1} + 0,65 * \text{B1}^2 / \text{B2}$ (More details on model selection in Table S2 in Supplementary
265 material). The map was consistent with the observed flood susceptibility pattern ($F_{1,33} = 25.09$;
266 $p < 0.001$, adjusted $R^2 = 0.43$) (Fig. 3a). The resulting trees for the classification of vegetation classes
267 provided the following error rates C_1 : 9%, C_2 : 14%, C_3 : 16%, C_4 : 16%, C_5 : 10%. Plant species
268 composition was significantly different between classes as defined by Twinspan ($F_{1,106} = 5.62$;
269 $p = 0.001$). Using transect data, we found that average vegetation class was strongly correlated with
270 Ellenberg moisture index ($F_{1,42} = 116.1$; $p < 0.001$; adjusted $R^2 = 0.73$) (Fig. 3b) confirming the value of
271 these two methods to map wetness gradients. We noticed that the cover of class 5 is probably
272 overestimated on a part of the wettest area. This area was submerged by water several weeks before
273 the image capture and vegetation reflectance was modified by alluvium deposits. It was classified as
274 belonging both to the wettest classes (1-3) and class 5 but field data confirm that this latter class was
275 not observed there. This classification error causes an underestimation of avoidance of class 5 by
276 birds because this area exhibit high bird densities. Nevertheless, we detected an avoidance of this
277 class for 4 species. Thus, this limit does not alter the conclusion of this study.

278 All bird species except the Corn bunting settled in meadows with Ellenberg values higher than
279 random locations in the study area (Bootstrap based on Wilcoxon test; $p < 0.001$) (Fig. 4 and Table 2).
280 Yet, this species also tended to select plots with high Ellenberg values ($p = 0.10 \pm 0.16$). Generally,
281 birds selected wetter hay meadows than available in the study area (Fig. 4 and Table 2). The analysis


282 of vegetation classes provided supplementary information (Fig. 5). Except for the Corn Bunting, the
283 three wettest vegetation classes represented a larger proportion of the buffers around bird locations
284 than available (Table 3). For all species but the Whinchat, the driest vegetal class (5) represented a
285 smaller proportion of the buffer area around bird locations than expected by chance. For all species,
286 vegetation class 3 was more represented around bird location than available. Results were more
287 contrasting for vegetation class 4. It was more represented around bird locations for the Corn
288 Bunting and the Whinchat, whereas it was less represented for the Reed Bunting and the Corncrake.
289 More globally, the pattern of habitat selection for vegetation classes was different between species

290 The proportion of area covered by each vegetation class differed between AES contract types
291 ($\text{Chi}^2 = 350.75$, $\text{df} = 4$, $p < 0.001$) (Fig.5). The driest vegetal classes (5) covered a larger proportion of
292 the meadows available for mowing in June (47 %) than in July (23%). Logically, we observed the
293 reverse pattern for the wettest vegetal classes 1, 2 and 3 which represented a lower proportion of
294 the meadows mown in June (31%) than in July (55%). The proportion of the area covered by
295 vegetation class 4 was constant between both periods (22 %). Regarding mowing date, birds of all
296 species settled more frequently in parcels available for mowing in July than expected by chance (Chi^2
297 = 521.10, $\text{df}=4$, $p < 0.001$). In the study area only 22 % of meadow area was mown in July whereas
298 between 45 % and 66 % of birds settled in this parcels (Fig.5).

299

300 **Figure 2:** Maps extract of the study area representing with the wetness gradient as determined
301 by Ellenberg moisture index and vegetation classes (1 the wettest and 5 the driest). Grassland areas
302 were used as a mask prior to mapping.

303



304


305

306

307


308

Figure 3: a. Linear relationship between Ellenberg moisture index and flood susceptibility index as measured on 35 plots during a flood event in 2013. b. Linear relationship between Ellenberg moisture index and average vegetation class (1 is the driest, 5 the wettest class) in hay meadows (n=44).



309

310 **Figure 4:** Boxplots of distribution of Ellenberg moisture index calculated for 100m buffers around
 311 species occurrences and in the study area (available). * indicates that a species did not settle at
 312 random across the wetness gradient and significantly preferred wetter meadows (Wilcoxon signed-
 313 rank test).




314

315

316

317 **Figure 5:** Distribution of bird occurrences across vegetation classes in floodplain grasslands. The
 318 first histogram represents the proportion of each class in the study area. All other histograms
 319 represent the differences between the proportion of each vegetation class available in the study area
 320 and the proportion of these classes in a 100m-buffer around bird occurrences. Filled symbols indicate
 321 whether birds significantly selected or avoided this class (Wilcoxon signed-rank tests). Open symbols
 322 indicate that no selection or avoidance was detected (see values in Table 2). Vegetation classes were
 323 ranked from the wettest (1) to the driest (5).

324




325

326

327 **Figure 6:** (left panel) Comparison between the distribution of the five vegetation classes (from C₁
328 the wettest to C₅ the driest) observed across the grasslands of the study area and in the parcels
329 under AES contracts. Mowing dates were allowed either in June or in July. (right panel) Comparison
330 between the proportion of area available for mowing in June and July, and the proportion of the
331 population for each bird species on these two management options.

332



333

334

335

336 **Table 2:** Test of habitat selection comparing Ellenberg index in 100m buffers around bird
 337 occurrences and random background locations, and using bootstrap results of Wilcoxon
 338 tests. Selection is considered to occur when p-value < 0.05, in this case the value is in bold.
 339
 340

	p average	2.50%	97.50%
Yellow Wagtail	<0.001	<0.001	<0.01
Corn Bunting	0.10	<0.001	0.62
Reed Bunting	<0.001	<0.001	<0.001
Corncrake	<0.001	<0.001	<0.001
Whinchat	<0.001	<0.001	<0.001

341

342

343 **Table 2:** Cover of each vegetation class within a 100m buffer around species occurrences and
 344 random background locations. A species is considered to settle more often than by chance in a given
 345 class when p-value < 0.05 (value in bold). Vegetation classes were classified for the wettest (1) to the
 346 driest (5). V is the Wilcoxon statistic and p is the corresponding p-value.

	1		2		3		4		5	
	V	p	V	p	V	p	V	P	V	p
Whinchat	62531	p <0.001	72221	p <0.001	66832	<0.001	57279	p <0.05	46081	0.078
Yellow Wagtail	6593	0.112	8265	p <0.001	9339	<0.001	5082	0.223	3786	p <0.001
Corn Bunting	3279	0.998	3615	0.341	4744	<0.001	4404	p <0.01	1297	p <0.001
Reed Bunting	10291	p <0.001	13992	p <0.001	14941	<0.001	6178	p <0.05	3609	p <0.001
Corncrake	68357	p <0.001	94337	p <0.001	108846	<0.001	47725	p <0.01	10081	p <0.001

347

348

349 **4. Discussion**

350 Floodplain grasslands, the location the work was conducted, do not represent a uniform habitat.
351 Soil moisture gradient generates spatial variation in the composition of plant community (Fig. 2).
352 Using remote sensing, we characterized Ellenberg index at the pixel scale (2.5 m) so that we could
353 map the moisture gradient at a high spatial resolution. The ability of this index to represent wetness
354 was confirmed by the positive relationship with flooding susceptibility, a variable that we could
355 extract from direct observations in the field during a flood event (Fig 3a.). The vegetation community
356 showed some discontinuity across the soil moisture gradient though using the TWINSpan method
357 (Hill, 1979) we could define five phytosociological classes spread across the soil moisture range.
358 Average level of association in a plot was strongly related to Ellenberg wetness index (Fig 3b.). These
359 remote sensing data provided us with a layer of an informative predictor, the moisture gradient, to
360 analyse fine habitat selection of birds.

361 Basing on these data, we found that no bird species settled randomly across the grasslands of the
362 study area. They rather selected parcels depending on vegetation composition, which itself reflected
363 the moisture gradient. The wettest meadows (high values of Ellenberg index) hosted generally more
364 grassland birds in 2011 (Fig. 4), a distribution pattern observed in other years (Noël, 2003, A. Besnard
365 unpublished data). Even the Corn Bunting, which exploits a large range of open habitats (Donald and
366 Evans, 1995) showed a tendency to select the wettest meadows. Consistently, birds did not settle
367 randomly according to vegetation class. This approach provided additional information to the
368 wetness index. The Reed Bunting and the Corncrake, specialists of grasslands and wetlands with tall
369 sward (Brickle and Peach, 2004; Green et al., 1997a), settled in areas containing a higher proportion
370 of the wettest vegetation classes (1-3) than available across the floodplain (Fig. 3b). They also
371 avoided plots with a high proportion of the driest classes (4-5). Similarly, the Whinchat which is a

372 specialist of extensively managed hay meadows (Müller et al., 2005), and the Yellow Wagtail which
373 can breed in alternative habitats like crops (Gilroy et al., 2011) preferentially selected the wettest
374 meadows. Finally, the Corn Bunting selected grasslands with a higher proportion of classes 3 and 4,
375 and avoided both wettest and driest classes. Overall, no species selected areas covered by a large
376 proportion of the driest class (5), confirming the results obtained by the method based on Ellenberg
377 index. Interestingly, this latter class is the most represented in the study area (36%) whereas classes
378 1, 2 and 3 cover respectively only 10%, 9% and 21%. Species' responses to class 4 were more diverse.
379 The Corn Bunting settled in areas offering a large cover of this class, and the Whinchat tended to
380 slightly show the same pattern. In contrast, the Reed Bunting and the Corncrake tended to avoid
381 areas where this type of vegetation was more abundant than observed in average in the study area.
382 These results highlight interspecific differences in habitat selection and are consistent with the
383 ecology of these birds. Yet, differences were usually described for a larger range of ecological
384 conditions (Jacobs et al., 2012). It is thus particularly interesting to observe that it is possible to
385 determine species-specific habitat selection profiles within a much narrower ecological range.
386 Furthermore, species tended to keep their relative position in the wetness gradient with the
387 restricted ecological range of conditions encountered within the grasslands of these floodplains.

388 Our results suggest that the wettest grasslands (high Ellenberg index and large proportion of
389 classes 1-3) were the most suitable for birds. Intermediate class (4) were suitable for some species
390 and avoided by others but selection or avoidance was lower. In contrast, the driest grasslands (class
391 5) tended to be avoided by all species. We can draw hypotheses to explain this pattern. The wettest
392 classes can be characterized by a higher forb cover than drier class. We expect trophic resources (e.g.
393 pollinator insects) (Oppermann, 1990; Pywell et al., 2011) and vegetation structure to depend on the
394 grass/forbs ratio (Oppermann, 1990). All bird species are mainly insectivorous during the breeding
395 season (Brickle, 1999; Brickle and Peach, 2004; Britschgi et al., 2006; Davies, 1977; Green et al.,

396 1997a). Therefore, the quantity or quality of trophic resources may be higher on the wet end of the
397 moisture gradient than on its dry end. Vegetation structure may be of importance too as a higher
398 proportion of forbs may provide better sites for building nest and more perches for foraging and
399 social activities as well as reducing predation risk.

400

401 *Flood, meadow management and bird conservation*

402 Vegetation mapping based on remote sensing technique appears as a helpful tool to investigate
403 habitat selection in grassland habitats and to guide conservation actions. In our study area, the
404 proportion of any given vegetation class drastically differed between AES contract types (Fig. 5). Early
405 mown parcels (June) were mostly covered by the driest vegetal classes whereas parcels mown later
406 (July) were mostly covered by the wettest classes. This is a logical outcome of a scheme based on
407 volunteering. Farmers tend to select the contract level which minimizes the loss of hay quality. On
408 the driest meadows, that are mature early, farmers preferentially choose contracts with early
409 mowing dates. It results that the spatial pattern of mowing dates matches fairly well the wetness
410 gradient across the study area.

411 It is possible to describe the wetness gradient of an area using only topographical data (Besnard
412 et al., 2013; Beven and Kirkby, 1979). However, the low altitudinal variation in floodplains may not be
413 grasped by the resolution of available many digital elevation models, that are the raw data for
414 computing topographic methods, which limit our ability to map accurately the wetness gradient. Also
415 pixel size of these models may not be appropriate or relevant in species with small territories for
416 which small-scale features of the habitat may matter, for their nest site for instance. In contrast, the
417 high resolution provided by satellite imagery (2.5 m here) allows adjusting the spatial scale of the
418 analysis to the scale relevant for any bird species. Digital elevation models can be available with very

419 high altitudinal resolution in some areas, in which case it could be interesting to carry out a
420 comparative assessment of these two methods. However, it is important to outline that unlike
421 topographical methods the local plant community is an integrative measure of various processes
422 generating soil moisture (topography, soil composition and climate). Vegetation mapping takes also
423 into account anthropogenic modifications (like dams or drainages) which cannot be integrated by
424 topographic methods, for instance in cultivated lands where vegetation depends mainly on
425 agricultural processes. Finally, the remote sensing method is more ecologically realistic in naturally
426 sown grasslands where environmental factors, namely flood, largely influence vegetation. This
427 approach requires more resources for image acquisition, fieldwork, and botanical expertise than
428 topographical approaches. However, these constraints are balanced by deeper insight into habitat
429 selection of species, even in homogeneous landscapes, and probably the providing of more robust
430 information to base on to delineate areas for conservation.

431

432 **Acknowledgement**

433 The study was funded by Plan Loire Grandeur Nature, European Regional Development Fund (ERDF),
434 Région des Pays de la Loire and Agence de l'eau Loire-Bretagne. We are grateful to the LPO Anjou for
435 providing us Corncrake data. This project is part of Zone Atelier Loire. The Spot 5 imagery was
436 acquired thanks to the ISIS program. We also thank Xavier Solé and Antoine Body for their help in the
437 field.

438

439 **References**

- 440 Barton, K., 2013. MuMIn: Multi-model inference, R package version 1.9.13.
- 441 Anderson, M., 2001. A new method for non-parametric multivariate analysis of variance. *Austral*
442 *Ecol.* 32–46.
- 443 Azpiroz, A.B., Isacch, J.P., Dias, R. a., Di Giacomo, A.S., Fontana, C.S., Palarea, C.M., 2012. Ecology and
444 conservation of grassland birds in southeastern South America: A review. *J. F. Ornithol.* 83, 217–
445 246. doi:10.1111/j.1557-9263.2012.00372.x
- 446 Besnard, A.G., Jeunesse, I. La, Pays, O., Secondi, J., 2013. Topographic wetness index predicts the
447 occurrence of bird species in floodplains. *Divers. Distrib.* 19, 955–963. doi:10.1111/ddi.12047
- 448 Besnard, A.G., Secondi, J., 2014. Hedgerows diminish the value of meadows for grassland birds:
449 Potential conflicts for agri-environment schemes. *Agric. Ecosyst. Environ.* 189, 21–27.
450 doi:10.1016/j.agee.2014.03.014
- 451 Beven, K., Kirkby, M.M.J., 1979. A physically based, variable contributing area model of basin
452 hydrology/Un modèle à base physique de zone d'appel variable de l'hydrologie du bassin
453 versant. *Hydrol. Sci. J.* 24, 43–69.
- 454 Biondi, E., 2011. Phytosociology today: Methodological and conceptual evolution. *Plant Biosyst.* - An
455 *Int. J. Deal. with all Asp. Plant Biol.* 145, 19–29. doi:10.1080/11263504.2011.602748
- 456 Braun-Blanquet, J., 1964. *Pflanzensoziologie: grundzüge der vegetationskunde.*
- 457 Breiman, L., Friedman, J., Stone, C., Olshen, R., 1984. *Classification and regression trees*, Chapman & .
458 ed. New York.
- 459 Brennan, L., Kuvlesky, W., 2005. North American Grassland Birds: An Unfolding Conservation Crisis? *J.*
460 *Wildl. Manage.* 69, 1–13. doi:10.2193/0022-541X(2005)069<0001:NAGBAU>2.0.CO;2
- 461 Brickle, N., 1999. Diet of nestling Corn Buntings *Miliaria calandra* in southern England examined by
462 compositional analysis of faeces. *Bird Study* 46, 319–329.
- 463 Brickle, N.W., Peach, W.J., 2004. The breeding ecology of Reed Buntings *Emberiza schoeniclus* in
464 farmland and wetland habitats in lowland England. *Ibis (Lond. 1859)*. 146, 69–77.
- 465 Britschgi, A., Spaar, R., Arlettaz, R., 2006. Impact of grassland farming intensification on the breeding
466 ecology of an indicator insectivorous passerine, the Whinchat *Saxicola rubetra*: Lessons for
467 overall Alpine meadowland management. *Biol. Conserv.* 130, 193–205.
468 doi:10.1016/j.biocon.2005.12.013
- 469 Broyer, J., 2007. Définition d'un calendrier des fenaison compatible avec la reproduction des
470 passereaux. *Alauda* 75, 145–158.

- 471 Burnham, K. P., D. R. Anderson., 2002. Model selection and multimodel inference : a practical
472 information-theoretic approach. Second edition. Springer, New York, New York, USA.
- 473 Chytrý, M., Otýpková, Z., 2003. Plot sizes used for phytosociological sampling of European
474 vegetation. *J. Veg. Sci.* 563–570.
- 475 Corillon, R., 1981. Flore et végétation de la vallée de la Loire, Imprimerie. ed. Paris.
- 476 Davies, C.E., Moss, D., Hill, M.O., 2004. EUNIS habitat classification revised 2004.
- 477 Davies, N.B., 1977. Prey selection and social behaviour in wagtails. *J. Anim. Ecol.* 46, 37–57.
- 478 Davis, S., 2005. Nest-site selection patterns and the influence of vegetation on nest survival of mixed-
479 grass prairie passerines. *Condor* 605–616.
- 480 Davis, S.K., Brittingham, M., 2004. Area sensitivity in grassland passerines: effects of patch size, patch
481 shape, and vegetation structure on bird abundance and occurrence in southern Saskatchewan.
482 *Auk* 121, 1130–1145.
- 483 Davranche, A., Lefebvre, G., Poulin, B., 2010. Wetland monitoring using classification trees and SPOT-
484 5 seasonal time series. *Remote Sens. Environ.* 114, 552–562. doi:10.1016/j.rse.2009.10.009
- 485 Davranche, A., Poulin, B., Lefebvre, G., 2013. Mapping flooding regimes in Camargue wetlands using
486 seasonal multispectral data. *Remote Sens. Environ.* 138, 165–171.
487 doi:10.1016/j.rse.2013.07.015
- 488 Donald, P., Evans, A., 1995. Habitat selection and population size of Corn Buntings *Miliaria calandra*
489 breeding in Britain in 1993. *Bird Study* 42, 190–204.
- 490 Eggenberg, S., Möhl, A., 2013. Flora Vegetativa [Relié], 2th editio. ed. Bern.
- 491 Ejsmond, J., 2008. The effect of mowing on next-year predation of grassland bird nests: experimental
492 study. *Polish J. Ecol.* 56, 299–307.
- 493 Ellenberg, H., Weber, H.E., Düll, R., Wirth, V., Werner, W., Paulißen, D., 1992. Zeigerwerte von
494 Pflanzen in Mitteleuropa. *Scr. Geobot* 18, 1–258.
- 495 Esposito, F., Malerba, R.D., Semeraro, G., Tamma, V., 1999. The effects of pruning medods on the
496 predictive accuracy of indices. *Appl. Stoch. Model. Buisness Ind.* 15, 277–299.
- 497 Fischer, K., Busch, R., Fahl, G., Kunz, M., Knopf, M., 2012. Habitat preferences and breeding success
498 of Whinchats (*Saxicola rubetra*) in the Westerwald mountain range. *J. Ornithol.* 154, 339–349.
499 doi:10.1007/s10336-012-0898-z
- 500 Fisher, R.J., Davis, S.K., 2010. From Wiens to Robel: A Review of Grassland-Bird Habitat Selection. *J.*
501 *Wildl. Manage.* 74, 265–273. doi:10.2193/2009-020

- 502 Foucault, B. De, 1984. Systématique, structuralisme et synsystématique des prairies hygrophiles des
503 plaines atlantiques françaises. Univ. Rouen.
- 504 Gilroy, J.J., Anderson, G.Q. a., Vickery, J. a., Grice, P. V., Sutherland, W.J., 2011. Identifying
505 mismatches between habitat selection and habitat quality in a ground-nesting farmland bird.
506 Anim. Conserv. 14, no–no. doi:10.1111/j.1469-1795.2011.00480.x
- 507 Goward, S.N., Markham, B., Dye, D.G., Dulaney, W., Yang, J.L., 1991. Normalized Difference
508 Vegetation Index Measurements From The Advanced Very High-Resolution Radiometer.
509 Remote Sens. Environ. 35, 257–277.
- 510 Green, R., Rocamora, G., Schaffer, N., 1997a. Populations, ecology and threats to the Corncrake *Crex*
511 *crex* in Europe. Vogelwelt 118, 117–134.
- 512 Green, R., Tyler, G., Stowe, T., Newton, A., 1997b. A simulation model of the effect of mowing of
513 agricultural grassland on the breeding success of the corncrake (*Crex crex*). J. Zool. 243, 81–115.
- 514 Guo, X., 2004. Measuring Spatial and Vertical Heterogeneity of Grasslands Using Remote Sensing
515 Techniques. J. Environ. Informatics 3, 24–32. doi:10.3808/jei.200400024
- 516 Helzer, C.J., Jelinski, D.E., 1999. The relative importance of patch area and perimeter-area ratio to
517 grassland breeding birds. Ecol. Appl. 9, 1448–1458.
- 518 Hesselink, A., Weerts, H., Berendsen, H., 2003. Alluvial architecture of the human-influenced river
519 Rhine, The Netherlands. Sediment. Geol.
- 520 Hill, M.O., 1979. TWINSpan: A FORTRAN Program for Arranging Multivariate Data in an Ordered
521 Two-way Table by Classification of the Individuals and Attributes.
- 522 Hill, M.O., Mountford, J.O., Roy, D.B., Bunce, R.G.H., 1999. Ellenberg's indicator values for British
523 plants. ECOFACT Volume 2 Technical Annex.
- 524 Jacobs, R.B., Thompson, F.R., Koford, R.R., La Sorte, F. a., Woodward, H.D., Fitzgerald, J. a., 2012.
525 Habitat and landscape effects on abundance of Missouri's grassland birds. J. Wildl. Manage. 76,
526 372–381. doi:10.1002/jwmg.264
- 527 Kleijn, D., Baquero, R. a, Clough, Y., Díaz, M., De Esteban, J., Fernández, F., Gabriel, D., Herzog, F.,
528 Holzschuh, a, Jöhl, R., Knop, E., Kruess, a, Marshall, E.J.P., Steffan-Dewenter, I., Tscharrntke, T.,
529 Verhulst, J., West, T.M., Yela, J.L., 2006. Mixed biodiversity benefits of agri-environment
530 schemes in five European countries. Ecol. Lett. 9, 243–54; discussion 254–7. doi:10.1111/j.1461-
531 0248.2005.00869.x
- 532 Martinez, J., Letoan, T., 2007. Mapping of flood dynamics and spatial distribution of vegetation in the
533 Amazon floodplain using multitemporal SAR data. Remote Sens. Environ. 108, 209–223.
534 doi:10.1016/j.rse.2006.11.012
- 535 McNeely, J., 1994. Protected areas for the 21st century: working to provide benefits to society.
536 Biodivers. Conserv. 405, 390–405.

- 537 Moeslund, J., Arge, L., Bøcher, P., 2013. Topographically controlled soil moisture is the primary driver
538 of local vegetation patterns across a lowland region. *Ecosphere* 4, 1–26.
- 539 Moilanen, A., Wilson, K., Possingham, H., 2009. Spatial conservation prioritization: quantitative
540 methods and computational tools.
- 541 Morris, A.J., Gilroy, J.J., 2008. Close to the edge: predation risks for two declining farmland
542 passerines. *Ibis (Lond. 1859)*. 150, 168–177.
- 543 Müller, M., Spaar, R., Schifferli, L., Jenni, L., 2005. Effects of changes in farming of subalpine
544 meadows on a grassland bird, the whinchat (*Saxicola rubetra*). *J. Ornithol.* 146, 14–23.
545 doi:10.1007/s10336-004-0059-0
- 546 Noël, F., 2003. Étude de l'avifaune nicheuse des prairies inondables de fauche dans les Basses Vallées
547 Angevines. *Crex* 7, 53–58.
- 548 Oppermann, R., 1990. Suitability of different vegetation structure types as habitat for the whinchat
549 (*Saxicola rubetra*). *Vegetatio* 109–116.
- 550 Poulin, B., Davranche, A., Lefebvre, G., 2010. Ecological assessment of *Phragmites australis* wetlands
551 using multi-season SPOT-5 scenes. *Remote Sens. Environ.* 114, 1602–1609.
552 doi:10.1016/j.rse.2010.02.014
- 553 Price, E., 2002. *Lowland Grassland and Heathland Habitats (Habitat Guides)*. Routledge.
- 554 Pywell, R.F., Meek, W.R., Hulmes, L., Hulmes, S., James, K.L., Nowakowski, M., Carvell, C., 2011.
555 Management to enhance pollen and nectar resources for bumblebees and butterflies within
556 intensively farmed landscapes. *J. Insect Conserv.* 15, 853–864. doi:10.1007/s10841-011-9383-x
- 557 Schaffers, A., Raemakers, I., Sýkora, K., Braak, C. Ter, 2008. Arthropod assemblages are best
558 predicted by plant species composition. *Ecology* 89, 782–794.
- 559 Schaffers, A., Sýkora, K., 2000. Reliability of Ellenberg indicator values for moisture, nitrogen and soil
560 reaction: a comparison with field measurements. *J. Veg. Sci.* 11, 225–244.
- 561 Suzuki, R., Xu, J., Motoya, K., 2006. Global analyses of satellite-derived vegetation index related to
562 climatological wetness and warmth. *Int. J. Climatol.* 26, 425–438. doi:10.1002/joc.1256
- 563 Therneau, T.M., Atkinson, E.J., 1997. *An Introduction to Recursive Partitioning Using the RPART*
564 *Routines*.
- 565 Tichý, L., 2002. JUICE, software for vegetation classification. *J. Veg. Sci.* 451–453.
- 566 Tryjanowski, P., Hartel, T., Báldi, A., Szymański, P., Tobolka, M., Herzon, I., Goławski, A., Konvička, M.,
567 Hromada, M., Jerzak, L., Kujawa, K., Lenda, M., Orłowski, G., Panek, M., Skórka, P., Sparks, T.H.,
568 Tworek, S., & A.W., Żmihorski, M., 2011. Conservation of Farmland Birds Faces Different
569 Challenges in Western and Central-Eastern Europe. *Acta Ornithol.*
570 doi:10.3161/000164511X589857

571 Tucker, M., Heath, M.F., Tomialojc, L., Grimmett, R., 1994. Birds in Europe: Their Conservation Status
572 (Birdlife Conservation). BirdLife International.

573 Whittingam, M.J., Evans, K.L., 2004. The effects of habitat structure on predation risk of birds in
574 agricultural landscapes. *Ibis* (Lond. 1859). 146, 210–220.

575

576 SUPPLEMENTARY MATERIAL

577 **Table S1.** Model selection for mapping the ewetness index using the MuMin R package. Only models with $\Delta AICc < 1.5$ are shown.

578 Description of variables are given in Table 1 (1-1_ refer to ID index.

579

Intercept	Coefficient estimates																		Df	logLik	AICc	$\Delta AICc$	Weight	
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18						
114.10	0.19			49.73			1.65	-164.96												7	-110.97	237.08	0.00	0.00048
115.33				49.95		0.19	1.48	-167.03												7	-111.02	237.18	0.10	0.00046
105.32				42.22			1.36	-155.15		0.17										7	-111.13	237.40	0.32	0.00041
107.07		0.17		44.44			1.55	-156.43												7	-111.17	237.47	0.39	0.00040
84.54		-1.12		25.56				-131.68		1.25										7	-111.36	237.85	0.77	0.00033
60.41	3.29	-6.90						-109.44		3.71		-2.71								8	-110.40	238.27	1.19	0.00027
78.70		-1.57				1.67		-159.23			45.54									7	-111.67	238.48	1.40	0.00024
-19.01	5.30	-8.37								2.98		-2.36				-1.57	3.85			8	-110.52	238.51	1.44	0.00023
139.51	1.12	-0.96	-56.04				1.70	-185.53												8	-110.53	238.52	1.45	0.00023
139.25	1.11	-0.95					1.70	-185.38	-37.46											8	-110.53	238.53	1.45	0.00023
139.19	1.11	-0.95			-56.21		1.70	-185.35												8	-110.53	238.53	1.45	0.00023
100.51	0.10				-32.83		1.29	-142.17												7	-111.71	238.54	1.47	0.00023
100.49	0.10						1.29	-142.13	-21.86											7	-111.71	238.55	1.47	0.00023

100.38 0.10

-32.55

1.29 -141.93

-0.43

7 -111.72 238.56

1.49

0.00023

580 Richardson, A. J., & Everitt, J. H. (1992). Using spectra vegetation indices to estimate rangeland
581 productivity. *Geocarto International*, 1, 63-69.

582 Adell, C., & Puech, C. (2003). L'analyse spatiale des plans d'eau extraits par télédétection satellitale
583 permet-elle de retrouver la marque cynégétique en Camargue? *Bulletin de la société française*
584 *de photogrammétrie et de télédétection*, 172, 76-86.

585 Rouse, J. W., Haas, R. H., Schell, J. A., & Deering, D. W. (1973). Monitoring vegetation systems in the
586 great plains with ERTS. Third ERTS Symposium, NASA SP-351,1.

587 Rondeaux, G., Steven, & M, Baret, F. (1996). Optimization of Soil-Adjusted Vegetation Indices.
588 *Remote Sensing of Environment*, 55, 95-107.

589 McFeeters, S. K. (1996). The use of the normalised difference water index (NDWI) in the delineation
590 of open water features. *International Journal of Remote Sensing*, 17, 1425-1432.

591 Symonds, MRE., & Moussalli, A., (2011). A brief guide to model selection, multimodel inference and
592 model averaging in behavioural ecology using Akaike's Information Criterion, *Behavioral Ecology and*
593 *Sociobiology*, 65, 13-21.

594 Huete, A. R. (1988). A Soil-Adjusted Vegetation Index (SAVI). *Remote sensing of Environment*, 25,
595 295-309.

596 Pearson, R.L., & Miller, L.D. (1972). *Remote mapping of standing crop biomass for estimation of the*
597 *productivity of the short-grass Prairie, Pawnee National Grasslands, Colorado*. Processing of the
598 8th International Symposium on Remote Sensing of Environment, ERIM, Ann Arbor, MI, 1357-
599 1381.

600 Lillesand, T. M., & Kiefer, R. W. (1987). *Remote sensing and image interpretation*, 2e édition, John
601 Wiley and Sons, New York, 721 p.

602 Caillaud, L., Guillaumont, B., & Manaud, F. (1991). *Essai de discrimination des modes d'utilisation*
603 *des marais maritimes par analyse multitemporelle d'images SPOT. Application aux marais maritimes*
604 *du Centre Ouest*. IFREMER report (H4.21) 485, 24 p.

605