



**HAL**  
open science

# Efficiency of Industrially Relevant Atropisomeric Diphosphines in Copper-Catalyzed 1,4-Asymmetric Conjugate Addition of Dialkylzincs to Cyclic or Acyclic Enones or Dienones

Marie S. T. Morin, Thomas Vives, Olivier Baslé, Christophe Crévisy, Virginie Ratovelomanana-Vidal, Marc Mauduit

## ► To cite this version:

Marie S. T. Morin, Thomas Vives, Olivier Baslé, Christophe Crévisy, Virginie Ratovelomanana-Vidal, et al.. Efficiency of Industrially Relevant Atropisomeric Diphosphines in Copper-Catalyzed 1,4-Asymmetric Conjugate Addition of Dialkylzincs to Cyclic or Acyclic Enones or Dienones. *Synthesis: Journal of Synthetic Organic Chemistry*, 2015, 47 (17), pp.2570–2577. 10.1055/s-0034-1378813 . hal-01188236

**HAL Id: hal-01188236**

**<https://univ-rennes.hal.science/hal-01188236v1>**

Submitted on 19 Oct 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

---

# Efficiency of Industrially Relevant Atropisomeric Diphosphines in Copper-Catalyzed 1,4-Asymmetric Conjugate Addition of Dialkylzincs to (a)Cyclic (di)Enones

Marie S. T. Morin<sup>a</sup>, Thomas Vives<sup>a</sup>, Olivier Baslé<sup>a</sup>, Christophe Crévisy<sup>\*a</sup>, Virginie Ratovelomanana-Vidal<sup>b\*</sup>, Marc Mauduit<sup>a\*</sup>

<sup>a</sup>Ecole Nationale Supérieure de Chimie de Rennes, CNRS UMR 6226  
11 Allée de Beaulieu, CS 50837, 35708, Rennes Cedex 7, France

<sup>b</sup>PSL Research University, Chimie ParisTech - CNRS, Institut de Recherche de Chimie Paris, 75005 Paris, France

Fax: +33 22 323 8112

E-mail: marc.mauduit@ensc-rennes.fr

**Received:** The date will be inserted once the manuscript is accepted.


---

**Abstract:** Industrially relevant atropisomeric diphosphines such as BINAP, SYNPHOS and DIFLUORPHOS have demonstrated their efficiency in the copper-catalyzed asymmetric conjugate addition of various dialkylzincs to  $\alpha$ -aryl enones,  $\alpha$ -aryl dienones and cyclic dienones. Excellent 1,4- or 1,6-regioselectivities and enantioselectivities (up to 97% ee) were reached, even with challenging sterically hindered Michael acceptors.

**Key words:**  $\alpha$ -Aryl enones; Dienones; Conjugate addition; Copper; Dialkylzinc reagents; BINAP; SYNPHOS, DIFLUORPHOS

---


In the field of modern asymmetric synthesis, the metal-catalyzed Asymmetric Conjugate Addition (ACA) represents one of the most powerful methods to build valuable optically active compounds for the pharmaceutical and agrochemical industries.<sup>1</sup> In this regard, intensive research in the ligand design has been performed over the last two decades, notably for copper-catalyzed ACA.<sup>2</sup> Among the plethora of phosphorus-based ligands described in this area, atropisomeric chiral phosphoramidites and phosphines have been deeply studied affording the expected 1,4-adducts in good yields and remarkable enantioselectivities.<sup>2</sup> Curiously, although atropisomeric diphosphines were successfully used in the Cu-catalyzed addition of trialkylaluminium or Grignard reagents to various cyclic and/or acyclic  $\alpha,\beta$ -unsaturated carbonyls (ketones,<sup>3</sup> esters<sup>4</sup>, thioesters<sup>5</sup> and  $\alpha$ -ketoesters<sup>6</sup>) (Scheme 1, A), to the best of our knowledge, no report concerned the addition of less reactive hard nucleophiles such as dialkylzinc reagents. Very recently, we serendipitously discovered the efficiency of (*R*)-BINAP (**L-1**) as a ligand in the Cu-catalyzed 1,4-ACA of diethylzinc to linear enones (ees ranging from 82 to 99%).<sup>7</sup> Remarkably, the Cu/BINAP catalytic system appeared particularly efficient with  $\delta$ -branched  $\alpha,\beta$ -unsaturated aryl ketones. The value of these catalytic processes allowed the efficient development of a highly regio- and enantioselective sequential 1,6/1,4 addition involving acyclic dienones (de >97%) (Scheme 1, B). As a result of these pioneering results, we were prompted to extend our study and to evaluate a series of industrially relevant atropisomeric diphosphine ligands such as BINAP (**L-1**), (*S*)-SYNPHOS (**L-2**)<sup>8</sup> and (*R*)-DIFLUORPHOS (**L-3**)<sup>9</sup> in the selective conjugate addition of dialkylzincs to various  $\alpha$ -aryl enones,  $\alpha$ -aryl dienones and cyclic dienones (Scheme 1, C).



**Scheme 1** Copper-catalyzed conjugate additions involving atropisomeric ligands

The screening of atropisomeric ligands **L1-3** was started with the conjugate addition of diethylzinc to the sterically hindered cyclohexyl-substituted  $\alpha$ -phenyl enone **1a** (Table 1), a challenging Michael acceptor that has been scarcely studied to date.<sup>10</sup> Under optimized reaction conditions (copper-thiophene-2-carboxylate (CuTc)/**L-X** 5 mol%, THF, room temperature, Table 1, entry 1), (*R*)-BINAP (**L-1**) afforded the expected 1,4-adduct **2a** in 92% isolated yield and high enantioselectivity (97% ee). At lower catalyst loading, the enantioselectivity of the addition decreased to 88% ee (Table 1, entry 2).

Table 1 Screening of atropisomeric ligands in 1,4 Cu-ACA of dialkylzinc to  $\alpha$ -aryl enones<sup>a</sup>



Entry	<b>1</b>	Ar	R	Alk	L-X	Product	Yield (%) <sup>b</sup>	ee (%) <sup>c</sup>		
1	<b>1a</b>	Ph	Cy	Et	<b>L-1</b>		<b>92</b>	<b>97 (S)</b>		
2 <sup>d</sup>					<b>L-1</b>				91	88 (S)
3					<b>L-2</b>				92	90 (R)
4					<b>L-3</b>				98	93 (S)
5 <sup>e</sup>	<b>1b</b>	Ph	Me	Et	<b>L-1</b>		74	99		
6	<b>1c</b>	2-Nph	Cy	Et	<b>L-1</b>		89	94		
7	<b>1a</b>	Ph	Cy	<i>i</i> -Pr	<b>L-1</b>		54	12		
8	<b>1d</b>	Ph	CH <sub>2</sub> <i>t</i> -Bu	Et	<b>L-1</b>		<b>96</b>	<b>91</b>		
9					<b>L-2</b>				96	-89
10					<b>L-3</b>				98	83
11 <sup>e</sup>	<b>1e</b>	Ph	<i>i</i> -Bu	Et	<b>L-1</b>		85	94		
12	<b>1d</b>	Ph	CH <sub>2</sub> <i>t</i> -Bu	Me	<b>L-1</b>		44	93		
13 <sup>f</sup>	<b>1e</b>	Ph	<i>i</i> -Bu	Me	<b>L-1</b>		30	91		
14	<b>1f</b>	Ph	<i>t</i> -Bu	Et	<b>L-1</b>		22 <sup>g</sup>	0		

<sup>a</sup> Reaction conditions: **1a-f** (0.5 mmol), Et<sub>2</sub>Zn (1.5 mmol), CuTc (0.025 mmol), **L-X** (0.025 mmol), THF (0.5 mL), rt, 14 to 36 h.

<sup>b</sup> Isolated yield. <sup>c</sup> Determined by GC on **2** (see ESI).

<sup>d</sup> Reaction performed with 2 mol% of CuTc/**L-1**. <sup>e</sup> A 1:1.5 MeTHF/Cyclohexane ratio was used at -40°C, see ref. 7.

<sup>f</sup> A 1:1.5 MeTHF/Cyclohexane ratio was used at 0°C, see ref. 7.


<sup>g</sup> <sup>1</sup>H NMR yield with internal standard (see ESI).

In the case of (*S*)-SYNPHOS (**L-2**) and (*R*)-DIFLUORPHOS (**L-3**), similar excellent yields were reached (up to 98%) but with slightly lower enantioselectivity (90% and 93% ee, respectively, entries 2-3). One should note that the results obtained for **2a** are equivalent to the best results reported in the literature both in term of isolated yield and enantioselectivity.<sup>11</sup> Moreover, in comparison with the less hindered substrate **1b** bearing a methyl group,<sup>7</sup> the erosion of enantioselectivity remained exceptionally quite low (entry 4, 99% ee vs entry 1, 97% ee). On the other hand, the replacement of the phenyl ketone function by the more bulky 2-naphthyl ketone fragment only led to a slight decrease of the efficiency to form the corresponding adduct **2c** with excellent 89% isolated yield and 94% enantiomeric excess (entry 6). In view of these studies, we decided to extend the scope by attempting the addition of a bulkier alkyl nucleophile such as *i*-Pr<sub>2</sub>Zn (entry 7). The corresponding 1,4-adduct **2d** was isolated for the first time with a moderate isolated yield of 54%, despite a prolonged reaction time (36 h). Unfortunately, the enantioselectivity of the addition was quite low (12% ee). The latter result illustrated a limit of the BINAP ligand (**L-1**) toward sterically-demanding ACA involving organozinc reagents as nucleophiles. We pursued our ligand screening in the addition of Et<sub>2</sub>Zn to the newly synthesized  $\alpha$ -phenyl enone **1d** bearing a *tert*-butylCH<sub>2</sub> side chain (entries 8-10). Again, the BINAP ligand (**L-1**) appeared to be the most efficient one, affording the 1,4-adduct **2e** in excellent 96% isolated yield and 91% ee, while (*S*)-SYNPHOS (**L-2**) and (*R*)-DIFLUORPHOS (**L-3**) reached 89 and 83% ee, respectively. The steric hindrance from

the CH<sub>2</sub>*t*-Bu group had slightly altered the enantioselectivity of the addition in comparison with adduct **2f** bearing an *isobutyl* side chain (91 vs 94% ee, entries 8 and 11). Noteworthy, the addition of the less reactive dimethylzinc to enone **1d** was less productive, affording the corresponding 1,4-adduct **2g** in 44% isolated yield, while the enantioselectivity remained excellent reaching 93% ee, in the same range as the previously reported adduct **2h** (entries 12-13). Unfortunately, the addition of diethylzinc to the more sterically-demanding enone **1f** bearing a *t*-Bu moiety was unsuccessful leading only to racemic product **2i** in 22% yield (entry 14).<sup>11</sup> In comparison with previously reported copper-catalytic systems involving BINAP-type ligands for the 1,4-ACA of alkyl Grignard<sup>3,4</sup> or trialkylaluminum<sup>6</sup> reagents to electron deficient unsaturated systems, our methodology gave similar efficiency both in term of isolated yields and enantioselectivities.

The capacity to control both the regio- and the enantioselectivity in metal-catalyzed conjugate addition involving electrophilic extended unsaturated systems remains a challenge for organic chemists and the development of new catalytic systems is highly desirable.<sup>12</sup> Therefore, having proved the efficiency of BINAP (**L-1**) toward simple aliphatic  $\alpha$ -aryl enones, we then decided to extend this method to the more challenging acyclic and cyclic dienic Michael acceptors to produce either 1,6- or 1,4-adducts.<sup>13</sup> First, we tried the addition of diethylzinc to  $\alpha$ -phenyl dienone **1g** (Table 2). We were pleased to observe that BINAP (**L-1**) promoted an excellent regioselectivity in favor of the 1,4-adduct **2j** (96/4 ratio) with 82% isolated yield after 3 h at room temperature and up to 92% ee (entry 1). This 1,4-selectivity was also recently observed with **L-1** for the addition of AlMe<sub>3</sub> on an  $\alpha$ -ketoester analogue, as reported by Alexakis and Gremaud (98% ee).<sup>6</sup> When (*S*)-SYNPHOS (**L-2**) and (*R*)-DIFLUORPHOS (**L-3**) were used, the additions were completed after 5 to 6 h to furnish **2j** with a similarly high level of regioselectivity (97/3 and 95/5 respectively, entries 2-3) but with lower ees (54% and 80% respectively). It is important to notice that enantioenriched 1,4-adducts such as **2j** could be easily transformed into valuable synthetic intermediates through the post-oxidation of the phenylketone as reported by Coates and Sowerby during the synthesis of Zizaene<sup>14</sup> or the styrenyl side chain as described by Zhang and co-workers.<sup>13f</sup>

Table 2 Screening of atropisomeric ligands in 1,4 Cu-ACA of Et<sub>2</sub>Zn to  $\alpha$ -aryl dienone **1g**<sup>a</sup>



Entry	L-X	Time (h)	Ratio <b>2j</b> / <b>2j'</b> <sup>b</sup>	Yield (%) <sup>c</sup>	ee (%) <sup>d</sup>
1	<b>L-1</b>	3	96/4	82	92
2	<b>L-2</b>	5	97/3	70	-54
3	<b>L-3</b>	6	95/5	66	80

<sup>a</sup> Reaction conditions: **1g** (0.5 mmol), Et<sub>2</sub>Zn (1.5 mmol), CuTc (0.025 mmol), **L-X** (0.025 mmol), THF (0.5 mL), 0 °C, 3 to 6 h.

<sup>b</sup> Determined by <sup>1</sup>H NMR spectroscopy. <sup>c</sup> Isolated yield of **2j**/**2j'**.


<sup>d</sup> Determined by GC on **2** (see ESI).

We finally focused our attention on the enantioselective addition of diethylzinc to the cyclic dienone **1h** (Table 3). Although the relative steric hindrances at the  $\beta$  and  $\delta$  positions should favour the 1,6-addition, it was previously observed that the regioisomeric outcome of such a reaction is also under strong dependence of the nature of both the nucleophile and the chiral ligand.<sup>15</sup> Using a Cu/diphosphine **L1-3** catalytic system, a complete regioselectivity in favour of the 1,6-adduct **2k** was observed (after re-conjugation in the presence of DBU). With (*R*)-BINAP (**L-1**), complete conversion occurred after 5 h at room temperature, furnishing **2k** in 60% isolated yield and 82% ee (entry 1). At lower temperature, the enantioselectivity was not improved and the yield decreased significantly to 26% despite 21 h of reaction (entry 2). A similar level of enantio-induction was reached with (*S*)-SYNPHOS (**L-2**), however the addition was less productive after 14 h of reaction at room temperature (54% of yield, entry 3). (*R*)-DIFLUORPHOS (**L-3**) appeared also less efficient under the same reaction conditions, yielding the 1,6-adduct in only 14% yield with 74% of enantioselectivity (entry 4).

In conclusion, we have demonstrated the efficiency of atropisomeric diphosphines ligands in the enantioselective conjugate addition of dialkylzinc reagents onto  $\alpha$ -aryl enones,  $\alpha$ -aryl dienones as well as cyclic dienones by using CuTc as a catalyst. The industrially relevant (*R*)-BINAP proved superior to both (*S*)-SYNPHOS and (*R*)-DIFLUORPHOS as a ligand for this transformation both in terms of productivity and enantioselectivity, providing valuable 1,4- and 1,6-

adducts in relatively good isolated yields and up to 97% *ee*. Unfortunately, highly sterically demanding enones or bulky organozinc reagents remain problematic leading only to moderate yields and low enantioselection.

Table 3 Screening of atropisomeric ligands in 1,6 Cu-ACA of Et<sub>2</sub>Zn to cyclic dienone **1h**<sup>a</sup>



Entry	L-X	T (°C)	Time (h)	Ratio <b>2k/2k'</b> <sup>b</sup>	Yield (%) <sup>c</sup>	<i>ee</i> (%) <sup>d</sup>
1	<b>L-1</b>	rt	5	>98/2	60	82
2	<b>L-1</b>	0	21	>98/2	26	82
3	<b>L-2</b>	rt	14	>98/2	54	-82
4	<b>L-3</b>	rt	14	>98/2	14 <sup>e</sup>	74

<sup>a</sup> Reaction conditions: **1h** (0.5 mmol), Et<sub>2</sub>Zn (1.5 mmol), CuTc (0.025 mmol), **L-X** (0.025 mmol), THF (0.5 mL), rt, 5 to 21 h.

<sup>b</sup> Determined by <sup>1</sup>H NMR spectroscopy. <sup>c</sup> Isolated yield of **2k**.

<sup>d</sup> Determined by GC on **2** (see ESI). <sup>e</sup> <sup>1</sup>H NMR yield with internal standard (see ESI).

Further developments for such challenging reactions are currently under investigation in our group and will be applied in the enantioselective synthesis of bio-relevant targets. In that context, BINAP is a commercially available and rather inexpensive chiral ligand that has already demonstrated its potential for industrial applications.<sup>16</sup> Consequently, the fact that a Cu/BINAP system catalyses efficiently and selectively various ACA of dialkylzincs to both enones and dienones is of a great interest in organic synthesis.

All reactions were performed under an argon atmosphere using oven-dried glassware. Et<sub>2</sub>Zn and *i*-Pr<sub>2</sub>Zn were purchased from Aldrich and used without further purification. Me<sub>2</sub>Zn was purchased from Acros and used without further purification. Substrates **1a**<sup>17</sup>, **1f**<sup>18</sup>, **1g**<sup>19</sup> and **1h**<sup>13b</sup> were synthesized as reported in the literature. THF was distilled from Na/benzophenone under argon and dichloromethane was distilled from calcium hydride under nitrogen. All other chemical reagents and solvents were obtained from commercial sources and used without further purification. <sup>1</sup>H (400 MHz), <sup>13</sup>C (100 MHz), NMR spectra were recorded on a Bruker ARX400 spectrometer with complete proton decoupling for nuclei other than <sup>1</sup>H. Chemical shifts are reported in ppm with the solvent resonance as the internal standard (CDCl<sub>3</sub>, <sup>1</sup>H: δ 7.26 ppm, <sup>13</sup>C: δ 77.16 ppm). Data are reported as follows: chemical shift (δ in ppm, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, quint = quintuplet, sept = septuplet, m = multiplet), coupling constants (Hz) and integration. Optical rotations were recorded using a polarimeter Perkin-Elmer 341. High resolution mass spectrometry analyses were performed at the Centre Régional de Mesures Physiques de l'Ouest (CRMPO), Université de Rennes 1. Enantiomeric excesses were measured using gas chromatography (capillary column - Chiraldex Beta-dex, 30 m x 0.25 mm id, 0.25 μm or G-TA, 30 m x 0.25 mm id, 0.12 μm) or HPLC (Chiralcel OJ-H or IA columns).

#### (*E*)-3-cyclohexyl-1-(naphthalene-2-yl)prop-2-en-1-one (**1c**)

According to the literature procedure<sup>17</sup> and starting from cyclohexanecarboxaldehyde (5 mmol), the desired enone **1c** was obtained in 61% yield (806 mg).

<sup>1</sup>H NMR (400 MHz; CDCl<sub>3</sub>): δ 8.44 (s, 1H), 8.02 (dd, *J* = 8.6 Hz, 1.7 Hz, 1H), 7.97 (d, *J* = 8.0 Hz, 1H), 7.90 (t, *J* = 9.0 Hz, 2H), 7.62-7.53 (m, 2H), 7.09 (dd, *J* = 15.5 Hz, 6.5 Hz, 1H), 6.99 (dd, *J* = 15.5 Hz, 1.0 Hz, 1H), 2.35-2.26 (m, 1H), 1.90-1.71 (m, 5H), 1.41-1.19 (m, 5H). <sup>13</sup>C NMR (101 MHz; CDCl<sub>3</sub>): δ 191.2, 155.0, 135.6, 135.5, 132.6, 130.0, 129.6, 128.5, 128.3, 127.9, 126.8, 124.7, 123.4, 41.2, 32.0, 26.11, 25.9.

HRMS (ESI) : *m/z* calcd for C<sub>19</sub>H<sub>21</sub>O: 265.15924; [M+H]<sup>+</sup> found 265.1593 (0 ppm).

#### (*E*)-5,5-dimethyl-1-phenylhex-2-en-1-one (**1d**)

According to the literature procedure<sup>17</sup> and starting from 3,3-dimethylbutyraldehyde (4 mmol), the desired enone **1d** was obtained in 56% yield (457 mg).

<sup>1</sup>H NMR (400 MHz; CDCl<sub>3</sub>): δ 7.95-7.92 (m, 2H), 7.57-7.53 (m, 1H), 7.49-7.44 (m, 2H), 7.10 (dt, *J* = 15.4 Hz, 7.7 Hz, 1H), 6.86 (dt, *J* = 15.3 Hz, 1.3 Hz, 1H), 2.21 (dd, *J* = 7.8 Hz, 1.3 Hz, 2H), 0.97 (s, 9H).

<sup>13</sup>C NMR (101 MHz; CDCl<sub>3</sub>): δ 190.6, 147.4, 138.0, 132.5, 128.4<sub>8</sub>, 128.4<sub>6</sub>, 127.9, 47.3, 31.6, 29.5.

HRMS (ESI) : *m/z* calcd for C<sub>14</sub>H<sub>18</sub>O: 225.12554; [M+Na]<sup>+</sup> found 225.1253 (1 ppm).

### 1,4-asymmetric conjugate addition on $\alpha$ -aryl enones: Typical Procedure

A flame dried Schlenk flask, under an argon atmosphere, was charged with (*R*)-BINAP ligand (12.5 mg, 0.02 mmol, 5 mol%), CuTc (3.8 mg, 0.02 mmol, 5 mol%) and dry THF (0.5 mL). The resulting mixture was stirred for 10 min at room temperature and to this solution was added dropwise a solution of dialkylzinc reagent (3 equiv.) The reaction mixture was stirred for 10 min at room temperature and a solution of substrate **1** (0.4 mmol, 1 equiv.) in 0.5 mL of dry THF was added. The reaction mixture was stirred at room temperature for 14 h for the product **2a**, 16 h for **2c** and **2e**, 36 h for **2d** and 24 h for **2g** and **2i**. The reaction was quenched by the addition of ethanol (4 mL). The mixture was stirred for 15 min and filtered through a short column of silica gel with EtOAc. The solvents were removed under reduced pressure. The crude mixtures were purified by flash chromatography on silica gel (pentane/Et<sub>2</sub>O: 98/2) to afford the corresponding products as colorless oils. Ee was obtained by GC or HPLC analysis of the purified product.

#### 3-cyclohexyl-1-phenylpentan-1-one (**2a**)

Using the typical procedure with (*R*)-BINAP, (*S*)-**2a** was obtained in 92% yield (90 mg) and 97% ee (HPLC: Chiralcel IA column, *n*-Hexane/*i*-PrOH: 99/1, 0.5 mL/min., R<sub>T1(major)</sub> = 11.8 min, R<sub>T2(minor)</sub> = 13.2 min.). [ $\alpha$ ]<sub>D</sub><sup>20</sup> = -3.4 (*c* 1, CHCl<sub>3</sub>). Data were in accordance with the literature.<sup>10b,20</sup>

#### 3-cyclohexyl-1-(naphthalen-2-yl)pentan-1-one (**2c**)

Using the typical procedure with (*R*)-BINAP, **2c** was obtained in 89% yield (105 mg) and 94% ee (HPLC: Chiralcel OJ-H column, *n*-Hexane, 0.5 mL/min., R<sub>T1(major)</sub> = 38.2 min., R<sub>T2(minor)</sub> = 44.3 min.). [ $\alpha$ ]<sub>D</sub><sup>20</sup> = -15.6 (*c* 1, CHCl<sub>3</sub>).

<sup>1</sup>H NMR (400 MHz; CDCl<sub>3</sub>):  $\delta$  8.47 (s, 1H), 8.04 (dd, *J* = 8.6 Hz, 1.8 Hz, 1H), 7.98 (d, *J* = 7.9 Hz, 1H), 7.90 (t, *J* = 7.8 Hz, 2H), 7.63-7.54 (m, 2H), 3.11 (dd, *J* = 16.1 Hz, 5.6 Hz, 1H), 2.91 (dd, *J* = 16.1 Hz, 7.5 Hz, 1H), 2.07-2.00 (m, 1H), 1.78-1.53 (m, 5H), 1.49-1.44 (m, 2H), 1.38-1.05 (m, 6H), 0.90 (t, *J* = 7.4 Hz, 3H).

<sup>13</sup>C NMR (101 MHz; CDCl<sub>3</sub>):  $\delta$  201.1, 135.6, 135.0, 132.7, 129.6<sub>9</sub>, 129.6<sub>7</sub>, 128.5, 128.4, 127.9, 126.8, 124.2, 41.4, 40.5, 40.4, 30.5, 29.5, 27.0, 26.9<sub>5</sub>, 26.9<sub>2</sub>, 24.2, 12.1.

HRMS (ESI) : *m/z* calcd for C<sub>21</sub>H<sub>26</sub>ONa: 317.18814; [M+Na]<sup>+</sup> found 317.1877 (1 ppm).

#### 3-cyclohexyl-4-methyl-1-phenylpentan-1-one (**2d**)

Using the typical procedure with (*R*)-BINAP, **2d** was obtained in 54% yield (56 mg) and 12% ee (HPLC: Chiralcel IA column, *n*-Hexane/*i*-PrOH: 99/1, 0.5 mL/min., R<sub>T1(major)</sub> = 11.6 min, R<sub>T2(minor)</sub> = 13.4 min.). [ $\alpha$ ]<sub>D</sub><sup>20</sup> = -0.7 (*c* 1, CHCl<sub>3</sub>).

<sup>1</sup>H NMR (400 MHz; CDCl<sub>3</sub>):  $\delta$  8.00-7.97 (m, 2H), 7.57-7.53 (m, 1H), 7.49-7.44 (m, 2H), 2.88 (dd, *J* = 17.7 Hz, 5.9 Hz, 1H), 2.80 (dd, *J* = 17.7 Hz, 5.1 Hz, 1H), 2.04 (quint, *J* = 5.6 Hz, 1H), 1.92-1.80 (m, 1H), 1.75-1.66 (m, 3H), 1.65-1.52 (m, 2H), 1.43-1.34 (m, 1H), 1.27-0.93 (m, 5H), 0.90 (d, *J* = 6.8 Hz, 3H), 0.81 (d, *J* = 6.8 Hz, 3H).

<sup>13</sup>C NMR (101 MHz; CDCl<sub>3</sub>):  $\delta$  200.7, 137.6, 132.8, 128.6, 128.1, 44.1, 40.3, 37.7, 31.8, 30.1, 28.8, 26.9<sub>4</sub>, 26.9<sub>1</sub>, 26.8, 21.6, 18.9.

HRMS (ESI) : *m/z* calcd for C<sub>18</sub>H<sub>26</sub>ONa: 281.18814; [M+Na]<sup>+</sup> found 281.1883 (1 ppm).

#### 3-ethyl-5,5-dimethyl-1-phenylhexan-1-one (**2e**)

Using the typical procedure with (*R*)-BINAP, **2e** was obtained in 96% yield (84 mg) and 91% ee (HPLC: Chiralcel OJ-H column, *n*-Hexane, 0.2 mL/min, R<sub>T1(major)</sub> = 38.4 min., R<sub>T2(minor)</sub> = 43.9 min.). [ $\alpha$ ]<sub>D</sub><sup>20</sup> = -16.3 (*c* 1, CHCl<sub>3</sub>).

<sup>1</sup>H NMR (400 MHz; CDCl<sub>3</sub>):  $\delta$  7.97-7.94 (m, 2H), 7.57-7.52 (m, 1H), 7.48-7.43 (m, 2H), 2.99 (dd, *J* = 16.3 Hz, 7.5 Hz, 1H), 2.83 (dd, *J* = 16.3 Hz, 5.6 Hz, 1H), 2.13-2.05 (m, 1H), 1.40-1.33 (m, 2H), 1.29 (dd, *J* = 13.6 Hz, 5.2 Hz, 1H), 1.19 (dd, *J* = 14.2 Hz, 5.3 Hz, 1H), 0.91-0.88 (m, 12H).

<sup>13</sup>C NMR (101 MHz; CDCl<sub>3</sub>):  $\delta$  200.6, 137.7, 132.9, 128.7, 128.2, 47.8, 45.4, 32.3, 31.2, 30.1, 29.3, 11.3.

HRMS (ESI) : *m/z* calcd for C<sub>16</sub>H<sub>24</sub>ONa: 255.17249; [M+Na]<sup>+</sup> found : 255.1726 (0 ppm).

#### 3,5,5-trimethyl-1-phenylhexan-1-one (**2g**)

Using the typical procedure with (*R*)-BINAP, **2g** was obtained in 44% yield (38 mg) and 93% ee (GC: G-TA column, helium (40 cm/sec), 100 °C -150 min. - 5 °C/min. - 160 °C - 10 min., R<sub>T1(minor)</sub> = 135.3 min., R<sub>T2(major)</sub> = 140.9 min.). [ $\alpha$ ]<sub>D</sub><sup>20</sup> = -13.2 (*c* 1, CHCl<sub>3</sub>).

<sup>1</sup>H NMR (400 MHz; CDCl<sub>3</sub>):  $\delta$  7.96-7.93 (m, 2H), 7.57-7.53 (m, 1H), 7.48-7.44 (m, 2H), 2.92 (dd, *J* = 15.9, 5.7 Hz, 1H), 2.82 (dd, *J* = 15.9, 8.0 Hz, 1H), 2.32-2.21 (m, 1H), 1.32 (dd, *J* = 14.0, 4.0 Hz, 1H), 1.18 (dd, *J* = 14.0, 6.5 Hz, 1H), 1.00 (d, *J* = 6.6 Hz, 3H), 0.91 (s, 9H).

<sup>13</sup>C NMR (101 MHz; CDCl<sub>3</sub>):  $\delta$  200.4, 137.6, 132.9, 128.7, 128.2, 51.2, 48.3, 31.3, 30.2, 26.5, 23.1.

HRMS (ESI) : *m/z* calcd for C<sub>15</sub>H<sub>22</sub>ONa: 241.15684; [M+Na]<sup>+</sup> found : 241.1567 (1 ppm).

#### 3-ethyl-4,4-dimethyl-1-phenylpentan-1-one (**2i**)

Using the typical procedure with (*R*)-BINAP, **2i** was obtained in 22% yield (determined by <sup>1</sup>H NMR using mesitylene as an internal standard) and 0% ee (HPLC: Chiralcel OJ column, *n*-Hexane, 0.2 mL/min., R<sub>T1</sub> = 28.9 min., R<sub>T2</sub> = 38.1 min.). Data were in accordance with the literature.<sup>12</sup>

### 1,4-asymmetric conjugate addition on $\alpha$ -aryl dienone (**1g**): Typical Procedure

A flame dried Schlenk flask, under an argon atmosphere, was charged with diphosphine ligand (0.025 mmol, 5 mol%), CuTc (4.8 mg, 0.025 mmol, 5 mol%) and dry THF (0.5 mL). The resulting mixture was stirred for 10 min at room temperature and to this solution was added dropwise a 1M solution of diethylzinc in hexanes (1.5 mL, 1.5 mmol, 3 equiv.). The reaction mixture was stirred for 10 min at room temperature and a solution of substrate **1g** (117 mg, 0.5 mmol, 1 equiv.) in 0.5 mL of dry THF was added. The reaction mixture was stirred at 0 °C until the completion. The reaction was quenched by the addition of ethanol (4 mL). The mixture was stirred for 15 min and filtered through a short column of silica gel with EtOAc. The solvents were removed under reduced pressure. The crude mixtures were purified by flash chromatography on silica gel (pentane/Et<sub>2</sub>O: 98/2) to afford the corresponding mixture of adducts **2j/2j'** as colorless oils. Data were in accordance with literature.<sup>7</sup> Ee was obtained by chiral HPLC analysis of the purified product (Chiralcel OJ-H column, *n*-Hexane/*i*-PrOH: 99/1, 1 mL/min, R<sub>T1(1,4 adduct, minor)</sub> = 18.1 min, R<sub>T2(1,4 adduct, major)</sub> = 20.0 min.).

With (*R*)-BINAP: After 3h of reaction, a 96/4 mixture of adducts **2j**/**2j'** was isolated in 82% yield (109 mg) and 92% ee.  
With (*S*)-SYNPHOS: After 5h of reaction, a 97/3 mixture of adducts **2j**/**2j'** was isolated in 70% yield (92 mg) and -54% ee.  
With (*R*)-DIFLUORPHOS: After 6h of reaction, a 95/5 mixture of adducts **2j**/**2j'** was isolated in 66% yield (87 mg) and 80% ee.

### 1,6-asymmetric conjugate addition on dienone (**1h**): Typical Procedure

A flame dried Schlenk flask, under an argon atmosphere, was charged with diphosphine ligand (0.025 mmol, 5 mol%), CuTc (4.8 mg, 0.025 mmol, 5 mol%) and dry THF (1 mL). The resulting mixture was stirred for 10 min. A 1M solution of diethylzinc in hexanes (1.5 mL, 1.5 mmol, 3 equiv.) was added and the reaction mixture was stirred for 10 min. Finally, a solution of substrate **1h** (90 mg, 0.5 mmol, 1 equiv.) in 0.5 mL of THF was added. The reaction mixture was stirred at room temperature until the completion. The reaction was quenched by the addition of solid NH<sub>4</sub>Cl (500 mg). The solution was stirred for 1 h and then filtered on a small pad of silica, washed with EtOAc, and concentrated under vacuo. The residue was dissolved in 4 mL of dry CH<sub>2</sub>Cl<sub>2</sub> and DBU (100 mL, 0.7 mmol, 1.4 equiv) was added to the solution. The reaction was stirred for 5 h and then filtered on a small pad of silica, washed with EtOAc, and concentrated under vacuo. The crude product was purified by flash chromatography on silica gel (pentane/Et<sub>2</sub>O: 85/15) to afford **2k**. Data were in accordance with literature.<sup>13b</sup> Ee was obtained by GC analysis of the purified product (Beta-dex column, helium (40 cm/sec), 100 °C-400 min-5 °C/min-170 °C-10 min, Rt<sub>(minor)</sub> = 380.5 min, Rt<sub>(major)</sub> = 385.4 min.)

With (*R*)-BINAP: After 5 h of reaction, **2k** was isolated in 60% yield (63 mg) and 82% ee.

With (*S*)-SYNPHOS: After 14 h of reaction, **2k** was isolated in 54% yield (56 mg) and -82% ee.

With (*R*)-DIFLUORPHOS: After 14 h of reaction, **2k** was isolated in 14% yield (determined by <sup>1</sup>H NMR using mesitylene as an internal standard) and 74% ee.

### Acknowledgement

This work was supported by the Agence Nationale de la Recherche (SCATE, n° ANR-12-B507-0009-01; grant to MSTM). MM and OB also thank the CNRS.

### Supporting Information

Supporting information for this article is available online at <http://dx.doi.org/>.

### References

- (1) For metal-catalysed asymmetric nucleophile addition to electron-deficient alkenes, see: (a) Mauduit, M.; Baslé, O.; Clavier, H.; Crévisy, C.; Denicourt-Nowicki, A. In *Comprehensive Organic Synthesis II, vol. 4* (Knochel, P., Molander G. A.) Eds, Elsevier, **2014**, pp 186; For an excellent comprehensive review, see also: (b) Hawner, C.; Alexakis, A. *Chem. Commun.* **2010**, 46, 7295.
- (2) For Cu-catalysed ACA, see: (a) *Copper-Catalyzed Asymmetric Synthesis*; Alexakis, A.; Krause, N.; Woodward, S., Eds.; Wiley-VCH: Weinheim, Germany **2014**. (b) Jerphagnon, T.; Pizzuti, M. G.; Minnaard, A. J.; Feringa, B. L. *Chem. Soc. Rev.* **2009**, 38, 1039. (c) Alexakis, A.; Bäckvall, J.; Krause, N.; Pàmies, O.; Diéguez, M. *Chem. Rev.* **2008**, 108, 2796.
- (3) den Hartog, T.; Rudolph, A.; Macia, B.; Minnaard, A. J.; Feringa, B. L., *J. Am. Chem. Soc.*, **2010**, 132, 14349.
- (4) (a) Wang, S.-Y.; Ji, S.-J.; Loh, T.-P. *J. Am. Chem. Soc.* **2007**, 129, 276-277. (b) Wang, S.-Y.; Lum, T.-K.; Ji, S.-J.; Loh, T.-P. *Adv. Synth. Catal.* **2008**, 350, 673. (c) Wang, S.-Y.; Song, P.; Loh, T.-P. *Adv. Synth. Catal.* **2008**, 350, 3185. (d) Wang, S.-Y.; Loh, T.-P. *Chem. Commun.* **2010**, 46, 8694; see also ref.3
- (5) (a) Maciá Ruiz, B.; Geurts, K.; Fernández-Ibáñez, M. A.; ter Horst, B.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2007**, 9, 5123. (b) Lee, J. C. H.; Hall, D. G. *J. Am. Chem. Soc.* **2010**, 132, 5544.
- (6) Gremaud, L.; Alexakis, A. *Angew. Chem. Int. Ed.* **2012**, 51, 794.
- (7) Magrez-Chiquet, M.; Morin, M. S. T.; Wencel-Delord, J.; Drissy Amraoui, S.; Baslé, O.; Alexakis, A.; Crévisy, C.; Mauduit, M. *Chem. Eur. J.* **2013**, 19, 13663.
- (8) (a) Duprat de Paule, S.; Champion, N.; Ratovelomanana-Vidal, V.; Genet, J.-P.; Dellis, P. FR2830254, EP1436304, **2001**, WO Patent 03029259, **2003**. (b) Duprat de Paule, S.; Jeulin, S.; Ratovelomanana-Vidal, V.; Genêt, J.-P.; Champion, N.; Dellis, P. *Eur. J. Org. Chem.* **2003**, 1931. (c) Duprat de Paule, S.; Jeulin, S.; Ratovelomanana-Vidal, V.; Genet, J. P.; Champion, N.; Deschaux, G.; Dellis, P. *Org. Process Res. Dev.* **2003**, 7, 399.
- (9) (a) Jeulin, S.; Duprat de Paule, S.; Ratovelomanana-Vidal, V.; Genêt, J.-P.; Champion, N.; Dellis, P. *Angew. Chem. Int. Ed.* **2004**, 43, 320. (b) Jeulin, S.; Duprat de Paule, S.; Ratovelomanana-Vidal, V.; Genet, J.-P.; Champion, N.; Dellis, P. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, 101, 5799. (c) Genêt, J.-P.; Ayad, T.; Ratovelomanana-Vidal, V. *Chem. Rev.* **2014**, 114, 2824.
- (10) To the best of our knowledge, only two reports disclose the 1,4-adduct **2a** with a maximum of 77% yield and 96% ee, with Cu-ACA processes involving phosphino-based BINOL or SPINOL ligands: (a) Endo, K.; Ogawa, M.; Shibata, T. *Angew. Chem. Int. Ed.* **2010**, 49, 2410. (b) Endo, K.; Hamada, D.; Yakeishi, S.; Ogawa, M.; Shibata, T. *Org. Lett.* **2012**, 14, 2342.
- (11) To the best of our knowledge, only one report mentioned this challenging ACA with lower yield of 5% but higher ee up to 74%, see: Takahashi, Y.; Yamamoto, Y.; Katagari, K.; Danjo, H.; Yamaguchi, K.; Imamoto, T. *J. Org. Chem.* **2005**, 70, 9009.
- (12) For a review dealing with ACA on extended Michael acceptors, see: Csaky, A. G.; de la Herran, G.; Murcia, M. C. *Chem. Soc. Rev.* **2010**, 39, 4080.

- (13) For examples of Cu-catalyzed ACA on conjugated dienones, see: (a) Fillion, E.; Wilsily, A.; Liao, E.-T. *Tetrahedron: Asymmetry* **2006**, *17*, 2957. (b) Henon, H.; Mauduit, M.; Alexakis, A. *Angew. Chem. Int. Ed.* **2008**, *47*, 9122. (c) Wencel-Delord, J.; Alexakis, A.; Crévisy, C.; Mauduit, M. *Org. Lett.* **2010**, *12*, 4335. (d) Tissot, M.; Poggiali, D.; Henon, H.; Müller, D.; Guénee, L.; Mauduit, M.; Alexakis, A. *Chem Eur. J.* **2012**, *18*, 8731. (e) Magrez, M. Wencel-Delord, J.; Alexakis, A.; Crévisy, C.; Mauduit, M. *Org. Lett.* **2012**, *14*, 3576. (f) Ma, Z.; Xie, F.; Yu, Han.; Zhang, Y.; Wu, X.; Zhang, W., *Chem. Commun.* **2013**, *49*, 5292.
- (14) Coates, R. M.; Sowerby, R. L. *J. Am. Chem. Soc.*, **1972**, *94*, 5386.
- (15) For instance, it was observed that the Cu/phosphoramidite catalysed addition of Et<sub>2</sub>Zn or Et<sub>3</sub>Al to a cyclic dienone structurally close to **1h** gave the 1,6-adduct, whereas the Cu/NHC catalysed addition of a Grignard reagent led to the 1,4-adduct. See reference 13b.
- (16) For a specific review on BINAP, see: (a) Noyori, R.; Takaya, H. *Acc. Chem. Res.* **1990**, *23*, 345. For industrial enantioselective metal-catalytic processes using BINAP, see: (b) Akutagawa, S. *Appl. Catal. A: General* **1995**, *128*, 171.
- (17) Liu, D.-N.; Tian, S.-K. *Chem. Eur. J.* **2009**, *15*, 4538
- (18) Pinto, Diana C. G. A.; Silva, Artur M. S.; Lévai, A.; Cavaleiro, José A. S.; Patonay, T.; Elguero, J. *Eur. J. Org. Chem.* **2000**, *2000*, 2593
- (19) Rana, N. K.; Selvakumar, S.; Singh, V. K. *J. Org. Chem.* **2010**, *75*, 2089
- (20) Shintani, R.; Fu, G. C. *Org. Lett.* **2002**, *4*, 3699

#### Graphical Abstract

