



**HAL**  
open science

# Influence of discharge, hydraulics, water temperature, and dispersal on density synchrony in brown trout populations (*Salmo trutta*)

Victor Bret, Benjamin Bergerot, Hervé Capra, Véronique Gouraud, Nicolas Lamouroux

## ► To cite this version:

Victor Bret, Benjamin Bergerot, Hervé Capra, Véronique Gouraud, Nicolas Lamouroux. Influence of discharge, hydraulics, water temperature, and dispersal on density synchrony in brown trout populations (*Salmo trutta*). *Canadian Journal of Fisheries and Aquatic Sciences*, 2016, 73 (3), pp.319–329. 10.1139/cjfas-2015-0209 . hal-01324705

**HAL Id: hal-01324705**

**<https://univ-rennes.hal.science/hal-01324705>**

Submitted on 7 Jul 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Influence of discharge, hydraulics, water temperature and dispersal on  
2 density synchrony in brown trout populations (*Salmo trutta*)

3 Bret Victor<sup>1</sup>, Bergerot Benjamin<sup>2</sup>, Capra Hervé<sup>3</sup>, Gouraud Véronique<sup>1</sup>, Lamouroux Nicolas<sup>3</sup>

4

5 <sup>1</sup> EDF R&D, LNHE Department, HYNES (Irstea – EDF R&D), 6 Quai Watier, Chatou Cedex  
6 78401, France

7

8 <sup>2</sup> Hepia Geneva, University of Applied Sciences Western Switzerland, Technology,  
9 Architecture and Landscape, Centre de Lullier, route de Presinge 150, CH-1254 Jussy,  
10 Switzerland

11

12 <sup>3</sup> IRSTEA Lyon, UR MALY, HYNES (Irstea – EDF R&D), centre de Lyon-Villeurbanne, F-  
13 69626 Villeurbanne, France

14

15

16 E-mails :

17 victor.bret@edf.fr, benjamin.bergerot@hesge.ch, herve.capra@irstea.fr,  
18 veronique.gouraud@edf.fr, nicolas.lamouroux@irstea.fr

19

20 Correspondence: Victor Bret, Tel +33130878459, victor.bret@edf.fr

21

## 22 **Abstract**

23 Environmental factors may cause synchronous density variations between populations. A  
24 better understanding of the processes underlying synchrony is fundamental to predicting  
25 resilience loss in metapopulations subject to environmental change. The present study  
26 investigated the determinants of synchrony in density time series of three age-groups of  
27 resident brown trout (0+, 1+ and adults) in 36 stream reaches. A series of Mantel tests were  
28 implemented to disentangle the relative effects on trout synchrony of geographical proximity,  
29 environmental synchrony in key environmental variables affecting trout dynamics (discharge,  
30 water temperature, hydraulics and spawning substrate mobility) and density-dependent  
31 dispersal. Results indicated that environmental synchrony strongly explained trout synchrony  
32 over distances less than 75km. This effect was partly due to a negative influence on 0+ trout  
33 of strong discharges during the emergence period and a more complex influence of substrate  
34 mobility during the spawning period. Dispersal between reaches had a weak influence on  
35 results. Juvenile and adult densities were strongly driven by survival processes and were not  
36 influenced by environmental synchrony. The results suggest that the environment can have  
37 general effects on population dynamics that may influence the resilience of metapopulations.

## 38 **Keywords:**

39 Moran effect; Freshwater fish; Population dynamics; Density-dependent dispersal; Mantel  
40 tests; Bypassed section

## 41 **Résumé**

42 Les facteurs environnementaux peuvent causer des fluctuations synchrones de densités entre  
43 populations. Une meilleure compréhension des processus expliquant la synchronie est  
44 fondamentale pour prédire des pertes de résilience des métapopulations sujettes à des  
45 changements environnementaux. Nous étudions la synchronie des chroniques de densités de

46 trois classes d'âge de la truite commune (0+, 1+ et adultes) entre 36 tronçons de cours d'eau.  
47 Nous utilisons des tests de Mantel pour discriminer les effets relatifs de la proximité  
48 géographique, de la synchronie de variables environnementales clefs (débit, température de  
49 l'eau, conditions hydrauliques et mobilité du substrat) et de la dispersion densité-dépendante.  
50 La synchronie environnementale expliquait fortement la synchronie de la truite jusqu'à des  
51 distances de 75km. Cet effet était dû en partie à l'influence négative sur les 0+ des hauts  
52 débits pendant l'émergence et une influence de la mobilité du substrat pendant la période de  
53 ponte. La dispersion entre tronçon influençait faiblement nos résultats. Les densités de  
54 juvéniles et d'adultes étaient fortement structurées par des processus de survie, mais n'étaient  
55 pas influencées par la synchronie des conditions environnementales. Les résultats suggèrent  
56 que l'environnement peut avoir des effets généraux sur la dynamique de population qui  
57 peuvent influencer la résilience des métapopulations.

58 **Mots-clefs:**

59 Effet Moran; Poissons d'eau douce ; Dynamique de populations; Dispersion densité-  
60 dépendante; Test de Mantel ; Tronçon court-circuité

61

62 **Introduction**

63 Temporal variations in population density depend on both abiotic (e.g., environmental) and  
64 biotic processes (e.g., density-dependent dispersal) operating at various spatial scales (Alonso  
65 et al. 2011; Richard et al. 2013). Environmental factors may cause synchronous fluctuations  
66 in density in populations with similar density-dependence structures (Moran 1953). These  
67 "Moran effects" may explain the synchronous variation in abundance of herbivorous insects  
68 and mammals in sites located as much as 1,000 km apart (Koenig 2002; Liebhold et al.  
69 2004). Density-dependent regulation (through dispersal of individuals) can also induce

70 synchrony between connected populations, but these effects are generally weaker (Ripa 2000)  
71 and occur over shorter distances (Ranta et al. 1998). In many previous synchrony analyses,  
72 the influence of environmental variables (e.g., Holyoak and Lawler 1996) or density-  
73 dependent regulation (e.g., Tedesco et al. 2004) could be neglected. However, their relative  
74 influence on population synchrony needs to be analyzed to identify the general drivers of  
75 population dynamics and to better quantify the resilience of metapopulations subject to  
76 environmental change. When the main drivers of synchrony are environmental factors,  
77 metapopulations can be threatened by synchronous environmental disturbances. In contrast,  
78 populations in which synchrony is driven by density-dependent dispersal may be highly  
79 resilient to environmental disturbance. Analyzing data sets combining close and distant sites  
80 can contribute to better understanding the relative influence of the environment and of  
81 density-dependent regulation on population synchrony.

82 Spatial synchrony has been frequently studied and observed in riverine fish populations.  
83 Chevalier et al. (2014) studied 27 freshwater fish species commonly found throughout  
84 France. They found low but significant levels of synchrony (average correlation between  
85 pairs of reaches generally  $<0.1$ ) that were related to the life-history strategies and the upper  
86 thermal tolerance limits of species. Of these species, stream-resident brown trout (*Salmo*  
87 *trutta*) has been especially well documented, making it useful for the study of synchrony.  
88 This species has a wide native range and is known to be sensitive to environmental conditions  
89 (e.g., high flow rates, extreme water temperatures). Synchrony in such environmental  
90 conditions may lead to synchrony in trout density. Cattaneo et al. (2003) showed the  
91 influence of hydrologic synchrony (high discharge levels during the emergence period) on the  
92 density synchrony of young-of-the-year trout in 37 stream reaches. One limitation of existing  
93 analyses of trout synchrony (e.g., Cattaneo et al. 2003; Zorn and Nuhfer 2007) is that they did  
94 not involve key habitat factors for trout population dynamics (e.g., hydraulics and water

95 temperature; Armstrong et al. 2003). They generally used proxies (e.g., air temperature,  
96 discharge rate, geographic proximity) that are spatially correlated, which makes it difficult to  
97 distinguish their relative influences. Studying pairs of geographically close reaches with  
98 contrasting environmental characteristics (e.g., contrasting levels of discharge regulation)  
99 would be particularly useful to better disentangle the relative influence of environmental  
100 factors and density-dependent dispersal on synchrony. Another difficulty in interpreting  
101 synchrony in fish populations is that synchrony in a given age-group may be inherited from  
102 the previous age-groups (Grenouillet et al. 2001; Lobón-Cerviá 2009). Ideally, age-group  
103 successions should be taken into account in analyzing synchrony.

104 The present study provides an analysis of spatial synchrony in density time series of  
105 three age-groups of brown trout. The originality of the analysis lies in addressing certain  
106 important limitations of previous synchrony studies. In particular, the present analyses  
107 involved geographically close sites with differing environmental characteristics (disconnected  
108 by dams or not, bypassed by hydroelectric plants or not), to refine analysis of the relative  
109 influence of environmental factors and density-dependent dispersal on synchrony. In  
110 addition, key quantitative environmental variables influencing population dynamics  
111 (hydraulics, water temperature) were taken into account, as were relationships between  
112 successive age-groups.

### 113 **Materials and methods**

114 In brief, the dataset covered 36 stream reaches in which trout abundance was assessed  
115 annually (320 surveys: i.e., reach×year combinations) and discharge, hydraulics and water  
116 temperature were described. These data were transformed into distance matrices describing  
117 geographic distance, trout synchrony and environmental synchrony between pairs of reaches.  
118 Series of Mantel tests were then implemented to characterize the spatial scales of trout

119 synchrony and environmental synchrony (step 1), and to focus on the influence on trout  
120 synchrony of environmental synchrony on the one hand (step 2) and density-dependent  
121 dispersal on the other (step 3).

## 122 **Study reaches and geographic distances**

123 The 36 reaches belonged to 22 French rivers distributed across continental France (Fig. 1)  
124 and had a wide range of environmental characteristics (e.g., width ranging from 2.9 to 15.5m,  
125 median streambed particle diameter from 0.1 to 64cm; Table 1). They were selected based on  
126 the availability of hydraulic and water temperature data. In each reach, fish were sampled for  
127 at least four pairs of consecutive years (consecutive years being needed in order to take  
128 account of age-group successions). We checked that brown trout was the dominant species  
129 (relative density > 80% on at least one survey) in the eight reaches where other species were  
130 sampled. Each reach included one or several sequences of pools, runs and/or riffles. Due to  
131 changes in sampling teams (consulting firms) or harsh hydraulic conditions during some  
132 surveys, sampled length was slightly modified in half of the reaches during the study period,  
133 affecting 18% of surveys (maximum length change: 25%; median change: 13%). One reach  
134 had its length divided by two at the middle of the time series, but was kept as a single reach  
135 for analysis as its hydraulic characteristics remained unchanged. Groups of close reaches  
136 selected in the same river (Fig. 1) might or might not be disconnected by dams or bypassed  
137 by hydroelectric plants. A total of 18 reaches were bypassed, and therefore showed decreased  
138 low-flow and flood frequency. Consequently, the data set included pairs of very close reaches  
139 characterized by different hydraulic conditions. No chemical pollution was reported in these  
140 reaches.

141 Two kinds of geographic distance were computed between pairs of reaches: Euclidean and  
142 river network distances. Network distance is potentially more relevant to describe density-  
143 dependent dispersal, and was computed for the 93 out of 630 pairs of reaches that were not

144 separated by the sea, using a theoretical hydrographic network developed for France (Pella et  
145 al. 2012). All distances were log-transformed to approximate normality.

#### 146 **Trout data and synchrony in trout time series**

147 Between 4 and 19 surveys (mean: 8.9) were conducted per reach between 1991 and 2012.  
148 Reaches were sampled by wading, using two-pass removal electrofishing sampling meeting  
149 European Committee for Standardization guidelines (CEN 2003). Fish densities were  
150 estimated on the Carle and Strub (1978) method. Sampling was performed without blocking  
151 nets, in summer or early autumn (median date: September 13). Sampled area (between 175  
152 and 2,295 m<sup>2</sup>) was computed as sampled length × reach width at median flow. All fish were  
153 measured (to the nearest 1mm) and length-frequency histograms were used to distinguish  
154 three age-groups: 0+ (young-of-the-year), 1+ (older than one year, generally juveniles) and  
155 adult (all fish older than two years). Scales were available for 10 reaches only, but confirmed  
156 the suitability of using length-frequency distributions (see Sabaton et al. 2008). Adults were  
157 considered as the potential reproductive pool. Age-group densities (number of individuals per  
158 100m<sup>2</sup>) were log-transformed to normalize their distributions.

159 Due to the strength of the relationship between densities of successive age-groups for brown  
160 trout (Zorn and Nuhfer 2007; Lobón-Cerviá 2009), a global model of age-group succession  
161 (averaged across reaches) was determined. All synchrony analyses were performed on  
162 residuals of this global model, in order to reduce serial correlation and better identify the  
163 causes of synchrony (e.g., Buonaccorsi et al. 2001; Santin-Janin et al. 2014). Specifically,  
164 linear regressions were fitted for all age-groups (0+, 1+, Ad) relating log-transformed density  
165 at year  $y$  ( $D_{0+,y}$ ;  $D_{1+,y}$ ;  $D_{Ad,y}$ ) to the density of previous age-group at year  $y-1$   
166 (respectively:  $D_{Ad,y-1}$ ;  $D_{0+,y-1}$ ;  $D_{1+,y-1}$  and  $D_{Ad,y-1}$ ). Adults at year  $y$  depended on both  
167 adults and 1+ fish at year  $y-1$ , as the adult group combined fish of several age-groups. Slopes



168 significantly lower than 1 were taken to indicate global density-dependence in population  
169 dynamics (for  $D_{0+,y}$  and  $D_{1+,y}$ ).

170 To quantify the potential limits of this approach, mixed-effect linear models with a reach-  
171 level random effect (i.e., with regression coefficients that could vary across reaches) were  
172 also fitted to the data and compared with the global model to appreciate the generality of our  
173 global models across reaches. Importantly, residuals of the mixed models were not analyzed,  
174 even when they fitted better than the global models, because they could not be interpreted  
175 together, not being calculated from the same regression model in all reaches. In addition, data  
176 for a given individual reach were often insufficient to provide a robust model of age-group  
177 succession. In other words, analyzing the residuals of the global models was a means of  
178 removing average serial correlation while calculating density descriptors similarly in all  
179 reaches.

180 All further synchrony analyses were made on residual densities of the global succession  
181 models, hereafter noted  $r_{0+}$ ,  $r_{1+}$  and  $r_{Ad}$ . For each of the three age-groups, synchrony  
182 between pairs of reaches was described by  $36 \times 36$  distance matrices. The elements of these  
183 distance matrices were dissimilarity measures, calculated as  $1 - \rho$ , where  $\rho$  was the Pearson  
184 correlation between age-group residual density for the corresponding pair of reaches (varying  
185 between 0 and 2). Synchrony values were transformed into distance values using  $1 - \rho$ , as  
186 distance values are conventionally used with Mantel tests ( $1 - \rho$  decreases with synchrony).  
187 Pairs of reaches with less than three years of simultaneous density and environmental  
188 information (33% of cases) were excluded from all analyses. Finally, the average number of  
189 years per reach used to calculate  $\rho$  values was 5.8.

190 **Environmental data and synchrony in environmental time series**

191 According to the literature, trout population dynamics may be influenced by discharge,  
192 hydraulic and thermal conditions during key periods of the trout life cycle. Therefore, for  
193 each year preceding fish sampling, environmental conditions were described for four key  
194 periods, using five environmental variables (Table 2). Only the 10 environmental descriptors  
195 (period×variable combinations) for which a causal relationship with some age-group  
196 densities was expected were considered (Table 2).

197 These four key periods were: (i) adult spawning migration (September 1<sup>st</sup> to January 31<sup>st</sup>); (ii)  
198 egg development (November 1<sup>st</sup> to February 29<sup>th</sup>); (iii) fry emergence (March 1<sup>st</sup> to April  
199 30<sup>th</sup>); and (iv) the summer growth period (July 1<sup>st</sup> to September 13<sup>th</sup>, the median date of the  
200 320 trout surveys). The dates for the first three periods (hereafter: ‘spawning’, ‘egg’,  
201 ‘emergence’) were estimated for France by a group of fourteen experts from several  
202 organizations on the basis of numerous trout monitoring campaigns (Gouraud et al. 2014).  
203 The ‘summer’ period was defined to describe low-flow conditions preceding sampling.

204 Two of the five environmental variables (Table 2) were daily discharge percentiles, two were  
205 hydraulic variables (flow velocity and substrate mobility), and the fifth was the frequency of  
206 low temperature. The field data and models used to calculate these environmental variables  
207 are detailed in Appendix A. In brief, field data involved daily discharge and daily water  
208 temperature measured in most reaches. Missing values were estimated using extrapolation  
209 models. For water temperature, extrapolation model tests generally indicated errors of the  
210 order of 1°C. Hydraulic conditions were derived from numerical hydraulic models or detailed  
211 hydraulic measurements (N>100) made throughout each reach.

212 The two daily discharge percentiles described low- and high-flow magnitude. Low-flow  
213 magnitude ( $Q_{90}$ , defined as daily discharge exceeded 90% of the time during the period,  $m^3 \cdot s^{-1}$ )

214 <sup>1</sup>) was used to test the effect of summer low-flows on all age-groups (see Nislow and  
215 Armstrong 2012). High-flow magnitude ( $Q_{10}$ , defined as daily discharge exceeded 10% of the  
216 time during the period,  $m^3 \cdot s^{-1}$ ) was used to test the effect of spates on  $rAd$  during spawning  
217 (spawners may be more sensitive to spates during their migration), on  $r0+$  during egg  
218 development (e.g., Unfer et al. 2011) or on residual density in all age-groups during  
219 emergence. All age-groups were considered as potentially influenced by high spring floods  
220 occurring during this period because spates have a major impact on 0+ fish (Jensen and  
221 Johnsen 1999; Cattaneo et al. 2003; Unfer et al. 2011) and may be strong enough to influence  
222 the survival and dispersal of older cohorts (Young et al. 2010).

223 The other three variables, describing hydraulics and thermal conditions, were not percentiles  
224 but indicated the frequency above or below quantitative thresholds of events that could  
225 influence trout life cycle. Frequency of high daily velocities ( $fV0.5$ ) was defined by a  
226 threshold of  $0.5 m \cdot s^{-1}$ , corresponding to the upper end of the preferred range of current  
227 velocity for 0+ (Heggenes 1996; Roussel and Bardonnnet 2002). The influence of  $fV0.5$  on  
228  $r0+$  was tested during emergence, due to the reduced swimming ability of recently hatched  
229 juveniles (Armstrong et al. 2003). The influence of frequency of spawning substrate mobility  
230 ( $fMob$ , frequency of daily discharge > critical discharge; see Appendix A) during spawning,  
231 egg development and emergence was tested on  $r0+$ , due to a potential direct influence of bed  
232 mobility on mortality in early life stages (Unfer et al. 2011). Regarding the thermal threshold,  
233 sub-lethal temperatures for brown trout (<0°C for all age-stages, >13°C during egg  
234 development, and >22°C for older stages; Elliott and Elliott 2010) were exceptional in the  
235 study reaches (only 0.7% of daily water temperatures were < 1°C and none were >19.6°C);  
236 cold periods were therefore defined by temperature thresholds that occurred more frequently,  
237 corresponding to the first third of the thermal preference range reported by Elliott and Elliott  
238 (2010): <4.3°C for egg development and <7.3°C for older age-stages. The influence of sub-

239 optimal temperature ( $fTlow$ , the frequency of days with  $T_{water,d}$  below threshold) on  $r0+$  was  
240 tested during egg development ( $T_{water,d} < 4.3^{\circ}C$ ) and emergence ( $T_{water,d} < 7.3^{\circ}C$ ). Influence  
241 on older age-groups was not tested as these fish can actively seek thermal refuge (Cunjak et  
242 al. 2013).

243 Regarding biological synchrony, environmental synchrony between pairs of reaches was  
244 assessed for each environmental variable on  $36 \times 36$  distance matrices with dissimilarity  
245 measures equal to  $1 - \rho$ , where  $\rho$  was the Pearson correlation of the environmental variable  
246 between the corresponding pair of reaches.

#### 247 **Data analyses**

248 Several Mantel tests (Mantel 1967) were used to analyze the relative influence of  
249 environmental synchrony and density-dependent dispersal on trout synchrony. All these  
250 Mantel tests analyzed the probability of the observed relationship (Mantel R) between  
251 dissimilarity values of two or three distance matrices (geographic distance, trout synchrony  
252 and environmental synchrony) occurring randomly (significance threshold: 0.05; 4,000  
253 random permutations; “vegan” R package; Oksanen et al. 2013).

#### 254 *Step 1: Spatial scales of synchrony*

255 The spatial scales of synchrony were first analyzed by correlating trout synchrony and  
256 environmental synchrony to Euclidean distance. A strong relationship between trout  
257 synchrony and geographic distance could be due to the combined influence of density-  
258 dependent dispersal and environmental synchrony. However, density-dependent dispersal  
259 could be expected to generate synchronous trout responses over smaller geographic distances  
260 than environmental synchrony. The tests relating trout synchrony to Euclidean distance were  
261 repeated using network distance: a stronger link between trout synchrony and network  
262 distance than Euclidean distance would suggest an effect of dispersal.

263 *Step 2: Influence of environmental synchrony on trout synchrony*

264 To better analyze the influence of environmental synchrony on trout synchrony, trout  
265 synchrony was first correlated to environmental synchrony (Mantel tests for all relationship  
266 hypotheses in Table 2) and results were compared to those between geographic distance and  
267 trout synchrony (at step 1).

268 When univariate Mantel tests were significant with both an environmental variable and  
269 geographic distance for a given age-group, partial (multivariate) Mantel tests were performed  
270 to better distinguish their relative effects (Smouse et al. 1986). To avoid having to perform  
271 numerous tests, partial Mantel tests were performed for only one environmental variable  
272 within each type of environmental group (discharge, hydraulics, temperature). The variable  
273 selected was the most significant one found on univariate Mantel testing.

274 Finally, to help interpret the synchronous influence of the environment on trout density, time  
275 series of trout residual density were co-plotted against environmental variables. These time  
276 series (trout and environment) were standardized by reach and averaged between groups of  
277 reaches with synchronous trout series. The groups were obtained by hierarchical cluster  
278 analysis based on the distance matrices of trout residual density (Ward algorithm; as in  
279 Cattaneo et al. 2003).

280 *Step 3: Influence of density-dependent dispersal on trout synchrony*

281 To better analyze the influence of density-dependent dispersal between reaches on observed  
282 synchrony, tests that were significant at steps 1 and 2 were repeated on two subsets of data  
283 characterized by reduced possibilities of dispersal. In the first subset, 30 reaches in which  
284 dispersal was limited were selected: reaches more than 50km apart or separated by a dam  
285 preventing upstream passage; the 50 km threshold was, to the best of our knowledge, greater  
286 than the maximum distance reported for brown trout displacement (Young et al. 2010).

287 Dispersal between these reaches was possible only during spates (e.g., drift through the dam).  
 288 The second data subset comprised only 22 reaches between which dispersal was impossible  
 289 (more than 50km apart or separated by impassable dams in both directions). As there were  
 290 various ways of removing reaches, tests on the two subsets were repeated 100 times with  
 291 differing random selection of reaches to be removed. Thus, for each significant result of steps  
 292 1 and 2, the percentage of cases (in the 100 repetitions) in which the Mantel test remained  
 293 valid on the data subsets was quantified.

## 294 **Results**

### 295 **Trout data and synchrony in trout times series**

296 Median trout density per reach across surveys varied between 10.3 and 51.3 individuals per  
 297 100m<sup>2</sup>. On average, 38% of sampled individuals were 0+, 34% were 1+ and 28% were adult.  
 298 Linear regressions indicated a significant relationship between the densities of successive  
 299 age-groups ( $P < 0.001$ ; Fig. 2, plain lines). Model coefficients [ $\pm$  standard deviation] were:

$$300 \quad (1) \quad \log(0+y) = 1.37 [\pm 0.17] + 0.34 [\pm 0.10]. \log(Ad_{y-1}) \quad (R^2 = 0.04)$$

$$301 \quad (2) \quad \log(1+y) = 0.80 [\pm 0.06] + 0.55 [\pm 0.03]. \log(0+y_{-1}) \quad (R^2 = 0.54)$$

$$302 \quad (3) \quad \log(Ad_y) = 0.27 [\pm 0.08] + 0.25 [\pm 0.04]. \log(1+y_{-1}) \\
 + 0.57 [\pm 0.03]. \log(Ad_{y-1}) \quad (R^2 = 0.54)$$

303 The global model  $R^2$  for 1+ and adults indicated that the global linear model appropriately  
 304 reflected an average age-group succession across reaches (>50% of variability explained).  
 305 For the 0+ age-group, the model showed low explained variance. The slopes of Eqs (1) and  
 306 (2) and the sum of slopes in Eq (3) were significantly less than 1, suggesting some degree of  
 307 apparent density-dependence regulation in age-group successions.

308 Mixed models (Fig. 2, dashed lines), significantly improved fit, with  $R^2$  values of 0.30, 0.70  
309 and 0.71 for the three age-groups.  $\Delta AIC$  between models with random effects ('mixed  
310 model') and fixed effect ('global model') were respectively 14, 23 and 9 for 0+, 1+ and  
311 adults.

312 Median Pearson  $\rho$  between all pairs of reaches was close to 0 ( $\rho_{r0+}=0.16$ ,  $\rho_{r1+}=-0.01$ ,  
313  $\rho_{rAd}=0.11$ ), indicating that there was no obvious global synchrony at the spatial scale of the  
314 whole dataset.

### 315 **Step 1: Spatial scales of synchrony**

316 Euclidean distance between pairs of reaches varied between 1.2 and 1,029.0 km, with 39  
317 pairs of reaches less than 15 km apart. For the 93 out of 630 pairs of reaches that were not  
318 separated by the sea, distance via the river network ranged from 1.2 to 640.0km, (mean:  
319 262.0 km), with 25 pairs of reaches less than 15 km apart.

320 Mantel tests showed that trout synchrony of  $r0+$  and  $r1+$  was significantly related to  
321 Euclidean distance (Fig. 3). Although significant, these tests revealed low Mantel R  
322 ( $R^2 < 0.08$ ). For  $r0+$ , however, the degree of synchrony was strong (half of correlations  $> 0.5$ )  
323 for reaches less than ~75 km apart, and even stronger (75% of correlations  $> 0.5$ ) for reaches  
324 less than 5 km apart. This effect was weaker for  $r1+$  (Fig. 3). Focusing on reaches for which  
325 network distance could be computed, Mantel R between  $r0+$  and geographic distance was  
326 lower for network distance than Euclidean distance ( $R^2 = 0.08$  vs. 0.16; Fig. 4). Correlations  
327 between  $r1+$  and geographic distance were no longer significant when analysis focused on  
328 these reaches.

329 Environmental synchrony was also significantly related to Euclidean distance for most  
330 descriptors (7 out of 10;  $R^2$  between 0.04 and 0.24), with particularly strong relationships for  
331 variables and periods related to high flow (see one example for each environmental group in

332 Fig. 3<sup>1</sup>). No spatial synchrony was found for only three environmental descriptors: *fTlow*  
333 during the egg period, *fMob* during the emergence period and  $Q_{90}$  during the summer. Other  
334 environmental synchronies were strong (generally  $>0.5$ ) up to distances around  $\sim 75$  km,  
335 except for *fTlow* during emergence (Fig. 3), which showed strong synchronies over longer  
336 distances ( $> 200$  km).

### 337 **Step 2: Influence of environmental synchrony on trout synchrony**

338 Mantel tests relating trout synchrony to the environment were significant in 4 out of 14 tests  
339 (Table 2; Fig. 5), all concerning  $r_{0+}$  residual density. Synchrony in  $r_{0+}$  was related to  $Q_{10}$   
340 (mainly during emergence and secondarily during the egg period) and *fMob* (mainly during  
341 spawning and secondarily during the egg period). Overall Mantel R was lower ( $R^2 \leq 0.05$ ) than  
342 for Euclidean distance. Mantel test results partly depended on a substantial number of pairs of  
343 reaches in which both physical and biological synchrony were strong (Fig. 5).

344 Partial Mantel tests (Table 3) were made for combinations of Euclidean distance and each of  
345 the two main environmental descriptors that influenced  $r_{0+}$ . The effect of Euclidean distance  
346 on  $r_{0+}$  synchrony was significant when the environmental effect was removed, whereas the  
347 reverse was not significant ( $P \geq 0.06$ ).

348 Co-plots of time series for trout residual density and the two main significant environmental  
349 descriptors in the four groups of synchronous reaches (Fig. 6) showed that  $Q_{10}$  during the  
350 emergence period was negatively associated with  $r_{0+}$  (at least in the three most synchronous  
351 groups), whereas there was no clear positive or negative direction of an annual effect of  
352 *fMob*. The four groups identified by cluster analysis did not necessarily involve  
353 geographically close reaches.

---

<sup>1</sup> Relations for all environmental descriptors are presented in the supplementary material (Fig. S1)



### 354 **Step 3: Influence of density-dependent dispersal on trout synchrony**

355 Four of the 6 significant Mantel tests relating trout density to Euclidean distance or the  
356 environment remained significant (in more than 75% of trials) when tested on the first data  
357 subset, in which dispersal was limited. These concerned the relationships with Euclidean  
358 distance and the two main environmental descriptors mentioned above ( $Q_{10}$  during  
359 emergence and  $fMob$  during spawning). By contrast, only one of the six tests remained  
360 significant (in 95% of trials) when tested on the second subset, in which dispersal was  
361 impossible. This test concerned the relationship between  $r0+$  and Euclidean distance.  
362 Dispersal between reaches may have influenced other results, which remained significant in  
363 <35% of tests on the second data subset.

364 The results of partial Mantel tests were similar whether performed on all reaches or on the  
365 limited dispersal subset, but were seldom significant when performed on the subset in which  
366 dispersal was impossible (Table 3).

## 367 **Discussion**

### 368 **A Moran effect on 0+ trout**

369 The present study supports the notion that salmonid populations are frequently synchronous  
370 (Copeland and Meyer 2011) and contributes to disentangling the relative influence of  
371 environmental factors and density-dependent dispersal on trout synchrony. The results  
372 principally suggest that a Moran effect is responsible for 0+ synchronies between  
373 geographically close reaches. Synchronies in older age-groups (1+ and adults) were weaker  
374 and not linked to environmental synchronies. Four elements supported the notion that 0+  
375 synchrony is due to a Moran effect. Firstly, synchrony in  $r0+$  was particularly strong over a  
376 distance of ~75km, a distance consistent with the spatial scale of environmental synchrony.  
377 This distance of ~75km is greater than the 50km reported for freshwater populations in the

378 meta-analysis by Myers et al. (1997). We were also considering a larger geographic scale in  
379 the present study, compared with reaches < 70km apart in Hayes 1995 or <25km apart in  
380 Lobón-Cervía 2004. Secondly, synchrony was related less to network distance than Euclidean  
381 distance, suggesting a weak influence of dispersal. Thirdly, several significant Mantel tests  
382 related 0+ synchrony to environmental synchrony (high flow during emergence and substrate  
383 mobility during spawning). And fourthly, many tests on  $r0+$  synchrony remained significant  
384 on the data subset where density-dependent dispersal between reaches was unlikely.

385 The Mantel tests showed low Mantel R values, but this statistic alone does not reflect the  
386 strength of synchrony. Mantel R is expected to be low in data sets collected over large spatial  
387 areas and with relatively short time series (see also Cattaneo et al. 2003; Chevalier et al.  
388 2014). Pairs of reaches more than 75km apart (Fig. 3) may not be synchronized due to a  
389 variety of environmental characteristics not considered here. This inevitably generates noise  
390 in the relation between geographic distance and trout synchrony. In the present study, plots  
391 relating 0+ synchrony to Euclidean distance or environmental synchrony indicated that 0+  
392 synchrony between geographically close reaches was frequently very strong (e.g., trout  
393 synchrony  $\rho$  was > 0.5 in 75% of pairs of reaches less than 5km apart; Fig. 3). More than the  
394 Mantel R value itself, these plots and the P-value of the Mantel tests indicated a strong,  
395 biologically significant level of synchrony between geographically close reaches.

#### 396 **Accounting for a global age-group succession model**

397 A first strength of the present approach was to consider age-groups individually rather than  
398 pooled. This can increase the observed degree of synchrony (Grenouillet et al. 2001), as  
399 suggested by the present median synchrony levels ( $\rho_{r0+}=0.16$ ,  $\rho_{r1+}=-0.01$ ,  $\rho_{rAd}=0.11$ ),  
400 which were generally higher than those obtained by Chevalier et al. (2014) after pooling age-  
401 groups ( $\rho=0.038$ ).

402 A second strength of the present approach was to reduce serial dependence between  
403 successive age-groups, by using global models of age-group succession. Serial dependence is  
404 one of the two main statistical issues in synchrony analysis (Liebhold et al. 2004), together  
405 with the influence of temporal trends (not found in the present dataset). Global succession  
406 models for 1+ and adults explained more than 50% of density variability, confirming the  
407 strength of serial dependence and the importance of taking it into account in analyzing  
408 synchrony with the environment. The global model relating 0+ to adults explained a smaller  
409 part of variability. Nevertheless, analyses were performed on  $r0+$  rather than directly on 0+  
410 density, in order to be consistent with the analyses for other age-groups. The weak serial  
411 dependence of 0+ on adults was consistent with other findings (e.g., Lobón-Cerviá 2013).  
412 Overwhelming environmental drivers may tend to make 0+ dependence on adults difficult to  
413 detect (Daufresne and Renault 2006). This relationship can also be affected by the lower  
414 catchability of small fish (e.g., Ruiz and Laplanche 2010) or by confusion between sampled  
415 adults and the actual parental stock. Actual spawning stock can depend on variability in  
416 maturity age (Olsen and Vøllestad 2005), on potential stocking issues of which the details are  
417 not well established, or on migration of adults between sampled reaches and spawning areas  
418 (Young et al. 2010).

419 The present global model of age-group succession for all reaches was the strongest  
420 assumption involved in accounting for serial dependence. This hypothesis was rarely tested  
421 explicitly in synchrony studies but is essential to investigating Moran effects. Liebhold et al.  
422 (2004) pointed out that the hypothesis probably does not hold in many systems, as spatial  
423 variation in population dynamics is frequent. In the present study, comparing the global  
424 models with reach-dependent mixed models suggested that a large part of population  
425 dynamics was taken into account by the global models. However, the mixed models indicated  
426 that variations in population dynamics did occur across reaches (higher explanatory power of

427 mixed models). This may partly lower the level of synchrony ( $\rho$  values and Mantel R)  
428 observed, because spatial variation in density-dependent dynamics reduces the synchrony  
429 caused by environmental stochasticity (Liebhold et al. 2006). Sampling error may also have  
430 lowered the observed levels of synchrony (Santin-Janin et al. 2014). However, sampling  
431 errors were unlikely to have influenced the main results, because the magnitude of sampling  
432 error was probably much lower than the magnitude of annual density variation, which can be  
433 as great as 10-fold between certain years.

434 The slopes of the global models for 1+ and adults were significantly less than 1, indicating an  
435 apparent global density-dependence survival for these age-groups, which was rarely  
436 previously documented (but see Richard et al. 2013). The present results also suggested a  
437 density-dependent regulation on 0+, but we remain cautious about this finding as the global  
438 0+ model showed very low explanatory power ( $R^2=4\%$ ). Density-dependence mechanisms on  
439 0+ have often been discussed in the literature, being difficult to identify (e.g., Elliott 1984;  
440 Nicola et al. 2008; Lobón-Cerviá 2013), mainly due to their high annual variability  
441 influenced by environmental conditions.

#### 442 **Environmental drivers of 0+ synchrony**

443 The correlation between  $r_{0+}$  and Euclidean distance was stronger than that between  $r_{0+}$  and  
444 environmental synchrony, although the dataset included geographically close reaches with  
445 differing characteristics due to dams. Thus, close reaches are likely to be synchronous, even  
446 if they are separated by dams and have different flow regimes. Moreover, partial tests  
447 revealed that environmental variables did not explain  $r_{0+}$  synchrony when the effect of  
448 Euclidean distance was removed. Therefore, Euclidean proximity probably accounted for a  
449 combined effect of several environmental variables including those studied here (e.g., high  
450 flow during emergence, or substrate mobility during spawning) and others not included in  
451 analysis.

452 Nevertheless,  $r_{0+}$  synchrony correlated significantly with  $Q_{10}$  during emergence and with  
453  $fMob$  during spawning. These results are consistent with the observation of a negative  
454 influence of high flow on small individuals during emergence (e.g. Hayes 1995; Cattaneo et  
455 al. 2003; Nicola et al. 2009), due to higher mortality and/or drift. Effects of spawning  
456 substrate mobility on  $0+$  were more rarely mentioned in the literature (but see Jensen and  
457 Johnsen 1999). Unfer et al. (2011) suggested that these effects could be positive (reshaping  
458 spawning grounds due to substrate turnover) or negative (scouring redds and destroying eggs)  
459 according to their timing. This could explain why the positive or negative direction of the  
460 annual effect of  $fMob$  was harder to identify.

461 The other environmental descriptors implemented in analysis, including the frequency of high  
462 current velocity, did not explain trout synchrony. However, using velocity percentiles instead  
463 of discharge ( $Q_{10}$ ) percentiles would have led to comparable results, due to the monotonic  
464 relationship between discharge and velocity in reaches. Therefore, an influence of hydraulics  
465 on trout synchrony cannot be ruled out. The non-significant effect of temperature may be due  
466 to local adaptation to the thermal regime (Filipe et al. 2013). It is also possible that the  
467 present dataset covered a larger range of hydraulic than thermal conditions (e.g., median flow  
468 velocity ranged from 0.1 to 0.7 m.s<sup>-1</sup> while median water temperature ranged from 6.6° to  
469 11.5°C; Table 1).

470 Including additional environmental descriptors could have increased understanding of trout  
471 synchrony. For example, food availability and oxygen concentration were not monitored in  
472 the study reaches, but may contribute to synchrony. Other descriptors of the available  
473 environmental regimes could also have been used (relating to magnitude, frequency, duration,  
474 timing or rate of change; Poff et al. 1997). For example, other thresholds, such as 2-year high  
475 seasonal discharge as used by Cattaneo et al. (2002), could have been tested. Finally,  
476 averaged physical conditions within a given reach could have been translated into hydraulic

477 habitat values that account for the heterogeneity of microhabitat conditions within the reach  
478 (Lamouroux and Capra 2002). However, to ensure statistical power, we only considered the  
479 variables that were most likely to explain density synchrony.

#### 480 **Influence of density-dependent dispersal on 0+ synchrony**

481 In previous papers, density-dependent dispersal between study populations was often ruled  
482 out *a priori* because pairs of populations were totally disconnected (Tedesco et al. 2004) or  
483 too distant (network distance >50km for 95% of the pairs of reaches in Cattaneo et al. 2003).  
484 In contrast, the present results on  $r0+$  synchrony remained significant when density-  
485 dependent dispersal between reaches was unlikely (tests on the first subset, where reaches  
486 could be connected by downstream drift only). Thus,  $r0+$  synchrony was linked to  
487 environmental synchrony and not to density-dependent dispersal. The limited density-  
488 dependent dispersal of 0+ reported in the literature also supports this conclusion. For  
489 example, Vollestad et al. (2012) mentioned a scale of dispersal of 200m while Dieterman and  
490 Hoxmeier (2011) and Vatland and Caudron (2015) estimated that only a small proportion of  
491 the 0+ population was involved in emigration from reaches due to density-dependence  
492 ( $\leq 10\%$ ).

493 By contrast, except for the test linking  $r0+$  and Euclidean distance, the present tests were no  
494 longer significant when dispersal between reaches was totally impossible, even by drift (tests  
495 on the second subset). This may partly be due to reduced statistical power (i.e., fewer reaches  
496 involved). However, this suggests that the possibility of drifting from one reach to another  
497 can explain synchronous emigrations from reaches that are not explained by high flow  
498 variables. Drift between reaches would potentially explain synchronous immigration, but  
499 cannot reasonably explain synchronous emigration of 0+. Therefore, this result again  
500 suggests that synchrony between geographically close reaches is not perfectly explained by  
501 the present high flow variables.

502 **Synchrony in 1+ and adults**

503 For older age-groups, except for a weak relation linking  $r1+$  synchrony and geographic  
504 distance, the present results were not able to explain trout synchrony. Authors often failed to  
505 identify constraining abiotic conditions for juveniles and adults (e.g., Cattaneo et al. 2002)  
506 except after exceptional events (e.g., a 50-year flood in Young et al. 2010). As they grow,  
507 stream-resident salmonids show increased swimming ability and may move more easily to  
508 avoid stressful conditions, reducing the influence of the environment (Unfer et al. 2011;  
509 Nislow and Armstrong 2012). Accordingly, movements of juveniles and adults toward  
510 sheltered areas during high flows (Bunt et al. 1999) or toward cold waters during droughts  
511 (Elliott 2000) have been reported. Moreover, movements of older individuals can occur at  
512 distances much larger than the reach (Ovidio et al. 1998), partially masking the links between  
513 trout density and the environment through sink/source recolonization processes (Zorn and  
514 Nuhfer 2007). The synchrony of  $r1+$  was related to Euclidean distance only, suggesting that  
515 environmental variables other than those studied here are likely involved.

516 In summary, the present study confirmed a Moran effect on 0+ trout density, operating  
517 mainly over distances <75km. A negative influence of high discharge was identified during  
518 emergence and a more complex relationship with spawning substrate mobility during the  
519 spawning period. Nevertheless, other environmental variables than those tested are likely  
520 involved. By contrast, dispersal between reaches had a weak influence on 0+ synchrony. The  
521 synchrony analyses provide useful information for building models of brown trout population  
522 dynamics integrating both biotic aspects (e.g., density-dependent and density-independent  
523 survival, dispersal) and abiotic mechanisms (e.g., the effects of high flow and spawning  
524 substrate movement). The results also suggest that brown trout populations may show low  
525 resilience in case of more frequent high flows over a given area (e.g., watershed). In a context

526 of global environmental change, further synchrony analyses are needed to better quantify the  
527 risk of extinction and potential resilience of freshwater fish metapopulations.

## 528 **Acknowledgments**

529 We thank the numerous people (working at ONEMA, Electricité de France, IRSTEA,  
530 ECOGEA, angling associations and other consulting firms) who contributed to electrofishing  
531 and habitat measurements. We also thank the organizations which provided environmental  
532 data (ONEMA, Météo France, and angling associations), Hervé Pella (IRSTEA Lyon) who  
533 provided the network distances and two anonymous reviewers who provided helpful  
534 suggestions.

## 535 **References**

- 536 Alonso, C., García de Jalón, D., Álvarez, J., and Gortázar, J. 2011. A large-scale approach  
537 can help detect general processes driving the dynamics of brown trout populations in  
538 extensive areas. *Ecol. Freshw. Fish* **20**(3): 449-460. doi:10.1111/j.1600-  
539 0633.2011.00484.x.
- 540 Armstrong, J. D., Kemp, P. S., Kennedy, G. J. A., Ladle, M., and Milner, N. J. 2003. Habitat  
541 requirements of Atlantic salmon and brown trout in rivers and streams. *Fish. Res.*  
542 **62**(2): 143-170. doi:doi:10.1016/S0165-7836(02)00160-1.
- 543 Bunt, C. M., Cooke, S. J., Katopodis, C., and McKinley, R. S. 1999. Movement and summer  
544 habitat of brown trout (*Salmo trutta*) below a pulsed discharge hydroelectric  
545 generating station. *Regul. River.* **15**(5): 395-403. doi:10.1002/(SICI)1099-  
546 1646(199909/10)15:5<395::AID-RRR556>3.0.CO;2-1.
- 547 Buonaccorsi, J. P., Elkinton, J. S., Evans, S. R., and Liebhold, A. M. 2001. Measuring and  
548 testing for spatial synchrony. *Ecology* **82**(6): 1668-1679. doi:10.1890/0012-  
549 9658(2001)082[1668:matfss]2.0.co;2.



- 550 Carle, R. T., and Strub, M. R. 1978. A new method for estimating population size from  
551 removal data. *Biometrics* **34**(4): 621-630.
- 552 Cattaneo, F., Lamouroux, N., Breil, P., and Capra, H. 2002. The influence of hydrological  
553 and biotic processes on brown trout (*Salmo trutta*) population dynamics. *Can. J. Fish.*  
554 *Aquat. Sci.* **59**: 12-22. doi:10.1139/f01-186.
- 555 Cattaneo, F., Hugueny, B., and Lamouroux, N. 2003. Synchrony in brown trout, *Salmo trutta*,  
556 population dynamics: a 'Moran effect' on early-life stages. *Oikos* **100**(1): 43-54.  
557 doi:doi:10.1034/j.1600-0706.2003.11912.x.
- 558 CEN (2003). Water quality – sampling of fish with electricity. European Standard.
- 559 Chevalier, M., Laffaille, P., and Grenouillet, G. 2014. Spatial synchrony in stream fish  
560 populations: influence of species traits. *Ecography* **37**(10): 960-968.  
561 doi:10.1111/ecog.00662.
- 562 Copeland, T., and Meyer, K. A. 2011. Interspecies Synchrony in Salmonid Densities  
563 Associated with Large-Scale Bioclimatic Conditions in Central Idaho. *Trans. Am.*  
564 *Fish. Soc.*(140:4): 928-942. doi:10.1080/00028487.2011.599261.
- 565 Cunjak, R. A., Linnansaari, T., and Caissie, D. 2013. The complex interaction of ecology and  
566 hydrology in a small catchment: a salmon's perspective. *Hydrol. Process.* **27**(5): 741-  
567 749. doi:10.1002/hyp.9640.
- 568 Daufresne, M., and Renault, O. 2006. Population fluctuations, regulation and limitation in  
569 stream-living brown trout. *Oikos* **113**(3): 459-468. doi:doi:10.1111/j.2006.0030-  
570 1299.14295.x.
- 571 Dieterman, D. J., and Hoxmeier, R. J. H. 2011. Demography of Juvenile and Adult Brown  
572 Trout in Streams of Southeastern Minnesota. *Trans. Am. Fish. Soc.* **140**(6): 1642-  
573 1656. doi:10.1080/00028487.2011.641883.

- 574 Elliott, J. M. 1984. Growth, size, biomass and production of young migratory trout *Salmo*  
575 *trutta* in a lake district stream, 1966-83. *J. Anim. Ecol.* **53**: 979-994.
- 576 Elliott, J. M. 2000. Pools as refugia for brown trout during two summer droughts: trout  
577 responses to thermal and oxygen stress. *J. Fish Biol.* **56**(4): 938-948.  
578 doi:doi:10.1111/j.1095-8649.2000.tb00883.x.
- 579 Elliott, J. M., and Elliott, J. A. 2010. Temperature requirements of Atlantic salmon *Salmo*  
580 *salar*, brown trout *Salmo trutta* and Arctic charr *Salvelinus alpinus*: predicting the  
581 effects of climate change. *J. Fish Biol.* **77**(8): 1793-1817. doi:10.1111/j.1095-  
582 8649.2010.02762.x.
- 583 Filipe, A. F., Markovic, D., Pletterbauer, F., Tisseuil, C., De Wever, A., Schmutz, S., Bonada,  
584 N., and Freyhof, J. 2013. Forecasting fish distribution along stream networks: brown  
585 trout (*Salmo trutta*) in Europe. *Diversity and Distributions* **19**(8): 1059-1071.  
586 doi:10.1111/ddi.12086.
- 587 Gouraud, V., Baran, P., Bardonnnet, A., Beaufrère, C., Capra, H., Caudron, A., Delacoste, M.,  
588 Lescaux, J. M., Naura, M., Ovidio, M., Poulet, N., Tissot, L., Sebaston, C., and  
589 Baglinière, J.-L. 2014. Sur quelles connaissances se baser pour évaluer l'état de santé  
590 des populations de truite commune (*Salmo trutta*)? *Hydroécologie Appliquée*: 1-28.  
591 doi:10.1051/hydro/2014001.
- 592 Grenouillet, G., Hugueny, B., Carrel, G. A., Olivier, J. M., and Pont, D. 2001. Large-scale  
593 synchrony and inter-annual variability in roach recruitment in the Rhône River: the  
594 relative role of climatic factors and density-dependent processes. *Freshw. Biol.* **46**(1):  
595 11-26. doi:10.1046/j.1365-2427.2001.00637.x.
- 596 Hayes, J. W. 1995. Spatial and temporal variation in the relative density and size of juvenile  
597 brown trout in the Kakanui River, North Otago, New Zealand. *New Zeal. J. Mar.*  
598 *Fresh.* **29**(3): 393-407. doi:10.1080/00288330.1995.9516674.

- 599 Heggenes, J. A. N. 1996. Habitat selection by brown trout (*Salmo trutta*) and young atlantic  
600 salmon (*S. salar*) in streams : Static and dynamic hydraulic modelling. Regulated  
601 Rivers: Research & Management **12**(2-3): 155-169. doi:10.1002/(SICI)1099-  
602 1646(199603)12:2/3<155::AID-RRR387>3.0.CO;2-D.
- 603 Holyoak, M., and Lawler, S. P. 1996. Persistence of an Extinction-Prone Predator-Prey  
604 Interaction Through Metapopulation Dynamics. Ecology **77**(6): 1867.  
605 doi:10.2307/2265790.
- 606 Jensen, A. J., and Johnsen, B. O. 1999. The functional relationship between peak spring  
607 floods and survival and growth of juvenile Atlantic Salmon (*Salmo salar*) and Brown  
608 Trout (*Salmo trutta*). Funct. Ecol. **13**(6): 778-785. doi:10.1046/j.1365-  
609 2435.1999.00358.x.
- 610 Koenig, W. D. 2002. Global patterns of environmental synchrony and the Moran effect.  
611 Ecography **25**(3): 283-288. doi:10.1034/j.1600-0587.2002.250304.x.
- 612 Lamouroux, N., and Capra, H. 2002. Simple predictions of instream habitat model outputs for  
613 target fish populations. Freshw. Biol. **47**: 1543–1556. doi:10.1046/j.1365-  
614 2427.2002.00879.x.
- 615 Liebhold, A., Koenig, W. D., and Bjørnstad, O. N. 2004. Spatial synchrony in population  
616 dynamics. Annu. Rev. Ecol. Evol. Syst. **35**(1): 467-490.  
617 doi:doi:10.1146/annurev.ecolsys.34.011802.132516.
- 618 Liebhold, A. M., Johnson, D. M., and Bjørnstad, O. N. 2006. Geographic variation in density-  
619 dependent dynamics impacts the synchronizing effect of dispersal and regional  
620 stochasticity. Popul. Ecol. **48**(2): 131-138. doi:10.1007/s10144-005-0248-6.
- 621 Lobón-Cerviá, J. 2004. Discharge-dependent covariation patterns in the population dynamics  
622 of brown trout (*Salmo trutta*) within a Cantabrian river drainage. Can. J. Fish. Aquat.  
623 Sci. **61**(10): 1929-1939. doi:10.1139/F04-118.

- 624 Lobón-Cerviá, J. 2009. Why, when and how do fish populations decline, collapse and  
625 recover? The example of brown trout (*Salmo trutta*) in Rio Chaballos (northwestern  
626 Spain). *Freshw. Biol.* **54**(6): 1149-1162. doi:10.1111/j.1365-2427.2008.02159.x.
- 627 Lobón-Cerviá, J. 2013. Recruitment and survival rate variability in fish populations: density-  
628 dependent regulation or further evidence of environmental determinants? *Can. J. Fish.*  
629 *Aquat. Sci.* **71**(2): 290-300. doi:10.1139/cjfas-2013-0320.
- 630 Mantel, N. 1967. The detection of disease clustering and a generalized regression approach.  
631 *Cancer Res.* **27**: 209-220.
- 632 Moran, P. A. P. 1953. The statistical analysis of the Canadian Lynx cycle. *Australian Journal*  
633 *of Zoology* **1**(3): 291. doi:10.1071/zo9530291.
- 634 Myers, R. A., Mertz, G., and Bridson, J. 1997. Spatial scales of interannual recruitment  
635 variations of marine, anadromous, and freshwater fish. *Can. J. Fish. Aquat. Sci.* **54**(6):  
636 1400-1407 doi:10.1139/f97-045.
- 637 Nicola, G. G., Almodóvar, A., Jonsson, B., and Elvira, B. 2008. Recruitment variability of  
638 resident brown trout in peripheral populations from southern Europe. *Freshw. Biol.*  
639 **53**(12): 2364-2374. doi:10.1111/j.1365-2427.2008.02056.x.
- 640 Nicola, G. G., Almodóvar, A., and Elvira, B. 2009. Influence of hydrologic attributes on  
641 brown trout recruitment in low-latitude range margins. *Oecologia* **160**(3): 515-524.  
642 doi:10.1007/s00442-009-1317-x.
- 643 Nislow, K. H., and Armstrong, J. D. 2012. Towards a life-history-based management  
644 framework for the effects of flow on juvenile salmonids in streams and rivers.  
645 *Fisheries Management and Ecology* **19**(6): 451-463. doi:10.1111/j.1365-  
646 2400.2011.00810.x.

- 647 Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Simpson,  
648 G. L., Solymos, P., Stevens, M. H. H., and Wagner, H. (2013). vegan: Community  
649 Ecology Package. R package version 2.0-10.
- 650 Olsen, E. M., and Vøllestad, L. A. 2005. Small-scale spatial variation in age and size at  
651 maturity of stream-dwelling brown trout, *Salmo trutta*. Ecol. Freshw. Fish **14**(2): 202-  
652 208. doi:10.1111/j.1600-0633.2005.00094.x.
- 653 Ovidio, M., Baras, E., Goffaux, D., Birtles, C., and Philippart, J. C. 1998. Environmental  
654 unpredictability rules the autumn migration of brown trout (*Salmo trutta* L.) in the  
655 Belgian Ardennes. Hydrobiologia **371-372**(0): 263-274.  
656 doi:10.1023/A:1017068115183.
- 657 Pella, H., Lejot, J., Lamouroux, N., and Snelder, T. H. 2012. Le réseau hydrographique  
658 théorique (RHT) français et ses attributs environnementaux. Geomorphologie **3/2012**:  
659 317-336. doi:10.4000/geomorphologie.9933.
- 660 Poff, N. L., Allan, J. D., Bain, M. B., Karr, J. R., Prestegard, K. L., Richter, B. D., Sparks,  
661 R. E., and Stromberg, J. C. 1997. The natural flow regime : a paradigm for river  
662 conservation and restoration. Bioscience **47**(11): 769-784
- 663 Ranta, E., Kaitala, V., and Lundberg, P. 1998. Population variability in space and time: the  
664 dynamics of synchronous population fluctuations. Oikos **83**: 376-382.  
665 doi:10.2307/3546852
- 666 Richard, A., Cattaneo, F., and Rubin, J.-F. 2013. Biotic and abiotic regulation of a low-  
667 density stream-dwelling brown trout (*Salmo trutta*) population: effects on juvenile  
668 survival and growth. Ecol. Freshw. Fish: n/a-n/a. doi:10.1111/eff.12116.
- 669 Ripa, J. 2000. Analysing the Moran effect and dispersal: their significance and interaction in  
670 synchronous population dynamics. Oikos **89**(1): 175-187. doi:10.1034/j.1600-  
671 0706.2000.890119.x.

- 672 Roussel, J. M., and Bardonnnet, A. 2002. Habitat de la truite commune (*Salmo trutta L.*)  
673 pendant la période juvénile en ruisseau : préférences, mouvements, variations  
674 journalières et saisonnières. Bull. Fr. Pêche Piscic. **365/366**: 435-454.  
675 doi:10.1051/kmae:2002044.
- 676 Ruiz, P., and Laplanche, C. 2010. A hierarchical model to estimate the abundance and  
677 biomass of salmonids by using removal sampling and biometric data from multiple  
678 locations. Can. J. Fish. Aquat. Sci. **67**(12): 2032-2044. doi:10.1139/F10-123.
- 679 Sabaton, C., Souchon, Y., Capra, H., Gouraud, V., Lascaux, J. M., and Tissot, L. 2008. Long-  
680 term brown trout populations responses to flow manipulation. River. Res. Applic.  
681 **24**(5): 476-505. doi:10.1002/rra.1130.
- 682 Santin-Janin, H., Hugueny, B., Aubry, P., Fouchet, D., Gimenez, O., and Pontier, D. 2014.  
683 Accounting for sampling error when inferring population synchrony from time-series  
684 data: a Bayesian state-space modelling approach with applications. PloS one **9**(1):  
685 e87084. doi:10.1371/journal.pone.0087084.
- 686 Smouse, P. E., Long, J. C., and Sokal, R. R. 1986. Multiple regression and correlation  
687 extensions of the Mantel test of matrix correspondence. Syst. Zool. **35**(4): 627-632.
- 688 Tedesco, P. A., Hugueny, B., Paugy, D., and Fermon, Y. 2004. Spatial synchrony in  
689 population dynamics of West African fishes: a demonstration of an intraspecific and  
690 interspecific Moran effect. J. Anim. Ecol. **73**(4): 693-705. doi:10.1111/j.0021-  
691 8790.2004.00843.x.
- 692 Unfer, G., Hauer, C., and Lautsch, E. 2011. The influence of hydrology on the recruitment of  
693 brown trout in an Alpine river, the Ybbs River, Austria. Ecol. Freshw. Fish **20**(3):  
694 438-448. doi:10.1111/j.1600-0633.2010.00456.x.
- 695 Vatland, S., and Caudron, A. 2015. Movement and early survival of age-0 brown trout.  
696 Freshw. Biol.: n/a-n/a. doi:10.1111/fwb.12551.

- 697 Vollestad, L. A., Serbezov, B., Bass, A., Bernatchez, L., Olsen, E. M., and Taugbøl, A. 2012.  
 698 Small-scale dispersal and population structure in stream-living brown trout (*Salmo*  
 699 *trutta*) inferred by mark–recapture, pedigree reconstruction, and population genetics.  
 700 *Can. J. Fish. Aquat. Sci.* **69**(9): 1513–1524. doi:10.1139/F2012-073.
- 701 Young, R. G., Hayes, J. W., Wilkinson, J., and Hay, J. 2010. Movement and mortality of  
 702 adult brown trout in the Motupiko River, New Zealand: effects of water temperature,  
 703 flow, and flooding. *Trans. Am. Fish. Soc.* **139**(1): 137-146. doi:10.1577/T08-148.1.
- 704 Zorn, T. G., and Nuhfer, A. J. 2007. Regional Synchrony of Brown Trout and Brook Trout  
 705 Population Dynamics among Michigan Rivers. *Trans. Am. Fish. Soc.* **136**(3): 706-  
 706 717. doi:10.1577/t06-275.1.

707

708 **Tables**709 **Table 1.** Physical characteristics of the 36 stream reaches.

Physical characteristics	Min	Mean	Max
Width at median discharge (m)	2.9	8.2	15.5
Reach slope (%)	0.3	3.4	13.2
Elevation (m)	15.0	787.7	1370.0
Distance from source (km)	3.0	17.9	49.0
Basin area (km <sup>2</sup> )	9.0	135.1	605.0
Median daily discharge Q <sub>50</sub> (m <sup>3</sup> .s <sup>-1</sup> )	0.1	1.0	2.7
Median diameter of streambed particle (cm)	0.1	16.6	64.0
Reach flow velocity (m.s <sup>-1</sup> ) at Q <sub>50</sub>	0.1	0.4	0.7
Median daily water temperature (°C)	6.6	8.3	11.5

710

711 **Table 2.** Environmental variables used in Mantel tests relating trout synchrony to environmental  
 712 synchrony (discharge, hydraulics or temperature regime).

<i>Variable group</i>		<b>Periods</b>			
<b>Code</b>	<b>Definition</b>	<b>Spawning</b>	<b>Egg</b>	<b>Emergence</b>	<b>Summer</b>
<i>Discharge</i>					
$Q_{90}$	daily discharge exceeded 90% of the time				$r0+$ ; $r1+$ ; $rAd$
$Q_{10}$	daily discharge exceeded 10% of the time	$rAd$	<b><math>r0+</math></b>	<b><u><math>r0+</math></u></b> ; $r1+$ ; $rAd$	
<i>Hydraulics</i>					
$fV0.5$	Frequency of current velocity $> 0.5\text{m}\cdot\text{s}^{-1}$			$r0+$	
$fMob$	Mobility frequency of the spawning substrate	<b><u><math>r0+</math></u></b>	<b><math>r0+</math></b>	$r0+$	
<i>Temperature</i>					
$fTlow$	Frequency of $T_{water,d}$ below threshold		$r0+$ ( $<4.3^{\circ}\text{C}$ )	$r0+$ ( $<7.3^{\circ}\text{C}$ )	

713 **Note:** Univariate Mantel tests relating synchrony in residual density of each age-group ( $r0+$ ,  
 714  $r1+$ ,  $rAd$ ) and environmental synchrony were made for a subset of environmental descriptors  
 715 (variables $\times$ periods). Bold values correspond to significant associations (p-value  $<0.05$ ) and  
 716 the most significant association for a given variable group is underlined.



717 **Table 3.** Results of Partial Mantel tests analyzing the association of  $r0+$  synchrony, geographic  
 718 distance and synchrony of one environmental descriptor.

Effect tested	Effect removed	Partial p-value	Partial $r^2$	% tests still significant on first subset	% tests still significant on second subset
$\text{Log}(\text{Euclid dist})$	$Q_{10}$ Emergence	<0.01	0.06	100	35
$Q_{10}$ Emergence	$\text{Log}(\text{Euclid dist})$	0.20			
$\text{Log}(\text{Euclid dist})$	$f\text{Mob}_{\text{Spawning}}$	<0.01	0.07	100	11
$f\text{Mob}_{\text{Spawning}}$	$\text{Log}(\text{Euclid dist})$	0.06			

719 **Note:** For significant partial tests, partial  $R^2$  was computed and we indicate the percentage of  
 720 significant tests when the Mantel test was repeated on two subsets of reaches (first subset:  
 721 limited dispersal between reaches; second subset: no dispersal).

722 **Figures**

723 **Fig. 1.** Locations of the 36 reaches (18 in bypassed sections).

724 **Fig. 2.** Relationships between raw densities of successive age-groups. Recruitment is linked to the  
725 adult density of the previous year (1<sup>st</sup> column) and 1+ are linked to previous density of 0+ (2<sup>nd</sup>  
726 column). Adult densities are linked to previous density of 1+ and adults (3<sup>rd</sup> and 4<sup>th</sup> columns); they are  
727 represented here after setting one of the two explanatory variables at its mean value across reaches.  
728 These relations are shown for all reaches pooled (1<sup>st</sup> row) and 5 randomly selected reaches (2<sup>nd</sup> to 5<sup>th</sup>  
729 rows). The solid lines correspond to a global linear regression and the dashed lines to linear mixed-  
730 effects models with a reach-level random effect.

731 **Fig. 3.** Trout synchrony (1<sup>st</sup> column) or environmental synchrony (2<sup>nd</sup> column) related to geographic  
732 distance. Linear relationships (full lines) correspond to the Mantel tests. Local polynomial fittings  
733 (dashed lines) present the smoothed spatial evolution of synchronies. For significant Mantel tests,  
734 Mantel  $R^2$  and p-values are given. The environmental variables showing the strongest relation to  
735 geographic distance (with highest  $R^2$ ) are shown for each environmental group (Table 2).

736 **Fig. 4.** Synchrony of  $r0+$  related to geographic distance (Euclidean or network distance), as in Figure  
737 3 but restricted to 93 pairs of reaches for which a network distance could be computed.

738 **Fig. 5.** Significant relationships (on Mantel tests) between the synchrony of  $r0+$  and the synchrony of  
739 environmental descriptors.

740 **Fig. 6.** Time series of  $r0+$  trout residual densities and environmental descriptors illustrating the most  
741 significant Mantel tests. Series were standardized by reach and grouped according to the results of a  
742 cluster analysis on trout synchrony. The cluster analysis was performed on the 18 (out of 36) reaches  
743 with more than 5 sampling year between 1999 and 2006.  $Q_{10}$  time series were multiplied by -1 to  
744 facilitate interpretation. The histograms in the right column represent the frequency distributions of  
745  $\rho_{r0+}$  between all pairs of reaches within the groups.




Fig. 1. Locations of the 36 reaches (18 in bypassed sections).  
169x159mm (300 x 300 DPI)




Fig. 2. Relationships between raw densities of successive age-groups. Recruitment is linked to the adult density of the previous year (1st column) and 1+ are linked to previous density of 0+ (2nd column). Adult densities are linked to previous density of 1+ and adults (3rd and 4th columns); they are represented here after setting one of the two explanatory variables at its mean value across reaches. These relations are shown for all reaches pooled (1st row) and 5 randomly selected reaches (2nd to 5th rows). The solid lines correspond to a global linear regression and the dashed lines to linear mixed-effects models with a reach-level random effect.

179x179mm (300 x 300 DPI)




Fig. 3. Trout synchrony (1st column) or environmental synchrony (2nd column) related to geographic distance. Linear relationships (full lines) correspond to the Mantel tests. Local polynomial fittings (dashed lines) present the smoothed spatial evolution of synchronies. For significant Mantel tests, Mantel  $R^2$  and  $p$ -values are given. The environmental variables showing the strongest relation to geographic distance (with highest  $R^2$ ) are shown for each environmental group (Table 2).

159x142mm (300 x 300 DPI)




Fig. 4. Synchrony of  $r_{0+}$  related to geographic distance (Euclidean or network distance), as in Figure 3 but restricted to 93 pairs of reaches for which a network distance could be computed.  
79x79mm (300 x 300 DPI)




Fig. 5. Significant relationships (on Mantel tests) between the synchrony of  $r_{0+}$  and the synchrony of environmental descriptors.  
199x500mm (300 x 300 DPI)




Fig. 6. Time series of  $r_{0+}$  trout residual densities and environmental descriptors illustrating the most significant Mantel tests. Series were standardized by reach and grouped according to the results of a cluster analysis on trout synchrony. The cluster analysis was performed on the 18 (out of 36) reaches with more than 5 sampling year between 1999 and 2006.  $Q_{10}$  time series were multiplied by -1 to facilitate interpretation. The histograms in the right column represent the frequency distributions of  $\rho_{r0+}$  between all pairs of reaches within the groups.

129x211mm (300 x 300 DPI)



## 1 **Appendix A: Daily data series**

### 2 **A.1 Discharge**

3 Daily discharge ( $\text{m}^3 \cdot \text{s}^{-1}$ ) was continuously gauged in 23 reaches (SP2T pressure probes). For  
4 seven reaches close to a gauging station (<40 km via the stream network), daily discharge  
5 was spatially extrapolated after correcting for drainage area. In the remaining six reaches,  
6 which were all below dams, daily discharge was calculated from the upstream discharge and  
7 the operating schedule of the dam. Less than 5% of the selected surveys had missing daily  
8 discharge values within the preceding year (with < 33% of missing values for key periods  
9 used in analysis).

### 10 **A.2 Hydraulic conditions: flow velocity and substrate mobility**

11 Reach daily current velocities ( $\text{m} \cdot \text{s}^{-1}$ ) were estimated as the ratio between daily discharge and  
12 the corresponding cross-section area (average depth  $\times$  average width). This required  
13 estimating depth-discharge and width-discharge relationships for the reach (i.e., at-a-reach  
14 hydraulic geometry relationships; Stewardson 2005). For this purpose and for 24 reaches,  
15 hydraulic geometry relationships were fitted to conventional power laws (Leopold and  
16 Maddock 1953) using estimates of depth and width made at two distinct discharges (median  
17 ratio between the two discharges: 5.1). For each measured discharge, the wetted widths of  
18 regularly spaced cross-sections ( $n > 15$ ) and the water depth at regularly spaced points along  
19 cross-sections ( $n > 100$  across the reach) were measured. For five reaches, the same method  
20 was used, except that cross-sections were not regularly spaced but weighted by the length of  
21 streams they represented. In the seven remaining reaches, the mean velocity was obtained  
22 from a numerical hydraulic model calibrated in the reach for habitat modeling purposes  
23 (Ginot et al. 1998).

24 Spawning substrate mobility was defined as the number of days with discharge above the  
 25 critical discharge theoretically moving particles of diameter 0.02 m (typical size for  
 26 spawning; Kondolf and Wolman 1993). This critical discharge was estimated using the  
 27 classical Shields' criterion (Shields 1936).

### 28 **A.3 Water temperature time series**

29 Depending on the availability of water temperature data for a given reach, three procedures  
 30 were used to estimate missing water temperature values.

#### 31 *Twenty-two reaches with available water temperature measurements*

32 These reaches had at least one year of daily water temperature data measured in the reach (on  
 33 average, 5.4 years of data per reach). To predict missing values in these reaches (29% of time  
 34 series), daily water temperature on day  $d$  ( $T_{water,d}$ ) was modeled from air temperature during  
 35 the three previous days ( $T_{air,d-2}$  to  $T_{air,d}$ ), partly accounting for inertia and hysteresis effects.  
 36 A logistic model (used on a weekly time-step in Mohseni et al. 1998) was implemented:

$$37 \quad (A1) \quad T_{water,d} = \mu + \frac{(A-\mu)}{(1+\exp[\Gamma \cdot (B-(0.5 \times T_{air,d} + 0.3 \times T_{air,d-1} + 0.2 \times T_{air,d-2}))])}$$

38  $T_{air}$  was recorded at meteorological stations close to our reaches (median Euclidean distance:  
 39 6.8 km, maximum: 31.1 km). Parameters corresponded to minimal ( $\mu$ ) and maximal ( $A$ ) water  
 40 temperature and air temperature at inflection point ( $B$ ) or were linked to the slope at  
 41 inflection point ( $\Gamma$ ). Each reach model was fitted using least-squares criteria (“nls” function  
 42 of R 3.1.1 software; R Core Team 2014).

43 Following Mohseni et al. (1998), Root Mean Squared Error (RMSE) and Nash-Sutcliffe  
 44 coefficients (NSC; Nash and Sutcliffe 1970) were computed for each reach to estimate  
 45 accuracy and goodness of fit. Water temperature models calibrated on these 22 reaches with

46 calibration data had a low RMSE (min: 0.7; mean: 1.0; max: 1.5 °C) and explained a large  
47 part of variability (min NSC: 0.82; mean: 0.90; max: 0.95). Fitting the parameter  $\mu$  (minimal  
48 water temperature) or setting it at 0 hardly changed the results (maximal variation: +2.8% in  
49 RMSE and -0.54% in NSC); it was therefore set at 0 in all reaches. Using lagged air  
50 temperature over three days instead of daily air temperature reduced the RMSE of the model  
51 by 14.6% on average.

52 *Five reaches close to another reach with water temperature data*

53 Five additional reaches were close to another reach (< 5km; mean: 2.6 km, with no tributary  
54 between reaches) with similar physical properties (<15% difference in flow or width; <10%  
55 difference in median grain size); the two were then considered to have similar water  
56 temperatures.

57 *Nine reaches without water temperature data*

58 For the nine remaining reaches, water temperature was predicted from air temperature using a  
59 global logistic model, fitted simultaneously on the 22 reaches with water temperature data.  
60 The global logistic model used had the same formulation as in Eq. (A1) but  
61 parameters  $\mu$ ,  $A$ ,  $B$  and  $\Gamma$  depended linearly on parameters that could explain differences  
62 between water and air temperatures (longitude, latitude, distance from the source, and  
63 difference in altitude between the reach and the air temperature recorder).

64 The RMSE of this global model was calculated when fitted on all available data. The global  
65 water temperature model had an RMSE of 1.3 and an NSC of 0.87. Cross-validation was then  
66 used (leaving one reach out in turn) to estimate the average residual absolute error on water  
67 temperature using the global model. Cross-validations indicated a mean absolute error of less  
68 than 1°C in 59% of cross-validations (maximum: 2.3°C for one reach).

69 **Appendix : References**

- 70 Ginot, V., Souchon, Y., Capra, H., Breil, P., and Valentin, S. (1998). Logiciel EVHA 2.0.  
71 Evaluation de l'habitat physique des poissons en rivière, Cemagref BEA/LHQ et  
72 Ministère de l'Aménagement du Territoire et de l'Environnement: 130.
- 73 Kondolf, G. M., and Wolman, M. G. 1993. The sizes of salmonid spawning gravels. *Water*  
74 *Resour. Res.* **29**(7): 2275-2285. doi:10.1029/93WR00402.
- 75 Leopold, L. E., and Maddock, T. 1953. The hydraulic geometry of stream channels and some  
76 physiographic implications. *USGS*(252): 57.
- 77 Mohseni, O., Stefan, H. G., and Erickson, T. R. 1998. A nonlinear regression model for  
78 weekly stream temperatures. *Water Resour. Res.* **34**(10): 2685-2692.  
79 doi:10.1029/98WR01877.
- 80 Nash, J. E., and Sutcliffe, J. V. 1970. River flow forecasting through conceptual models part I  
81 — A discussion of principles. *J. Hydrol.* **10**(3): 282-290.  
82 doi:http://dx.doi.org/10.1016/0022-1694(70)90255-6.
- 83 R Core Team (2014). *R: A language and environment for statistical computing.*, R  
84 Foundation for Statistical Computing, Vienna, Austria.
- 85 Shields, A. 1936. *Anwendung der Aehnlichkeitsmechanik und der Turbulenzforschung auf*  
86 *die Geschiebebewegung.* Technical University Berlin.
- 87 Stewardson, M. 2005. Hydraulic geometry of stream reaches. *J. Hydrol.* **306**(1-4): 97-111.  
88 doi:http://dx.doi.org/10.1016/j.jhydrol.2004.09.004.
- 89