



HAL
open science

Geographical distribution and morphometry of *Heimyscus fumosus* (Brosset et al., 1965)

Violaine Nicolas, Wim Wendelen, Walter Verheyen, Marc Colyn

► **To cite this version:**

Violaine Nicolas, Wim Wendelen, Walter Verheyen, Marc Colyn. Geographical distribution and morphometry of *Heimyscus fumosus* (Brosset et al., 1965). *Revue d'Écologie*, 2003, 58 (2), pp.197-208. hal-01346042

HAL Id: hal-01346042

<https://univ-rennes.hal.science/hal-01346042>

Submitted on 6 Sep 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GEOGRAPHICAL DISTRIBUTION AND MORPHOMETRY OF *HEIMYSCUS FUMOSUS* (BROSSET *ET AL.*, 1965)

Violaine NICOLAS¹, Wim WENDELEN², Walter VERHEYEN³ & Marc COLYN¹

RÉSUMÉ

Plus de 35 ans après sa description, le Muridé africain *Heimyscus fumosus* reste mal connu et peu de spécimens sont déposés dans les musées. Depuis 1992 nous avons collecté 239 spécimens dans dix localités d'Afrique centrale, nous permettant de confirmer sa distribution géographique. Cette espèce est présente dans les forêts de plaine situées entre les rivières Sanaga et Oubangui-Congo. De plus, l'étude d'une population du sud-ouest du Gabon nous a permis d'analyser les variations morphologiques intra-populationnelles chez cette espèce. Nos résultats mettent en évidence une augmentation avec l'âge de la plupart des mesures externes et crâniennes; en revanche, le dimorphisme sexuel est faible.

SUMMARY

More than 35 years after its original description, our knowledge of the De Balsac's mouse (*Heimyscus fumosus*) remains poor, and few specimens are deposited in museums. We collected 239 specimens of this species in ten central African localities. We therefore had the opportunity to confirm its geographical distribution, which covers lowland forests between the Sanaga and the Oubangui-Congo rivers. Moreover, based upon specimens collected from a single locality, we analysed intra-populational morphological variations of the De Balsac's mouse. We showed that most external and cranial measurements increased with age; while sexual dimorphism was low.

INTRODUCTION

De Balsac's Mouse (*Hylomyscus fumosus*) was described in 1965 from Gabon by Brosset *et al.* While the validity of the species is not questionable, its affiliation

¹ Laboratoire Ethologie-Evolution-Ecologie, C.N.R.S. U.M.R. 6552, Station Biologique, Université de Rennes 1, F-35380 Paimpont, France. Tel (33) 02.99.61.81.62 (81.63). E-mail marc.colyn@univ-rennes1.fr.

² Departement Vertebraten, Koninklijk Museum voor Midden-Afrika, Tervuren, Belgium.

³ Departement Biologie, R.U.C.A., Universitair Centrum Antwerpen, Groenenborgerlaan 171, 2020 Antwerpen, Belgium. E-mail: walter.verheyen@ua.ac.be.

to the genus *Hylomyscus* gives rise to controversy. Misonne (1969) suggested the creation of a new genus *Heimyscus* based on its cranial characteristics. He was followed by Meester & Setzer (1971), and Musser & Carleton (1993). Although Duplantier (1982), emphasized that the external morphology of *H. fumosus* differs significantly from that of other *Hylomyscus*, he kept it within the genus *Hylomyscus*. In this paper, we use the nomenclature proposed by Musser & Carleton (1993).

In the original description, the external and cranial morphology of *H. fumosus* was carefully described. However, no information on cranial measurements were given and few specimens were deposited in museums.

Our knowledge of *H. fumosus* remains poor. It is mentioned from five localities in Gabon (Brosset *et al.*, 1965), Central African Republic (Petter & Genest, 1970), Cameroon (Robbins *et al.*, 1980), Republic of Congo (Granjon, 1991), and Equatorial Guinea (Lasso, 1995), and one paper deals with its ecology (Duplantier, 1982).

In the context of several international projects on the tropical forest biodiversity, we carried out surveys on small mammals communities in four central African countries. During these field works we collected 239 *H. fumosus* from ten localities. In this paper, we precise the distribution of this species and its habitat preferences, and, for the first time, analyse morphological variations of one population from Gabon.

STUDY AREAS AND METHODS

From 1992 to 2001, we captured *H. fumosus* in ten localities (Table I) located in Cameroon, Central African Republic (C.A.R.), Republic of Congo and Gabon. They were trapped with Sherman and snap traps arranged in lines or grids, and with pitfall lines with drift fences (see details in Colyn *et al.*, 1996). In order to define the geographical distribution of *H. fumosus*, we combined data from our surveys, museums collections (Table I) and bibliographical data.

At most localities, several types of habitat were surveyed (primary forest, secondary forest, fallow land, savannahs). Details on habitat characteristics are given in Nicolas & Barrière (2001) for Monts Doudou, in Barrière & Nicolas (2000) for Batouri, and in Colyn *et al.* (1996) for other localities. Habitat preferences and spatial variations in trapping success (number of specimens caught per 100 trap-nights) were evaluated by Chi-square tests.

Intra-population variability of external and cranial measurements of *H. fumosus* was studied on 207 specimens trapped in Monts Doudou (Gabon). Five standard external measures were taken on freshly killed mice: weight (W), head+body length (HB), tail length (TL), hindfoot length with nail (HF), ear length (EL). Specimens were preserved in 10 % formalin and skulls were later on removed and cleaned at the lab. Twenty three craniometrical and dental dimensions were measured on every skull (Tab. II). All specimens were age-classified using the following stages of tooth eruption and tooth-wear patterns of the maxillary toothrows: age-class 0: M³ not yet fully erupted; age-class 1: all cheekteeth fresh but fully erupted, the exposed dentine of the three cusps of the first row of M¹ are not yet fused; age-class 2: wear is light, dentine-bridge between the cusps of the first row of M¹ becomes progressively wider as the cusps wear down; age-class 3: wear is obvious, the bottom of the groove between row 2 and 3 is reached and the groove is

TABLE I

Number of specimens of Heimyscus fumosus collected by our team in ten Central African localities, and number of specimens deposited in museum and reviewed for our analysis of the geographical distribution of this species. Localities of collect, name of the collector, year of capture, and museum where the specimens are deposited are precised. The following acronyms identify the museum: CMNH: Carnegie Museum of Natural History, MHNP: Museum d'Histoire Naturelle de Paris, MRAC: Musée Royal de l'Afrique Centrale de Tervuren, SBP: Station Biologique de Paimpont

| Country | Locality | Co-ordinates | Collector | Year of collect | Museum | Nb of skulls examined |
|---|-----------------|---------------|-----------|-----------------|--------|-----------------------|
| Specimens collected by our team: | | | | | | |
| Cameroon | Bodjouo | 03.21N-13.02E | | 1995 | SBP | 2 |
| C.A.R. | Grima | 04.02N-17.05E | | 1996 | SBP | 2 |
| | Bambio | 03.57N-16.58E | | 1994 | SBP | 1 |
| | Batouri | 03.54N-17.02E | | 1994 1999 | SBP | 6 |
| | Salo | 03.11N-16.06E | | 1994 | SBP | 2 |
| Rep. Congo | Mbomo | 00.24N-14.44E | | 1995 to 1997 | SBP | 8 |
| Gabon | La Makandé | 00.41S-11.55E | | 1995 | SBP | 7 |
| | Gongué | 00.47S-11.55E | | 1992 | SBP | 3 |
| | Monts Doudou | 02.09S-10.30E | | 2000 to 2001 | SBP | 207 |
| | Malounga | 03.09S-10.45E | | 1998 | SBP | 1 |
| Specimens in museums: | | | | | | |
| Cameroon | Efulen (1) | 02.47N-10.32E | Good | 1944 | CMNH | 1 |
| | Eseka (2) | 03.38N-10.47E | Robbins | 1974 | CMNH | 1 |
| | Yaoundé (3) | 03.52N-11.31E | Mouchet | 1956 | MHNP | 1 |
| C.A.R. | Kongana (10) | 02.47N-16.25E | Ray | 1993 | MRAC | 6 |
| | Boukoko (11) | 03.54N-17.55E | Pujol | 1962 | MHNP | 1 |
| Congo | Kouboutchi (13) | 04.19S-11.47E | Granjon | 1990 | MHNP | 2 |
| Gabon | Makokou (14) | 00.34N-12.52E | Dubost | 1962 to 1964 | MHNP | 18 |

TABLE II

Recapitulation and short description of the measurements as used in this study. For a full description we refer to Verheyen et al. (1996)

| Numbers | Acronyms | Morphometrical characters |
|---------|-------------------|--|
| M1 | GRLS | greatest length of skull |
| M2 | PRCO | condylobasal length |
| M3 | HEBA | henselion-basion |
| M4 | HEPA | henselion-palation |
| M5 | PAFL | length of palatal foramen |
| M6 | DIA1 | length of diastema |
| M7 | DIA2 | distance between alveolus M ¹ and cutting edge of upper incisor |
| M8 | INTE | smallest interorbital breadth |
| M9 | ZYGO | zygomatic breadth |
| M10 | PALA | smallest palatal breadth |
| M11 | UPTE | length of upper cheekteeth |
| M12 | UPDA | breath of upper dental arch |
| M13 | M ¹ BR | greatest breadth of first upper molar |
| M14 | ZYPL | smallest breadth of zygomatic plate |
| M15 | BNAS | greatest breath of nasals |
| M16 | LNAS | greatest length of nasals |
| M17 | LOTE | length of mandibular teeth |
| M18 | BULL | length of auditory bulla |
| M19 | BRCA | greatest breadth of braincase |
| M20 | DINC | depth of upper incisor |
| M21 | ROHE | mediosagittal projection of rostrum height |
| M22 | ROBR | greatest rostrum breadth |
| M23 | PCPA | distance between coronoid and angular processes |

interrupted on the lingual side of the molar; age-class 4: wear is extensive, the dentine of row 2 and 3 make contact with each other; age-class 5: wear is severe, M¹ is heavily eroded and all rows are communicating. Sexual dimorphism and growth were statistically evaluated by Student t-test on the basis of both external and cranial measurements.

RESULTS

GEOGRAPHICAL DISTRIBUTION AND HABITAT PREFERENCE (Fig. 1)

H. fumosus, described from Makokou (Gabon, Brosset *et al.*, 1965), is also mentioned from Eseka (Cameroon, Robbins *et al.*, 1980), La Maboké (C.A.R., Peter & Genest, 1970), Monte Alen (Equatorial Guinea, Lasso, 1995) and Kouboutchi (Republic of Congo, Granjon, 1991); specimens from Yaoundé and Efulen (Cameroon), and from Kongana and Boukoko (C.A.R.) are also deposited in different museums. We collected this species in ten additional localities: Bodjouo (Cameroon), Grima, Bambio, Batouri and Salo (C.A.R.), Mbomo (Republic of Congo), La Makandé, Gongué, Monts Doudou and Malounga (Gabon).

Further to the West, in Nigeria (Gambari forest, Happold, 1977; Gotel Mountains and Mambilla Plateau, Hutterer *et al.*, 1992), Ghana (Jeffrey, 1977), Ivory Coast (Adam, 1977; Dosso, 1983; Collinet *et al.*, 1984), Liberia (Mont Nimba, Verschuren, 1986) and Guinea (Mont Nimba, Gautun, 1986) *H. fumosus* was never captured. The same applies to the North for the C.A.R (Bamingui-Brendja, Colyn, unpub. data), and to the East for the Democratic Republic of Congo (Kikwit, Leirs *et al.*, 1999; Ituri forest, Gubista, 1999; Masako, Dudu, 1991; Kisangani left and right bank of the Congo river, Colyn unpub. data). More eastward, this species was




Figure 1. — Geographical distribution of *Heimyscus fumosus*. The symbol + refers to the type locality, * refers to specimen deposit in a museum, and ● refers to our own data. The numbers refer to the following localities: (1) Efulen, (2) Eseka, (3) Yaoundé, (4) Bodjouo, (5) Grima, (6) Bambio, (7) Batouri, (8) La Maboké, (9) Boukoko, (10) Salo, (11) Kongana, (12) Monte Alen, (13) Makokou, (14) Mbomo, (15) La Makandé, (16) Gongué, (17) Monts Doudou, (18) Malounga, (19) Kouboutchi.

never found in Uganda (Mayanja, Delany, 1971; Kibale, Struhsaker, 1997) and Malawi (Zomba Plateau, Nyika Plateau and Mulanje Mountain, Happold & Happold, 1989).

All these data confirm that *H. fumosus* is only present in lowland forests, between the Sanaga River (West) and the Oubangui-Congo River (East).

In Monts Doudou, all specimens were captured in primary forest, and at the nine other localities, 28 out of 32 specimens were collected in primary forest. The four others were collected in fallow land (two in Gougué and one in Bodjouo) or in secondary forest (one in Batouri). *H. fumosus* seems to prefer primary forest, rather than secondary forest or open areas. Moreover, it would prefer *terra firme* forest rather than riparian or swamp forest, as indicated by captures realized in Monts Doudou forest: trapping success of 0.17, 0.07 and 0.07 in *terra firme*, riparian and swamp forests respectively ($\chi^2 = 6.822$, $df = 2$, $P < 0.05$). Finally, this species seems to be mainly terrestrial since no capture occurred on lianas or branches.

For a given habitat (*terra firme* forest), the abundance of *H. fumosus* varied depending on sites. For example, its trapping success was of 0.17, 0.02 and 0.02 in Monts Doudou, Mbomo and Batouri respectively ($\chi^2 = 61.400$, $df = 2$, $P < 0.001$) and its relative abundance among the murid community was of 9, 4.3 and 1.1 % respectively at the same localities.

INTRA-POPULATIONAL MORPHOLOGICAL VARIATIONS

The dorsal pelage is dark-grey, a little reddish for old individuals; the ventral colour, from the cheeks to the beginning of the tail, is whitish. The belly hairs are grey basally and white in the terminal half. The coloration of the belly passes gradually into that of the back. The dorsal sides of the fore and hindfeet are white, and the morphology of the hindfoot agrees with the description given by Brosset *et al.* (1965). The ear, black-grey, is well developed (mean length 16.6 mm). The tail is slightly longer than the head and body length. The mean length of the tail for adult specimens is 95 mm (70 mm – 115 mm); the head and body length are 90 mm (68 mm – 104 mm). The mean hindfoot length for adult specimens is 20.2 mm (16.3 mm – 22.1 mm). Females have 3 pairs of nipples (1 pectoral and 2 inguinal).

Figure 2 illustrates the variation present in cusp and ridge patterns of the right maxillary teeth of four representative specimens among the 207 specimens examined. Cusp and ridge patterns are rather stable from one individual to another. In 40 % of the specimens (e.g. GA1984 and GA2982), the t_3 cusp of M^1 shows a beginning of splitting in two. The two most variable characters are the length and breadth of the molars.

Sexual dimorphism was tested for each age-class separately, and for all age-classes combined. Whatever the age-class, mice of both sexes do not differ significantly for most of the considered variables. When all age-classes are pooled together in analysis, males have a larger interorbital breadth (M8) and rostrum breadth (M22) than females (Table III).

The influence of growth on cranial and external measures was analysed for each sex separately, and for both sexes combined. Whatever the sex, most measurements increase significantly with age. Only five measurements out of 28 did not vary significantly: hindfoot length (HF), upper cheekteeth length (M11), upper molar breadth (M13), mandibular cheekteeth length (M17) and auditory bulla length (M18; Table IV). The increase with age, of skull length (M1), zygomatic breadth




Figure 2. — Comparative drawings of the right maxillary teeth of *Heimyscus fumosus*. Left to right: GA3440, GA1984, GA1437 and GA2982.

(M9), nasals breadth (M15), length of nasals (M16), braincase breadth (M19) and rostrum breadth (M22) are visualized in Figure 3.

DISCUSSION

H. fumosus is present in a large area of lowland forests, between the Sanaga River and the Oubangui-Congo River. Its low trap success could explain its rarity in museum collections.

It seems to prefer primary forest, and particularly *terra firme* forest. However, it can also be present occasionally in adjacent habitats. Thus, it was collected in manioc fields, at the immediate vicinity of a marshland, and in riparian and secondary forest by Brosset *et al.* (1965) and by Lasso (1995). The preference of *H. fumosus* for *terra firme* forest could be related to the fact that this species digs burrows (Duplantier, 1982), has a poor swimming ability (V.N. pers. obs.), and is terrestrial (Brosset *et al.*, 1965; Duplantier, 1982; our data).

Misonne (1969) suggested the creation of the genus *Heimyscus* for this species, owing to its cranial morphology. However, no thorough comparative effort was made to justify his proposition, and a comparative revision of *Hylomyscus* and *Heimyscus* is still needed. According to Verheyen & Bracke (1966), in order to make a systematic revision of a given Murid-species, one must make first a thorough morphological study of one “interbreeding population”. Our study on the intra-popula-

TABLE III

Comparison between males and females of *Heimyscus fumosus* for 5 external measurements and 23 cranial measurements. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. To provide general trends results for all age-classes combined are given. Body weights are in grams and other measurements are in millimetres

| | females | | | | | | males | | | | | |
|-----|---------|-------|-------|--------|-------|---------|-------|-------|-------|--------|------|------|
| | n | mean | min | max | SD | t-value | n | mean | min | max | SD | |
| W | 83 | 17.19 | 7.00 | 27.00 | 3.97 | -0.25 | 102 | 17.33 | 7.80 | 23.00 | 3.62 | |
| BL | 90 | 86.82 | 62.00 | 102.00 | 8.95 | -0.39 | 112 | 87.28 | 64.00 | 104.00 | 7.68 | |
| TL | 88 | 91.11 | 37.00 | 115.00 | 11.16 | -1.62 | 109 | 93.40 | 70.00 | 111.00 | 8.73 | |
| HF | 90 | 19.87 | 17.00 | 22.00 | 0.92 | -0.18 | 112 | 19.91 | 9.40 | 22.10 | 1.46 | |
| EL | 80 | 16.37 | 12.00 | 25.00 | 1.59 | -0.37 | 106 | 16.45 | 13.30 | 19.20 | 1.22 | |
| M1 | 67 | 25.42 | 22.65 | 27.70 | 0.98 | 0.39 | 88 | 25.36 | 22.30 | 27.05 | 1.08 | |
| M2 | 71 | 23.35 | 20.05 | 25.45 | 1.05 | -0.17 | 99 | 23.38 | 19.90 | 25.20 | 1.10 | |
| M3 | 72 | 19.99 | 16.95 | 22.15 | 0.98 | -0.13 | 99 | 20.01 | 16.65 | 21.65 | 1.02 | |
| M4 | 77 | 11.15 | 9.75 | 12.25 | 0.53 | 0.23 | 102 | 11.13 | 9.70 | 12.05 | 0.53 | |
| M5 | 77 | 3.95 | 3.15 | 4.55 | 0.27 | 0.52 | 102 | 3.92 | 3.10 | 4.75 | 0.30 | |
| M6 | 77 | 7.16 | 5.80 | 8.20 | 0.44 | 0.01 | 102 | 7.16 | 5.90 | 7.95 | 0.43 | |
| M7 | 77 | 7.84 | 6.40 | 8.85 | 0.49 | -0.17 | 102 | 7.86 | 6.40 | 8.85 | 0.49 | |
| M8 | 77 | 4.56 | 3.95 | 5.00 | 0.19 | -2.62 | ** | 101 | 4.64 | 4.20 | 5.10 | 0.21 |
| M9 | 73 | 11.68 | 10.50 | 12.95 | 0.44 | 0.41 | 101 | 11.66 | 10.45 | 12.75 | 0.44 | |
| M10 | 77 | 2.73 | 2.35 | 3.35 | 0.18 | 0.22 | 102 | 2.73 | 2.20 | 3.10 | 0.17 | |
| M11 | 77 | 3.61 | 3.20 | 3.80 | 0.13 | -0.78 | 102 | 3.63 | 3.30 | 4.05 | 0.15 | |
| M12 | 77 | 5.07 | 4.70 | 5.70 | 0.20 | -0.24 | 102 | 5.08 | 4.60 | 5.50 | 0.19 | |
| M13 | 77 | 1.18 | 1.05 | 1.25 | 0.04 | 0.95 | 102 | 1.18 | 1.10 | 1.35 | 0.05 | |
| M14 | 77 | 2.21 | 1.75 | 2.55 | 0.16 | 0.82 | 102 | 2.20 | 1.75 | 2.55 | 0.15 | |
| M15 | 77 | 2.49 | 1.90 | 3.10 | 0.19 | -1.00 | 102 | 2.52 | 2.15 | 2.85 | 0.16 | |
| M16 | 72 | 9.89 | 7.70 | 11.20 | 0.66 | 0.16 | 95 | 9.88 | 7.85 | 11.25 | 0.69 | |
| M17 | 77 | 3.42 | 3.15 | 3.70 | 0.13 | -0.03 | 102 | 3.42 | 3.20 | 3.70 | 0.12 | |
| M18 | 75 | 4.07 | 3.75 | 4.50 | 0.18 | 1.19 | 100 | 4.04 | 3.55 | 4.60 | 0.18 | |
| M19 | 75 | 10.54 | 9.80 | 11.30 | 0.32 | 0.15 | 102 | 10.54 | 9.55 | 11.20 | 0.30 | |
| M20 | 77 | 1.14 | 0.85 | 1.35 | 0.09 | -1.52 | 102 | 1.16 | 0.90 | 1.40 | 0.10 | |
| M21 | 77 | 5.19 | 4.50 | 5.85 | 0.27 | -1.02 | 101 | 5.24 | 4.50 | 5.80 | 0.29 | |
| M22 | 77 | 4.37 | 3.70 | 5.15 | 0.30 | -2.09 | * | 102 | 4.47 | 3.80 | 5.10 | 0.26 |
| M23 | 76 | 6.51 | 5.30 | 7.60 | 0.43 | -0.05 | 102 | 6.52 | 5.45 | 7.50 | 0.43 | |

TABLE IV

Comparison between *Heimyscus fumosus* of age-classes 1 and 2 for 5 external measurements and 23 cranial measurements. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Body weights are in grams and other measurements are in millimetres

| | age-class 1 | | | | | | | age-class 2 | | | | |
|-----|-------------|-------|-------|--------|------|---------|-----|-------------|-------|-------|--------|------|
| | n | mean | min | max | SD | t-value | | n | mean | min | max | SD |
| W | 76 | 15.91 | 7.80 | 25.00 | 3.62 | -5.37 | *** | 75 | 18.57 | 13.30 | 23.00 | 2.31 |
| HB | 83 | 84.36 | 64.00 | 98.00 | 7.32 | -5.75 | *** | 80 | 90.33 | 70.00 | 104.00 | 5.80 |
| TL | 80 | 89.93 | 70.00 | 108.00 | 8.46 | -4.16 | *** | 80 | 95.74 | 37.00 | 111.00 | 9.19 |
| HF | 82 | 19.71 | 9.40 | 22.10 | 1.58 | -1.80 | | 81 | 20.07 | 17.60 | 22.10 | 0.89 |
| EL | 77 | 16.19 | 13.30 | 18.70 | 1.13 | -3.12 | * | 78 | 16.82 | 14.20 | 25.00 | 1.40 |
| M1 | 66 | 25.02 | 22.30 | 27.05 | 1.03 | -4.19 | *** | 73 | 25.66 | 22.60 | 26.85 | 0.78 |
| M2 | 71 | 22.91 | 20.05 | 24.95 | 1.04 | -5.20 | *** | 82 | 23.67 | 20.75 | 25.00 | 0.74 |
| M3 | 72 | 19.58 | 17.25 | 21.55 | 0.94 | -5.24 | *** | 82 | 20.28 | 17.55 | 21.60 | 0.70 |
| M4 | 77 | 10.92 | 9.70 | 11.80 | 0.54 | -5.02 | *** | 85 | 11.30 | 10.20 | 12.00 | 0.40 |
| M5 | 77 | 3.85 | 3.10 | 4.55 | 0.29 | -3.95 | *** | 85 | 4.02 | 3.40 | 4.75 | 0.25 |
| M6 | 77 | 6.96 | 5.90 | 7.65 | 0.40 | -5.93 | *** | 85 | 7.29 | 6.30 | 7.95 | 0.31 |
| M7 | 77 | 7.62 | 6.50 | 8.40 | 0.45 | -6.33 | *** | 85 | 8.01 | 7.00 | 8.75 | 0.33 |
| M8 | 77 | 4.57 | 3.95 | 5.10 | 0.22 | -1.91 | | 84 | 4.63 | 4.30 | 5.00 | 0.17 |
| M9 | 75 | 11.51 | 10.50 | 12.30 | 0.45 | -3.52 | *** | 84 | 11.73 | 10.45 | 12.60 | 0.34 |
| M10 | 77 | 2.66 | 2.20 | 3.10 | 0.16 | -4.52 | *** | 85 | 2.76 | 2.50 | 3.05 | 0.13 |
| M11 | 77 | 3.62 | 3.35 | 4.00 | 0.14 | 0.20 | | 85 | 3.61 | 3.20 | 4.05 | 0.15 |
| M12 | 77 | 4.99 | 4.60 | 5.30 | 0.16 | -4.78 | *** | 85 | 5.11 | 4.75 | 5.50 | 0.16 |
| M13 | 77 | 1.18 | 1.10 | 1.25 | 0.04 | -0.55 | | 85 | 1.18 | 1.05 | 1.35 | 0.05 |
| M14 | 77 | 2.18 | 1.75 | 2.45 | 0.15 | -1.91 | | 85 | 2.22 | 1.80 | 2.55 | 0.14 |
| M15 | 77 | 2.45 | 1.90 | 2.85 | 0.18 | -4.17 | *** | 85 | 2.56 | 2.25 | 2.90 | 0.15 |
| M16 | 73 | 9.63 | 7.70 | 10.95 | 0.71 | -4.65 | *** | 79 | 10.09 | 8.20 | 11.25 | 0.51 |
| M17 | 77 | 3.43 | 3.20 | 3.70 | 0.11 | 0.86 | | 85 | 3.41 | 3.15 | 3.70 | 0.13 |
| M18 | 75 | 4.05 | 3.55 | 4.60 | 0.19 | 0.12 | | 83 | 4.05 | 3.75 | 4.50 | 0.17 |
| M19 | 76 | 10.47 | 9.55 | 11.30 | 0.33 | -2.27 | * | 84 | 10.58 | 9.90 | 11.20 | 0.26 |
| M20 | 77 | 1.11 | 0.85 | 1.30 | 0.09 | -5.60 | *** | 85 | 1.18 | 0.90 | 1.35 | 0.07 |
| M21 | 77 | 5.11 | 4.50 | 5.75 | 0.28 | -4.77 | *** | 84 | 5.30 | 4.50 | 5.85 | 0.23 |
| M22 | 77 | 4.31 | 3.70 | 4.95 | 0.27 | -5.15 | *** | 85 | 4.51 | 3.90 | 5.00 | 0.21 |
| M23 | 77 | 6.34 | 5.40 | 7.05 | 0.42 | -5.19 | *** | 84 | 6.64 | 5.55 | 7.50 | 0.31 |




Figure 3. — Drawings of the dorsal view of two *Heimyscus fumosus*, illustrating the variability in skull morphology relative to age. Left to right: GA3119 (age-class 1) and GA2542 (age-class 4).

tional morphological variations of *H. fumosus* in Monts Doudou could provide a baseline for an incoming systematic revision of the genus *Heimyscus*.

ACKNOWLEDGEMENTS

We wish to express our gratitude to the curators of the mammal collections of a number of museums who allowed us to study the type-specimen and collections in their care; particularly C. Denys and M. Tranier. Our gratitude goes to the numerous colleagues and local collaborators who assisted us in our collecting effort (P. Barrière, O. Perpète, D. Cornélis, S. Guimondou, J.N. Assoume Ondo, A. Essono Bigang,....), and the local authorities who made our field work possible. Field studies were supported by (1) EU-DGVIII-Ecofac program “Conservation et Utilisation Rationnelle des Ecosystèmes Forestiers en Afrique centrale” (managed by AGRECO, GEIE, BDPA, SCETAGRI, SECA, CIRAD-FORET); (2) PRGIE-WWF Gabon and (3) EU-DGVIII-Biofac program “Origine et maintien de la biodiversité en Afrique Centrale” (University of Rennes 1 / CNRS, UMR 6552). V. Nicolas received a fellowship from the Ministère de l’Éducation Nationale, de l’Enseignement Supérieur et de la Recherche (France). We are grateful to A. Gautier-Hion who provided numerous helpful comments on this manuscript.

REFERENCES

- ADAM, F. (1977). — Données préliminaires sur l'habitat et la stratification des rongeurs en forêt de Basse Côte d'Ivoire. *Mammalia*, 41 : 283-290.
- BARRIÈRE, P. & NICOLAS, V. (2000). — *Écologie et structuration des peuplements de micro-mammifères musaraignes et rongeurs*. Rapport ECOFAC.
- BROSSET, A., DUBOST, G. & HEIM DE BALSAC, H. (1965). — Mammifères inédits récoltés au Gabon. *Biol. Gab.*, 1 : 148-174.
- COLLINET, J., COUTURIER, G., GUILLAUMET, J.L., KAHN, F., MOREAU, R. & SANGARÉ, Y. (1984). — Le système cultural et ses contraintes. Pp.113-184, in : J.L. Guillaumet, G. Couturier & H. Dosso (eds). *Recherche et aménagement en milieu forestier tropical humide : le projet Taï de Côte d'Ivoire*. UNESCO.
- COLYN, M., CORNÉLIS, D. & PERPÈTE, O. (1996). — *Synthèse "Micro-mammifères" Muridae et Soricidae ; structure des peuplements : richesse et diversité spécifiques et indices d'abondance*. Programme ECOFAC.
- DELANY, M.J. (1971). — The biology of small rodents in Mayanja Forest, Uganda. *J. Zool. Lond.*, 165: 85-129.
- DOSSO, H. (1983). — *Étude des rongeurs de forêts hygrophiles conservées et de zones anthropisées de la Côte d'Ivoire méridionale*. Thèse, Université d'Abidjan, Côte d'Ivoire.
- DUDU, A. (1991). — *Étude du peuplement d'insectivores et de rongeurs de la forêt ombrophile de basse altitude du Zaïre (Kisangani, Masako)*. Thèse, Université d'Anvers, Belgique.
- DUPLANTIER, J.M. (1982). — *Les rongeurs myomorphes forestiers du nord-est du Gabon : peuplements, utilisation de l'espace et des ressources alimentaires, rôle dans la dispersion et la germination des graines*. Thèse, Université des Sciences et techniques du Languedoc, France.
- GAUTUN, J.C., SANKHON, I. & TRANIER, M. (1986). — Nouvelle contribution à la connaissance des rongeurs du massif guinéen des monts Nimba (Afrique occidentale). Systématique et aperçu quantitatif. *Mammalia*, 50 : 205-217.
- GRANJON, L. (1991). — Les rongeurs myomorphes du bassin du Kouilou (Congo). Pp. 265-278, in : R.J. Dowsett & F. Dowsett-Lemaire (eds). *Flore et faune du bassin du Kouilou, Congo et leur exploration*. Tauraco Research Report n°4, Tauraco Press, Belgique.
- GUBISTA, K.R. (1999). — Small mammals of the Ituri Forest, Zaire: diversity and abundance in ecologically distinct habitats. *J. Mammal.*, 80: 252-262.
- HAPPOLD, D.C.D. (1977). — A population study on small rodents in the tropical rain forest on Nigeria. *Rev. Ecol. (Terre et Vie)*, 31: 385-455.
- HAPPOLD, D.C.D. & HAPPOLD, M. (1989). — Biogeography of montane small mammals in Malawi, Central Africa. *J. Biogeogr.*, 16: 353-367.
- HUTTERER, R., DIETERLEN, F. & NIKOLAUS, G. (1992). — Small mammals from forest islands of eastern Nigeria and adjacent Cameroon, with systematical and biogeographical notes. *Bonn. Zool. Beitr.*, 43: 393-414.
- JEFFREY, S.M. (1977). — Rodent ecology and land use in western Ghana. *J. Appl. Ecol.*, 14: 741-755.
- LASSO, C. (1995). — *Biodiversidad animal des parque nacional de Monte ALEN*. Proyecto ECOFAC, AGRECO-CTFT.
- LEIRS, H., MILLS, J.N., KREBS, J.W., CHILDS, J.E., AKAIBE, D., WOOLLEN, N., LUDWIG, G., PETERS, C. & KSIAZK, T.G. (1999). — Search for the ebola virus reservoir in Kikwit, Democratic Republic of the Congo: reflections on a vertebrate collection. *The Journal of Infectious Diseases*, 179: S155-S163.
- MEESTER, J. & SETZER, H.W. (1971). — *The mammals of Africa, an identification manual*. Smithsonian Institution Press, Washington.
- MISONNE, X. (1969). — African and Indo-Australian Muridae, evolutionary trends. *Annales du Musée Royal de l'Afrique centrale, Tervuren Belgique, série IN-8, Sciences Zoologiques*, 172: 125.
- MUSSER, G. & CARLETON, D. (1993). — Family Muridae, in: D. Wilson, & D.A Reeder (eds). *Mammal species of the world: a taxonomic and geographic reference*. Smithsonian Institution Press, Washington and London.
- NICOLAS, V. & BARRIÈRE, P. (2001). — *Biodiversité, biogéographie et écologie de la conservation des Monts Doudou : modèles fauniques et dynamique évolutive*. Programme WWF *Diversité biologique du refuge forestier pléistocène des Monts Doudou (Gabon)*.

- PETTER, F. & GENEST, H. (1970). — Liste préliminaire des rongeurs myomorphes de République Centrafricaine. Description de deux souris nouvelles : *Mus oubanguii* et *Mus goundae*. *Mammalia*, 34 : 451-458.
- ROBBINS, L., CHOATE, J.R. & ROBBINS R. (1980). — Nongeographic and interspecific variation in four species of *Hylomyscus* (Rodentia: Muridae) in southern Cameroon. *Annals of Carnegie Museum*, 49: 31-48.
- STRUHSAKER T. (1997). — *Ecology of an African rain forest: logging in Kibale and the conflict between conservation and exploitation*. University Press of Florida, Gainesville.
- VERHEYEN, W. & BRACKE, E. (1966). — The influence of aging on the craniometrical characters of *Pracomys jacksoni* (De Winton, 1897). *Ann. Mus. Roy. Afr. Centr.*, 144: 91-110.
- VERHEYEN, W., COLYN, M. & HULSELMANS, J. (1996). — Re-evaluation of the *Lophuromys nudicaudus* Heller, 1911 species-complex with a description of a new species from Zaire (Muridae – Rodentia). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, 66: 241-273.
- VERSCHUREN, J. (1986). — Perspectives pour la conservation des habitats, de la grande faune et des parcs nationaux en Afrique. *Rev. Zool. Afr.*, 100 : 73-83.