



HAL
open science

Temperature-Induced Neutral-Ionic Phase Transition in the (EDT-TTF-I₂)₂TCNQF Mixed-Stack Charge-Transfer Salt

A. Frackowiak, I. Olejniczak, R. Świetlik, O. Jeannin, M. Fourmigué

► **To cite this version:**

A. Frackowiak, I. Olejniczak, R. Świetlik, O. Jeannin, M. Fourmigué. Temperature-Induced Neutral-Ionic Phase Transition in the (EDT-TTF-I₂)₂TCNQF Mixed-Stack Charge-Transfer Salt. *Journal of Physical Chemistry C*, 2016, 120 (41), pp.23740–23747. 10.1021/acs.jpcc.6b07642 . hal-01398047

HAL Id: hal-01398047

<https://univ-rennes.hal.science/hal-01398047v1>

Submitted on 5 Jan 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Temperature Induced Neutral-Ionic Phase Transition in the (EDT-TTF-I₂)₂TCNQF Mixed-Stack Charge-Transfer Salt

Arkadiusz Frąckowiak,[†] Iwona Olejniczak,[†] Roman Świetlik,^{*,†} Olivier
Jeannin,[‡] and Marc Fourmigué[‡]

[†]*Institute of Molecular Physics, Polish Academy of Sciences, Smoluchowskiego 17, 60-179
Poznań, Poland*

[‡]*Institut des Sciences Chimiques de Rennes, Université Rennes 1, CNRS UMR 6226, Campus de
Beaulieu, 35042 Rennes, France*

E-mail: roman.swietlik@ifmpan.poznan.pl

Phone: +48 61 8695 165. Fax: +48 61 8684 524

Abstract

We report the infrared (IR) and Raman study of the 2:1 mixed-stack charge transfer salt (EDT-TTF-I₂)₂TCNQF, where EDT-TTF = ethylenedithiotetrathiafulvalene and TCNQ = tetracyanoquinodimethane, which undergoes a temperature induced neutral-ionic phase transition. Polarized infrared (IR) and Raman spectra of single crystals were measured in the 8-293 K temperature range. Temperature variations of vibrational modes that are assigned to both donors and acceptors show that the average degree of charge-transfer is growing continuously from about zero at room-temperature to about $1e$ at $T = 8$ K, nevertheless at about 100 K a regime change is observed. The coexistence of molecular species of different ionicity is detected in the whole temperature range. The IR vibrational features due to electron-molecular vibrational coupling increase their intensities considerably upon cooling giving an evidence of relatively small gradual distortions of crystal structure. Existence of ferroelectric domains is suggested. A scheme of the neutral-ionic phase transition in a 2:1 complex is proposed and discussed.

Introduction

Low dimensional organic charge-transfer (CT) salts provide a variety of structural and physical properties, which originate from strong electron correlations, intrinsic instabilities and an interplay between charge/spin and lattice degrees of freedom¹⁻³. In particular, quasi-one dimensional mixed-stack CT crystals composed of π -electron donor (D) and acceptor (A) molecules alternating along a stack (...DADA...) attract much attention due to a possible neutral-ionic phase transition (NIT)^{4,5}. Temperature- or pressure-induced increase of molecular ionicity (ρ) in these materials is the result of the competition between effective ionization energy of a DA pair and the long range electrostatic energy gained by the ionized lattice (Madelung energy). Passage from quasi-neutral ($\rho < 0.5e$) to quasi-ionic state ($\rho > 0.5e$) manifests itself by a change of CT degree and a dimerization distortion (...D^{+ ρ} A^{- ρ} D^{+ ρ} A^{- ρ} D^{+ ρ} A^{- ρ} ...) along the stacking axis. Therefore, the NIT transition can be presented as a complex interplay between the average site ionicity ρ and the stack dimerization δ . The ionicity may display a discontinuous change at critical temperature (T_c). Such a jump of ρ is favored by inter-site energy (V) and electron-molecular vibration (EMV) coupling which modulates the on-site energy (U). On the other hand, hopping integral (t) along the stack favors continuous ionicity change. The stack dimerization observed in the wide temperature range is then a result of coupling of electrons to lattice phonons. The NIT is associated to many intriguing phenomena like a dramatic increase in electrical conductivity and dielectric constant⁵. Among the methods used for investigations of the NIT, vibrational spectroscopy is one of the most reliable because it allows one to directly determine the charge on the donor and acceptor molecules based on its stretching modes^{6,7}. The spectroscopy is also sensitive to the symmetry breaking associated with the stack distortions, that yields the intensity increasing of totally symmetric modes due to the EMV-coupling^{8,9}.

Tetrathiafulvalene-*p*-chloranil (TTF-QCl₄) is the prototypical material which exhibits the neutral-ionic phase transition¹⁰. There is a number of known NIT crystals with 1:1 stoichiometry^{5,11}. Some of them are variants of TTF-QCl₄ like dimethyltetrathiafulvalene-chloranil (DMTTF-QCl₄),¹² DMTTF-QBr_{*x*}Cl_{4-*x*}¹³, and TTF_{1-*x*}TSF_{*x*}QCl₄¹⁴ where TSF = tetraselenofulvalene. Tetramethylbenzidine-

TCNQ^{15,16}, is another example of material with the discontinuous ionicity change. On the contrary to the TTF-QCl₄ salt with a first order NIT transition¹⁷, the DMTTF-QCl₄ salt exhibits a second order phase transition with continuous change of ρ upon cooling¹⁸. A continuous ionicity change with T at ambient pressure has been observed in 2-chloro-5-methyl-p-phenylenediamine-2,5-dimethyl-dicyanoquinonediimine (ClMePD-DMeDCNQi)¹⁹ where the enhancement of the stack dimerization under cooling is regarded as a driving force of the phase transition²⁰.




Figure 1: Projection view along b axis of the extended unit cell of (EDT-TTF-I₂)₂TCNQF, showing the alternated ...ADDADDA... stacks. The pink dotted lines indicate the halogen bond interaction between iodine and nitrogen atoms. The single fluorine atom is disordered on two positions related by inversion center.

Recently, for the first time the temperature-induced neutral-ionic phase transition has been reported in the (EDT-TTF-I₂)₂TCNQF mixed-stack charge-transfer complex characterized by 2:1 stoichiometry, formed by iodinated ethylenedithiotetrathiafulvalene (EDT-TTF-I₂) and tetracyanoquinodimethane derivative (TCNQF)²¹. The complex crystallizes in the triclinic system, space group $P\bar{1}$ with two TCNQF molecules on inversion centers and four EDT-TTF-I₂ molecules in general positions in the unit cell. Dyads of donor molecules in the "head-tail" configuration alternate with acceptor molecules along a - b axis. The resulting stacks (...ADDADDA...) interact with each other to form layers (Fig. 1). On the other hand, EDT-TTF-I₂ molecules link with TCNQF molecules along the c axis, by highly directional C-I...N≡C halogen bonds. Infrared spectra of (EDT-TTF-I₂)₂TCNQF in the C≡N stretching vibration range of TCNQF molecule provided an evidence that the salt undergoes NIT transition with lowering temperature²¹. Based on the X-ray data, it has

1
2
3
4 been shown that the molecular ionicity increasing in $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ does not induce a
5 structural transition such as the loss of an inversion center, but is associated with a halogen bond
6 strengthening and unit-cell contraction.
7

8
9 Here, we present a detailed spectral analysis of temperature induced neutral-ionic phase tran-
10 sition in the $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ salt of 2:1 stoichiometry. In the first section of Results and
11 Discussion, we compare the reflectance and absorbance spectra of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ in the
12 broad frequency range at 8 and 293 K. In the second section, we concentrate on the origin of the
13 temperature induced neutral-ionic phase transition in $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$. The totally symmet-
14 ric donor and acceptor modes, which are activated in the infrared spectra, show that molecular
15 charge enhancement is associated with structure distortion. In the last section, we estimate the ion-
16 icity ρ based on the charge-sensitive stretching vibrations of the EDT-TTF-I₂ molecule. We show
17 that ρ in $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ gradually increases upon cooling, with remarkable enhancement
18 below about 100 K. Our overall goal is to clarify the unknown aspects of the neutral-ionic phase
19 transition in this mixed-stack material of 2:1 stoichiometry.
20
21
22
23
24
25
26
27
28
29
30
31

32 33 34 35 **Experimental**

36
37 Single crystals of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ were synthesized and crystallized according to a pro-
38 cedure described previously²¹. Typical dimensions of the elongated plates used in measurements
39 were $0.5 \times 0.2 \times 0.02 \text{ mm}^3$. The optical axes of the sample were determined as those displaying the
40 largest reflectance anisotropy. The infrared spectra were measured along the *a-b* and *a+b* axes, that
41 are nearly parallel and nearly perpendicular to the stacking direction, respectively. The polarized
42 reflectance (from 600 to 12000 cm^{-1}) and transmittance (from 600 to 4000 cm^{-1}) spectra were
43 recorded with a resolution of 4 cm^{-1} using a Bruker Equinox 55 FT-IR spectrometer equipped
44 with a Hyperion 1000 infrared microscope and a set of suitable polarizers. The absolute values of
45 reflectance were obtained using an aluminum mirror. The optical conductivity was calculated us-
46 ing Kramers-Krönig transformation of the reflectance data²². In the low frequency range, the data
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 were extrapolated assuming constant value appropriate for insulators, and in the high frequency
5 range, the data were extrapolated assuming a ω^{-2} behavior (up to 10^6 cm^{-1}) and a ω^{-4} behavior
6 (for higher frequencies). Raman spectra for the electric field vector of the laser beam polarized
7 parallel to $a+b$ axis were measured from $100 - 2300 \text{ cm}^{-1}$ in a backscattering geometry with a
8 Raman LABRAM HR800 spectrometer equipped with a microscope. Laser line of the 632.8 nm
9 (He-Ne) was used with power reduced below 0.1 mW to avoid sample overheating. The spectra
10 were recorded with a spectral resolution of 2 cm^{-1} .
11

12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Variable temperature IR and Raman spectra were measured with the help of a continuous-flow
cold-finger cryostat manufactured by the Oxford Instruments. A good thermal contact between
cold-finger and the sample was fulfilled using vacuum grease. The spectra were collected at sev-
eral temperatures from 296 to 8 K. The rate of cooling/heating was 1 K/min. The complex vibra-
tional features in the optical conductivity and transmittance spectra were decomposed by standard
peak fitting techniques that allow to extract the center frequency and the integral intensity (oscil-
lator strength). The oscillators were fitted using the Voigt spectroscopy function in the PeakFit
program²³.

61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161

Results and discussion

Infrared spectra at 8 and 293 K

Figure 2 shows room-temperature and 8 K infrared spectra of single crystals of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ polarized along two mutually perpendicular directions in the frequency range $700 - 9000 \text{ cm}^{-1}$.

The optical response of the material is characterized by a significant reflectance anisotropy. In the




Figure 2: (a) Infrared reflectance spectra of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ at 293 K and 8 K polarized parallel to the stack; inset displays a schematic view of a single stack; (b) polarized optical conductivity spectra derived from the reflectance by the Kramers-Krönig transformation; ν_{CT} mark the position of the mid-infrared charge-transfer band; (c) infrared absorption spectra of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ at 293 K and 8 K polarized perpendicular to the stack (note the logarithmic frequency scale).

1
2
3 direction of the highest reflectance value, that is nearly parallel to a - b axis (stacking direction), we
4 show the reflectance spectra (Fig. 2a), and in the $a+b$ direction, we show the transmittance spectra
5 (Fig. 2c).
6
7

8
9 The room-temperature reflectance spectrum polarized $E||a$ - b displays semiconducting behav-
10 ior, with the reflectance level of about 0.25 at low frequency and the reflectance edge at about 6500
11 cm^{-1} (Fig. 2a). In the 700 - 2300 cm^{-1} frequency range we observe rather weak and broad vi-
12 brational modes of EDT-TTF-I₂ and TCNQF molecules, activated due to the EMV-coupling²⁷⁻²⁹.
13
14 Between 293 K and 8 K, the overall reflectance level increases slightly in the whole frequency
15 range, and the reflectance edge is shifted about 1500 cm^{-1} toward higher frequencies. At the
16 same time, we see a significant intensity enhancement of vibrational modes. The optical conduc-
17 tivity spectra of (EDT-TTF-I₂)₂TCNQF, obtained from the $E||a$ - b reflectance spectra through the
18 Kramers-Krönig transformation are shown in Fig. 2b. The room-temperature spectrum polarized in
19 the stacking direction is dominated by the broad mid-infrared electronic band centered at around
20 3000 cm^{-1} that is attributed to a superposition of two charge transfer excitations, between the
21 dyads of EDT-TTF-I₂ and TCNQF molecules and between the EDT-TTF-I₂ molecules themselves
22 (Fig. 2b). Frequency of the maximum of the electronic CT band along stacking direction does
23 not change much between 293 K and 8 K (Fig. 2b); a relatively small intensity increase and band
24 broadening is found. On the other hand, in the spectra polarized perpendicularly to the stacking
25 direction ($E||a+b$), the electronic band attributed to a side-by-side CT between EDT-TTF-I₂ and
26 TCNQF is partially observed between 3000 and 4000 cm^{-1} (Fig. 2c). The measured crystals were
27 not thin enough to perform transmittance investigations in higher frequency range.
28
29

30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Vibrational features observed both in the optical conductivity ($E||a$ - b) and the transmittance
spectrum ($E||a+b$) of (EDT-TTF-I₂)₂TCNQF in the range between 700 and 2300 cm^{-1} are re-
lated to both TCNQF and EDT-TTF-I₂ molecules. In the optical conductivity spectrum in the
stacking direction, vibrational bands are rather broad and weak at room-temperature. These bands
are mostly the consequence of the EMV-coupling. Between 293 K and 8 K, the most prominent
EMV-coupled modes of EDT-TTF-I₂ and TCNQF molecules dramatically increase their intensity

1
2
3
4 and shift toward lower frequencies, as expected for a compound undergoing the neutral-ionic tran-
5 sition. On the other hand, vibrational bands in the absorbance spectrum in the interstack $E||a+b$
6 direction are relatively narrow. These are mostly IR active modes of both TCNQF and EDT-TTF-I₂
7 but bands related to coupling with side-by-side CT transition are also possible. Temperature evolu-
8 tion of vibrational modes in (EDT-TTF-I₂)₂TCNQF, that is related with the changes of molecular
9 ionicity and stack distortion, will be discussed in detail in the following sections.

17 **Neutral-ionic phase transition**

20 The structure of the mixed-stack CT materials is intrinsically unstable towards lattice distortion
21 upon approaching an ionic phase with lowering temperature^{2,3}. Such structural distortions are
22 usually evidenced by intensity enhancement of the totally symmetric molecular vibrations, that are
23 activated in the infrared spectra due to EMV-coupling^{8,9}. Here, we focus on the C≡N stretching
24 TCNQF modes in the spectral range 2100-2250 cm⁻¹, where no EDT-TTF-I₂ vibrational modes
25 are present. It is known, that the C≡N stretching modes are charge-sensitive, so they are useful
26 to molecular charge estimation³⁰⁻³². As shown in our previous paper²¹, investigations of these
27 modes give an unambiguous evidence of the neutral-ionic phase transition. However, it should be
28 also taken into account, that the C≡N stretching modes are sensitive to molecular environment like
29 electrostatic potential and hydrogen (halogen) bonding^{20,33}. This is particularly important from the
30 point of view of relatively strong halogen bond interaction in (EDT-TTF-I₂)₂TCNQF²¹.

31
32
33
34
35
36
37
38
39
40
41
42
43 Figure 3a shows the optical conductivity spectra of (EDT-TTF-I₂)₂TCNQF in the C≡N stretch-
44 ing region, polarized along stacking direction ($E||a-b$) at selected temperatures. The relatively
45 weak and broad mode observed at about 2190 cm⁻¹ at room-temperature is the counterpart of to-
46 tally symmetric TCNQ mode $\nu_2(a_g)$, that is activated in the infrared spectra due to EMV-coupling.
47 Schematic view of the mode, that we denote here as ν_g^A , is shown as the inset in Fig. 3b. During
48 cooling from room-temperature down to about 100 K, maximum of the ν_g^A band centered at 2192
49 cm⁻¹ at 293 K shifts continuously toward lower frequencies of about 25 cm⁻¹ (Fig. 4a); below
50 about 100 K, the band position changes slightly reaching 2166 cm⁻¹ at 8 K. Softening of the ν_g^A
51
52
53
54
55
56
57
58
59
60




Figure 3: (a) IR optical conductivity and (b) absorbance spectra of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ in the frequency range of the stretching $\text{C}\equiv\text{N}$ vibration at selected temperature (adopted from Ref.²¹).




Figure 4: (a) Temperature evolution of the frequency of the TCNQF ν_g^A band observed in the optical conductivity spectra; (b) temperature dependence of the ν_g^A intensity normalized to the low-temperature value; (c) temperature evolution of the normalized intensity of the 2175 and 2185 cm^{-1} components of the $\text{C}\equiv\text{N}$ spectral features observed in the absorbance spectra.

mode is mostly caused by the ionicity variation and the EMV-coupling. At the same time, we observe significant intensity enhancement of the ν_g^A band. As shown in Fig. 4b, intensity of ν_g^A (normalized to the 8 K value) increases gradually with lowering temperature. Such a temperature dependence of the mode intensity in $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$, suggests a continuous increase of structure distortion in the a - b direction due to the neutral-ionic phase transition.

Figure 3b shows the absorbance spectra of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ in the $\text{C}\equiv\text{N}$ stretching range, polarized perpendicular to the stacking direction ($E||a+b$) at selected temperatures. In this frequency range, we observe a broad multi-component spectral feature, with four strongest components at about 2175, 2185, 2207 and 2223 cm^{-1} at room-temperature. Two bands observed at 2207 and 2223 cm^{-1} are assigned to two components of normally infrared active $\text{C}\equiv\text{N}$ mode of TCNQF, that is counterpart of the unsymmetrical TCNQ $\nu_{19}(b_{1u})$ mode. Schematic view of the TCNQF mode, that we denote here as ν_u^A , is shown as the inset in Fig. 3b. The higher-frequency feature at 2223 cm^{-1} is assigned to ν_u^A mode of neutral TCNQF molecules. The band at 2207 cm^{-1} is related to ν_u^A which corresponds to small number of ionized TCNQF^- molecules that can exist in the crystal at room-temperature. When temperature decreases the intensity of the TCNQF^- band at 2207 cm^{-1} strongly grows giving thus a clear evidence of the neutral-ionic transition²¹. With decreasing temperature, two other bands grow at around 2175 and 2185 cm^{-1} , which can be related to the ν_g^A mode activated due to EMV-coupling. Fig. 4c shows temperature depen-

1
2
3
4 dence of the normalized integral intensity of the 2175 and 2185 cm^{-1} components of the broad
5 $\text{C}\equiv\text{N}$ stretching band in the absorbance spectra. As one can notice, we observe similar behavior
6 as shown in Fig. 4b for the $\nu_{\text{g}}^{\text{A}}$ mode in $\text{E}\parallel a-b$ polarization. Therefore, the intensity enhancement
7 of the $\nu_{\text{g}}^{\text{A}}$ components of the $\text{C}\equiv\text{N}$ band in the spectra polarized perpendicular to the stacking di-
8 rection is also related with a structure distortion that appears parallel to stacking direction ($\text{E}\parallel a-b$).
9 In this case the $\nu_{\text{g}}^{\text{A}}$ mode interacts most probably with the side-by-side CT, therefore temperature
10 dependence of the $\nu_{\text{g}}^{\text{A}}$ suggests that it exists a relatively small distortion in the direction perpendic-
11 ular to the stacks but it is significantly smaller than in the $a-b$ direction. In fact, CT interaction in
12 the mixed-stack materials is highly sensitive to a subtle structural change³⁰. The doublet structure
13 of the band can be interpreted as an indication that in the crystal exist molecules with different
14 charges. The two components at 2175 and 2185 cm^{-1} can be related to TCNQF^{-} and TCNQF^0
15 molecules, respectively.
16
17

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Figures 5a, 5b and 6 show optical conductivity ($\text{E}\parallel a-b$), absorbance ($\text{E}\parallel a+b$) and Raman
($\text{E}\parallel a+b$) spectra of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ in the frequency range 1000-1700 cm^{-1} , where both
donor and acceptor bands are seen. Here, we analyze the stretching vibrations of TCNQF that are
activated in the infrared spectra due to EMV-coupling. We assign the band at about 1170 cm^{-1}
in the optical conductivity spectrum at room-temperature (Fig. 5a) to C–F and C–C stretching
vibration of TCNQF, as shown in the inset in Fig. 7b. Here, we denote this mode as $\nu_{\text{a}}^{\text{A}}$. For this
polarization the 1170 cm^{-1} band has a clear doublet structure in the whole temperature range but
the intensity of lower-frequency component grows stronger on cooling down. Analogously, the
 $\nu_{\text{a}}^{\text{A}}$ band has also a doublet structure in the absorption spectra polarized in the direction $\text{E}\parallel a+b$,
centered at 1177 and 1204 cm^{-1} (Fig. 5b) and in the Raman spectrum at 1177 and 1194 cm^{-1} at
room-temperature (Fig. 6). Similar behavior to the $\nu_{\text{a}}^{\text{A}}$ band is also observed in the case of another
TCNQF mode at 1605 cm^{-1} in the optical conductivity spectrum ($\text{E}\parallel a-b$), which we assign to the
totally symmetric C=C stretching vibration of the TCNQF ring, as shown in the inset in Fig. 7d.
Here, we denote this band as $\nu_{\text{b}}^{\text{A}}$. In the spectra polarized perpendicularly to the stack, the $\nu_{\text{b}}^{\text{A}}$
features are centered at 1607 and 1614 cm^{-1} in the absorbance spectrum (Fig. 5b) and at 1608




Figure 5: (a) IR optical conductivity, (b) absorbance spectra of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ in the 1000-1700 cm^{-1} frequency range at selected temperatures.

and 1612 cm^{-1} in the Raman spectrum (Fig. 6). We suggest that the doublet structure of the TCNQF bands in $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ is due to coexistence of two species of TCNQF molecules of different ionicity.




Figure 6: Raman spectra of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ in the $1000\text{-}1700\text{ cm}^{-1}$ frequency range at selected temperatures.

As results from our DFT calculations³⁴, when TCNQF molecule is ionized the frequencies of the ν_a^A and ν_b^A modes shift towards lower frequencies by 16 and 21 cm^{-1} , respectively. Upon cooling, maximum of the EMV-coupled ν_a^A mode shifts gradually from 1171 cm^{-1} at 293 K to 1107 cm^{-1} at 8 K (Fig. 7a) due to molecular ionicity variation and lattice distortion. At the same time, the ν_a^A band increases intensity (Fig. 7b), what is an indication of the structure distortion in $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$. On the other hand, the ν_a^A band in the Raman spectrum exhibits moderate temperature changes down to about 100 K and then it shifts from 1176 cm^{-1} at 100 K to 1155 cm^{-1} at 8 K (Fig. 7a). The red-shift of ν_a^A in the Raman spectra is associated mostly with the ionicity change of TCNQF molecules. Frequency of the ν_b^A mode is also sensitive to ionicity variation in $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$. The ν_b^A band shifts from 1607 cm^{-1} at 293 K to 1598 cm^{-1} at 8 K in the optical conductivity spectra ($E\parallel a-b$). Only the 1607 cm^{-1} component of ν_b^A is temperature dependent in the absorbance spectra ($E\parallel a+b$). Upon cooling, it shifts 6 cm^{-1} toward lower frequencies (Fig. 7c). At the same time, the ν_b^A band increases the intensity with lowering temperature (Fig. 7d). The temperature dependence of a band intensity that is similar to ν_a^A and

ν_b^A bands is also observed for C–S stretching band of EDT-TTF-I₂ at around 1000 cm⁻¹ (see Fig. 9c).

In the whole temperature range we observe bands related to TCNQF⁻ ions and to TCNQF⁰ neutral molecules but on temperature decreasing the TCNQF⁻ bands considerably grow as expected for neutral-ionic transition. Our analysis of the C≡N, C=C and C–S stretching modes shows that the transition in (EDT-TTF-I₂)₂TCNQF is related with the structure distortions both in the direction parallel (E||*a-b*) and perpendicular (E||*a+b*) to the stacks. The ionicity and lattice distortion change continuously with temperature decreasing, however these changes become stronger below about 100 K, i. e. there is an important regime change at this temperature. In case of the EDT-TTF-I₂ modes, δ_5^D is related to the bending vibration of CH₂ groups of EDT-TTF-I₂ molecules. Temperature dependence of the frequency and intensity of δ_5^D and other EDT-TTF-I₂ modes, like ν_3^D , ν_2^D and ν_{27}^D (corresponding to stretching C=C vibrations) are similar to the temperature behavior of TCNQF modes (Figs. 5a, 5b, 6).




Figure 7: (a) Temperature evolution of the frequency of the TCNQF ν_a^A band observed in the optical conductivity spectra (■) and in the Raman spectra (□); (b) temperature dependence of the intensity (normalized to 8 K value) of the TCNQF ν_a^A band observed in the optical conductivity spectra; (c) temperature evolution of the frequencies of the two components of the TCNQF ν_b^A band observed in the absorbance spectra; (d) temperature dependence of the intensity (normalized to 8 K value) of the TCNQF ν_b^A doublet band observed in the absorbance spectra.

Figure 8 shows Raman spectra of (EDT-TTF-I₂)₂TCNQF polarized in E||*a+b* in the frequency




Figure 8: Raman spectra of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ in the range of the deformation vibrations of the donor and acceptor molecules at selected temperatures.

range $250\text{-}350\text{ cm}^{-1}$ at selected temperatures. We assign the bands at 268 and 290 cm^{-1} at room-temperature to deformation vibrations of EDT-TTF- I_2 rings⁶. The band at 320 cm^{-1} , that can be related to deformation modes of the TCNQF or EDT-TTF- I_2 splits in two components at 309 and 326 cm^{-1} with lowering the temperature. Significant changes of vibrational features in the low frequency Raman spectra of low-dimensional CT salts are usually related to structural transition. Thus, we suggest that the band splitting at around 100 K in $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ is due to structure distortion.

Estimation of the charge on EDT-TTF- I_2

The most important parameter to discuss the neutral-ionic phase transition is the ionicity ρ of donor and acceptor molecules in the whole temperature range. In Ref.[²¹], the actual degree of charge transfer ρ was tentatively estimated from the evolution of the intramolecular bond lengths within the TCNQF molecule to exceed $0.5e$. The evolution of the $\text{N}\cdots\text{I}$ halogen bond distances with temperature indicate a larger charge transfer, closer to $0.7\text{-}0.8e$. A more quantitative analysis based on the evolution of the $\text{C}\equiv\text{N}$ stretching frequencies was not attempted at that time since, as mentioned above, these modes can be perturbed by the molecular environment and particularly

by the strong halogen bonding interaction taking place with the iodine atom of the EDT-TTF-I₂ molecules. In order to circumvent this problem, we have focused our attention on the charge localized on the (EDT-TTF-I₂)₂ dimer, which should be less sensitive to the influence of environment. Figure 9a shows the infrared spectra of (EDT-TTF-I₂)₂TCNQF, polarized perpendicular to the stacks (E||a+b) in the spectral region of C–S, C–I vibrational modes of EDT-TTF-I₂. These modes are well separated from TCNQF vibrational modes. Based on the well-known behavior of the tetrathiafulvalene derivatives^{35,36}, we assume that an ionization of EDT-TTF-I₂ molecule leads to contraction of C–S bonds length, and therefore hardening of the corresponding charge-sensitive modes. Here, we assign the single peak observed at 914 cm⁻¹ at room-temperature (Fig. 9a) as the IR active C–S, C–I stretching mode of EDT-TTF-I₂, marked as $\nu_{C-S,C-I}^D$.




Figure 9: (a) IR absorption spectra of (EDT-TTF-I₂)₂TCNQF in the C–S stretching frequency range at selected temperatures; (b) temperature evolution of the frequency of the EDT-TTF-I₂ stretching $\nu_{C-S,C-I}^D$ mode; (c) the intensity change of the two components at 999 cm⁻¹ (□) and 1011 cm⁻¹ (■) of the EDT-TTF-I₂ stretching ν_{C-S}^D mode, that is related to the stack dimerization.

We compare the frequency of the $\nu_{C-S,C-I}^D$ mode in the absorbance spectra of (EDT-TTF-I₂)₂TCNQF polarized perpendicularly to the stack (E||a+b) with frequency of $\nu_{C-S,C-I}^D$ in other two isostructural complexes: (EDT-TTF-I₂)₂TCNQ and (EDT-TTF-I₂)₂TCNQF₂. As results from our previous studies there is almost no charge-transfer in the (EDT-TTF-I₂)₂TCNQ salt, while the (EDT-TTF-I₂)₂TCNQF₂ salt is fully ionized, i. e. in these salts the (EDT-TTF-I₂)₂ dimers possess charges

0 and $1e$, respectively²¹. It is to be emphasized that both salts are isostructural with $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$. In the absorbance spectra ($E\parallel a+b$) of the $(\text{EDT-TTF-I}_2)_2\text{TCNQ}$ neutral complex the $\nu_{\text{C-S,C-I}}^{\text{D}}$ mode is observed at 914 cm^{-1} at 293 K (See the Supporting Information, Figure S1, Table S1). In fact, for the neutral EDT-TTF-I₂ molecule⁶, the experimental frequency of the mode is 917 cm^{-1} . On the other hand, in the absorbance spectra ($E\parallel a+b$) of the $(\text{EDT-TTF-I}_2)_2\text{TCNQF}_2$ ionic complex, the $\nu_{\text{C-S,C-I}}^{\text{D}}$ band has a doublet structure and the frequencies of the mode are 919 and 929 cm^{-1} (See the Supporting Information, Figure S1, Table S1); this is an indication that the charge $1e$ is distributed non-uniformly among molecules forming $(\text{EDT-TTF-I}_2)_2$ dimers. The $\nu_{\text{C-S,C-I}}^{\text{D}}$ mode in the $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ shows a small shift towards higher frequencies upon cooling down to 110 K (Fig. 9a). Then below 110 K, the $\nu_{\text{C-S,C-I}}^{\text{D}}$ mode splits into two peaks centered at 916 and 929 cm^{-1} at 8 K. Hence, frequencies of the C–S, C–I stretching mode components of EDT-TTF-I₂ in $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ indicate a change of ionicity ρ from about zero at room-temperature to about $1e$ at 8 K. Splitting of the $\nu_{\text{C-S,C-I}}^{\text{D}}$ band at 100 K suggests the simultaneous presence of quasi-neutral and quasi-ionic molecules in the ionic phase, which means that the charge distribution among molecules in EDT-TTF-I₂ dyad components is not uniform.

Let us focus on the possible charge distribution in the $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ complex. According to our results, at room-temperature in the neutral ground state all the molecules are mostly neutral (Fig. 10a). Upon cooling the complex becomes ionic and two types of charge distributions in the 2:1 complex are possible, as shown in Figs. 10b and 10c. When the charge is uniformly distributed among molecules in the dimer (Fig. 10b), apparent inversion centers remain, but it is not the case in $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ because we do not observe donors with charge $+0.5$. On the other hand, when the charge is localized on one of the donors in the dimeric unit (Fig. 10c), inversion centers no longer exist and the lattice should be deformed. The coexistence of molecules with different charges is indeed observed in our IR spectra. Therefore, we suggest that the charge distribution in the ionic state of $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ can be represented by the model shown in the Fig. 10c. However, the structural investigations show that TCNQF molecules are located on an inversion center and no indication of any structural phase transition, such as the loss of




Figure 10: Schematic representation of the neutral-ionic phase transition (NIT) in the 2:1 material; mixed-stack in the neutral phase (a); two possibilities of charge distribution in the ionic phase (b), (c).

this center, was found down to $T = 20 \text{ K}$ ²¹. Two possibilities depicted here in Fig. 10c most probably involve the presence of oppositely polarized ferroelectric domains in the structure, and also the domain walls that can be considered as solitons. It can be assumed that the polarization of domains fluctuates between opposite directions in the time scale which is larger than that one appropriate for IR spectroscopy. In this picture the distortion inside domains disappears in the time scale of the X-ray experiment, therefore the stack distortion and thus the lost of the inversion center is not seen in the structure data. It is also important that unpaired spins in the ionic state interact antiferromagnetically giving rise to magnetic phase transitions²¹. Such a scenario has been already discussed in case of 1:1 mixed stacked materials^{37,38}. In case of $(EDT-TTF-I_2)_2TCNQF$ characterized by stoichiometry 2:1, the situation is obviously more complex taking into account both charge-transfer between dimer of donor molecules and acceptors and non-uniform charge distribution within dimers. In fact, the non-uniform distribution of charge evidenced here also explains why the structural characteristics deduced from the X-ray diffraction studies (intramolecular bond lengths and halogen bond distances) give an averaged charge transfer less than $1e$. The

1
2
3
4 non-uniform charge distribution among the EDT-TTF-I₂ molecules in the dimer is connected with
5
6 the neutral-ionic transition observed here, involving a continuous and smooth increase of charged
7
8 species with decreasing the temperature. The non-uniform charge distribution is also associated
9
10 with the strong charge localization as shown by the insulating character of the salt down to the
11
12 lowest temperatures.
13
14
15

16 Conclusion

17
18
19 In summary, using infrared and Raman spectroscopy we have investigated the neutral-ionic phase
20
21 transition in (EDT-TTF-I₂)₂TCNQF, the first mixed-stack salt of 2:1 stoichiometry showing such
22
23 transition. We estimated the molecular ionicity ρ in (EDT-TTF-I₂)₂TCNQF based on charge-
24
25 sensitive stretching vibrations of EDT-TTF-I₂ molecules; ρ changes continuously from about zero
26
27 at room-temperature to about $1e$ at 8 K but coexistence of two species of different ionicity persists
28
29 down to the lowest temperature. Our spectroscopic data indicate that at about 100 K a regime
30
31 change is found. The nominally totally symmetric TCNQF and EDT-TTF-I₂ modes, activated
32
33 in the infrared spectra due to EMV-coupling, provide an evidence that stacks ...ADDADDA...
34
35 are slightly distorted already at room-temperature. Intensity increasing of such modes indicates
36
37 that ionicity enhancement in (EDT-TTF-I₂)₂TCNQF is associated with gradual increase of the
38
39 structure distortion in the direction parallel to the stacks; in the direction perpendicular to the stacks
40
41 a subtle structure distortion is also possible. We demonstrated and discussed a scheme of the charge
42
43 distribution in the mixed-stack charge-transfer salt of a 2:1 stoichiometry that undergoes a neutral-
44
45 ionic phase transition. We suggest that ferroelectric domains exist in (EDT-TTF-I₂)₂TCNQF and
46
47 their polarization fluctuates between the opposite directions.
48
49
50
51

52 Acknowledgement

53
54
55 We thank J. Lieffrig for the preparation of the crystalline samples used in these studies. This
56
57 work was supported by the Institute of Molecular Physics PAS and the ANR (Paris, France) under
58
59
60

contract No. 08-BLAN-0091-02.

Supporting Information Available

The following files are available free of charge.

The following files are available free of charge.

- Figure S1 shows the IR absorbance spectra of $(\text{EDT-TTF-I}_2)_2\text{TCNQ}$, $(\text{EDT-TTF-I}_2)_2\text{TCNQF}$ and $(\text{EDT-TTF-I}_2)_2\text{TCNQF}_2$ in the C-S, C-I stretching frequency range with a brief description.
- Table S1 shows the frequency of the bands related to the C-S, C-I stretching vibrations at room-temperature.

References

- (1) Horiuchi, S.; Okimoto, Y.; Kumai, R.; Tokura, Y. Quantum Phase Transition in Organic Charge-Transfer Complexes. *Science* **2003**, *299*, 229–232.
- (2) Dressel, M. Ordering Phenomena in Quasi-One-Dimensional Organic Conductors. *Naturwissenschaften* **2007**, *94*, 527–541.
- (3) Tomić, S.; Dressel, M. Ferroelectricity in Molecular Solids: a Review of Electrodynamic Properties. *Rep. Prog. Phys.* **2015**, *78*, 096501–096526.
- (4) Torrance, J. B.; Vazquez, J. E.; Mayerle, J. J.; Lee, V. Y. Discovery of a Neutral-To-Ionic Phase Transition in Organic Materials. *Phys. Rev. Lett.* **1981**, *46*, 253–257.
- (5) Horiuchi, S.; Kumai, R.; Okimoto, Y.; Tokura, Y. Chemical Approach to Neutral–Ionic Valence Instability, Quantum Phase Transition, and Relaxor Ferroelectricity in Organic Charge-Transfer Complexes. *Chem. Phys.* **2006**, *325*, 78–91.

- 1
2
3
4 (6) Łapiński, A.; Ouahab, L.; Imakubo, T. Spectroscopic and Theoretical Study on Organic
5 Donors: DIET, DIEDO, and DIETS. *Vib. Spectrosc.* **2010**, *52*, 22–30.
6
7
8 (7) Murata, T.; Nakamura, K.; Yamochi, H.; Saito, G. Crystal Structure and Properties of Charge-
9 Transfer Complex of N-butylguanine and FTCNQ. *Synth. Met.* **2009**, *159*, 2375–2377.
10
11 (8) Meneghetti, M.; Pecile, C.; Yakushi, K.; Yamamoto, K.; Kanoda, K.; Hiraki, K. Study
12 of the Phase Transitions of (DI-DCNQI)₂M (M=Ag, Li, Cu) Through the Analysis of the
13 Temperature-Dependent Vibronic and Vibrational Infrared Absorptions. *J. Solid State Chem.*
14 **2002**, *168*, 632–638.
15
16 (9) Meneghetti, M.; Bozio, R.; Pecile, C. Electron-Molecular Vibration Coupling in 2-D Organic
17 Conductors: High and Low Temperature Phases of alpha-(BEDT-TTF)₂I₃. *J. Physique* **1986**,
18 *47*, 1377–1387.
19
20 (10) Horiuchi, S.; Kobayashi, K.; Kumai, R.; Ishibashi, S. Ionic Versus Electronic Ferroelectricity
21 in Donor-Acceptor Molecular Sequences. *Chem. Lett.* **2014**, *43*, 26–35.
22
23 (11) Girlando, A.; Painelli, A.; Bewick, S. A.; Soos, Z. G. Charge Fluctuations and Elec-
24 tron-Phonon Coupling in Organic Charge-Transfer Salts with Neutral-Ionic and Peierls
25 Transitions. *Synth. Met.* **2004**, *141*, 129–138.
26
27 (12) Aoki, S.; Nakayama, T.; Miura, A. Temperature-Induced Neutral-Ionic Transition in
28 dimethyltetrathiafulvalene-p-chloranil. *Phys. Rev. B* **1993**, *48*, 626–629.
29
30 (13) Horiuchi, S.; Okimoto, Y.; Kumai, R.; Tokura, Y. Ferroelectric Valence Transition and Phase
31 Diagram of a Series of Charge-Transfer Complexes of 4,4'-Dimethyltetrathiafulvalene and
32 Tetrahalo-p-benzoquinones. *J. Am. Chem. Soc.* **2001**, *123*, 665–670.
33
34 (14) Horiuchi, S.; Kumai, R.; Tokura, Y. Chemical Control of Ferroelectric Neutral-Ionic Tran-
35 sition in Charge-Transfer Complexes, TTF_{1-x}TSF_xQC1₄ [TTF = Tetrathiafulvalene; TSF =
36 Tetraselenafulvalene; QC1₄ = p-Chloranil]. *J. Am. Chem. Soc.* **1998**, *120*, 7379–7380.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 (15) Iwasa, Y.; Koda, T.; Tokura, Y.; Kobayashi, A.; Iwasawa, N.; Saito, G. Temperature-Induced
5 Neutral-Ionic Transition in tetramethylbenzidine-tetracyanoquinodimethane (TMB-TCNQ).
6 *Phys. Rev. B* **1990**, *42*, 2374–2377.
7
8
9
10 (16) Iwasa, Y.; Watanabe, N.; Koda, T.; Saito, G. Two-Step Neutral-Ionic Phase Transition in
11 Organic Charge-Transfer Compounds: Possible Staging Effect. *Phys. Rev. B* **1993**, *47*, 2920–
12 2923.
13
14
15
16 (17) Tokura, Y.; Koda, T.; Mitani, T.; Saito, G. Neutral-To-Ionic Transition in tetrathiafulvalene-
17 p-chloranil as Investigated by Optical Reflection Spectra. *Solid State Commun.* **1982**, *43*,
18 757–760.
19
20
21
22
23 (18) Ranzieri, P.; Masino, M.; Girlando, A.; Lemée-Cailleau, M. H. Temperature-Induced Valence
24 and Structural Instability in DMTTF-CA: Single-Crystal Raman and Infrared Measurements.
25 *Phys. Rev. B* **2007**, *76*, 134115.
26
27
28
29
30 (19) Aoki, S.; Nakayama, T. Temperature-Induced Neutral-Ionic Transition in 2-chloro-5-methyl-
31 p-phenylenediamine-2,5-dimethyl-dicyanoquinonediimine. *Phys. Rev. B* **1997**, *56*, R2893–
32 R2896.
33
34
35
36 (20) Masino, M.; Girlando, A.; Farina, L.; Brillante, A. A New Type of Neutral-Ionic Interface
37 in Mixed Stack Organic Charge Transfer Crystals: Temperature Induced Ionicity Change in
38 CIMePDMeDCNQI. *Phys. Chem. Chem. Phys.* **2001**, *3*, 1904–1910.
39
40
41
42
43 (21) Liefbrig, J.; Jeannin, O.; Frąckowiak, A.; Olejniczak, I.; Świetlik, R.; Dahaoui, S.;
44 Aubert, E.; Espinosa, E.; Auban-Senzier, P.; Fourmigué, M. Charge-Assisted Halogen Bond-
45 ing: Donor-Acceptor Complexes with Variable Ionicity. *Chem. Eur. J.* **2013**, *19*, 14804–
46 14813.
47
48
49
50 (22) Wooten, F. *Optical Properties of Solids*, 1st ed.; Academic Press: New York, USA, 1972.
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 (23) The Voigt spectroscopy function is a four-parameter theoretical model of the spectral line
5 including the Gaussian and Lorentzian types of broadening.
6
7
8
9 (24) Frisch, M. J. et al. Gaussian 03 (Revision B.03). Gaussian, Inc.: Pittsburgh PA, 2003.
10
11 (25) Sun, R.; Yao, J.; Li, S.; Gu, R. Raman Spectroscopic and Density Functional Theory Studies
12 on a benzothiazole-2-thione Derivative. *Vib. Spectrosc.* **2008**, *47*, 38–43.
13
14
15
16 (26) Scott, A. P.; Random, L. Harmonic Vibrational Frequencies: An Evaluation of Hartree-
17 Fock, Møller-Plesset, Quadratic Configuration Interaction, Density Functional Theory, and
18 Semiempirical Scale Factors. *J. Phys. Chem.* **1996**, *100*, 16502–16513.
19
20
21
22 (27) Rice, M. J. Organic Linear Conductors as Systems for the Study of Electron-Phonon Interac-
23 tions in the Organic Solid State. *Phys. Rev. Lett.* **1976**, *37*, 36–39.
24
25
26
27
28 (28) Girlando, A.; Marzola, F.; Pecile, C.; Torrance, J. B. Vibrational Spectroscopy of Mixed
29 Stack Organic Semiconductors: Neutral and ionic Phases of tetrathiafulvalene- U chloranil
30 (TTF- U CA) Charge Transfer Complex. *J. Chem. Phys.* **1983**, *79*, 1075–1085.
31
32
33
34 (29) Yartsev, V. M.; Graja, A. Electron- U Intramolecular Vibration Coupling in Charge-Transfer
35 Salts Studied by Infrared Spectroscopy. *Int. J. Mod. Phys. B* **1998**, *12*, 1643–1672.
36
37
38
39 (30) Hasegawa, T.; Mochida, T.; Kondo, R.; Kagoshima, S.; Iwasa, Y.; Akutagawa, T.; Naka-
40 mura, T. T.; Saito, G. Mixed-Stack Organic Charge-Transfer Complexes with Intercolumnar
41 Networks. *Phys. Rev. B* **2000**, *62*, 10059–10066.
42
43
44
45
46 (31) Bozio, R.; Girlando, A.; Pecile, C. Vibrational Analysis of Spectra of quinonoid Molecu-
47 lar Ions. Part 3. U Vibrational Spectra and Assignment of 7,7,8,8-tetracyanoquinodimethane
48 Radical Anion. *J. Chem. Soc., Faraday Trans. 2* **1975**, *71*, 1237–1254.
49
50
51
52
53 (32) Okamoto, H.; Tokura, Y.; Koda, T. Optical Study of Structural Phase Transition in Or-
54 ganic Charge-Transfer Crystals K- and Rb-tetracyanoquinodimethane. *Phys. Rev. B* **1987**,
55
56
57
58
59
60

- 1
2
3
4 (33) Maienschein-Cline, M. G.; Londergan, C. H. The CN Stretching Band of Aliphatic Thio-
5 cyanate is Sensitive to Solvent Dynamics and Specific Solvation. *J. Phys. Chem. A* **2007**,
6 *111*, 10020–10025.
7
8
9
10 (34) Frąckowiak, A. (to be published).
11
12
13 (35) Kozlov, M. E.; Pokhodnia, K. I.; Yurchenko, A. A. Electron Molecular Vibration Coupling
14 in Vibrational Spectra of BEDT-TTF Based Radical Cation Salts. *Spectrochim. Acta A* **1989**,
15 *45*, 437–444.
16
17
18
19 (36) Meneghetti, M.; Bozio, R.; Zanon, I.; Pecile, C.; Ricotta, C.; Zanetti, M. Vibrational Behavior
20 of Molecular Constituents of Organic Superconductors: TMTSF, its Radical Cation and the
21 Sulphur Analogs TMTTF and TMTTF+. *J. Chem. Phys.* **1984**, *80*, 6210–6224.
22
23
24
25
26 (37) Tokura, Y.; Koshihara, S.; Iwasa, Y.; Okamoto, H.; Komatsu, T.; Koda, T.; Iwasawa, N.;
27 Saito, G. Domain-Wall Dynamics in Organic Charge-Transfer Compounds with One-
28 Dimensional Ferroelectricity. *Phys. Rev. Lett.* **1989**, *63*, 2405–2408.
29
30
31
32
33 (38) Kishida, H.; Takamatsu, H.; Fujinuma, K.; Okamoto, H. Ferroelectric Nature and Real-
34 Space Observations of Domain Motions in the Organic Charge-Transfer Compound
35 tetrathiafulvalene-p-chloranil. *Phys. Rev. B* **2009**, *80*, 205201.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Graphical TOC Entry

