

# Supplementary Material

## Activation of persulfate by Fe(III) species: Implications for 4-tert-butylphenol degradation

Yanlin Wu<sup>a,b,c,d</sup>, Romain Prulho<sup>a,b</sup>, Marcello Brigante<sup>a,b\*</sup>, Wenbo Dong<sup>e</sup>, Khalil Hanna<sup>c\*</sup>, Gilles Mailhot<sup>a,b</sup>

<sup>a</sup> Université Clermont Auvergne, Université Blaise Pascal, Institut de Chimie de Clermont-Ferrand, BP 10448, F-63000 Clermont-Ferrand, France


<sup>b</sup> CNRS, UMR 6296, ICCF, F-63171 Aubière, France

<sup>c</sup> Ecole Nationale Supérieure de Chimie de Rennes UMR CNRS 6226, 11 Allée de Beaulieu, CS 50837, F-35708 RENNES Cedex 7, France

<sup>d</sup> State Key Laboratory of Pollution Control and Resources Reuse, College of Environmental Science and Engineering, Tongji University, Shanghai 200092, China

<sup>e</sup> Shanghai Key Laboratory of Atmospheric Particle Pollution and Prevention, Department of Environmental Science and Engineering, Fudan University, Shanghai 200433, China

*Corresponding authors: marcello.brigante@univ-bpclermont.fr ; khalil.hanna@ensc-rennes.fr*



SM1: Effect of pH on the different forms of ferric iron present in aqueous solution