



## **Lipids in maternal diet influence yolk hormone levels and post-hatch neophobia in the domestic chick**

Elske N. de Haas, Ludovic Calandreau, Elisabeth Baéza, Pascal P. Chartrin,  
Rupert Palme, Anne-Sophie Darmaillacq, Ludovic Dickel, Sophie Lumineau,  
Cécilia Houdelier, Isabelle Denis, et al.

### **► To cite this version:**

Elske N. de Haas, Ludovic Calandreau, Elisabeth Baéza, Pascal P. Chartrin, Rupert Palme, et al..  
Lipids in maternal diet influence yolk hormone levels and post-hatch neophobia in the domestic chick.  
Developmental Psychobiology, 2017, 9999 (3), pp.1-10. 10.1002/dev.21504 . hal-01483726

**HAL Id: hal-01483726**

**<https://univ-rennes.hal.science/hal-01483726>**

Submitted on 5 Sep 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License



## Lipids in maternal diet influence yolk hormone levels and post-hatch neophobia in the domestic chick

Journal:	<i>Developmental Psychobiology</i>
Manuscript ID	DEV-16-124.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	de Haas, Elske; INRA-CNRS-Université de Tours François Rabelais, INRA Val de Loire Calandreau, Ludovic; INRA-CNRS-Université de Tours François Rabelais, INRA Val de Loire Baéza, Elisabeth; INRA Centre Val de Loire, Unité de Recherches Avicoles Chartrin, Pascal; INRA Centre Val de Loire, Unité de Recherches Avicoles Palme, Rupert; University of Veterinary Medicine, 1210 Vienna, Unit of Physiology, Pathophysiology, and Experimental Endocrinology, Institute of Medical biochemistry, Departement of Biomedical Sciences Darmaillacq, Anne-Sophie; University of caen Basse-Normandy, Groupe Memoire et plasticité comortementale Dickel, Ludovic; University of caen Basse-Normandy, Groupe Memoire et plasticité comortementale Lumineau, Sophie; University of Rennes 1, UMR CNRS 6552 Ethologie animale et humaine Houdelier, Cecilia; University of rennes, UMR6552; Denis, Isabelle; INRA-U1197, Unité de Neurobiologie de l'Olfaction Arnould, Cécile; INRA-CNRS-Université de Tours François Rabelais, INRA Val de Loire Meurisse, Maryse; INRA-CNRS-Université de Tours François Rabelais, UMR 85 Physiologie de la Reproduction et des Comportements Bertin, Aline; INRA-CNRS-Université de Tours François Rabelais, INRA Val de Loire
Keywords:	Gallus gallus domesticus, maternal effects, yolk hormones, poly unsaturated fatty acids, neophobia

SCHOLARONE™  
Manuscripts

**1 Lipids in maternal diet influence yolk hormone levels and post-hatch**  
**2 neophobia in the domestic chick.**

3 Elske N. de Haas<sup>1,2,3,4</sup>, Ludovic Calandreau<sup>1,2,3,4</sup>, Elisabeth Baéza<sup>5</sup>, Pascal Chartrin<sup>5</sup>, Rupert  
4 Palme<sup>6</sup>, Anne-Sophie Darmaillacq<sup>7</sup>, Ludovic Dickel<sup>7</sup>, Sophie Lumineau<sup>8</sup>, Cécilia Houdelier<sup>8</sup>,  
5 Isabelle Denis<sup>9</sup>, Cécile Arnould<sup>1,2,3,4</sup>, Maryse Meurisse<sup>1,2,3,4</sup>, Aline Bertin<sup>1,2,3,4</sup>

6  
7 <sup>1</sup>INRA Centre Val de Loire, UMR85 Physiologie de la Reproduction et des Comportements,  
8 Laboratoire de Comportement, Neurobiologie et Adaptation, 37380 Nouzilly, France ; <sup>2</sup>  
9 CNRS, UMR7247 Physiologie de la Reproduction et des Comportements, 37380 Nouzilly,  
10 France; <sup>3</sup>Université François Rabelais de Tours, 37041, Tours, France; <sup>4</sup>Institut Français du  
11 Cheval et de l'Equitation, 37380, Nouzilly, France; <sup>5</sup>INRA Centre Val de Loire, Unité de  
12 Recherches Avicoles, 37380 Nouzilly, France ; <sup>6</sup>Unit of Physiology, Pathophysiology, and  
13 Experimental Endocrinology, Institute of Medical Biochemistry, Department of Biomedical  
14 Sciences, University of Veterinary Medicine, 1210 Vienna, Austria ; <sup>7</sup>Groupe Mémoire et  
15 Plasticité comportementale, EA 4259, Université de Caen Normandie UNICAEN, GMPc  
16 EA4259, Caen, France ; <sup>8</sup>UMR CNRS 6552, Ethos, Université de Rennes1, Rennes, France; <sup>9</sup>  
17 INRA, Unité de Neurobiologie de l'Olfaction (NBO, INRA-U1197), 78352 Jouy-en-Josas,  
18 France

19 *Correspondence to be sent to:* [aline.bertin@tours.inra.fr](mailto:aline.bertin@tours.inra.fr)

20 Equipe " Comportement, Neurobiologie et Adaptation"

21 INRA Val de Loire, UMR 85 Physiologie de la Reproduction et des Comportements

22 CNRS-UMR 7247 - Université de Tours - IFCE

23 F-37380 NOUZILLY FRANCE

25 **Running title:** maternal diet engenders maternal effects

26

We assessed whether the ratio of dietary n-6/n-3 polyunsaturated fatty acids (PUFA) during egg formation engenders transgenerational maternal effects in domestic chicks. We analysed yolk lipid and hormone concentrations, and HPA-axis activity in hens fed a control diet (high n-6/n-3 ratio) or a diet enriched in n-3 PUFAs (low n-6/n-3 ratio) for six consecutive weeks. Their chicks were tested for neophobia during the first week of life. We found higher corticosterone metabolites in droppings of hens fed the diet enriched in n-3 and significantly higher concentrations of yolk progesterone, androstenedione and estradiol in their eggs compared to controls. Chicks of hens fed the n-3 enriched diet showed a lower body mass at hatch than controls and expressed higher neophobia when exposed to a novel object. These results add support to the hypothesis that the nutritional state of female birds produces variation in yolk hormone levels and engender maternal effects.

**Keywords:** *Gallus gallus domesticus*, maternal effects, yolk hormones, poly unsaturated fatty acids, neophobia

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

52           Understanding the reasons for phenotypic variation is one of the cornerstones of  
53   behavioural biology. Non-genetic maternal effects play a key role in generating phenotypic  
54   variation in many taxa (Ledón-Rettig, Richards, & Martin, 2013). In birds, prenatal maternal  
55   effects can affect the morphology, behaviour and physiology of offspring (Ruuskanen, 2015).  
56   Such effects are partly mediated via variation in yolk-hormone concentrations of maternal  
57   origin according to different environmental conditions (Groothuis, Muller, von Engelhardt,  
58   Carere, & Eising, 2005). Yolk androgens (mainly testosterone and its precursor  
59   androstenedione) have been widely studied in wild bird populations. Differential allocation of  
60   androgens into eggs is presumed to be a way to adapt offspring to their hatching environment  
61   (Gil, 2008; Groothuis, et al., 2005). Androgen concentrations can vary substantially within  
62   and between egg clutches as a consequence of the maternal environment; factors such as  
63   social conditions (density, aggressive interactions, mate quality) (Kingma et al., 2009; Pilz &  
64   Smith, 2004; Schwabl, 1997), parasitic infection (Müller, Heylen, Eens, Rivera-Gutierrez, &  
65   Groothuis, 2013) and predation risk (Coslovsky, Groothuis, de Vries, & Richner, 2012) can  
66   influence hormone production. Because yolk hormone levels show such a strong context-  
67   dependency within species (Ruuskanen, 2015), the specific environmental cues triggering  
68   maternal effects must be elucidated further.

69           Recently, food resource availability has been identified as an environmental factor  
70   engendering differential maternal hormone allocation. In the lesser black-backed gull (*Larus*  
71   *fuscus*), pied flycatcher (*Ficedula hypoleuca*) and black-legged kittiwake (*Rissa tridactyla*)  
72   food supplementation prior to, and during oviposition affected androgen quantity in the yolk  
73   (Verboven et al., 2003; Verboven, Monaghan, Nager, & Evans, 2010). While in great tits  
74   (*Parus major*), no effect of food supplementation on yolk androgens has been recorded (S.  
75   Ruuskanen, Darras, de Vries, Visser, & Groothuis, 2016). Manipulation of food resources  
76   also shows to increase the within clutch variation of yolk-hormones in canaries (*Serinus*

1  
2  
3 77 *canaria*) (Jonas Vergauwen, Goerlich, Groothuis, Eens, & Müller, 2012) and zebra finches  
4  
5 78 (*Taeniopygia guttata*) (Sandell, Adkins-Regan, & Ketterson, 2007). These studies  
6  
7 79 demonstrate that the nutritional state of maternal birds during oviposition contributes to  
8  
9 80 differences in yolk hormone concentrations between or within a clutch. In the reported  
10  
11 81 studies, enrichment of the maternal diet was achieved by providing additional protein and/or  
12  
13 82 lipids from different sources (e.g. seeds, boiled eggs). This procedure leaves open the question  
14  
15 83 on the influence of the nutritional composition of the food provided to the mother. To our  
16  
17 84 knowledge the effect of the quality of maternal diet on egg hormone levels and offspring  
18  
19 85 phenotype hasn't been investigated so far in birds.  
20  
21  
22

23  
24 86 This question is of particular importance for factory-farmed birds which are fed an  
25  
26 87 unique diet. The domestic chicken (*Gallus gallus domesticus*) is the most abundant bird  
27  
28 88 species on the earth but, despite this importance, many avenues of research remained to be  
29  
30 89 explored in order to advance our knowledge on maternal effects and behavioral development  
31  
32 90 (Dixon, Sparks, & Rutherford, 2016). Housing conditions (Janczak, Torjesen, &  
33  
34 91 Rettenbacher, 2009), unpredictable access to food (Janczak, Torjesen, Palme, & Bakken,  
35  
36 92 2007; Nätt et al., 2009), maternal social status (Muller, Eising, Dijkstra, & Groothuis, 2002),  
37  
38 93 early social stress (Goerlich, Nätt, Elfving, Macdonald, & Jensen, 2012) and thermal  
39  
40 94 environment (Bertin et al., 2013) cause variation in yolk androgens and progesterone levels in  
41  
42 95 domestic hens. This caused modifications in growth, feeding behaviors, and emotional  
43  
44 96 reactivity of their chicks (Goerlich, Nätt, Elfving, Macdonald, & Jensen, 2012; Nätt et al.,  
45  
46 97 2009; Bertin et al., 2013). In domestic hens, maternal stress during egg formation —mimicked  
47  
48 98 with subcutaneous corticosterone implantation— increased plasma corticosterone levels and  
49  
50 99 decreased the synthesis of reproductive hormones which accumulate in the yolk (Henriksen,  
51  
52 100 Groothuis, & Rettenbacher, 2011). In addition, on-farm longitudinal observations showed that  
53  
54 101 parental stress physiology correlate with offspring's anxiety and expression of damaging  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

behavior (de Haas, Bolhuis, Kemp, Groothuis, & Rodenburg, 2014; Rodenburg & de Haas, 2016).

In domestic hens we previously showed that adding fish oil to the maternal diet induced food neophobia in the offspring (Aigueperse, Calandreau, & Bertin, 2013). As fish oil is particularly rich in n-3 poly-unsaturated fatty acids (PUFA), this result suggested that PUFA in the maternal diet may directly affect the development of behaviour in the offspring. The aim of the present study was to test this hypothesis and assess whether the ratio between n-6 and n-3 PUFA in the maternal diet during egg formation could engender maternal effects. N-3 and n-6 PUFA unambiguously influence the development of the central nervous system in both mammalian and avian species (Noble & Cocchi, 1989). But these effects are better-understood for mammals. In mammals, PUFAs and their mediators are known to regulate many processes within the brain, such as neurotransmission, cell survival and neuro-inflammation, and also affect hypothalamic–pituitary–adrenal (HPA) axis responses, thereby possibly playing a role in anxiety-like behavior and cognition (e.g. Bazinet & Layé, 2014; Lafourcade et al., 2011). In the domestic chicken, PUFA ratios in the maternal diet are transferred to the egg yolk (Noble & Cocchi, 1990) and subsequently to the tissues of the developing chick embryo (Maldjian, Farkas, Noble, Cocchi, & Speake, 1995). Numerous studies show that the PUFA profile of the developing chick brain reflects the PUFA composition in the yolk (Cherian & Sim, 1993; Anderson, Connor, Corliss, & Lin, 1989) but the consequences on behavioral development remain to be investigated.

We analysed yolk lipid and hormone concentrations, and HPA-axis activity in hens fed a control diet (high n-6/n-3 ratio) or a diet enriched in n-3 PUFAs (low n-6/n-3 ratio). Development and neophobia (i.e. fear of novelty) of the hatchlings were analysed. Neophobia is considered to be a temperament trait which can impair an animal’s capacity to adapt to new resources, new habitats and express behavioural innovation (Greenberg & Mettke-hofmann,

2001; Webster & Lefebvre, 2001). As a high level of neophobia can reduce fitness (Brown, Ferrari, Elvidge, Ramnarine, & Chivers, 2013), this behavioural trait is a growing topic of interest for behavioural biologists (Camín, Martín-Albarracín, Jefferies, & Marone, 2016; Greggor, Thornton, & Clayton, 2015). The precocial domestic chick is a particularly appropriate model in which to analyse developmental influences on neophobia since chicks start food selection soon after hatching (Bertin et al., 2010). In addition, under commercial conditions the refusal to accept new foods leads to a major reduction in feed intake and, subsequently, in growth and animal welfare (Murphy, 1977).

## Methods

### *Ethics statement*

All birds were maintained at the Experimental Unit PEAT of INRA (Nouzilly, France). The Experimental Unit is registered by the ministry of Agriculture with the license number B-37-175-1 for animal experimentation. All experiments were approved by the Ethic Committee in Animal Experimentation of Val de Loire, CEEA Vdl (reference number 02153.02). The CEEA vdl is registered by the National Committee "Comité National de Réflexion Ethique sur l'Expérimentation Animale" under the number 19. All experiments were performed in accordance with the European Communities Council Directive 2010/63/UE. All animals were sold for rehabilitation at the end of the experiment.

### *Maternal hens and diets*

Hens (n=40) and diets were provided by the experimental unit PEAT of INRA (Nouzilly, France). Two groups of 20 one-year old White Leghorn laying hens were matched for body weight, and divided into two dietary treatment groups, a control diet and a diet enriched in n-3 FA. Diets were administered for six consecutive weeks. The control diet contained soya-oil



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

151 (3%) with a n-6:n-3 ratio of 7.99, a value within the range recommended for commercial  
152 laying hens (Van Elswyk, 1997). The n-3 diet contained 3% menhaden fish-oil (Sigma Life  
153 Science) having an n-6:n-3 ratio of 0.82. Diets were iso-caloric (2820 kcal ME/kg) and iso-  
154 nitrogenous (170 g CP/kg), differing only in the composition of PUFAs (electronic  
155 supplementary material, table S1). Hens were housed individually in wire-mesh cages of 100  
156 x 100 x 50 cm (*l x w x h*), with the possibility of tactile, visual, and vocal contact with  
157 neighbouring hens. Cages contained wood shavings and a wooden perch on the floor, a nest  
158 enclosed by red curtain flaps, a nipple drinker and a feed trough. Water and food were  
159 provided *ad libitum*. Four hens were excluded from the experiment, three due to a lack of  
160 habituation to the experimental diets (food conservatism) and one death unrelated to the diet  
161 treatment, resulting in n =18 hens per diet group.

162 *Morpho-physiological measurements of hens*

163 All hens were weighed once per week. 24 h-feed intake per hen was measured once a week by  
164 calculating the difference in weight of the feeder before and after 24 h. The number of eggs  
165 laid was recorded daily. Laying rate was calculated as a percentage of the number of eggs laid  
166 per female per day.

167 In order to evaluate HPA activity in chickens, faecal corticosterone metabolite (FCM)  
168 concentrations were measured (Rettenbacher, Mostl, Hackl, Ghareeb, & Palme, 2004). In the  
169 sixth week of diet treatment, one fresh faecal dropping per hen was collected from the home  
170 cage. Urine was removed from the sample as the concentrations of FCM in the liquid portions  
171 of droppings are more likely to represent acute HPA activity caused by handling or disturbing  
172 the animals during collection (Rettenbacher, et al., 2004). Each sample was homogenized and  
173 stored at -20°C. From each sample an aliquot (0.5 g) was extracted with 60% methanol

(Mostl, Spendier, & Kotrschal, 2001) and analysed by a cortisone enzyme immunoassay (EIA) validated for chickens and fully described in (Rettenbacher, et al., 2004).

*Analysis of egg quality, yolk hormones and yolk lipids*

As the vitellogenesis has an average length of 8 days (Lacassagne, 1960), we collected eggs for hormonal and lipid assays during the fifth and sixth weeks of treatment to ensure hens were fully habituated to the diets, and that differences in egg components were a result of the maternal diet treatments. One egg per female was weighed and stored at -20°C for hormonal assay. Eggshells were separated and dried for 24 h and weighed. Frozen yolk was separated from the albumen and weighed. The weight of albumen was calculated by subtracting the weight of the eggshell and yolk from that of the whole egg. We then determined the ratio of each component relative to the egg mass (yolk mass / egg mass; albumen mass / egg mass; shell mass / egg mass) for each female. A total of 18 eggs were collected per group. The concentrations of immunoreactive progesterone, testosterone, androstenedione and oestrogens were analysed by EIAs. Details of the extraction protocol are found in (Guesdon et al., 2011). For a full description of the assays including specific antibodies, see (Palme, Touma, Arias, Dominchin, & Lepschy, 2013). Intra- and inter-assay coefficients of variation were less than 10% and 15%, respectively.

In order to measure yolk lipid content, three yolks per female were pooled and homogenized. We then determined the total lipid content of the yolk and the percentages of saturated, mono-unsaturated and poly-unsaturated FAs within the lipid fraction. Yolk lipid analysis was conducted by trans-methylation with gas chromatography (Perkin Elmer Autosystem, St. Quentin en Yvelines, France) following the protocol described in (Chartrin, Berri, Lebihan-Duval, Quentin, & Baéza, 2005).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

During the fourth and fifth week of treatment, hens were artificially inseminated three times (same mix of sperm for all the hens). Two hundred and thirty nine eggs (mean of  $5.6 \pm 0.4$  eggs per female) produced offspring. Eggs were incubated at  $37.8^{\circ}\text{C}$  with 56% relative humidity for 21 days.

*Chicks and housing conditions*

Forty-eight chicks from hens fed the n-3 enriched diet (n-3 chicks) and 50 control chicks (C chicks) were used in this experiment. Chicks were identified with a metal leg-ring. Chicks were housed in non-sibling pairs but with birds from the same diet treatment. We chose not to mix treatments within each cage as potential unknown differences in chicks' social or emotional behaviour between treatments may mask prenatal effects. Chicks were housed in plastic cages (50 cm x 40 cm x 30 cm) placed in rows with treatments in alternation within a row (two adjacent cages were from different treatments). All the cages were equally distributed between two identical rooms. Cages had wood shavings on the floor and a wire-mesh lid. Food and water were provided *ad libitum*. Chicks were fed a postnatal starter mash with a PUFA ratio similar to the maternal diet. We used this diet so that the PUFA ratio of the chick's diet would not counteract the effects induced by the maternal diet on offspring's brain PUFA levels. This is especially important since precocial birds adapt their feeding behaviour to their mother's diet choices (Hess, 1964). Moreover, when the chick's brain experiences a deficiency in PUFAs due to restriction of the maternal diet, it is able to compensate for this deficiency by altering lipid metabolism and demonstrating preferential uptake of the PUFA lacking in the diet (Anderson, Van Winkle, & Connor, 1992). As chicks rely on the vitelline sac reserve for 2/3 days after hatching, any phenotypic differences during the first days of life between chicks of both maternal diets could therefore be attributed to maternal diet effects. The n-3 diet administered to experimental chicks had an n-6:n-3 ratio of 1.03, and the control diet had an n-6:n-3 ratio of 8.15. Both diets were iso-caloric (2900 kcal ME/kg) and iso-

nitrogenous (207.2 g CP/kg) differing solely in PUFA composition (electronic supplementary material, table S1). To determine the sex we looked for the development of the comb at 23 days of age. The body weight of each chick was measured at hatching and at seven days of age.

#### *Behavioural characterization of chicks*

To characterize neophobia, behavioural responses to two novel foods and a novel object were assessed with a protocol previously described by (Bertin et al., 2015). Each test was executed at the same age for all chicks. All tests lasted 180 s.

Because chicks become distressed when socially isolated (Hocking, Haldane, Davidson, Sandøe, & Kristensen, 2015), we tested cage mates together ( $n = 24$  pairs of n-3 chicks and  $n = 25$  pairs of C chicks). Testing took place in a separate room within an experimental cage that had the same features as their home cage. Testing commenced 90 min after the feeder was removed from their home cage. Pairs were transported to the test room in a  $15\text{ cm} \times 15\text{ cm} \times 15\text{ cm}$  plastic container, and were gently deposited in an opaque enclosure ( $20\text{ cm} \times 6\text{ cm} \times 20\text{ cm}$ ) within the testing cage, opposite the feeding trough. After 30 s, the enclosure was lifted and removed while chicks remained inside the testing cage. The behaviour of one marked focal chick of each pair was recorded by an unseen observer. Focal chicks were chosen randomly at two days of age and were tagged with a blue-coloured mark on the head. The latency (s) to eat (the moment swallowing was observed) and time spent eating (s) were recorded.

(i) *Habituation test/familiarization*: at three days of age, chicks were familiarized with the testing cage and handling procedure. As chicks rely on their vitelline sac reserve the first days of life, this test also measured food motivation. The metallic feeding trough of their home cage was placed in the test cage, filled with 200 g of their usual diet.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

246 (ii) *Food neophobia tests*: we used these tests to assess the immediate reaction to novel foods.  
247 Chicks were exposed to changes in sensory properties of the food. At four days of age, the  
248 first test was done with cracked corn-wheat (cereals contained in familiar pellets). Millet  
249 seeds (a novel food) was used at five days of age for the second test.

250 (iii) *Object neophobia test*: chicks were tested at six days of age. The novel object was an  
251 unfamiliar feeder containing the familiar diet of the test pair.

252 **Statistical analysis**

253 Data were analyzed with Statview (SAS Institute Inc., Cary, NC). The distribution of  
254 variables was tested with a Kolmogorov-Smirnov test. Hen and egg data were normally  
255 distributed. Groups were compared using a t-test (FCM, mass of egg components, yolk-  
256 hormones, yolk-PUFA concentrations) or repeated measures ANOVA (hen body weight, food  
257 intake and laying rate) with the main effect of diet, time, and the interaction between time and  
258 diet. The number of chicks hatched (hatching success) and sex ratio between diet groups were  
259 compared with a Chi-square test. The weight of chicks at hatching and their growth rate  
260 during the first week were analyzed via ANOVA. Offspring behavioural data did not follow a  
261 normal distribution even after transformation, so the non-parametric Mann-Whitney *U*-test  
262 was used to compare groups. Data are presented as mean  $\pm$  SEM, with significance accepted  
263 at  $p < 0.05$ .

264  
265 **Results**

266 *Dietary effects on laying performance of hens*

After five weeks of consuming their respective diets, n-3 enriched hens had significantly higher FCM levels than hens of the C diet ( $398 \pm 57$  ng/g vs.  $99 \pm 19$  ng/g,  $t = 2.98$ ,  $p = 0.006$ ).

No effect of diet or interaction between diet and time were observed for body weight, 24h-food intake, or laying rate (see Table 1). For food intake, a time effect was observed (time:  $F_{4,33} = 15.3$ ,  $p < 0.01$ ). Hen food intake was lower during the first week compared to the second week for both diets (Table 1).

>>> INSERT Table 1 <<<

#### *Dietary effects on egg components, yolk hormones and yolk lipids*

The eggs of n-3 hens were significantly lighter than those of C hens ( $54.9 \pm 1.25$  g vs.  $58.0 \pm 0.86$  g,  $t = -2.11$ ,  $p = 0.04$ ). N-3 eggs had a significantly lower yolk mass/egg mass ratio ( $0.28 \pm 0.04$  vs.  $0.29 \pm 0.06$ ,  $t = 2.47$ ,  $p = 0.02$ ), and higher albumin mass/egg mass ratio ( $0.64 \pm 0.05$  vs.  $0.62 \pm 0.04$ ,  $t = -2.07$ ,  $p = 0.05$ ) than C eggs. Eggshell mass/egg mass ratio did not differ between diets ( $0.09 \pm 0.02$  for both diets,  $t = -0.41$ ,  $p = 0.68$ ).

Compared to C hens, the egg yolk of n-3 hens contained a significantly higher concentration of immunoreactive progesterone ( $t = 2.96$ ,  $p = 0.006$ ), oestrogens ( $t = 5.97$ ,  $p < 0.0001$ ) and androstenedione ( $t = 2.01$ ,  $p = 0.05$ ) with a trend in the same direction for testosterone ( $t = 1.68$ ,  $p = 0.10$ ; Figure 1). If instead of yolk hormone concentrations (ng/g) we analyse hormone contents (ng/yolk), the results remained the same except for androstenedione (n-3 eggs vs. n-6 eggs: progesterone:  $17897.2 \pm 2241.07$  ng vs.  $12042.04 \pm 1635.8$  ng,  $t = 2.1$ ,  $p = 0.04$ ; oestrogens:  $162.5 \pm 4.9$  ng vs.  $134.05 \pm 5.4$  ng,  $t = 3.85$ ,  $p = 0.0005$ ; androstenedione:  $785.6 \pm 75.06$  ng vs.  $641.7 \pm 94.16$  ng,  $t = 1.2$ ,  $p = 0.23$ ; testosterone:  $42.7 \pm 4.7$  ng vs.  $30.4 \pm 5.2$  ng,  $t = 1.77$ ,  $p = 0.08$ ).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The percentage of total lipid content in the yolk did not differ between diets ( $34.7 \pm 0.6\%$  vs.  $33.1 \pm 1.0\%$ ,  $t = 1.36$ ,  $p = 0.18$ ). Within the lipid fraction of the yolk, the percentage of saturated fatty acids did not significantly differ between diets ( $33.6 \pm 0.47\%$  vs.  $33.7 \pm 0.37\%$ ,  $t = -0.28$ ,  $p = 0.78$ ). For n-3 hens, the percentage of mono-unsaturated fatty acids was higher, while the percentage of PUFAs was lower in comparison to eggs from the C hens (mono:  $49.5 \pm 0.62\%$  vs.  $44.0 \pm 0.65\%$ ,  $t = -6.2$ ,  $p < 0.001$ , PUFA:  $16.7 \pm 0.36\%$  vs.  $22.5 \pm 0.60\%$ ,  $t = 8.22$ ,  $p < 0.001$ ). The yolk of eggs from n-3 hens had significantly higher percentages of n-3 PUFAs with a significantly lower n-6/n-3 ratio, while the yolk of eggs from C hens had significantly higher percentages of n-6 PUFAs and a significantly higher n-6/n-3 ratio (Table 2).

>>> INSERT Figure 1 <<<

>>> INSERT Table 2 <<<

*Dietary effects on hatching success and weight of chicks*

Hatching success did not differ between diets (98 hatched out of 111 fertile eggs for the n-3 group vs. 125 out of 138 fertile eggs for the n-6 group;  $\chi^2$ -test,  $p = 0.67$ ), neither did sex-ratio (25 females and 23 males in the n-3 group and 28 females and 22 males in the C group,  $\chi^2$ -test,  $p = 0.69$ ). N-3 chicks were lighter at hatching than C chicks ( $39.4 \pm 0.4\text{ g}$  vs.  $40.9 \pm 0.5\text{ g}$ ,  $t = -2.29$ ,  $p = 0.02$ ) but did not differ significantly in growth rate during the first week of life ( $8.60 \pm 0.16\text{ g/week}$  vs.  $9.72 \pm 1.86\text{ g/week}$ ,  $F_{3,93} = 0.66$ ,  $p = 0.57$ ).

*Dietary effects on neophobic behaviour in the chicks*

Latencies to eat and the time spent eating did not differ significantly between n-3 and C chicks in the habituation/familiarization test and both food neophobia tests (Table 3). In the

novel object test, n-3 chicks demonstrated a longer latency to eat and spent less time eating from the novel object compared to C chicks (Table 3).

>>> INSERT Table 3 <<<

## Discussion

The present experiment was designed to test the hypothesis that PUFA in the maternal diet could engender differential allocation of yolk hormones of maternal origin. We found higher corticosterone metabolites, reflecting heightened HPA-activity, in the faecal droppings of hens fed the diet enriched in n-3 compared to control animals. As expected, yolk n-6:n-3 ratio strongly reflected that of the maternal diet. There were significantly higher concentrations of yolk progesterone, androstenedione and oestrogens, as well as a trend for higher yolk testosterone in eggs of hens fed the n-3 enriched diet compared to controls. Chicks of hens fed the n-3 enriched diet showed a lower body weight at hatching and higher neophobia when exposed to a novel object compared to chicks of hens fed the control diet.

After six weeks of consuming a diet enriched in n-3 PUFAs, hens had elevated concentrations of corticosterone metabolites in the faeces and elevated levels of yolk hormones compared to hens fed a control diet. Additionally, hens of the n-3 diet laid lighter eggs, with lower yolk/egg mass ratios than controls. Diets were equal in protein and energy content, and were selected to meet the dietary requirements of laying hens (Van Elswyk, 1997). Food intake and body weight followed a similar pattern for n-3 and control hens. Thus, differences between groups cannot be explained by differences in the energy content of the diet or differences in food intake. Rather, they demonstrate an influence of n-6/n3 PUFA ratio in the diet on the metabolism and physiology of hens.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

334           These results suggest a potential action for the n-6:n-3 PUFA ratio of diet on the HPA-  
335   axis of hens as reported in other species (Fisher et al., 2014 ; Larrieu et al., 2014; Lafourcade,  
336   et al., 2011). First, hens of the n-3 diet had FCM levels more than 3-times greater than  
337   controls. FCM levels have been shown to reflect HPA activity (Rettenbacher & Palme, 2009).  
338   Second, in chickens with elevated corticosterone levels due to a corticosterone implant, egg  
339   mass, yolk mass and hatchling's weight reduction have also been recorded (Rie Henriksen,  
340   Groothuis, & Rettenbacher, 2011; R. Henriksen, Rettenbacher, & Groothuis, 2011) which is  
341   comparable to our chicks from hens treated with n-3. Moreover, in birds, environmental stress  
342   can induce HPA axis activation, causing a decrease in egg and offspring weight (Hayward &  
343   Wingfield, 2004; R. Henriksen, et al., 2011; Hsu, Dijkstra, Darras, de Vries, & Groothuis,  
344   2016). Third, it is known that maternal environmental stress induces similar increases in hens'  
345   yolk hormones as we found in the hens fed the n-3 diet (Bertin, et al., 2013; Guibert et al.,  
346   2010). In mammals, the effects of n-3 PUFAs on HPA axis functioning are generally  
347   beneficial, but not always. As in mice, a deficiency of n-3 PUFA can alter corticosterone  
348   secretion via modifications in glucocorticoid receptors (GRs) of the prefrontal cortex, but not  
349   in the HPA-axis (Larrieu, Hilal, De Smedt-Peyrusse, Sans, & Laye, 2016). In the chicken,  
350   GRs are also found in the pituitary (Bossis, Nishimura, Muchow, & Porter, 2004) and frontal  
351   brain areas (Bordone, Schrott, & Sparber, 1997). Glucocorticoid receptors can be modified by  
352   stressful conditions in a variety of avian species (Japanese quail (Zimmer & Spencer, 2014),  
353   starlings (Dickens, Romero, Cyr, Dunn, & Meddle, 2009), house sparrow (Lattin & Romero,  
354   2014)) and play a role in the lipid metabolism of the white-crowned sparrow (Landys,  
355   Piersma, Ramenofsky, & Wingfield, 2004) (Landys, Ramenofsky, Guglielmo, & Wingfield,  
356   2004). Our results on HPA-functioning are a likely consequence of the surplus of n-3 in the  
357   experimental diet. This may have resulted in a lack of arachidonic acid (AA), which is  
358   essential for metabolic and hormonal functioning (Huang, Leibovitz, Lee, & Millar, 1990).

Based on our results, we hypothesize that the ratio of n-6:n-3 PUFAs in the diet influence HPA activity in birds, but the mechanisms require further investigation. It is important to notice that domestic chickens have been raised for food thousands of years ago and were mainly fed with wheat seeds. As wheat seeds are poor in n-3 and rich in n-6 fatty acids, these animals may have developed specific adaptations and, as a consequence, may cope better with high n-6:n-3 ratios than with low n-6:n3 ratios in the diet.

Egg mass and yolk proportion was reduced in n-3 hens. These results are in accordance with those obtained by administering diets enriched with 2%, 3% or 4% fish oil (Gonzalez-Esquerria & Leeson, 2000; Huang, et al., 1990) in which hens' egg mass was found to decrease linearly with increasing percentages of fish oil (high in n-3). Diets enriched with fish oil can increase albumin production and decrease yolk production compared to control diets (Huang, et al., 1990). The n-3 diet we applied contained little linoleic acid (a molecular precursor essential to the production of n-6 PUFA AA). Moreover, AA production appeared to be strongly inhibited by the dominance of docohexaenoic acid (DHA) in the n-3 diet as demonstrated by the amount measured in the yolk. This could severely hamper AA metabolism, resulting in metabolic changes in the hens receiving the n-3 diet, and thereby altering egg composition (Huang, et al., 1990). PUFAs in the maternal diet can thus have an important effect on the amount and constitution of the resources available for embryo development.

Moreover, our data showed that fish oil in the diet led to higher concentrations of yolk progesterone, androstenedione and oestradiol and a trend for higher concentrations of yolk testosterone. These results add support to the hypothesis that the nutritional state of female birds causes variation in yolk hormone levels. However, at the present stage, the interpretation of the mechanisms that mediate variations in yolk hormone levels is bound to be speculative. As mentioned previously, it could be that increased levels of yolk hormones are the result of

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

HPA axis activation in the mother. In accordance with this hypothesis; recent studies showed that maternal plasma corticosterone levels influence reproductive hormone concentrations in the yolk (Henriksen, Groothuis, & Rettenbacher, 2011). The underlying physiological mechanisms remain poorly understood and are barely addressed in the literature. As argued before, it might be possible that corticosterone levels affect the steroidogenic activity of follicles. Another hypothesis could be related to the direct influence of PUFAs and their oxygenated derivatives on steroid synthesis. Several factors of the steroidogenic machinery are directly regulated by AA and AA-derived eicosanoids (Wathes, Abayasekara, & Aitken, 2007). The n-3/n-6 PUFA ratio in the diet largely determines the balance between the various types of oxygenated metabolites deriving from AA or from EPA and DHA (Yates, Calder, & Rainger, 2014). Therefore, the decreased AA/DHA+EPA ratio and the resulting eicosanoid pattern in n-3 supplemented hens may have directly contributed to the observed differences in yolk hormone levels. Consistently, dietary fat (such as those derived from fish oil) in the maternal diet is known to influence egg weight and plasma oestrogen levels (Whitehead, 1995), a role for PUFAs in the hormonal metabolism of birds seems, therefore, a plausible explanation.

We found that n-3 chicks were more inhibited (longer latency to eat and less time spent eating) than C chicks in the novel object test. This result indicates a higher level of object neophobia in n-3 chicks than C chicks. On the other hand, n-3 and C chicks did not differ in food neophobia test. Since the n-3 maternal diet modified the embryonic environment in several ways (yolk mass, yolk hormones and fatty acids), it is difficult to determine which mechanisms and which hormones affected object neophobia. Little is currently known regarding the developmental influence of yolk progesterone and androgens on neophobia. The influence of progesterone is barely investigated. In Northern bobwhite quail (*Colinus virginianus*), a recent study showed that elevated yolk progesterone levels

1  
2  
3 409 elevate emotional reactivity in neonates (Herrington, Vallin, & Lickliter, 2015). Interestingly,  
4  
5 410 an artificial increase in yolk hormone levels has been found to alter object neophobia but not  
6  
7 411 food neophobia in domestic chicks (Bertin, et al., 2015), but the direction of effects was not  
8  
9 412 the same as in the present study. Lower object neophobia was found in chicks exposed *in ovo*  
10  
11 413 to increased yolk progesterone, androstenedione and estradiol levels compared to controls.  
12  
13 414 Discrepancies in the direction of yolk hormonal effects are also seen in latencies to approach  
14  
15 415 novel objects depending on species and the injected dose (Vergauwen, Eens, & Muller, 2012).  
16  
17 416 Nevertheless, all of these experiments suggest that variation in yolk hormone levels can alter  
18  
19 417 specific aspects of neophobia.  
20  
21  
22

23 418 The data concerning the effects of environmental conditions during ontogeny on the  
24  
25 419 development of object or food neophobia are inconclusive. In accordance with our findings,  
26  
27 420 zebra finch chicks (*Taeniopygia guttata*) from eggs with artificially elevated yolk testosterone  
28  
29 421 show no difference in latency to eat novel foods than control animals (Tobler & Sandell,  
30  
31 422 2007). Similarly, in canaries (*Serinus canaria*), increased levels of yolk testosterone was  
32  
33 423 found to not be the primary source of variation in the expression of food neophobia (  
34  
35 424 Vergauwen, et al., 2012). Our data highlight the importance of studying the quality of the  
36  
37 425 maternal diet in order to understand more fully the ontogeny of emotional and behavioural  
38  
39 426 traits involved in adapting to new resources. This could be of particular importance when food  
40  
41 427 resources become scarcer under the projected future of climate change. In captive bird  
42  
43 428 populations (for production or conservation purposes), no specific recommendation is made  
44  
45 429 regarding PUFA quantity and quality in the diet. This study highlights the importance of  
46  
47 430 carefully considering food quality in captive bird populations as they might engender  
48  
49 431 transgenerational effects on development and behaviour. In our study, chicks were all feeding  
50  
51 432 around two/three days after hatching, therefore we cannot totally exclude the possibility that  
52  
53 433 the behaviour of the chicks was influenced by their respective diet. Two points are of  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

importance when considering this hypothesis. First, even when chicks start feeding, the residual yolk continues to be taken up and thereby provides a source of fatty acids of maternal origin (Romanoff, 1960). Second, in the domestic chicken and king penguin (*Aptenodytes patagonicus*) it was consistently found that the proportion of fatty acids in the brain phospholipids was particularly refractory to the effects of the chicks' diet compared to other tissues (Thil, Speake, & Groscolas, 2003; Anderson, et al., 1992). More specifically, high levels of brain n-3 produced by exposure to a maternal diet rich in fish oil are not reversible on a short time period (Anderson, et al., 1992). Although caution must be exercised, we argue that maternal effects more likely contributed to the differences observed in hatchlings' mass and behaviour than the post-natal diet.

The present data show maternal diet composition as an overlooked pathway by which the environment could engender large variation in yolk hormones levels. This may play a role in the development of offspring behaviour and phenotype. In farm birds, as unresolved welfare issues are mediated by unknown cues, this study lends support to perspectives in the field of maternal nutrition. Further investigations to the mechanisms are required but this study suggests common principles regarding the regulative action of n-3 and n-6 PUFA on fear-related behavior across vertebrate taxa. Using precocial birds could yield new insights into the mechanism by which the prenatal resources in PUFA influences behavioral development of individuals in vertebrates

**Acknowledgements**

All birds were maintained at the PEAT experimental unit, INRA, Nouzilly, France. We are grateful to all members of the unit, particularly P. Ganier and O. Callut for taking care of the birds. The authors are grateful to Dr. C. O'Brien for English proofreading. This work was

458 supported by the French National Research Agency (PReSTO'Cog ANR-13-BSV7-0002-02)  
459 and the University François Rabelais of Tours (Post-doctoral grant).

460

461

For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**References**

Aigueperse, N., Calandreau, L., & Bertin, A. (2013). Maternal Diet Influences Offspring Feeding Behavior and Fearfulness in the Precocial Chicken. *PLoS ONE*, 8(10), e77583. doi: 10.1371/journal.pone.0077583

Anderson, G., Connor, W., Corliss, J., & Lin, D. (1989). Rapid modulation of the n-3 docosahexaenoic acid levels in the brain and retina of the newly hatched chick. *Journal of Lipid Research*, 30(3), 433-441.

Anderson, G. J., Van Winkle, S., & Connor, W. E. (1992). Reversibility of the effects of dietary fish oil on the fatty acid composition of the brain and retina of growing chicks. *Biochimica et Biophysica Acta (BBA)-Lipids and Lipid Metabolism*, 1126(2), 237-246.

Bazinet, R. P., & Layé, S. (2014). Polyunsaturated fatty acids and their metabolites in brain function and disease. *Nature Reviews Neuroscience*, 15(12), 771-785.

Bertin, A., Arnould, C., Moussu, C., Meurisse, M., Constantin, P., Leterrier, C., & Calandreau, L. (2015). Artificially Increased Yolk Hormone Levels and Neophobia in Domestic Chicks. *Animals*, 5(4), 0408.

Bertin, A., Calandreau, L., Arnould, C., Nowak, R., Levy, F., Noirot, V., . . . Leterrier, C. (2010). In Ovo Olfactory Experience Influences Post-hatch Feeding Behaviour in Young Chickens. *Ethology*, 116(11), 1027-1037. doi: 10.1111/j.1439-0310.2010.01820.x

Bertin, A., Chanson, M., Delaveau, J., Mercierand, F., Möstl, E., Calandreau, L., . . . Collin, A. (2013). Moderate heat challenge increased yolk steroid hormones and shaped offspring growth and behavior in chickens. *PLoS ONE*, 8(2), e57670.

Bordone, L., Schrott, L., & Sparber, S. (1997). Ontogeny of Glucocorticoid Receptors in the Hyperstriatum-Hippocampus-Parahippocampal Area and Optic Tectum of the

- Embryonic Chicken (*Gallus domesticus*) Brain. *Journal of Neuroendocrinology*, 9(10), 753-761.
- Bossis, I., Nishimura, S., Muchow, M., & Porter, T. E. (2004). Pituitary expression of type I and type II glucocorticoid receptors during chicken embryonic development and their involvement in growth hormone cell differentiation. *Endocrinology*, 145(7), 3523-3531.
- Brown, G. E., Ferrari, M. C. O., Elvidge, C. K., Ramnarine, I., & Chivers, D. P. (2013). Phenotypically plastic neophobia: a response to variable predation risk. *Proceedings of the Royal Society of London B: Biological Sciences*, 280(1756). doi: 10.1098/rspb.2012.2712
- Camín, S. R., Martín-Albarracín, V., Jefferies, M., & Marone, L. (2016). Do neophobia and dietary wariness explain ecological flexibility? An analysis with two seed-eating birds of contrasting habits. *Journal of Avian Biology*, 47(2), 245-251. doi: 10.1111/jav.00697
- Chartrin, P., Berri, C., Lebihan-Duval, E., Quentin, M., & Baéza, E. (2005). Lipid and fatty acid composition of fresh and cured-cooked breast meat of standard, certified and label chickens. *Archiv Fur Geflugelkunde*, 69(5), 219-225.
- Cherian, G., & Sim, J. S. (1993). Net transfer and incorporation of yolk n-3 fatty acids into developing chick embryos. *Poultry Science*, 72(1), 98-105.
- Coslovsky, M., Groothuis, T., de Vries, B., & Richner, H. (2012). Maternal steroids in egg yolk as a pathway to translate predation risk to offspring: Experiments with great tits. *General and Comparative Endocrinology*, 176(2), 211-214.
- de Haas, E. N., Bolhuis, J. E., Kemp, B., Groothuis, T. G., & Rodenburg, T. B. (2014). Parents and early life environment affect behavioral development of laying hen chickens. *PLoS ONE*, 9(3), e90577.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Dickens, M., Romero, L., Cyr, N., Dunn, I., & Meddle, S. (2009). Chronic stress alters glucocorticoid receptor and mineralocorticoid receptor mRNA expression in the European starling (*Sturnus vulgaris*) brain. *Journal of Neuroendocrinology*, 21(10), 832-840.

Dixon, L., Sparks, N., & Rutherford, K. (2016). Early experiences matter: a review of the effects of prenatal environment on offspring characteristics in poultry. *Poultry Science*, 95(3), 489-499.

Fisher, R., Or'Rashid, M., Quinton, M., AlZahal, O., Boermans, H., McBride, B., & Karrow, N. (2014). Maternal supplementation with fishmeal protects against late gestation endotoxin-induced fetal programming of the ovine hypothalamic-pituitary-adrenal axis. *Journal of developmental origins of health and disease*, 5(03), 206-213.

Gil, D. (2008). Hormones in avian eggs: Physiology, ecology and behavior *Advances in the Study of Behavior*, Vol 38 (Vol. 38, pp. 337-398). San Diego: Elsevier Academic Press Inc.

Goerlich, V. C., Nätt, D., Elfwing, M., Macdonald, B., & Jensen, P. (2012). Transgenerational effects of early experience on behavioral, hormonal and gene expression responses to acute stress in the precocial chicken. *Hormones and Behavior*, 61(5), 711-718.

Gonzalez-Esquerria, R., & Leeson, S. (2000). Effect of feeding hens regular or deodorized menhaden oil on production parameters, yolk fatty acid profile, and sensory quality of eggs. *Poultry Science*, 79(11), 1597-1602.

Greenberg, R., & Mettke-hofmann, C. (2001). Ecological aspects of neophobia and neophilia in birds *Current Ornithology* (pp. 119-178): Springer.

Greggor, A. L., Thornton, A., & Clayton, N. S. (2015). Neophobia is not only avoidance: improving neophobia tests by combining cognition and ecology. *Current Opinion in Behavioral Sciences*, 6, 82-89.

- 536 Groothuis, T. G. G., Muller, W., von Engelhardt, N., Carere, C., & Eising, C. (2005).  
537 Maternal hormones as a tool to adjust offspring phenotype in avian species.  
538 *Neuroscience and Biobehavioral Reviews*, 29(2), 329-352. doi:  
539 10.1016/j.neubiorev.2004.12.002
- 540 Guesdon, V., Bertin, A., Houdelier, C., Lumineau, S., Formanek, L., Kotrschal, K., . . .  
541 Richard-Yris, M.-A. (2011). A Place to Hide in the Home-Cage Decreases Yolk  
542 Androgen Levels and Offspring Emotional Reactivity in Japanese Quail. *PLoS ONE*,  
543 6(9), e23941. doi: 10.1371/journal.pone.0023941
- 544 Guibert, F., Richard-Yris, M.-A., Lumineau, S., Kotrschal, K., Guémené, D., Bertin, A., . . .  
545 Houdelier, C. (2010). Social Instability in Laying Quail: Consequences on Yolk  
546 Steroids and Offspring's Phenotype. *PLoS ONE*, 5(11), e14069.
- 547 Hayward, L. S., & Wingfield, J. C. (2004). Maternal corticosterone is transferred to avian  
548 yolk and may alter offspring growth and adult phenotype. *General and Comparative*  
549 *Endocrinology*, 135(3), 365-371.
- 550 Herrington, J., Vallin, C., & Lickliter, R. (2015). Increased yolk progesterone interferes with  
551 prenatal auditory learning and elevates emotional reactivity in bobwhite quail (*Colinus*  
552 *virginianus*) chicks. *Developmental Psychobiology*, 57(2), 255-262. doi:  
553 10.1002/dev.21274
- 554 Henriksen, R., Groothuis, T. G., & Rettenbacher, S. (2011). Elevated plasma corticosterone  
555 decreases yolk testosterone and progesterone in chickens: linking maternal stress and  
556 hormone-mediated maternal effects. *PLoS ONE*, 6(8), e23824.
- 557 Henriksen, R., Rettenbacher, S., & Groothuis, T. G. G. (2011). Prenatal stress in birds:  
558 Pathways, effects, function and perspectives. *Neuroscience and Biobehavioral*  
559 *Reviews*, 35(7), 1484-1501. doi: 10.1016/j.neubiorev.2011.04.010
- 560 Hess, E. H. (1964). Imprinting in birds. *Science*, 146(3648), 1128-1139.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

561 Hocking, P. M., Haldane, K.-A., Davidson, E. M., Sandøe, P., & Kristensen, H. H. (2015).  
562 Behavioural consequences of visual deprivation occurring at hatch or in the early life  
563 of chickens. *Applied Animal Behaviour Science*, 172, 33-43.

564 Hsu, B. Y., Dijkstra, C., Darras, V. M., de Vries, B., & Groothuis, T. G. G. (2016). Maternal  
565 adjustment or constraint: differential effects of food availability on maternal  
566 deposition of macronutrients, steroids and thyroid hormones in rock pigeon eggs.  
567 *Ecology and Evolution*, 6(2), 397-411. doi: 10.1002/ece3.1845

568 Huang, Z., Leibovitz, H., Lee, C. M., & Millar, R. (1990). Effect of dietary fish oil on.  
569 omega.-3 fatty acid levels in chicken eggs and thigh flesh. *Journal of Agricultural and*  
570 *Food Chemistry*, 38(3), 743-747.

571 Janczak, A. M., Torjesen, P., Palme, R., & Bakken, M. (2007). Effects of stress in hens on the  
572 behaviour of their offspring. *Applied Animal Behaviour Science*, 107(1-2), 66-77. doi:  
573 10.1016/j.applanim.2006.09.016

574 Janczak, A. M., Torjesen, P., & Rettenbacher, S. (2009). Environmental effects on steroid  
575 hormone concentrations in laying hens' eggs. *Acta Agriculturae Scandinavica Section*  
576 *A Animal Science*, 59(2), 80-84. doi: 10.1080/09064700903023348

577 Kingma, S. A., Komdeur, J., Vedder, O., von Engelhardt, N., Korsten, P., & Groothuis, T. G.  
578 G. (2009). Manipulation of male attractiveness induces rapid changes in avian  
579 maternal yolk androgen deposition. *Behavioral Ecology*, 20(1), 172-179. doi:  
580 10.1093/beheco/arn130

581 Lacassagne, L. (1960). Etude comparée des réserves vitellines et de la durée de la phase de  
582 grand accroissement de l'ovocyte chez la poule domestique. Influence de l'âge de  
583 l'animal et de la longueur de la série. *Annales de Zootechnie*, 9(1), 85-96.

- 584 Lafourcade, M., Larrieu, T., Mato, S., Duffaud, A., Sepers, M., Matias, I., . . . Matute, C.  
585 (2011). Nutritional omega-3 deficiency abolishes endocannabinoid-mediated neuronal  
586 functions. *Nature Neuroscience*, 14(3), 345-350.
- 587 Landys, M. M., Piersma, T., Ramenofsky, M., & Wingfield, J. C. (2004). Role of the low-  
588 affinity glucocorticoid receptor in the regulation of behavior and energy metabolism in  
589 the migratory red knot *Calidris canutus islandica*. *Physiological and Biochemical*  
590 *Zoology*, 77(4), 658-668.
- 591 Landys, M. M., Ramenofsky, M., Guglielmo, C. G., & Wingfield, J. C. (2004). The low-  
592 affinity glucocorticoid receptor regulates feeding and lipid breakdown in the migratory  
593 Gambel's white-crowned sparrow *Zonotrichia leucophrys gambelii*. *Journal of*  
594 *Experimental Biology*, 207(1), 143-154.
- 595 Larrieu, T., Hilal, L., Fourrier, C., De Smedt-Peyrusse, V., Sans, N., Capuron, L., & Laye, S.  
596 (2014). Nutritional omega-3 modulates neuronal morphology in the prefrontal cortex  
597 along with depression-related behaviour through corticosterone secretion.  
598 *Translational psychiatry*, 4(9), e437.
- 599 Larrieu, T., Hilal, M. L., De Smedt-Peyrusse, V., Sans, N., & Laye, S. (2016). Nutritional  
600 Omega-3 Deficiency Alters Glucocorticoid Receptor-Signaling Pathway and Neuronal  
601 Morphology in Regionally Distinct Brain Structures Associated with Emotional  
602 Deficits. *Neural Plasticity*. doi: 10.1155/2016/8574830
- 603 Lattin, C. R., & Romero, L. M. (2014). Chronic stress alters concentrations of corticosterone  
604 receptors in a tissue-specific manner in wild house sparrows (*Passer domesticus*).  
605 *Journal of Experimental Biology*, 217(14), 2601-2608.
- 606 Ledón-Rettig, C. C., Richards, C. L., & Martin, L. B. (2013). Epigenetics for behavioral  
607 ecologists. *Behavioral Ecology*, 24(2), 311-324. doi: 10.1093/beheco/ars145

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

608 Maldjian, A., Farkas, K., Noble, R. C., Cocchi, M., & Speake, B. K. (1995). The transfer of  
609 docosaehaenoic acid from the yolk to the tissues of the chick embryo. *Biochimica et*  
610 *Biophysica Acta (BBA)-Lipids and Lipid Metabolism*, 1258(2), 81-89.

611 Mostl, E., Spendier, H., & Kotrschal, K. (2001). Concentration of immunoreactive  
612 progesterone and androgens in the yolk of hens' eggs (*Gallus domesticus*). *Wiener*  
613 *Tierarztliche Monatsschrift*, 88(3), 62-65.

614 Muller, W., Eising, C. M., Dijkstra, C., & Groothuis, T. G. G. (2002). Sex differences in yolk  
615 hormones depend on maternal social status in Leghorn chickens (*Gallus gallus*  
616 *domesticus*). *Proceedings of the Royal Society of London Series B-Biological*  
617 *Sciences*, 269(1506), 2249-2255. doi: 10.1098/rspb.2002.2159

618 Müller, W., Heylen, D., Eens, M., Rivera-Gutierrez, H. F., & Groothuis, T. G. G. (2013). An  
619 experimental study on the causal relationships between (ecto-)parasites, testosterone  
620 and sexual signalling. *Behavioral Ecology and Sociobiology*, 67(11), 1791-1798. doi:  
621 10.1007/s00265-013-1586-6

622 Murphy, L. B. (1977). Responses of domestic fowl to novel food and objects. *Applied Animal*  
623 *Ethology*, 3(4), 335-349.

624 Nätt, D., Lindqvist, N., Stranneheim, H., Lundeberg, J., Torjesen, P. A., & Jensen, P. (2009).  
625 Inheritance of Acquired Behaviour Adaptations and Brain Gene Expression in  
626 Chickens. *PLoS ONE*, 4(7), e6405.

627 Noble, R., & Cocchi, M. (1989). The relationship between the supply and demand for  
628 essential polyunsaturated fatty acids during mammalian and avian embryonic  
629 development. *Research and Development in Agriculture (United Kingdom)*.

630 Noble, R., & Cocchi, M. (1990). Lipid metabolism and the neonatal chicken. *Progress in*  
631 *Lipid Research*, 29(2), 107-140.

- 632 Palme, R., Touma, C., Arias, N., Dominchin, M., & Lepschy, M. (2013). Steroid extraction:  
 633 get the best out of faecal samples. *Wiener Tierarztliche Monatsschrift*, 100(9-10), 238-  
 634 246.
- 635 Pilz, K. M., & Smith, H. G. (2004). Egg yolk androgen levels increase with breeding density  
 636 in the European Starling, *Sturnus vulgaris*. *Functional Ecology*, 18(1), 58-66.
- 637 Rettenbacher, S., Mostl, E., Hackl, R., Ghareeb, K., & Palme, R. (2004). Measurement of  
 638 corticosterone metabolites in chicken droppings. *British Poultry Science*, 45, 704-711.  
 639 doi: 10.1080/00071660400006156
- 640 Rettenbacher, S., & Palme, R. (2009). Biological validation of a non-invasive method for  
 641 stress assessment in chickens. *Berliner Und Munchener Tierarztliche Wochenschrift*,  
 642 122(1-2), 8-12. doi: 10.2376/0005-9366-122-8
- 643 Rodenburg, T. B., & de Haas, E. N. (2016). Of nature and nurture: the role of genetics and  
 644 environment in behavioural development of laying hens. *Current Opinion in*  
 645 *Behavioral Sciences*, 7, 91-94.
- 646 Ruuskanen, S. (2015). Hormonally-mediated maternal effects in birds: lessons from the  
 647 flycatcher model system. *General and Comparative Endocrinology*, 224, 283-293.
- 648 Ruuskanen, S., Darras, V. M., de Vries, B., Visser, M. E., & Groothuis, T. G. G. (2016).  
 649 Experimental manipulation of food availability leads to short-term intra-clutch  
 650 adjustment in egg mass but not in yolk androgen or thyroid hormones. *Journal of*  
 651 *Avian Biology*, 47(1), 36-46. doi: 10.1111/jav.00728
- 652 Sandell, M. I., Adkins-Regan, E., & Ketterson, E. D. (2007). Pre-breeding diet affects the  
 653 allocation of yolk hormones in zebra finches *Taeniopygia guttata*. *Journal of Avian*  
 654 *Biology*, 38(3), 284-290. doi: 10.1111/j.2007.0908-8857.03640.x
- 655 Schwabl, H. (1997). The contents of maternal testosterone in house sparrow *Passer*  
 656 *domesticus* eggs vary with breeding conditions. *Naturwissenschaften*, 84(9), 406-408.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

657 Tobler, M., & Sandell, M. I. (2007). Yolk testosterone modulates persistence of neophobic  
658 responses in adult zebra finches, *Taeniopygia guttata*. *Hormones and Behavior*, 52(5),  
659 640-645. doi: 10.1016/j.yhbeh.2007.07.016

660 Van Elswyk, M. E. (1997). Comparison of n-3 fatty acid sources in laying hen rations for  
661 improvement of whole egg nutritional quality: a review. *British Journal of Nutrition*,  
662 78(01), S61-S69.

663 Verboven, N., Monaghan, P., Evans, D. M., Schwabl, H., Evans, N., Whitelaw, C., & Nager,  
664 R. G. (2003). Maternal condition, yolk androgens and offspring performance: a  
665 supplemental feeding experiment in the lesser black-backed gull (*Larus fuscus*).  
666 *Proceedings of the Royal Society B-Biological Sciences*, 270(1530), 2223-2232. doi:  
667 10.1098/rspb.2003.2496

668 Verboven, N., Monaghan, P., Nager, R. G., & Evans, N. P. (2010). The Effect of Maternal  
669 State on the Steroid and Macronutrient Content of Lesser Black-Backed Gull Eggs.  
670 *Physiological and Biochemical Zoology*, 83(6), 1009-1022. doi: 10.1086/656568

671 Vergauwen, J., Eens, M., & Muller, W. (2012). Pre- and postnatal environmental effects as  
672 potential sources of variation in neophobic behaviour in canaries. *Behaviour*, 149(8),  
673 849-867. doi: 10.1163/1568539x-00003006

674 Vergauwen, J., Goerlich, V. C., Groothuis, T. G., Eens, M., & Müller, W. (2012). Food  
675 conditions affect yolk testosterone deposition but not incubation attendance. *General  
676 and Comparative Endocrinology*, 176(1), 112-119.

677 Webster, S. J., & Lefebvre, L. (2001). Problem solving and neophobia in a columbiform-  
678 passeriform assemblage in Barbados. *Animal Behaviour*, 62(1), 23-32.

679 Whitehead, C. C. (1995). Plasma oestrogen and the regulation of egg weight in laying hens by  
680 dietary fats. *Animal Feed Science and Technology*, 53(2), 91-98.



681 Yates, C. M., Calder, P. C., & Rainger, G. E. (2014). Pharmacology and therapeutics of  
682 omega-3 polyunsaturated fatty acids in chronic inflammatory disease. *Pharmacology  
683 & Therapeutics*, 141(3), 272-282.

684 Zimmer, C., & Spencer, K. (2014). Modifications of Glucocorticoid Receptors mRNA  
685 Expression in the Hypothalamic-Pituitary-Adrenal Axis in Response to Early-life  
686 Stress in Female Japanese Quail. *Journal of Neuroendocrinology*, 26(12), 853-860.

For Peer Review



Table 1: Mean ( $\pm$  SE) body mass, food intake and laying rate of adult n-3 hens ( $N = 18$ ) and control hens ( $N = 18$ ) during dietary treatment.

Parameters	Hens	Week 1	Week 2	Week 3	Week 4	Week 5
Body mass (g)	n-3 enriched	1958.7 $\pm$ 33.9	1854.5 $\pm$ 40.6	1863.8 $\pm$ 37.3	1835.3 $\pm$ 35.3	1856.6 $\pm$ 36.5
	controls	1966.6 $\pm$ 67.2	1880.9 $\pm$ 60.9	1974.1 $\pm$ 62.5	1865.4 $\pm$ 60.8	1873.5 $\pm$ 62.3
24h feed intake (g)	n-3 enriched	154.7 $\pm$ 23.4	144.7 $\pm$ 7.4	173.7 $\pm$ 16.7	194.6 $\pm$ 22.8	224.4 $\pm$ 28.1
	controls	133.7 $\pm$ 11.8	169.6 $\pm$ 21.1	161.8 $\pm$ 20.5	219.0 $\pm$ 24.5	265.4 $\pm$ 24.3
Laying rate (number per day)	n-3 enriched	0.78 $\pm$ 0.07	0.89 $\pm$ 0.03	0.91 $\pm$ 0.02	0.78 $\pm$ 0.04	0.69 $\pm$ 0.06
	controls	0.71 $\pm$ 0.07	0.82 $\pm$ 0.06	0.77 $\pm$ 0.07	0.69 $\pm$ 0.07	0.65 $\pm$ 0.06

Table 2: Mean ( $\pm$  SE) percentages of n-3 and n-6 poly-unsaturated fatty acids in the yolk of eggs of hens fed a diet enriched in n-3 ( $N = 18$ ) or a control diet ( $N = 18$ ).

Poly Unsaturated fatty acids	Diets	
	n-3 enriched	Control
<b><i>n-6 fatty acids</i></b>		
Linoleic acid (LA), C18:2 n-6	13.5 % $\pm$ 0.23 <sup>a</sup>	18.9 % $\pm$ 0.47 <sup>b</sup>
Arachidonic acid (AA), C20:4 n-6	0.64 % $\pm$ 0.05 <sup>a</sup>	1.81 % $\pm$ 0.18 <sup>b</sup>
Docosatetraenoic acid (DTA), C22:4 n-6	0.03 % $\pm$ 0.005 <sup>a</sup>	0.22 % $\pm$ 0.02 <sup>b</sup>
<b><i>n-3 fatty acids</i></b>		
$\alpha$ -linolenic acid (ALA), C18:3 n-3	0.47 % $\pm$ 0.02 <sup>a</sup>	0.71 % $\pm$ 0.03 <sup>b</sup>
Eicosapentaenoic acid (EPA), C20:5 n-3	0.22 % $\pm$ 0.02 <sup>a</sup>	0.00 % $\pm$ 0.0 <sup>b</sup>
Docosapentaenoic acid (DPA), C22:5 n-3	0.15 % $\pm$ 0.01 <sup>a</sup>	0.07 % $\pm$ 0.02 <sup>b</sup>
Docosahexaenoic acid (DHA), C22:6 n-3	1.72 % $\pm$ 0.17 <sup>a</sup>	0.74 % $\pm$ 0.07 <sup>b</sup>
Ratio n-6/n-3	6.15 % $\pm$ 0.53 <sup>a</sup>	13.9 % $\pm$ 0.42 <sup>b</sup>

Means between columns with a different superscript (a, b) differ ( $p < 0.05$ )


1  
2  
3  
4  
5728Table 3: Mean ( $\pm$  SE) Latency to eat and time spent eating during neophobia tests of n-3 chicks ( $N = 24$  pairs) and control chicks ( $N = 25$  pairs)  
6  
7729and Mann-Whitney  $U$ -test outcome.  
8  
9

10730  
11  
12  
13731  
14  
15

Neophobia Tests	Parameters Measured	n-3 chicks	control chicks	$Z$	$p$ - values
Habituation	latency to eat (s)	40.5 $\pm$ 3.3	48.3 $\pm$ 6.5	-0.66	0.51
	time spent eating (s)	48.9 $\pm$ 3.8	52.1 $\pm$ 5.2	-1.07	0.28
Novel food	latency to eat (s)	99.1 $\pm$ 13.2	85.4 $\pm$ 10.6	-0.30	0.76
Test 1	time spent eating (s)	18.4 $\pm$ 3.7	22.8 $\pm$ 3.8	-0.90	0.36
Novel food	latency to eat (s)	84.3 $\pm$ 13.9	107.8 $\pm$ 10.2	-0.96	0.33
Test 2	time spent eating (s)	32.0 $\pm$ 9.1	24.2 $\pm$ 5.9	-0.15	0.88
Novel Object	latency to eat (s)	174.1 $\pm$ 5.9	155.2 $\pm$ 8.5	-2.11	0.03
	time spent eating (s)	2.1 $\pm$ 2.0	14.11 $\pm$ 5.1	-2.27	0.02

27732  
28  
29  
30733  
31  
32734  
33  
34  
35  
36735  
37  
38  
39736  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

737 FIGURE 1



738

739

740

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

741 **Figure caption**

742 **FIGURE 1:** Mean ( $\pm$  SE) yolk progesterone (P4), androstenedione (A4), estrogens (E) and  
743 testosterone (T) concentrations (ng/g of yolk) in the eggs from n-3 enriched hens ( $N = 18$ ) and  
744 control hens ( $N = 18$ ). \*\*  $p < 0.01$ ; \*  $p \leq 0.05$ ; #  $0.1 \leq p \leq 0.05$

For Peer Review