

New Seco-limonoids from *Cipadessa baccifera*: Isolation, Structure Determination, Synthesis and Their Antiproliferative Activities.

Bandi Siva,^a A. Venkanna,^a B. Poornima,^a S. Divya Reddy,^a J. Boustie,^b Schnell Bastien,^{a,b} Nishant Jain,^c P.Usha Rani,^d K.Suresh Babu,^{a,*}

^aNatural Products Laboratory, Division of Natural Product Chemistry, CSIR-Indian Institute of Chemical Technology, Hyderabad 500 607, India

^bUMR CNRS 6226 ISCR PNSCM, Université de Rennes 1, 2 Avenue du Professeur Léon Bernard, 35043 Rennes, France.

^cCentre for Chemical Biology, CSIR-Indian Institute of Chemical Technology, Hyderabad 500 007, India

^d Biology division, CSIR-Indian Institute of Chemical Technology, Hyderabad-500 007.

Table of Contents

Experimental values of **9-13**

FIGURE1: HRESIMS,IR AND ¹HNMR SPECTRUM OF COMPOUND **1**

FIGURE2: ¹³C NMR & DEPT135° SPECTRUM OF COMPOUND **1**

FIGURE3: HSQC SPECTRUM OF COMPOUND **1**

FIGURE4: HMBC SPECTRUM OF COMPOUND **1**

FIGURE5: COSY SPECTRUM OF COMPOUND **1**

FIGURE6: NOESY SPECTRUM OF COMPOUND **1**

FIGURE7: HRESIMS,IR AND ¹HNMR SPECTRUM OF COMPOUND **2**

FIGURE8: ¹³C NMR & DEPT135° SPECTRUM OF COMPOUND **2**

FIGURE9: HSQC SPECTRUM OF COMPOUND **2**

FIGURE10: HMBC SPECTRUM OF COMPOUND **2**

FIGURE11: COSY SPECTRUM OF COMPOUND **2**

FIGURE12: NOESY SPECTRUM OF COMPOUND **2**

- FIGURE13:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **9**
- FIGURE14:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **10a**
- FIGURE15:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND**10b**
- FIGURE16:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **10c**
- FIGURE17:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **11a**
- FIGURE18:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **11b**
- FIGURE19:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **11c**
- FIGURE20:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **11d**
- FIGURE21:** ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **11e**
- FIGURE22:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **12**
- FIGURE23:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **13a**
- FIGURE24:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **13b**
- FIGURE25:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **13c**
- FIGURE26:** ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **13d**
- FIGURE27:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **13e**
- FIGURE28:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **13f**
- FIGURE29:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **13g**
- FIGURE30:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **13h**
- FIGURE31:** HRESIMS, ^1H NMR & ^{13}C NMR SPECTRUM OF COMPOUND **13i**

1. Experimental

1.1. Methyl 2-((4S,4aS,7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-10-hydroxy-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-8-yl)acetate (9). Yield: 70%, time: 4h; white amorphous powder; $[\alpha]_D^{25} - 26.1$ (*c* 1.1, CHCl₃) IR(KBr): ν_{max} 3426, 2978, 1732, 1728, 1683, 1372, 1245, 1037 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ : 7.46 (1H, s, H-21), 7.39 (1H, s, H-23), 6.41(1H, s, H-22), 5.79 (1H, s, H-17), 5.09 (1H, s, H_a-30), 4.84 (1H, s, H_b-30), 3.69 (1H, d, *J* = 10.9 Hz, H-3), 3.70(3H, s, OMe), 3.29 (1H, t, *J* = 3.2 & 3.0 Hz, H-1), 2.90 (1H, d, *J* = 17.9 Hz, H_a-15), 2.59 (1H, d, *J* = 17.9 Hz, H_b - 15), 2.55 (1H, dd, *J* = 11.8 & 5.0 Hz, H-5), 2.50 (1H, OL, H_a-6), 2.09 (1H, d, *J* = 15.2 Hz, H_b-6), 2.18 (1H, br s, H-9), 2.07 (1H, m, H-11), 1.84 (2H, m, H-2), 1.85(1H, m, H_a-12) & 1.15 (1H, m, H_b-12), 1.53 (1H, m), 0.95 (1H, s, H-29), 0.89 (3H, s,H-18), 0.87(3H, s, H-19), 0.80 (1H, s, H-28); ¹³C NMR (75 MHz, CDCl₃) δ : 174.5(C-7), 170.4 (C-16), 146.4 (C-8), 142.7 (C-23), 140.6 (C-21), 120.9 (C-20), 110.8 (C-22), 109.8 (C-30), 79.9 (C-14), 77.5 (C-17), 74.9(C-3), 72.6 (C-1), 51.9 (C-9), 50.4 (C-10), 43.8 (C-12), 41.4 (C-13), 39.7 (C-4), 33.9 (C-15), 32.6 (C-6), 32.4 (C-6), 30.9 (C-2), 29.5 (C-11), 27.0 (C-28), 23.8 (C-5), 21.9(C-19), 14.38 (C-29), 13.8 (C-18); HRMS (ESI+) *m/z*: 473.2527 [M+H]⁺ (calcd for C₂₇H₃₇O₇[M+H]⁺, 473.2533).

1.2.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl butyrate (10a): Yield: 90%, time: 3h, colorless gum; $[\alpha]_D^{25} - 34.2$ (*c* 1.1, CHCl₃); IR (KBr): ν_{max} 2968, 1737, 1718, 1685, 1325, 1235, 1027 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ : 7.49 (1H, s, H-21), 7.39 (1H, s, H-23), 6.43 (1H, s, H-22), 5.80 (1H, s, H-17), 5.00 (1H, dd, *J* = 10.0 & 6.6

Hz, H-3), 5.09 (1H, s, H_a-30), 4.85 (H, s, H_b-30), 3.69 (3H, s, OMe), 3.24 (1H, t, *J* = 3.2&2.8 Hz, H-1), 2.90 (1H, d, *J* = 18.1 Hz, H_a-15) & 2.57 (1H, d, *J* = 18.1Hz, H_b-15), 2.54 (1H, dd, *J* = 11.6 & 5.1 Hz, H-5), 2.51(1H, overlapping, H_a-6) & 2.15 (1H, d, *J* = 15.1Hz, H_b-6), 2.23 (2H, m, butyryl -2'), 2.08 (1H, br s,H-9), 2.02 (1H, m, H-11_a) & 1.54 (1H, m, H-11_b), 1.89 (2H, m, H-2), 1.87 (2H, m, butyryl -3¹), 1.86 (1H, m, H_a-12) &1.31 (1H, m, H_b-12), 1.15 (3H, t, *J* = 7.5 Hz, butyryl -4'), 0.95 (1H, s,H-29), 0.87 (3H, s, H-18), 0.83 (3H, s, H-19), 0.80 (1H, s,H-28); ¹³ C NMR (75 MHz, CDCl₃) δ: 174.2 (C-7), 172.7 (C- butyryl-1'), 170.3 (C-16), 146.4 (C-8), 142.5 (C-23), 140.8 (C-21), 120.9 (C-20), 110.9 (C-22), 110.0 (C-30), 79.97 (C-14), 79.79 (C-17), 74.70 (C-3), 74.14 (C-1), 51.91 (C-9), 50.43 (C-10), 43.8 (C-12), 41.5 (C-13), 38.7 (C-4), 33.8 (C-15), 32.3 (C-6) , 30.9 (butyryl -2¹), 29.6 (C-2), 29.3 (C-11), 26.9 (C-28) , 23.8 (C-5) , 22.6(butyryl -3¹), 21.8 (C-19), 18.5 (butyryl -4¹), 15.6 (C-29), 13.8(C-18); HRMS (ESI+) *m/z*: 565.2781 [M+Na]⁺(calcd for C₃₁H₄₂O₈Na [M+Na]⁺, 565.2771).

1.3 (7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12amethanobenzo[b]pyrano[3,4-g]oxocin-10-yl propionate (10b). Yield: 85%, time: 2h, colorless gum; [α]_D²⁵ – 48.5 (*c* 0.9, CHCl₃); IR(KBr): ν_{max} 2974, 1734, 1728, 1682, 1334, 1238, 1035 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ: 7.49 (1H, s, H-21), 7.39 (1H, s, H-23), 6.43 (1H, s, H-22), 5.80 (1H, s, H-17), 5.09 (1H, s, H_a-30), 5.00 (1H, dd, *J* = 10.0 & 6.6, H-3), 4.85 (1H, s, H_b-30), 3.69(3H, s, OMe), 3.23 (1H, t, *J* = 3.28 & 2.8 Hz, H-1), 2.90 (1H, d, *J* = 18.1Hz, H_a-15) & 2.57 (1H, d, *J* = 18.1Hz, H_b-15), 2.54 (1H, dd, *J* = 11.6 & 5.1Hz, H-5), 2.51(1H, overlapping, H_a-6) & 2.15 (1H, d, *J* = 15.1Hz, H_b-6), 2.23 (2H, q, *J* = 7.5 & 15.1Hz, propionyl H-2'), 2.08 (1H, br s, H-9), 2.02 (1H, m, H_a-11) & 1.54 (1H, m, H_b-11), 1.88 (2H, m, H-2), 1.86 (1H, m, Ha-12) & 1.31 (1H, m, H_b-12), 1.15 (3H, t, *J* = 7.5Hz, propionyl H-3'), 0.95 (1H, s, H-29), 0.87 (3H, s, H-18), 0.83(3H, s,H-19), 0.80 (1H, s, H-28); ¹³ C NMR

(75 MHz, CDCl₃) δ: 174.2 (C-7), 173.6 (propionyl C-1'), 170.4 (C-16), 146.4 (C-8), 142.5 (C-23), 140.8 (C-21), 120.8 (C-20), 110.9 (C-22), 110.0 (C-30), 79.9 (C-14), 79.7 (C-17), 74.7 (C-3), 74.2 (C-1), 51.9 (C-9), 50.4 (C-10), 43.8 (C-12), 41.5 (C-13), 38.7 (C-4), 33.8 (C-15), 32.3 (C-6), 29.5 (C-2), 29.4 (C-11), 27.8 (propionyl C-2'), 26.9 (C-28), 23.8 (C-5), 21.8 (C-19), 15.6 (C-28), 13.8 (C-18), 9.21 (propionyl C-3'); HRMS (ESI+) *m/z*: 529.2774 [M+H]⁺ (calcd for C₃₀H₄₁O₈ [M+H]⁺, 529.2795).

1.4 (7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl isobutyrate (10c): Yield: 90%, time: 3h; colorless gum, $[\alpha]_D^{25} - 54.2$ (*c* 0.7, CHCl₃); IR(KBr):

ν_{max} 2994, 1737, 1722, 1692, 1324, 1231, 1031 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ: 7.50 (1H, s, H-21), 7.39 (1H, s, H-23), 6.43(1H, s, H-22), 5.81 (1H, s, H-17), 5.09 (1H, s, H_a-30), 4.98 (1H, dd, *J* = 10.9 & 5.6Hz, H-3), 4.85 (1H, s, H_b-30), 3.69(3H, s, OMe), 3.23 (1H, t, *J* = 3.28 & 2.8 Hz, H-1), 2.55(1H, dd, *J* = 13.6 & 6.1 Hz, H-5), 2.91 (1H, d, *J* = 18.1Hz, H_a-15), 2.58 (1H, d, *J* = 18.1Hz, H_b-15), 2.51(1H, OL, H_a-6) & 2.14 (1H, d, *J* = 15.1Hz, H_b-6), 2.05 (1H, m, H-2¹), 2.04 (1H, m, H_a-11) & 1.56 (1H, m, H_b-11), 2.03 (1H, br s, H-9), 1.90 (2H, m, H-2), 1.88 (1H, m, H_a-12) & 1.32 (1H, m, H_b-12), 1.18 (6H, d, *J* = 6.9 Hz, isobutyryl -3' & 4'), 0.97 (1H, s, H-29), 0.87 (3H, s, H-18), 0.83 (3H, s, H-19), 0.80 (3H, s, H-28); ¹³C NMR (75 MHz, CDCl₃) δ: 176.1 (C-7), 174.2 (isobutyryl -1'), 170.3 (C-16), 146.4(C-8), 142.5 (C-23), 140.8 (C-21), 120.9 (C-20), 110.9 (C-22), 110.0 (C-30), 79.9 (C-14), 79.8 (C-17), 74.7 (C-3), 73.9 (C-1), 51.9 (C-9), 50.4 (C-10), 43.8 (C-12), 41.5 (C-13), 38.8 (C-4), 33.8 (C-15), 32.3 (C-6), 31.9 (isobutyryl -2'), 29.6 (C-2), 29.5 (C-11), 26.8 (C-26), 23.8 (C-5), 21.8 (C-21), 19.1 (C-3'), 18.8 (isobutyryl -3', 4'), 15.6 (C-29), 13.84 (C-18); HRMS (ESI+) *m/z*: 543.2934 [M+H]⁺ (calcd for C₃₁H₄₃O₈ [M+H]⁺, 543.2952).

1. 5. (7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 2-chloroacetate (12). Yield: 79%, time: 3h, colorless oil; $[\alpha]_D^{25} - 32.7$ (*c* 1.2, CHCl₃); IR(KBr): ν_{max} 2992, 1731, 1725, 1698, 1324, 1237, 1034 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ : 7.49 (1H, bs, H-21), 7.39 (1H, bs, H-23), 6.43(1H, bs, H-22), 5.79 (1H, s, H-17), 5.11(1H, H_b-30), 5.05 (1H, dd, *J* = 10.5 & 5.8Hz, H-3), 4.87(1H, s, H_a-30), 4.06 (2H, chloroacetyl-H-2¹), 3.69(3H, s, OMe), 3.25 (1H, t, *J* = 3.2 & 6.0 Hz, H-1), 2.91 (1H, d, *J* = 18.3Hz, H_a-15) & 2.5 (1H, d, *J* = 18.3Hz, H_b-15), 2.59(1H, d, *J* = 15.1Hz, Ha-6) & 2.12 (1H, d, *J* = 15.1Hz, H_b-6), 2.54 (1H, m, H-5), 2.22 (1H, m, Ha-12)& 1.18 (1H, m, H_b-12), 2.09 (1H, br s,H-9), 2.07 (2H, m, H-2), 1.94 (1H, m, H_a-11) & 1.52 (1H, m, H_b-11), 0.99 (1H, s, H-29), 0.87 (3H, s, H-18), 0.83 (1H, s, H-19), 0.83(1H, s, ol, H-28); ¹³C NMR (75 MHz, CDCl₃) δ : 174.1 (C-7), 170.6 (C-16), 166.5 (chloroacetyl-1¹), 146.1 (C-8), 142.6(C-23), 140.8 (C-21), 120.7 (C-20), 111.1 (C-22), 109.9 (C-30), 80.0 (C-14), 79.8 (C-17), 77.5 (C-3), 74.5 (C-1), 52.0(C-9), 43.8(C-10), 41.4(C-12), 41.0(C-13), 40.4(chloroacetyl-C-2¹), 38.9(C-14), 33.82 (C-15), 32.28 (C-6), 29.41 (C-2), 29.31 (C-11), 26.87 (C-28), 23.82 (C-5), 21.89 (C-19), 15.5(C-29), 13.8 (C-18); HRMS (ESI+) *m/z*: 571.2094M+Na]⁺ (calcd for C₂₉H₃₇Cl O₈Na [M+Na]⁺, 571.2069).

1. 6.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 4-cyanobenzoate (11a). Yield: 70%, time: 11h; light yellow gum; $[\alpha]_D^{25} - 19.2$ (*c* 0.6, CHCl₃); IR(KBr): ν_{max} 3094, 1736, 1723, 1698, 1336, 1239, 1036 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ : 8.11 (2H, d, *J* = 8.5 Hz, Ar-3''& Ar-5''), 7.75 (2H, d, *J* = 8.5Hz, Ar-2''&Ar-6''), 7.52(1H, s, H-21), 7.41 (1H, s, H-23), 6.45(1H, s, H-22), 5.84 (1H, s, H-17), 5.12(1H, H_b-30), 5.24 (1H, dd, *J* = 10.5 & 6.2 Hz, H-3), 4.87(1H, s, H_a-30), 3.70(3H, s, OMe), 3.30 (1H, t, *J* = 3.2 & 2.8 Hz, H-

1), 2.92 (1H, d, $J = 18.1$ Hz, H_a-15) & 2.59 (1H, d, $J = 18.1$ Hz, H_b-15), 2.54 (1H, dd, $J = 13.6$ & 6.1Hz, H-5), 2.63 (1H, d, $J = 15.1$ Hz, H_a-6), 2.01 (2H, m), 2.24 (1H, m, H_a-12) & 1.32 (1H, m, H_b-12), 2.12 (1H, d, $J = 15.1$, H_b-6), 2.05 (1H, m, H_a-11) & 1.24 (1H, m, H_b-11), 2.02 (1H, br s, H-9), 0.89 (3H, s, H-18), 0.88 (3H, s, H-19), 0.88 (3H, s, ol, H-28), 1.10 (1H, s, H-29); ¹³C NMR (75 MHz, CDCl₃) δ: 174.1 (C-7), 170.2 (C-16), 164.0 (ester carbonyl C-1'), 146.2 (C-8), 142.6 (C-23), 140.8 (C-21), 134.28 (Ar-4''), 132.26 (Ar-1''), 127.04 (Ar-3'' & Ar-5''), 120.9 (C-20), 116.36 (Ar-2''&Ar-6''), 111.1 (C-22), 109.9 (C-30), 80.1 (C-14), 79.8 (C-17), 77.4 (C-3), 74.6 (C-1), 52.0 (C-9), 50.4 (C-10), 43.9 (C-12), 41.5 (C-13), 39.0 (C-4), 33.8 (C-15), 32.3 (C-6), 29.41 (C-2), 29.68 (C-11), 27.08 (C-28), 23.88 (C-5), 21.94 (C-19), 15.9 (C-29), 13.8 (C-18); MS (ESI+) *m/z*: 602.2 [M+H]⁺.

1.7.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 4-chlorobenzoate (11b). Yield: 65%, Time: 12h; light yellow gum; $[\alpha]_D^{25} - 24.1$ (*c* 0.8, CHCl₃) IR(KBr): ν_{max} 3097, 1731, 1721, 1694, 1335, 1232, 1031 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ: 7.95 (4H, d, $J = 8.6$ Hz, Ar H-3''& H-5''), 7.42 (4H, d, $J = 8.5$ Hz, Ar H-2''& H-6''), 7.52 (1H, s, H-21), 7.40 (1H, s, H-23), 6.45(1H, s, H-22), 5.12(1H, H_b-30), 5.85 (1H, s, H-17), 5.21 (1H, dd, $J = 10.5$ & 6.2 Hz, H-3), 4.87(1H, s, H_a-30), 3.70(3H, s, OMe), 3.29 (1H, t, $J = 3.28$ & 2.30 Hz, H-1), 2.92 (1H, d, $J = 18.1$ Hz, H_b-15), 2.58 (1H, d, $J = 18.1$ Hz, H_a-15), 2.64(1H, d, $J = 15.1$ Hz, H_a-6) & 2.14 (1H, d, $J = 15.1$ Hz, H_b-6), 2.53 (1H, dd, $J = 13.6$ & 6.1Hz, H-5), 2.26 (1H, m, H_a-12) & 1.31 (1H, m, H_b-12), 2.06 (1H, m, H_a-11) & 1.22 (1H, m, H_b-11), 2.01 (1H, br s, H-9), 2.00 (2H, m, H-2), 1.09 (1H, s, H-29), 0.88 (3H, s, H-18), 0.87 (3H, s, H-19), 0.87(3H, s, ol, H-28); ¹³C NMR (75 MHz, CDCl₃) δ: 174.2 (C-7), 170.3 (C-16), 164.8 (ester carbonyl- C-1'), 146.3 (C-8), 142.6 (C-23), 140.8 (C-21), 139.3 (Ar-4''), 130.8 (Ar-3''& 5''), 128.8 (Ar-2'' & 6''), 127.8 (Ar-

1"), 120.9 (C-20), 111.0 (C-21), 110.0 (C-22), 80.0 (C-14), 79.8 (C-17), 75.6 (C-3), 74.7 (C-1), 51.9 (C-9), 50.4 (C-10), 43.8 (C-12), 41.5 (C-13), 39.0 (C-4), 33.8 (C-15), 32.3 (C-6), 29.6 (C-2), 29.3 (C-11), 27.0 (C-28), 23.8 (C-5), 21.9 (C-19), 15.9 (C-29), 13.8 (C-18); HRMS (ESI+) m/z 633.2249 [M+Na]⁺ (calcd for C₃₄H₃₉O₈ClNa [M+Na]⁺, 633.2225).

1. 8. (7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 4-(trifluoromethyl)benzoate (11c). Yield: 70%, time: 12h, Yellow gum; $[\alpha]_D^{25} = -41.3$ (*c* 0.9, CHCl₃); IR(KBr): ν_{max} 3087, 1734, 1728, 1691, 1331, 1233, 1033cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ : 8.13 (2H, d, *J* = 8.1Hz, Ar-3"&Ar-5"), 7.72 (2H, d, *J* = 8.1Hz, Ar-2"&Ar-6"), 7.53 (1H, s, H-21), 7.41 (1H, s, H-23), 6.45(1H, s, H-22), 5.86 (1H, s, H-17), 5.12(1H, H_b-30), 5.25 (1H, dd, *J* = 9.1 & 7.5Hz, H-3), 4.87(1H, s, H_a-30), 3.69(3H, s, OMe), 3.30 (1H, t, *J* = 3.2 & 2.8Hz, H-1), 2.92 (1H, d, *J* = 18.1Hz, H_a-15), 2.60 (1H, d, *J* = 18.1Hz, H_b-15), 2.57(1H, d, *J* = 16.0Hz, H_a-6) & 2.12 (1H, d, *J* = 16 Hz, H_b-6), 2.54 (1H, dd, *J* = 17.3& 7.1Hz, H-5), 2.22 (1H, m, H_a-12) & 1.33 (1H, m,H_b-12), 2.04 (1H, br s, H-9), 2.04 (2H, m, H-2), 2.03 (1H, m, H_a-11), 1.41(1H, m, H_b-11), 1.11 (1H, s, H-29), 0.89 (3H, s,H-18), 0.88 (3H, s,H-19), 0.88 (3H, s, OL, H-28); ¹³C NMR (75 MHz, CDCl₃) δ : 174.1(C-7), 170.2 (C-16), 164.4 (ester carbonyl-C-1¹), 146.2 (C-8), 142.6 (C-23), 140.8 (C-21), 133.7 (Ar-4"), 129.8 (Ar-3"& Ar-5"), 125.4 (Ar-2"& Ar- 6"), 125.4 (Ar-1"), 120.9 (C-20), 111.0 (C-22), 110.0 (C-30), 80.1 (C-14), 79.8 (C-17), 76.0 (C-3), 74.7 (C-1), 57.8 (C-9), 51.9 (C-10), 43.9 (C-12), 41.5 (C-13), 39.0 (C-4), 33.8 (C-15), 32.3 (C-6), 31.9 (C-2), 29.4 (C-11), 27.0 (C-28), 23.9 (C-5), 21.9 (C-19), 15.9 (C-29), 13.8 (C-18); HRMS (ESI+) m/z : 667.2519 [M+Na]⁺ (calcd for C₃₅H₃₉F₃O₈Na [M+Na]⁺, 667.2489).

1. 9. (7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 2,4-

dinitrobenzoate (11d). Yield: 85%, time: 10h; yellow gum $[\alpha]_D^{25} - 21.3$ (*c* 0.8, CHCl₃); IR(KBr): ν_{max} 2997, 1741, 1732, 1701, 1337, 1237, 1033cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ : 9.24 (1H, t, *J* = 2.2Hz, Ar-3"), 9.13 (1H, bs, Ar-5"), 9.13 (1H, bs, Ar-6"), 7.53 (1H, s, H-21), 7.42 (1H, s, H-23), 6.46(1H, s, H-22), 5.83(1H, s, H-17), 5.34 (1H, dd, *J* = 12 & 4.5 Hz, H-3), 5.14 (1H, H_b-30), 4.89 (1H, s, H_a-30), 3.70 (3H, s, OMe), 3.34 (1H, t, *J* = 3.28 & 2.3Hz, H-1), 2.94 (1H, d, *J* = 18.1Hz, H_a-15)& 2.61 (1H, d, *J* = 18.1Hz, H_b-15), 2.62(1H, d , *J* = 15.1Hz, H_a-6) & 2.14 (1H, d, *J* = 15.1Hz, H_b-6), 2.55 (1H, dd, *J* = 13.6 & 6 Hz, H-5), 2.28 (1H, m,Ha-12) & 1.35 (1H, m, H_b-12), 2.07 (2H, m, H-2), 2.06 (1H, m, Ha-11) &1.23 (1H, m, H_b-11), 2.05 (1H, br s H-9), 1.09 (1H, s,H_b-29), 0.88 (3H, s,H_a-18), 0.87 (3H, s, H_b-19), 0.87(1H, s, ol, H_a-28), ¹³C NMR (75 MHz, CDCl₃) δ : 174.0(C-7), 170.2(C-16), 161.6 (ester carbonyl C-1'), 153.1(Ar-4"), 148.6 (Ar-2"), 145.9(C-8), 142.7 (C-23), 140.0 (C-21), 134.08 (Ar-1"), 131.5 (Ar-6"), 129.3 (Ar-5"), 122.3 (Ar-3"), 120.8 (C-20), 111.3 (C-22), 109.9 (C-30), 80.2 (C-14), 79.8 (C-17), 77.9 (C-3), 74.5 (C-1), 52.0 (C-9), 50.3 (C-10), 43.8 (C-12), 41.4 (C-13), 39.1 (C-4), 33.8 (C-15), 32.2 (C-6), 29.6 (C-2), 29.4 (C-11), 27.1(C-28), 23.8 (C-5), 21.9 (C-19), 16.0 (C-29), 13.9 (C-18). HRMS (ESI+) *m/z*: 667.2475 [M+H]⁺ (calcd for C₃₄H₃₉N₂O₁₂ [M+H]⁺, 667.2497).

1.10.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl thiophene-2-carboxylate (11e). Yield: 80%, time:12h; light brown gum; $[\alpha]_D^{25} - 54.5$ (*c* 0.6, CHCl₃); IR(KBr): ν_{max} 3097, 1721, 1715, 1699, 1329, 1237, 1033cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ : 7.79 (1H, dd, *J* = 0.9, 3.5, Ar-4") , 7.55 (1H, dd, *J* = 1.1, 5.0 Hz, Ar-2") , 7.11 (1H, m, Ar-3"), 7.51(1H, bs, H-21), 7.39 (1H, bs, H-23), 6.44(1H, bs, H-22), 5.85 (1H, s, H-17), 5.14 (1H, dd, *J* = 16.8 & 9.4 Hz, H-3), 5.11 (1H, H_b-30), 4.86 (1H, s, H_a-30), 3.69 (3H, s, OMe), 3.29 (1H, t, *J* = 3.2 & 6.0 Hz, H-1), 2.91 (1H, d, *J* = 18.1Hz, H_a-15)& 2.61 (1H, d, *J* = 18.1Hz, H_b-

15), 2.54 (1H, dd, $J = 10$ & 4.9 Hz, H-5), 2.62 (1H, d, $J = 15.1$ Hz, H_a-6) & 2.14 (1H, d, $J = 15.1$ Hz, H_b-6), 2.28 (1H, m, H_a-12) & 1.35 (1H, m, H_b-12), 2.07 (2H, m, H-2), 2.06 (1H, m, H_a-11) & 1.23 (1H, m, H_b-11), 2.05 (1H, br s, H-9), 1.07 (3H, s, H-29), 0.89 (3H, s, H-18), 0.86 (3H, s, H-19), 0.8 (3H, s, oH, H-28); ^{13}C NMR (75 MHz, CDCl₃) δ : 174.2 (C-7), 170.2 (C-16), 161.4 (ester carbonyl- C-1'), 146.3 (C-8), 142.6 (C-23), 140.9 (C-21), 134.0 (Ar-1''), 133.24 (Ar-2''), 132.20 (Ar-4''), 127.76 (Ar-3''), 120.9 (C-20), 110.9 (C-22), 110.0 (C-30), 80.0 (C-14), 79.7 (C-17), 75.6 (C-3), 74.7 (C-1), 51.9 (C-9), 50.4 (C-10), 43.8 (C-12), 41.5 (C-13), 39.0 (C-4), 33.8 (C-15), 32.3 (C-6), 29.2 (C-2), 29.6 (C-11), 27.0 (C-28), 23.9 (C-5), 21.9 (C-19), 15.8 (C-29), 13.84 (C-18); ESI-MS: m/z 605.2 [M+Na]⁺.

1.11.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 2-morpholinoacetate (13a). Yield: 81%, Time: 4h; oil; $[\alpha]_D^{25} - 65.1$ (c 0.6, CHCl₃); IR(KBr): ν_{max} 2983, 1742, 1731, 1689, 1331, 1241, 1031 cm⁻¹. ^1H NMR (300 MHz, CDCl₃) δ : 7.48 (1H, bs, H-21), 7.39 (1H, bs, H-23), 6.43 (1H, bs, H-22), 5.79 (1H, s, H-17), 5.06 (1H, d, $J = 11.7$ & 5.7 Hz, H-3), 5.10 (1H, s, H_a-30), 4.86 (1H, s, H_b-30), 3.76 (4H, m, morpholine -5''&3''), 3.69 (3H, s, OMe), 3.26 (2H, s, chloroacetyl-H-2'), 3.24 (1H, t, $J = 3.2$ & 3.0 Hz, H-1), 2.90 (1H, d, $J = 18.1$ Hz, H_a-15) & 2.56 (1H, d, $J = 18.1$ Hz, H_b-15), 2.60 (4H, m, morpholine H-2''& H-6''), 2.56 (1H, m, H-5), 1.84 (2H, m, H-2), 2.54 (1H, OL, H_a-6) & 2.10 (1H, d, $J = 15.2$ Hz, H_b-6), 2.21 (1H, br s, H-9), 2.07 (1H, m, H_a-11)& 1.54 (1H, m, H_b-11), 1.86 (1H, m, H_a-12)& 1.16 (1H, m, H_b-12), 0.95 (1H, s, H-29), 0.88 (3H, s, H-18), 0.83 (3H, s, H-19), 0.80 (3H, s, H-28); ^{13}C NMR (75 MHz, CDCl₃) δ : 174.21 (C-7), 170.3 (C-16), 169.3 (chloroacetyl-1'), 146.2 (C-8), 142.6 (C-23), 140.9 (C-21), 120.8 (C-20), 111.0 (C-22), 109.9 (C-30), 80.0 (C-14), 79.7 (C-17), 74.9 (C-3), 74.6 (C-1), 66.77 (morpholine -3''& 5''), 59.1 (C-2'), 53.21 (morpholine -2''& 6''), 51.9 (C-9),

50.3 (C-10), 43.8 (C-12), 41.5 (C-13), 38.7 (C-4), 33.8 (C-15), 32.3 (C-6), 29.7 (C-2), 29.3 (C-11), 27.0 (C-28), 23.8 (C-5), 21.9 (C-19), 15.6 (C-29), 13.8 (C-18). HRMS (ESI+) *m/z*: 600.3167 [M+H]⁺ (calcd for C₃₃H₄₆NO₉Na [M+Na]⁺, 600.3167)

1.12.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl-2-(pyrrolidin-1-yl)acetate (13b). Yield: 85%, time: 4h; yellow gum; $[\alpha]_D^{25} - 54.8$ (*c* 0.4, CHCl₃); IR(KBr): ν_{max} 2992, 1732, 1734, 1699, 1321, 1241, 1034cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ : ¹H NMR (300Hz , CDCl₃): 7.50 (1H, bs, H-21), 7.39 (1H, bs, H-23), 5.80 (1H, s, H-17), 6.43 (1H, bs, H-22), 5.10 (1H, s, H_a-30), 5.06 (1H, d, *J* = 11.7 & 5.7 Hz, H-3), 4.85 (1H, s, H_b-30), 3.69(3H, s, OMe), 3.43 (4H, m, pyrrolidine H-2" & H-5"), 3.26 (2H, s, chloroacetyl H-2'), 3.23 (1H, t, *J* = 3.2 & 3.0 Hz, H-1), 2.90 (1H , d , *J* = 18.1Hz, H_a-15), 2.58 (1H, d , *J* = 18.1Hz, H_b-15), 2.54 (1H, m, H-5), 2.50 (1H, d, *J* = 15.6 Hz, H_a-6) & 2.09 (1H, d, *J* = 15.6 Hz, H_b-6), 2.18 (1H, br s, H-9), 2.06 (1H, m, H_a-11)&1.55 (1H, m, H_b-11), 2.86(4H, m, pyrrolidine H-3" & H-4"), 1.85(1H, m, H_a-12) & 1.15 (1H, m, H_b-12), 1.84 (2H, m, H-2), 0.95 (3H, s, H-19), 0.87 (3H, s, H-18), 0.82 (3H, s, H-19), 0.81 (3H, s, H-28); ¹³C NMR (75 MHz, CDCl₃) δ : 174.2(C-7), 170.4(C-16), 174.5 (chloroacetyl C-1'), 146.3 (C-8), 142.6 (C-23), 140.9 (C-21), 120.8 (C-20), 110.9 (C-22), 110.0 (C-30), 79.9 (C-14), 79.7 (C-17), 74.6 (C-3), 72.6 (C-1), 57.2 (chloroacetyl C-2'), 51.94 (pyrrolidine C-2"& C-5"), 50.3 (C-9), 47.3 (C-10), 43.8 (C-12), 41.5 (C-13), 38.7 (C-4), 33.8 (C-15), 32.3 (C-6), 29.6 (C-2), 29.3 (C-11), 26.9 (C-28), 24.58 (pyrrolidine C-3'& C-4'), 23.8 (C-5), 22.49, 21.8 (C-19), 15.6 (C-29), 13.8 (C-18) ; HRMS (ESI+) *m/z*: 584.3222 [M+H]⁺ (calcd for C₃₃H₄₆NO₈ [M+Na]⁺, 584.3217).

1.13.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 2-

(4-methylpiperazin-1-yl)acetate (13c). Yield: 85%, time: 5h; yellow oil; $[\alpha]_D^{25} - 52.6$ (*c* 0.6, CHCl₃); IR(KBr): ν_{max} 2988, 1731, 1741, 1698, 1331, 1242, 1038cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ : 7.48 (1H, bs, H-21), 7.39 (1H, bs, H-23), 5.79 (1H, s, H-17), 6.43 (1H, bs, H-22), 5.10 (1H, s, H_b-30), 5.05 (1H, d, *J* = 12 & 4.5 Hz, H-3), 4.85 (1H, s, H_a-30), 3.68 (3H, s, OMe), 3.25 (2H, m, chloroacetyl H-2'), 3.24 (1H, t, *J* = 3.2 & 3.0 Hz, H-1), 3.20 (4H, m, piperazine H-3''&H-5''), 2.90 (1H, d, *J* = 18.1Hz, H_a-15), 2.58 (1H, d, *J* = 18.1Hz, H_b-15), 2.54 (4H, m, piperazine H-2'' & H-6''), 2.53 (1H, dd, *J* = 11.8 & 5.0 Hz, H-5), 2.50 (1H, d, *J* = 16.0 Hz, H_a-6) & 2.10 (1H, d, *J* = 15.2 Hz, H_b-6), 2.32(3H, m), 2.26 (3H, s, N-Methyl) 2.08 (1H, br s, H-9), 2.21(1H, m, H_a-12) & 1.14 (1H, m,H_b-12), 2.01 (1H, m, H_a-11) &1.53 (1H, m,H_b-11), 1.84 (2H, m, H-2), 0.87 (3H, s, H-18), 0.82 (3H, s, H-19), 0.80 (3H, s, H-28), 0.95 (3H, s, H-29); ¹³C NMR (75 MHz, CDCl₃) δ : 174.22 (C-7), 170.38 (C-16), 169.51 (chloroacetyl C-1'), 146.28 (C-8), 142.62 (C-23), 140.89 (C-21), 120.8 (C-20), 110.9 (C-22), 109.9 (C-30), 79.9 (C-14), 79.7 (C-17), 74.6 (C-3), 74.8 (C-1),59.2 (C-2'), 54 (piperazine C-2'' & 6''), 52.68(piperazine C-3''&C-5''), 51.9(C-9), 50.3 (C-10), 45.74 (N-Methyl), 43.8 (C-12), 41.5 (C-13), 38.6 (C-4), 33.8 (C-15), 32.3 (C-6), 29.6 (C-2), 29.3 (C-11), 26.9 (C-28), 23.8 (C-5), 21.8 (C-19), 15.6 (C-29), 13.8 (C-18); HRMS (ESI+) *m/z*: 613.3484 [M+H]⁺ (calcd for C₃₄H₄₉N₂O₈ [M+H]⁺, 613.3483).

1.14.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 2-(4-ethylpiperazin-1-yl)acetate (13d). Yield: 83%, time: 5h; yellow gum. $[\alpha]_D^{25} - 69.2$ (*c* 0.7, CHCl₃); IR(KBr): ν_{max} 2997, 1737, 1742, 1697, 1327, 1242, 1037cm⁻¹; ¹H NMR (300 MHz, in CDCl₃) δ (in ppm): 7.50 (1H, bs, H-21), 7.40 (1H, bs, H-23), 6.43 (1H, bs, H-22), 5.79 (1H, s, H-17), 5.11 (1H, s, H_a-30), 5.00 (1H, d, *J* = 11.8 & 4.7 Hz, H-3), 4.86 (1H, s, H_b-30), 3.69 (3H,s, OMe),3.25 (2H, s, H-2') 3.24 (1H, t, *J* = 3.2 & 3.0 Hz, H-1), 3.32 (2H, s, chloroacetyl H-1'), 2.90

(1H, d, $J = 18.1$ Hz, H_a-15) & 2.58 (1H, d, $J = 18.1$ Hz, H_b-15), 2.55 (1H, dd, $J = 11.8$ & 5.0 Hz, H-5), 2.54 (4H, m, piperazine H-2" & H-6"), 2.51(4H, m, piperazine H-3" & H-5"), 2.50 (1H, OL, H_a-6) & 2.09 (1H, d, $J = 15.2$ Hz, H_b-6), 2.33(2H, m, ethyl H-1"'), 2.18 (1H, br s, H-9), 2.07 (1H, m, H_a-11) & 1.53 (1H, m, H_b-11), 1.84 (2H, m, H-2), 1.85(1H, m, H_a-12) & 1.15 (1H, m, H_b-12), 1.27 (3H, t, $J = 7.5$ Hz, ethyl H-2"'), 0.94 (1H, s, H-29), 0.87 (3H, s, H-18), 0.83 (3H, s, H-19), 0.80 (1H, s, H-28); ^{13}C NMR (75 MHz, CDCl₃) δ : 174.2(C-7), 170.9 (C-16), 170.4 (chloroacetyl C-1'), 146.2 (C-8), 142.6 (C-23), 140.8 (C-21), 120.8 (C-20), 111.0 (C-22), 109.9 (C-30), 80.0(C-14), 80.0 (C-17), 74.6 (C-3), 74.6(C-1), 59.1(C-2'), 52.80 (piperazine C-2" & C-6"), 51.97 (piperazine C-3" & 5"), 51.4 (C-9), 50.3(C-10), 43.8(C-12), 41.4(C-13), 38.7(C-4), 33.8 (C-15), 32.3 (C-6), 31.9 (C-1"'), 29.6 (C-2), 29.3 (C-11), 26.9 (C-28), 23.8(C-5), 22.6 (C-2"'), 21.9(C-19), 15.7 (C-29), 13.9 (C-18); ESI-MS: m/z 627.3 [M+H]⁺.

1.15.tert-butyl-4-((7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl)oxy)-2-oxoethyl)piperazine-1-carboxylate (13e). Yield: 86%, time: 4h; yellow gum; $[\alpha]_D^{25} - 38.1$ (c 0.5, CHCl₃); IR(KBr): ν_{\max} 2995, 1735, 1732, 1695, 1325, 1242, 1035cm⁻¹; ^1H NMR (300 MHz, CDCl₃) δ : 7.49 (1H, bs, H-21), 7.39 (1H, bs, H-23), 6.43 (1H, bs, H-22), 5.78 (1H, s, H-17), 5.10 (1H, s, H_a-30), 5.04 (1H, d, $J = 12$ & 4.5 Hz, H-3), 4.85 (1H, s, H_b-30), 3.69 (3H, s, OMe), 3.35 (4H, m, piperazine H-3" & H-5"), 3.24 (1H, t, $J = 3.2$ & 3.0 Hz, H-1), 3.23 (2H, m, chloroacetyl H-1'), 2.90 (1H, d, $J = 18.1$ Hz, H_a-15), 2.57 (1H, d, $J = 18.1$ Hz, H_b-15), 2.55(1H, m, H-5), 2.50 (1H, d, $J = 16.0$ Hz, H_a-6) & 2.10 (1H, d, $J = 15.8$ Hz, H_b-6), 2.28(4H, m, piperazine H-2" & H-6") 2.08 (1H, brs, H-9), 2.21 (1H, m, H_a-11) & 1.53 (1H, m, H_b-11), 1.85(1H, m, H_a-12) & 1.16(1H, m, H_b-12), 1.84 (2H, m, H-2), 1.47 (9H, s, boc-3Me), 0.94 (3H, s, H-29), 0.87 (3H, s, H-18), 0.82(3H, s, H-19), 0.80

(3H, s, H-28); HRMS (ESI+) m/z : 721.3672 [M+Na]⁺ (calcd for C₃₈H₅₄N₂O₁₀Na[M+H]⁺, 721.3670).

1.16.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl-2-(4-(2-hydroxyethyl)piperazin-1-yl)acetate (13f). Yield: 75%, time: 6h; yellow gum; $[\alpha]_D^{25} = -44.6$ (c 0.6, CHCl₃); IR(KBr): ν_{max} 3497, 1731, 1718, 1698, 1328, 1238, 1033cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ : 7.48(1H, bs, H-21), 7.39 (1H, bs, H-23), 6.43(1H, bs, H-22), 5.79 (1H, s, H-17), 5.10 (1H, s, H_a-30), 5.05 (1H, d, $J = 11.1 \text{ \& } 5.2$ Hz, H-3), 4.85 (1H, s, H_b-30), 3.69 (3H, s, OMe), 3.69 (2H, t, $J = 5.2$, O-CH₂-2''), 3.24(2H, m, H-2'), 3.26(2H, s, chloroacetyl, H-2') 3.23 (1H, t, $J = 3.2 \text{ \& } 3.0$ Hz, H-1), 2.4 (1H, m, H-5), 2.90 (1H, d, $J = 18.1$ Hz, H_a-15), 2.65 (4H, m, H-3'' & H-5''), 2.61(2H,t, $J = 5.2$ Hz, N-CH₂, H-1''), 2.57 (1H, d, $J = 18.1$, H_b-15), 2.51 (1H, d, $J = 15.2$, H_a-6) & 2.09 (1H, d, $J = 15.2$ Hz, H_b-6), 2.17 (1H, br, H-9), 2.14 (4H, m, piperazine H-2'' & H-6'') 2.07 (1H, m, H_a-11) & 1.56 (1H, m, H_b-11), 1.98 (1H, m, H_a-12), 1.86 (2H, m, H-2), 1.16(1H, m, H_b-12), 0.95 (3H, s, H-29), 0.87 (3H, s, H-18), 0.82 (3H, s, H-19), 0.80 (3H, s, H-28); ¹³C NMR (75 MHz, CDCl₃) δ : 174.1(C-7), 170.3 (C-16), 169.4 (chloroacetyl C-1¹), 146.3(C-8), 142.6 (C-23), 140.9 (C-21), 120.8 (C-20), 110.9 (C-22), 109.9 (C-30), 80.0(C-14), 79.7 (C-17), 74.9 (C-3), 74.6(C-1), 59.19 (O-CH₂-2''), 59.1 (C-2'), 57.5 (N-CH₂-1''), 52.5(piperazine C-3'' & C-5''), 51.9(C-9), 50.9 (piperazine C-2'' & C-6'') 50.4(C-10), 43.8 (C-12), 41.5 (C-13), 38.7(C-4), 33.8 (C-15), 32.3 (C-6), 29.6(C-11), 27.0 (C-28), 23.8 (C-5), 21.9 (C-19), 15.6 (C-29), 13.8 (C-18); HRMS (ESI+) m/z : calcd for C₃₅H₅₁N₂O₉ [M+H]⁺: 643.3589; found: 643.3599 [M+H]⁺.

1.17.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 2-


(4-(furan-3-carbonyl)piperazin-1-yl)acetate (13g). Yield: 88%, yellow gum; $[\alpha]_D^{25} - 53.9$ (*c* 1.0, CHCl_3); IR(KBr): ν_{max} 2997, 1731, 1719, 1698, 1337, 1237, 1034 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ : 7.48 (1H, bs, H-21), 7.47 (1H, bs, furan-H), 7.39 (1H, bs, H-23), 6.43 (1H, bs, H-22), 6.47 (1H, bs, furan-H), 6.99 (1H, bs, furan-H), 5.79 (1H, s, H-17), 5.07 (1H, d, *J* = 11.5 & 5.09 Hz, H-3), 5.10 (1H, s, H_a-30), 4.85 (1H, s, H_b-30), 3.87 (4H, m, piperazine H-3" & H-5"), 3.69 (3H, s, OMe), 3.26 (2H, s, chloroacetyl C-2'), 3.25 (1H, t, *J* = 3.2 & 3.0, H-1), 2.90 (1H, d, *J* = 17.9 Hz, H_a-15), 2.67 (4H, m, piperazine H-2" & H-6"), 2.57 (1H, d, *J* = 17.9 Hz, H_a-15), 2.54 (1H, dd, *J* = 11.2 & 5.0 Hz, H-5), 2.58 (1H, d, *J* = 14.8 Hz, H_a-6) & 2.09 (1H, d, *J* = 14.8 Hz, H_b-6), 2.17 (1H, brs, H-9), 2.11 (1H, m, H_a-11) & 1.53 (1H, m, H_b-11), 2.21 (1H, m, H_a-12) & 1.16 (1H, m, H_b-12), 1.86 (2H, m, H-2), 0.95 (1H, s), 0.87 (3H, s, H-18), 0.82 (3H, s, H-19), 0.81 (1H, s); ^{13}C NMR (75 MHz, CDCl_3) δ : 207(Keto C-1") 174.1(C-7), 170.3(C-16), 169.3(chloroacetyl-1), 159.04, 147.82(furan C), 146.2(C-8), 143.63 (furan C), 142.6(C-23), 140.8(C-21), 120.8(C-20), 116.37(furan C), 111.2(C-22), 109.9(C-30), 79.9 (C-14), 79.8 (C-17), 74.6 (C-3), 75.05 (C-1), 59.04 (C-2'), 52.9(piperazine C-3"&5"), 52.3(piperazine C-2"&6"), 51.9 (C-9), 50.3(C-10), 43.8 (C-12), 38.7(C-13), 33.8 (C-4), 32.3 (C-15), 30.9 (C-6), 29.6 (C-2), 29.3(C-11), 27.0 (C-28), 23.8 (C-5), 21.8 (C-19), 15.6 (C-29), 13.8 (C-18); HRMS (ESI+) *m/z*: 693.3³⁸⁹ [M+H]⁺ (calcd for $\text{C}_{38}\text{H}_{49}\text{N}_2\text{O}_{10}$ [M+H]⁺, 693.3381).

1.18.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 2-(4-benzoylpiperazin-1-yl)acetate (13h). Yield: 89%, time: 6h; yellow gum; $[\alpha]_D^{25} - 41.9$ (*c* 0.4, CHCl_3); IR(KBr): ν_{max} 3097, 1721, 1715, 1699, 1329, 1237, 1033 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ : 7.49 (1H, bs, H-21), 7.39 (1H, bs, H-23), 7.39 (5H, m, 5Ar-H), 6.39 (1H, O1, H-22), 5.79 (1H, s, H-17), 5.10 (1H, s, H_a-30), 5.05 (1H, d, *J* = 11.7 & 5.05 Hz, H-3), 4.85 (1H, s, H_b-

30), 3.85(4H, m, piperazine H-3" & H-5"), 3.69 (3H, s, OMe), 3.27(2H, S, chloroacetyl -2'), 3.26 (1H, t, J = 3.2 & 3.0 Hz, H-1), 2.91 (1H, d, J = 17.9 Hz, H_a-15), 2.71 (1H, d, J = 17.9 Hz, H_b-15), 2.55 (1H, dd, J = 11.2 & 5.0 Hz, H-5), 2.52 (1H, d, J = 14.8 Hz, H_a-6) & 2.15 (1H, d, J = 14.8 Hz, H_b-6), 2.66(4H, m, piperazine H-2" & H-6"), 2.17 (1H, brs, H-9), 2.11(1H, m, H_a-11) & 1.86(1H, m, H_b-11), 2.22(1H, m, H_a-12) & 1.17 (1H, m, H_b-12), 1.87 (2H, m, H-2), 0.95 (1H, s, H-29), 0.87 (3H, s, H-18), 0.82 (3H, s, H-19), 0.80 (3H, s, H-28): ^{13}C NMR (75 MHz, CDCl₃) δ : 207.0 (keto C-1''), 174.1(C-7), 176.2 (chloroacetyl C-1'), 170.3 (C-16), 146.2 (C-8), 140.8 (C-21), 135.6 (Ar-C), 129.6 (Ar-C), 128.4 (Ar-C), 127.0 (Ar-C), 120.8 (C-20), 111.0 (C-22), 109.9 (C-30), 79.9 (C-14), 79.8 (C-17), 74.6 (C-3), 75.0(C-1), 58.97 (chloroacetyl C-2'), 52.9 (piperazine C-3" & 5"), 52.4 (piperazine C-2" & 6"), 51.9 (C-9), 50.3 (C-10), 43.8 (C-12), 43.8 (C-13), 38.6 (C-4), 30.9 (C-15), 27.0 (C-6), 29.3 (C-2), 29.6 (C-11), 21.8 (C-19), 23.8 (C-5), 15.6 (C-29), 13.8 (C-18); HRMS (ESI+) m/z : 703.3591 [M+H]⁺ (calcd for C₄₀H₅₁N₂O₉ [M+H]⁺, 703.3589).

1.19.(7aS,8S,10S,11aS,12aS)-4-(furan-3-yl)-8-(2-methoxy-2-oxoethyl)-4a,7a,9,9-tetramethyl-13-methylene-2-oxododecahydro-1H-7,12a-methanobenzo[b]pyrano[3,4-g]oxocin-10-yl 2-(diethylamino)acetate (13i). Yield: 90%, Time: 4h; oil; $[\alpha]_D^{25} - 43.2$ (*c* 0.9, CHCl₃); IR(KBr): ν_{max} 3087, 1731, 1725, 1698, 1339, 1233, 1034cm⁻¹; ^1H NMR (300 MHz, CDCl₃) δ : 7.50 (1H, bs, H-21), 7.39 (1H, bs, H-23), 6.43(1H, bs, H-22), 5.81 (1H, s, H-17), 5.10 (1H, s, H_a-30), 5.04 (1H, d, J = 11.5 & 5.09 Hz, H-3), 4.86 (1H, s, H_b-30), 3.69 (3H, s, OMe), 3.24 (2H, s, chloroacetyl -2'), 3.23 (1H, t, J = 3.2 & 3.0, H-1), 2.90 (1H, d, J = 18.1Hz, H_a-15), 2.70 (4H, m, N,N,-Diethyl-2CH₂), 2.57 (1H, d, J = 18.1Hz, H_b-15), 2.55 (1H, m, H-5), 2.50 (1H, d, J = 16.0Hz, H_a-6) & 2.09 (1H, d, J = 15.2 Hz, H_b-6), 2.18 (1H, br s,H-9), 1.87(1H, m,H_a-11) & 1.53(1H, m, H_b-11), 2.21(1H, m, H_a-12), 1.86 (2H, m, H-2), 1.15 (1H, m, H_b-12), 1.07 (6H, t,

$J = 7.1 \text{ \& } 14.2 \text{ Hz}$, N,N,-Diethyl 2Me), 0.96 (1H, s, H-29), 0.87 (3H, s, H-18), 0.83 (3H, s, H-19), 0.80 (3H, s, H-28); ^{13}C NMR (75 MHz, CDCl_3) δ : 181.1 (chloroacetyl C-1¹), 174.2 (C-7), 170.4 (C-16), 146.2 (C-8), 142.6 (C-23), 140.8 (C-21), 120.8 (C-20), 111.0 (C-22), 109.9 (C-30), 80.0 (C-14), 74.6 (C-17), 59.9 (chloroacetyl-2'), 52.0 (N,N,-Diethyl -2CH₂-), 51.4 (C-9), 50.3 (C-10), 43.8 (C-12), 41.4 (C-13), 38.7 (C-4), 33.8 (C-15), 32.3 (-6), 31.9 (C-2), 29.3 (C-11), 26.9(C-28), 23.8 (C-5), 21.90 (C-19), 15.73 (C-29), 14.10 (N,N,-Diethyl-2Me), 13.9 (C-18); HRMS (ESI+) m/z : 586.3347 [M+H]⁺ (calcd for $\text{C}_{33}\text{H}_{48}\text{NO}_8$ [M+H]⁺, 586.3374).



NPL IICR FTIR REPORT

Date: Friday, January 11, 2013





Figure 1: HRESIMS, IR and ^1H NMR Spectrums of Compound 1




Figure 2: ^{13}C NMR & DEPT 135° Spectrums of Compound 1




Figure 3: HSQC Spectrum of Compound 1




Figure 4: HMBC Spectrum of Compound 1




Figure 5: COSY Spectrum of Compound 1




Figure 6: NOESY Spectrum of Compound 1




Figure 7: HRESIMS, IR and ¹H NMR Spectrums of Compound 2




Figure 8: ^{13}C NMR & DEPT 135° Spectrums of Compound 2




Figure 9: HSQC Spectrum of Compound 2




Figure 10: HMBC Spectrum of Compound 2




Figure 11: COSY Spectrum of Compound 2




Figure 12: NOESY Spectrum of Compound 2




Figure 13: HRESIMS, 1H & ^{13}C NMR of Compound 9




Figure 14: HRESIMS, ^1H & ^{13}C NMR of Compound **10a**

KSB-CBS-PROPINYL #4-28 RT: 0.03-0.12 AV: 25 NL: 7.79E6
T: FTMS {1,1} + p ESI Full ms [100.00-2000.00]




Figure 15: HRESIMS, ^1H & ^{13}C NMR of Compound 10b

KS-B-CBS-ISO-BUT #3-34 RT: 0.02-0.14 AV: 32 NL: 6.48E6
T: FTMS {1,1} + p ESI Full ms [100.00-2000.00]




Figure 16: HRESIMS, ^1H & ^{13}C NMR of Compound 10c




Figure 17: HRESIMS, ^1H & ^{13}C NMR of Compound 11a




Figure 18: HRESIMS, ^1H & ^{13}C NMR of Compound **11b**




Figure 19: HRESIMS, ^1H & ^{13}C NMR of Compound **11c**




Figure 20: HRESIMS, ¹H & ¹³C NMR of Compound 11d





Figure 21: ¹H & ¹³C NMR of Compound 11e






Figure 22: HRESIMS, ^1H & ^{13}C NMR of Compound 12




Figure 23: HRESIMS, ^1H & ^{13}C NMR of Compound **13a**




Figure 24: HRESIMS, ¹H & ¹³C NMR of Compound **13b**




Figure 25: HRESIMS, ^1H & ^{13}C NMR of Compound **13c**

Elemental Composition Report

Page 1


Single Mass Analysis

Tolerance = 20.0 PPM / DBE: min = -1.5, max = 30.0
 Element prediction: Off
 Number of isotope peaks used for i-FIT = 3

Monoisotopic Mass, Odd and Even Electron Ions
 31 formula(e) evaluated with 1 results within limits (all results (up to 1000) for each mass)
 Elements Used:

C: 0-35 H: 0-52 N: 0-2 O: 0-10
 20160904_024 (0.323)

1: TOF MS 627.31ES+
 3.20e+006



Minimum: -1.5
 Maximum: 5.0 20.0 30.0

Mass	Calc. Mass	mDa	PPM	DBE	i-FIT	Norm	Conf(%)	Formula
627.3632	627.3645	-1.3	-2.1	11.5	520.1	n/a	n/a	C35 H51 N2 O8




Figure 26:¹H & ¹³C NMR of Compound 13d




Figure 27: HRESIMS, ^1H & ^{13}C NMR of Compound **13e**




Figure 28: HRESIMS, ^1H & ^{13}C NMR of Compound **13f**





Figure 29: HRESIMS, 1H & ^{13}C NMR of Compound 13g






Figure 31: HRESIMS, ^1H & ^{13}C NMR of Compound **13i**