



HAL
open science

Elemental properties, hydrology, and biology interact to shape concentration-discharge curves for carbon, nutrients, sediment, and major ions

Florentina Moatar, Benjamin W. Abbott, Camille Minaudo, Florence Curie, Gilles Pinay

► To cite this version:

Florentina Moatar, Benjamin W. Abbott, Camille Minaudo, Florence Curie, Gilles Pinay. Elemental properties, hydrology, and biology interact to shape concentration-discharge curves for carbon, nutrients, sediment, and major ions. *Water Resources Research*, 2017, 53 (2), pp.1270-1287. 10.1002/2016WR019635 . hal-01507692

HAL Id: hal-01507692

<https://univ-rennes.hal.science/hal-01507692v1>


Submitted on 21 Jun 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



RESEARCH ARTICLE

10.1002/2016WR019635

Special Section:

Concentration-discharge Relations in the Critical Zone

Key Points:

- Only 2 or 3 of the possible 9 C-Q modalities occurred for any given element, and these shapes were stable through time
- Elemental properties set C-Q templates for each parameter, secondarily influenced by biology, catchment characteristics, and hydrology
- Nearly two-thirds of C-Q relationships were nonlinear, emphasizing information cost of monotonic C-Q analysis

Supporting Information:

- Supporting Information S1
- Data Set S1

Correspondence to:

F. Moatar,
Florentina.moatar@univ-tours.fr

Citation:

Moatar, F., B. W. Abbott, C. Minaudo, F. Curie, and G. Pinay (2017), Elemental properties, hydrology, and biology interact to shape concentration-discharge curves for carbon, nutrients, sediment, and major ions, *Water Resour. Res.*, 53, 1270–1287, doi:10.1002/2016WR019635.

Received 11 AUG 2016

Accepted 10 JAN 2017

Accepted article online 20 JAN 2017

Published online 3 FEB 2017

© 2017. The Authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Elemental properties, hydrology, and biology interact to shape concentration-discharge curves for carbon, nutrients, sediment, and major ions

F. Moatar¹, B. W. Abbott^{2,3} , C. Minaudo¹, F. Curie¹, and G. Pinay²

¹University François-Rabelais Tours, EA 6293 Géo-Hydrosystèmes Continentaux, Parc de Grandmont, Tours, France,

²OSUR-ECOBIO-CNRS University of Rennes1, campus de Beaulieu, avenue du general Leclerc, Rennes, France,

³Department of Earth and Environmental Sciences, Michigan State University, East Lansing, Michigan, USA

Abstract To investigate the prevalence and cause of concentration-discharge (C-Q) relationships for carbon, nutrients, major ions, and particulates, we analyzed 40 years of water quality data from 293 monitoring stations in France. Catchments drained diverse landscapes and ranged from 50 to 110,000 km², together covering nearly half of France. To test for differences during low and high flows, we calculated independent C-Q slopes above and below the median discharge. We found that 84% of all catchment-element combinations were chemodynamic for at least half of the hydrograph and 60% of combinations showed nonlinear C-Q curves. Only two or three of the nine possible C-Q modalities were manifest for each parameter, and these modalities were stable through time, suggesting that intrinsic and extrinsic elemental properties (e.g., solubility, reactivity, and source dynamics) set basic C-Q templates for each parameter, which are secondarily influenced by biological activity during low flows, and the interaction between hydrology and catchment characteristics at high flows. Several patterns challenged current C-Q views, including low-flow chemostasis for TSS in 66% of catchments, low-flow biological mediation of NO₃⁻ in 71% of catchments, and positive C-Q for dissolved organic carbon independent of catchment size in 80% of catchments. Efforts to reduce nutrient loading decreased phosphorus concentration and altered C-Q curves, but NO₃⁻ continued to increase. While C-Q segmentation requires more data than a single analysis, the prevalence of nonlinear C-Q slopes demonstrates the potential information loss associated with linear or monotonic analysis of C-Q relationships, and conversely, the value of long-term monitoring.

1. Introduction

Relationships between solute or particulate concentration and discharge have long been used as indicators of biological and hydrological functioning at catchment and subcatchment scales [Hem, 1948; Johnson et al., 1969; Edwards, 1973; Thompson et al., 2011]. The shape of a concentration-discharge (C-Q) curve and its changes in time can indicate solute or particulate abundance and location [Bowes et al., 2005; Thomas et al., 2016], rate and timing of biogeochemical production and uptake [Herndon et al., 2015; Monteith et al., 2015], and hydrological flowpath and residence time [Hooper et al., 1990; Genereux et al., 1993]. C-Q analysis has been used at the event scale for hydrograph separation and to assess solute stocks and production rates [Evans and Davies, 1998; Hornberger et al., 2001; Chanat et al., 2002; Dupas et al., 2015], and at seasonal and multiannual scales to test how catchment characteristics (e.g., topography, soil, vegetation, anthropogenic activity) influence export dynamics [Butturini et al., 2008; Godsey et al., 2009; Musolff et al., 2015]. Consequently, investigating C-Q relationships for multiple elements at multiple spatial and temporal scales can be a powerful tool to address three of ecohydrology's fundamental questions: where does water come from, how long does it stay in the catchment, and what happens to the solutes and particulates it carries along the way [Oldham et al., 2013; Pinay et al., 2015; Abbott et al., 2016]. Additionally, C-Q relationships are necessary for reconstructing elemental fluxes to assess compliance with water quality legislation [European Union, 2000; Raymond et al., 2013; Bowes et al., 2014; Hirsch, 2014], and can be used to detect long-term shifts in hydrological and biogeochemical functioning due to disturbance, land-use, or climate change [Monteith et al., 2007; Petrone et al., 2007; Guzman et al., 2013; Törnqvist et al., 2015]. Despite this large body of work on C-Q relationships, unanswered questions remain about the relative importance of biological and




Figure 1. The nine possible concentration-discharge (C-Q) relationships when segmenting the hydrograph at the median discharge. Conceptual schematic on the left and examples of the modalities from catchments in this study on the right. $b_{50\text{inf}}$ is the slope of the log-transformed C-Q relationship for flows lower than the median and $b_{50\text{sup}}$ for flows greater than the median [Meybeck and Moatar, 2012].

hydrological controls for different elements [Moatar and Meybeck, 2005; Basu et al., 2010; Thompson et al., 2011; Bierzoza and Heathwaite, 2015], and it is not clear to what extent observed C-Q patterns are due to inherent properties of the solute or particulate (e.g., solubility, reactivity, charge) or catchment characteristics (e.g., topography, land cover, weather and climate, hydrologic routing) [Aitkenhead and McDowell, 2000; Basu et al., 2011; Onderka et al., 2012; Gall et al., 2013; Herndon et al., 2015; Torres et al., 2015].

At a basic level, C-Q curves can be described as chemostatic (no change in concentration with discharge) or chemodynamic, yielding three general C-Q shapes: “flat,” “up,” or “down” [Godsey et al., 2009; Musolff et al., 2015]. Positive or “up” chemodynamic relationships (Figure 1a) have been observed for suspended sediment and some nutrients, attributed to enhanced erosion during high flows for suspended sediment and associated pollutants, and to reconnection of surface or subsurface catchment components where a given element is more abundant [Holmes et al., 2012; Thomas et al., 2016]. Up behavior has been described as transport limited, since delivery to the stream network is controlled by the spatial extent and transport capacity of the terrestrial-aquatic nexus rather than production or abundance of the element [Basu et al., 2011]. Chemostatic or flat behavior (Figure 1e) has been observed for some elements associated with geologic weathering [Godsey et al., 2009], dissolved organic carbon (DOC) [Creed et al., 2015], and some nutrients in agricultural catchments [Basu et al., 2010; Thomas et al., 2016]. Flat behavior has been attributed to homogenous and uniform distribution of elements in the catchment, meaning that changes in hydrological connectivity and flowpath do not affect solute concentration, or alternatively to temporal stability of flowpaths (i.e., nonvariant source area) [Godsey et al., 2009; Thompson et al., 2011; Musolff et al., 2015]. Negative or “down” chemodynamic relationships (Figure 1i) have been observed for nutrients and major ions, attributed to dilution during high flows [Shanley et al., 2011; Meybeck and Moatar, 2012; Thomas et al., 2016]. Elements with down behavior have been described as source limited, since delivery to the stream network is determined by their abundance or rate of production, rather than the transport capacity of the catchment [Basu et al., 2011].

Because C-Q analysis requires repeated measurements across the hydrograph, understanding the prevalence and meaning of C-Q modalities depends on gathering and analyzing appropriate data for multiple elements from diverse catchments. While high-frequency data are still relatively rare and tend to be short

term [Schwientek et al., 2013; Strohmeier et al., 2013; Bierzoza and Heathwaite, 2015], there are many large databases of low-frequency water chemistry and discharge from government agencies and research institutes [Alewell et al., 2004; Lischeid, 2008; Holmes et al., 2012; Meybeck and Moatar, 2012; Dupas et al., 2013; Minaudo et al., 2015, 2016; Musolff et al., 2015; Zhang et al., 2016]. The analysis of C-Q curves on decadal timescales and at larger spatial scales shows great promise, however, our ability to generate general understanding of catchment processes from these records is limited by the availability of appropriate analytical tools to digest and distill discriminatory data [Kirchner, 2003; McDonnell et al., 2007]. A host of metrics have been proposed to summarize salient characteristics of C-Q relationships, including unit-less indices of variability, polynomial description of nonlinear trends, and segmentation based on hydrological or chemical analysis [Godsey et al., 2010; Jawitz and Mitchell, 2011; Thompson et al., 2011; Meybeck and Moatar, 2012; Musolff et al., 2015; Zhang et al., 2016]. While each of these metrics has different applications and limitations, several simple indices have been proposed that could leverage long-term C-Q relationships to shed light on catchment biogeochemistry and hydrology. Based on 128 long-term daily records, Meybeck and Moatar [2012] proposed splitting the hydrograph at the median daily flow and calculating separate log-C log-Q slopes (Figure 1). They reasoned that having individual slopes for low and high flows ($b_{50\text{inf}} < \text{median flow} < b_{50\text{sup}}$) accommodated nonlinearity and potential shifts in hydrological and biogeochemical controls at different hydrological states [Doyle et al., 2005; Meybeck and Moatar, 2012]. Because 60 to 99% (median 90%) of annual flux for most elements occurs at flows greater than the median [Meybeck and Moatar, 2012], $b_{50\text{sup}}$ is the relevant metric for controls on elemental loads. Conversely, $b_{50\text{inf}}$ provides insight into low-flow periods, revealing potential biological and geochemical dynamics that are not detectable when hydrological flow is dominant [Moatar et al., 2013]. With two C-Q segments each with three degrees of freedom, nine possible C-Q modalities emerge (Figure 1).

In this study, we applied the split-hydrograph method of Meybeck and Moatar [2012] to a large, long-term, low-frequency data set covering nearly half of France. Our goal was to quantify the prevalence of different C-Q shapes and identify how catchment characteristics (catchment size, land use, lithology, and hydrology), biological activity (primary and secondary productivity), and inherent solute and particulate properties (ionic charge, solubility, and reactivity) interact to determine C-Q response. Specifically we asked: 1. How general are C-Q relationships across catchments and how many of the 9 possible C-Q shapes actually occur for each parameter, 2. How does catchment size alter C-Q curves for a particular element and can these differences be used to infer stream network dynamics or instream processes, 3. How do biologically-reactive parameters differ from nonreactive parameters with similar molecular properties, and 4. How has land-use change altered solute concentrations and C-Q modalities for catchments with long-term records? To address these questions, we analyzed nearly half a million measurements of DOC, nutrients, major ions, and suspended sediment, collected over 40 years from 293 water quality monitoring stations in western and central France.

2. Methods

2.1. Study Sites and Sampling Design

Water chemistry was collected at 293 stations at weekly to monthly frequencies by the water quality agencies of Seine-Normandy (ABSN) and Loire-Brittany (ABLB) as a part of mandatory national monitoring surveys from 1970 to 2000, and to comply with the European Union Water Framework Directive since 2000 [European Union, 2000]. Studied catchments vary from 50 km² to 110,000 km² and include most ecoregions of France, except Mediterranean and high mountain environments which do not occur in these regions (Figure 2). The catchments drain numerous geologic substrata, including metamorphic, volcanic, carbonate, and non-carbonate sedimentary deposits. Together, ABSN (76,000 km²) and ABLB (Loire River basin 110,000 km², Brittany 29,000 km²) cover 215,000 km², approximately half of the French territory. Climatic conditions in the study regions are oceanic in Brittany (mean annual temperature and precipitation in western Brittany: 11.1°C and 1200 mm yr⁻¹), temperate in the Seine catchment and middle reaches of the Loire basin (12.2°C and 810 mm yr⁻¹), and continental in the higher-altitude catchments of the upper Loire in the Massif Central (9°C and 1500 mm yr⁻¹). Agriculture is widespread, with intensive wheat and corn production in the Beauce and Brie basins surrounding the Paris metropolitan area and in the lower Loire Valley; intensive pig, dairy, and poultry production in Brittany; and less intensive agricultural activities in the upper




Figure 2. Study sites superimposed on a geologic map of France. 293 catchments had sufficient data (at least 300 sampling points across the hydrograph) for inclusion in this study.

Loire and Seine catchments with a mix of crops and livestock. Human population in the combined catchments is around 25 million inhabitants, 11 million of which reside in the Paris metropolitan area.

We analyzed segmented C-Q curves for dissolved and particulate parameters collected by the water quality agencies, representing more than 460,000 unique date and location combinations across the 293 stations. NH_4^+ was not included in the final analysis due to data-quality issues (i.e., large proportion of measurements at or below the detection limit and widespread analytical artifacts). Analyzed parameters included variables typically associated with geochemical weathering such as dissolved silica (SiO_2), sodium (Na^+), calcium (Ca^{2+}), and magnesium (Mg^{2+}); variables strongly influenced by agricultural and urban land use such as nitrate (NO_3^-), orthophosphate (PO_4^{3-}), unfiltered total phosphorous (TP), dissolved organic carbon (DOC), total suspended solids (TSS), and conductivity (Cond); and elements associated with multiple factors (e.g., weathering, land use, and atmospheric deposition) such as sulfate (SO_4^{2-}), potassium (K^+), and chloride (Cl^-). To ensure that C-Q slopes were robust and representative, we only included stations with at least 300 C-Q couples (samplings where both concentration and discharge were measured) over the last 25 years for a given parameter. Because sampling frequency differed by parameter and station, the number of eligible sites varied by parameter, with NO_3^- and TP being the most frequently measured (293 and 281 eligible stations, respectively) and silica the least (20 eligible stations; Table 1). We compared discharge recorded at the time of sampling by the water quality agencies with nearby continuous monitoring stations in the national hydrometric network and found good agreement (a maximum of 20% difference in the worst case). For all stations, the sampling spanned over 92% of the total observed flow regime, and for 99% of sites, more than 95% of the flow variability was sampled, indicating good representativity of flow conditions (supporting information Figure S1).

2.2. Concentration-Discharge Analysis

Following *Meybeck and Moatar* [2012], we segmented the hydrograph at the median daily discharge based on long-term hydrological records and calculated slopes of the $\log C - \log Q$ relationship. While changes in

Table 1. Sample Size and Summary Statistics for Measured Parameters^a

Parameter	N (Stations)	Measurements per Station		C _{Q50} (mg L ⁻¹)			b _{50inf}	R ² _{inf}		CV _{inf} (flat)	b _{50sup}	R ² _{sup}		CV _{sup} (Flat)
		Median	max	min	Median	max	Median	Median	max	Median	Median	Median	max	Median
DOC	102	186	322	2.38	4.45	9.44	-0.01	0.08	0.34	26	0.14	0.17	0.54	30
NO ₃ ⁻	293	232	1830	2.37	19.0	78.6	0.29	0.23	0.73	33	0.05	0.09	0.30	27
TP	281	217	1340	0.03	0.13	2.18	-0.18	0.08	0.44	67	0.16	0.10	0.43	75
PO ₄ ³⁻	293	231	1680	0.03	0.20	5.92	-0.23	0.10	0.46	75	0.04	0.08	0.48	68
SiO ₂	20	181	298	8.21	10.7	17.0	0.33	0.13	0.47	45	0.12	0.06	0.09	26
Cl ⁻	171	209	1080	4.37	23.3	215	-0.18	0.20	0.62	18	-0.15	0.20	0.64	26
SO ₄ ²⁻	64	213	1040	4.70	19.3	321	-0.15	0.12	0.34	31	-0.08	0.09	0.51	28
Mg ²⁺	161	223	972	1.06	5.41	17.4	-0.06	0.11	0.41	18	-0.10	0.14	0.63	20
Ca ²⁺	159	225	972	2.93	28.8	117	-0.05	0.13	0.57	14	-0.08	0.16	0.79	19
TSS	281	230	1120	2.21	9.46	59.17	0.06	0.07	0.30	70	0.81	0.36	0.79	75
Cond.*	290	228	978	51.61	462	1390	-0.06	0.13	0.60	12	-0.10	0.18	0.78	16

^aC_{Q50} is the average concentration at the median daily flow. b_{50inf} and b_{50sup} are respectively the slopes of the log-transformed C-Q relationship for flows inferior and superior to the median daily flow. R²_{inf} and R²_{sup} are respectively the regression coefficients of b_{50inf} and b_{50sup}. CV is the coefficient of variation for C-Q relationships that were not significant. *Conductivity is expressed in μS cm⁻¹.

C-Q slope do not necessarily occur at the median flow, selecting a fixed inflection point facilitates comparisons between large numbers of catchments, and in practice usually corresponds to observed nonlinearity [Meybeck and Moatar, 2012]. C-Q relationships were analyzed as follows:

$$C = aQ^b \tag{1}$$

where C is concentration, a is a coefficient with units of concentration, Q is discharge, and b is a unit-less exponent representing the slope of the log-transformed C-Q relationship. We calculated individual slopes for flows below and above the median daily flow (b_{50inf} and b_{50sup}, respectively, where “inf” = inferior and “sup” = superior). To situate the two C-Q slopes on the concentration axis (i.e., calculate the intercept as an estimate of central tendency for concentration), we averaged the predicted concentrations for b_{50inf} (equation (2)) and b_{50sup} (equation (3)) at the median daily flow (equation (4)):

$$C_{Q50(1)} = \left[\frac{Q_{50}}{10^{a_{50inf}}} \right]^{1/b_{50inf}} \tag{2}$$

$$C_{Q50(2)} = \left[\frac{Q_{50}}{10^{a_{50sup}}} \right]^{1/b_{50sup}} \tag{3}$$

$$C_{Q50} = \left[\frac{C_{Q50(1)} + C_{Q50(2)}}{2} \right] \tag{4}$$

where C_{Q50} is the average solute or particulate concentration at the median daily flow (Q₅₀) and other terms are as defined in equation (1) on a logarithmic basis. We tested statistical significance of C-Q coefficients by Pearson product-moment correlation, with significant relationships (α < 0.05) classed as up or down (chemodynamic) depending on the slope, and nonsignificant relationships classed as flat (chemostatic). As reported previously, most slopes were significant when b > 0.2 [Meybeck and Moatar, 2012], though the actual threshold varied depending on the number and variance of C-Q couples in the regression (supporting information Figure S2). We reported R² when correlations were significant and the coefficient of variation (CV) of concentration for flat C-Q relationships, since R² is uninformative as a measure of dispersion when slopes are near zero (i.e., when C-Q correlations are nonsignificant).

To test how catchment characteristics, hydrologic dynamics, and biological activity influence C-Q dynamics, we calculated Spearman’s rank correlation coefficients for b_{50inf}, b_{50sup} and C_{Q50} as functions of agricultural land use, population density, carbonated bedrock coverage, mean summer water temperature (T °C) and chlorophyll (Chl), specific discharge (L sec⁻¹ km⁻²), base flow index (BFI) [Sloto and Crouse, 1996], Q_{mna5} (the minimum monthly discharge with a 5 year return period [Nicolle et al., 2013]), W2 (an index of hydrologic reactivity representing the percentage of total discharge that occurs during the highest 2% of flows) [Walsh and Lawler, 1981; Birgand et al., 2010; Moatar et al., 2013], and an index of seasonality based on the sum of the difference between monthly and annual concentrations [Walsh and Lawler, 1981; Tian et al.,

2016]. Additionally, to test if background concentration of an element influenced the likelihood of transport or source limitation, we correlated b_{50sup} and b_{50inf} with C_{Q50} .

2.3. Trend Analysis

In the 1990s, in response to widespread eutrophication of freshwater and estuarine ecosystems, national and European legislation was implemented (e.g., the Water Framework Directive) to limit diffuse and point-source pollution from agriculture and waste water treatment plants [Lehmann and Rode, 2001; Hartmann et al., 2007; Minaudo et al., 2015]. We took advantage of differences in nutrient loading pre and post implementation to test for changes in C-Q dynamics. While there were not enough sampling points to test for differences in seasonality and multiple C-Q segments simultaneously [Hirsch et al., 2010; Zhang et al., 2016], we compared b_{50sup} , b_{50inf} , and C_{Q50} for parameters with sufficient data points (NO_3^- , TP, PO_4^{3-} , Cl^- , Mg^{2+} , Ca^{2+} , and TSS) from 1970–1990 and 1991–2010. Because this analysis required at least 150 C-Q couples for each of the sampling periods, it was only possible for 14 to 38 stations depending on the parameter, and was somewhat biased toward larger catchments since there were no eligible catchments smaller than 1,500 km². We quantified relative change of slopes (b_{50}) and intercepts (C_{Q50}) to assess whether the C-Q modality and background concentration changed between the two periods.

For analyses and data visualization, we used R version 3.3.0, Matlab 7.8, ArcGis 10.2.2, and Python for ArcGis.

3. Results

3.1. Prevalence of C-Q Types for Elements With Different Sources and Transport Dynamics

While all 9 potential C-Q modalities occurred for some parameters across the 293 catchments, they were not equally represented (Figure 3 and supporting information Figure S2). C-Q modality varied by element, with most elements having two dominant modalities accounting for over 75% of catchments, and none




Figure 3. Prevalence of up, flat, and down C-Q modalities below and above the median flow for water quality parameters and ratios in this study. All modalities that were present in more than 10% of catchments for each parameter are shown.

Table 2. Percentage of Catchments With Down, Flat, and Up C-Q Slopes Below the Median Flow (b_{50inf}) and Above the Median Flow (b_{50sup})

Parameter	b_{50inf}			b_{50sup}			b_{50inf} Different Than b_{50sup} ^a
	Down	Flat	Up	Down	Flat	Up	
DOC	14	66	20	1	18	81	72
NO_3^-	3	26	71	12	62	27	67
TP	39	54	8	7	51	43	68
PO_4^{3-}	44	47	9	20	58	23	52
SiO_2	40	47	13	19	56	25	51
Cl^-	50	39	11	50	12	37	44
SO_4^{2-}	21	61	18	10	21	69	66
Mg^{2+}	38	48	13	66	30	4	49
Ca^{2+}	39	44	17	61	34	5	56
TSS	12	66	23	1	7	93	76
Conductivity	50	39	11	70	26	4	47

^aPercentage of catchment-element combinations with statistically different b_{50inf} and b_{50sup} .

having more than three (Figure 3). Approximately half of all C-Q relationships for low flows (b_{50inf}) were flat, while high-flow C-Q coupling (b_{50sup}) was spread more equally between up (37%), flat (34%), and down (29%) modalities (Table 2). Three modalities did not occur in more than 15% of catchments for any given element: down-up, flat-down, and up-down (Figure 3). The flat-up pattern was the most common across parameters, accounting for 24% of all station-element combinations, and representing the dominant modality for TP, DOC, TSS, and SO_4^{2-} (Figures 3 and 4; supporting information Figure S2). Down-down and flat-flat (chemostatic) were the next most prevalent patterns, each accounting for 16% of all station-element combinations. Overall, 60% of catchment-element combinations had a bent C-Q curve (different b_{50inf} than b_{50sup}), but this varied widely by element, with over 70% of TSS and DOC catchments showing nonlinear C-Q curves and less than 50% for Mg^{2+} , conductivity, and Cl^- (Table 2).

There were several differences potentially associated with elemental characteristics (e.g., solubility, reactivity, and charge). Chemostatic modalities fell into two distinct categories, with elements associated with sediment transport (TP, PO_4^{3-} , and TSS) showing high variability (median CV for b_{50inf} and b_{50sup} between 67 and 75%), and more mobile elements (major ions, DOC, and NO_3^-) showing much lower variability (median CV from 12 to 45%; Table 1). CV was similar for all elements above and below the median flow (Table 1). While less distinct than the separation based on transport dynamics, there were several differences between biologically reactive (DOC, NO_3^- , TP, PO_4^{3-}) and less reactive elements (TSS and major ions). Though mobilization of TSS and phosphorus species is associated with high flow events, b_{50inf} and b_{50sup} were systematically lower for TP and PO_4^{3-} than TSS, with PO_4^{3-} in particular consistently expressing more flat or down slopes, suggesting source limitation or biological retention of phosphorus (Table 1 and Figure 4). In contrast with down behavior of less reactive ions, NO_3^- b_{50inf} was strongly up (median = 0.29), suggesting biological retention or removal during low flows. Finally, DOC and SO_4^{2-} showed similar dominant and secondary modalities (Table 1; Figures 3 and 4), implying shifts in source area, production, or consumption particularly during high flows.

3.2. Influence of Catchment Characteristics on C-Q Types

Relationships between slopes (b_{50inf} , b_{50sup}), and intercepts (C_{Q50}) with catchment characteristics (e.g., land use, hydrology, and biological parameters) are presented in Table 3 and Figures 5 and 6; supporting information Figures S4, S5, and S6. The next two paragraphs highlight some of the broader trends between explanatory variables and groups of elements, followed by one paragraph focused on differences for individual elements.

C_{Q50} was more sensitive than b_{50inf} and b_{50sup} to catchment characteristics (i.e., more significant correlations and patterns), and was particularly responsive to percentage arable land and specific discharge, which were correlated with C_{Q50} for all elements except DOC (Figures 5 and 6; supporting information Figure S4A; Table 3). There were fewer clear patterns explaining variability of C-Q slopes, though seasonality index and base flow index (BFI) were associated with most parameters (Table 3). b_{50inf} was most closely associated with seasonality index, mean summer chlorophyll and water temperature, and generally had more significant correlations than b_{50sup} , which was associated with seasonality index, population density and BFI (Figures 5 and




Figure 4. Frequency histograms of b_{50inf} and b_{50sup} for the 293 catchments in this study. (a) Solutes and particulates. (b) Elemental ratios.

6; supporting information Figures S4, S5, and S6; Table 3). Of all the explanatory variables, BFI and seasonality index were associated with the largest number of individual slopes and intercepts across elements. BFI had particularly clear relationships with b_{50inf} and C_{Q50} of DOC and b_{50sup} of TSS, though this was notably not the case for nonbiologically reactive ions such as chlorine, calcium or magnesium (Table 3). Seasonality index was strongly negatively associated with less-reactive elements (e.g., Cl^- , Mg^{2+} , Ca^{2+}), suggesting that seasonal dilution is a primary control of slope and intercept for these parameters. Conversely, the relationship between seasonality index and more reactive elements (e.g., DOC, NO_3^- , TP, PO_4^{3-}) was weaker and




Figure 5. Relationships between dissolved organic carbon (DOC), nitrate (NO_3^-), and total phosphorus (TP) with (a) spatial scale, (b) base flow index (the proportion of annual flow occurring during base flow conditions [Sloto and Crouse, 1996]), (c) mean summer chlorophyll a concentration, and (d) percent agricultural land use. $b_{50\text{inf}}$ and $b_{50\text{sup}}$ are the log C-Q slopes below and above the median discharge, respectively, and C_{Q50} is the concentration at the median discharge.

more varied (i.e., both positive and negative correlations), indicating a decoupling of seasonal hydrological variation with C-Q dynamics, potentially due to biological processes.

There were three general patterns that held for all elements, when relationships were significant. First, $b_{50\text{inf}}$, $b_{50\text{sup}}$, and C_{Q50} , were always negatively correlated with specific discharge, meaning that catchments with higher water yield had lower median concentrations and were more likely to express dilution (source limitation) dynamics (supporting information Figure S5A and Table 3). Second, with the exception of $b_{50\text{sup}}$ for NO_3^- , which was weakly negative, the percentage arable land was always positively correlated with slopes and intercepts, meaning that catchments with more agricultural land use had higher concentrations and steeper C-Q slopes (Figures 5d and 6d; supporting information Figure S4B). Third, C_{Q50} was always negatively correlated with $b_{50\text{sup}}$, meaning that catchments with higher concentrations of a given element were more likely to be chemostatic or experience source limitation during high flows (Table 3 and supporting information Figure S6D).

Multiple patterns emerged of catchment effects on individual element C-Q slopes and intercepts. While C_{Q50} for most elements was positively correlated with percentage of agricultural land, PO_4^{3-} and TP C_{Q50} were better correlated with population density (supporting information Figure S4 and Table 3). DOC C_{Q50} varied strongly with hydrological metrics, increasing with W2 and decreasing with BFI and Q_{MNA5} (supporting information Figure S5 and Table 3). Ca^{2+} C_{Q50} was positively correlated with carbonate lithology (chalk, marl, limestone) and secondarily with arable land (supporting information Figure S4). Mean summer chlorophyll, a proxy for primary production, was associated with lower NO_3^- C_{Q50} , higher NO_3^- $b_{50\text{inf}}$, and a convergence toward chemostasis for TP and PO_4^{3-} , suggesting algal modulation of NO_3^- at low flows and for phosphorus across the hydrograph (Figure 5c and supporting information Figure S6B). Variability of C_{Q50} between catchments decreased with increasing catchment size for all parameters except TSS (Figure 5a and supporting information Figure S4). C-Q slopes for NO_3^- and phosphorus species often showed opposite patterns, including: 1. convergence of NO_3^- $b_{50\text{sup}}$ and $b_{50\text{inf}}$ with increasing BFI versus diverging phosphorus slopes (Figure 5B and supporting information Figure S5B), 2. converging phosphorus slopes with summer chlorophyll, versus diverging NO_3^- slopes (Figure 5C and supporting information Figure S6B), and 3. NO_3^- $b_{50\text{inf}}$ was systematically higher than $b_{50\text{sup}}$, whereas the opposite was true for TP and PO_4^{3-} .




Figure 6. Relationships between Chloride (Cl^-), calcium (Ca^{2+}), and total suspended solids (TSS) with (a) spatial scale, (b) base flow index, (c) mean summer chlorophyll a concentration, and (d) Percent agricultural land use. $b_{50\text{inf}}$ and $b_{50\text{sup}}$ are the log C-Q slopes below and above the median discharge, respectively, and C_{Q50} is the concentration at the median discharge.

Table 3. Statistically Significant ($\alpha < 0.05$) Spearman's Correlations for C_{Q50} , $b_{50\text{inf}}$ and $b_{50\text{sup}}$ in Relation to Catchment, Hydrological, and Biological Characteristics

Element	Parameter	Catchment Area	Arable Land	Carbonate Bedrock	Population Density	Specific Discharge	Base Flow Index	$Q_{\text{mna}5}$	W2	Seasonality Index	Summer Temp. °C	Summer Chlorophyll	C_{Q50}
DOC	$b_{50\text{inf}}$				-0.20		0.37	0.29	-0.34	0.21		-0.26	-0.29
	$b_{50\text{sup}}$		0.21	0.30		-0.36	0.26						-0.51
	C_{Q50}	-0.38		-0.26			-0.55	-0.54	0.57			0.26	
NO_3^-	$b_{50\text{inf}}$	0.30				-0.19	-0.21	-0.40	0.23	0.40	0.54	0.48	0.14
	$b_{50\text{sup}}$		-0.13				-0.19	-0.16		0.34	0.18		-0.31
	C_{Q50}	-0.24	0.70	0.22	0.21	-0.37	0.29			-0.31	-0.22		
TP	$b_{50\text{inf}}$	-0.14			-0.20	-0.20	-0.13	-0.25	0.22	-0.38		0.20	
	$b_{50\text{sup}}$	-0.14			-0.33		-0.16	-0.15	0.24	-0.37	-0.14	0.17	-0.38
	C_{Q50}		0.36		0.77	-0.44	0.27		-0.15		0.17	0.19	
PO_4^{3-}	$b_{50\text{inf}}$				-0.16	-0.17	-0.15	-0.30	0.22	-0.31		0.31	-0.14
	$b_{50\text{sup}}$				-0.44		-0.29	-0.27	0.32			0.25	-0.56
	C_{Q50}	0.11	0.39	0.22	0.76	-0.48	0.35		-0.26				
Cl^-	$b_{50\text{inf}}$	-0.31	0.20		-0.34			-0.21	0.25	-0.52	-0.18		
	$b_{50\text{sup}}$	-0.21			-0.25					-0.47	-0.25	-0.22	-0.18
	C_{Q50}	-0.34	0.57		0.36	-0.38		-0.38	0.39			0.27	
SO_4^{2-}	$b_{50\text{inf}}$									-0.33			
	$b_{50\text{sup}}$												-0.51
	C_{Q50}		0.33	0.44	0.48	-0.57	0.36						
Mg^{2+}	$b_{50\text{inf}}$	-0.27	0.36			-0.23	-0.18	-0.34	0.35	-0.60		0.22	0.22
	$b_{50\text{sup}}$									-0.31			-0.33
	C_{Q50}	-0.28	0.51		0.38	-0.32		-0.34	0.35	-0.26		0.18	
Ca^{2+}	$b_{50\text{inf}}$	0.20	0.62	0.49		-0.47	0.31			-0.68	0.23	0.21	0.54
	$b_{50\text{sup}}$		0.19	-0.29						-0.21		-0.17	-0.31
	C_{Q50}	0.32	0.59	0.83	0.28	-0.75	0.28			-0.33	0.27		
TSS	$b_{50\text{inf}}$	-0.19	0.15	0.15			0.22	0.24	-0.20	0.26	0.37	-0.32	
	$b_{50\text{sup}}$		0.22	0.41	0.21	-0.22	0.37	0.36	-0.39	0.53	-0.13	-0.37	-0.14
	C_{Q50}	0.36	0.25	0.22	0.36	-0.28	0.37	0.17	-0.34	-0.26	0.27	0.23	

Table 4. Changes in Median Flow Concentration and Stationarity of C-Q Slopes Before and After Implementation of Water Quality Legislation

Element	N (stations)	Change Between 1970–1990 and 1991–2010 (%)			Stationarity of C-Q Modality (%)	
		ΔC_{Q50}	Δb_{50inf}	Δb_{50sup}	b_{50inf}	b_{50sup}
NO_3^-	38	+41	+82	-62	74	53
TP	14	-45	+51	-121	64	50
PO_4^{3-}	36	-46	+97	+13	72	61
Cl^-	16	+21	-6	-28	94	87
Mg^{2+}	22	+11	+104	+6	50	41
Ca^{2+}	22	+4	+85	+3	68	55
TSS	31	-18	-1561	-27	65	87

3.3. Effects of Water Quality Mitigation on C-Q Slopes and Intercepts

For the stations with adequate sampling for the two time periods (1970–1990 and 1991–2010), there was little change in C_{Q50} for Ca^{2+} and Mg^{2+} (Table 4). However, there were substantial increases in NO_3^- and Cl^- and substantial decreases in PO_4^{3-} , TP, and TSS (Figure 7 and Table 4). Changes in TP C_{Q50} were accompanied by shifts in C-Q shape, with

approximately half of catchments changing modality, primarily becoming more chemostatic (Table 4), indicating substantial change in how TP was delivered to waterways. In contrast, Cl^- , NO_3^- , PO_4^{3-} , and TSS C-Q modalities were largely invariant despite substantial changes in C_{Q50} , except the b_{50sup} of NO_3^- where almost half of catchments changed modality, primarily from flat to up (Table 4). Mg^{2+} modality was the most variable in terms of percentage of catchments experiencing change, though this was due to a large number of catchments with C-Q slopes on the cusp of statistical significance, and actual changes were minor (Table 4). Altogether, C-Q shape remained the same between the two time periods for approximately




Figure 7. Examples of change in C-Q patterns between 1970–1990 and 1990–2010 for (a) NO_3^- and (b) PO_4^{3-} for two streams in the Loire catchment, the Seine River drainage basin and Brittany. These sites are representative of the direction and magnitude of change observed for most sites but see Table 4 for a summary of change across the data set.

70% of all element-basin combinations. For the 30% that changed, flat-flat to up-flat was the most common shift.

4. Discussion

The causes and prevalence of C-Q curves for different elements across catchments remains unknown, despite decades of research [Hem, 1948; Edwards, 1973; Thompson *et al.*, 2011]. To address this issue, we analyzed chemistry and discharge data collected over 40 years from 293 water-monitoring stations draining half of France. Our analysis of segmented C-Q slopes [Meybeck and Moatar, 2012] revealed several surprising patterns, including flat (chemostatic) behavior of TSS for low flows in 66% of catchments, up behavior of NO_3^- during low flows for 71% of catchments, and up behavior of DOC in 80% of catchments during high flows independent of catchment size. While dilution effects dominated C-Q behavior for nonreactive elements, we found evidence for biological modulation of nitrogen, phosphorus, and TSS, particularly during low flows. Long-term efforts to reduce nutrient loading in the studied catchments effectively decreased phosphorus concentration and altered phosphorus C-Q curves, but NO_3^- increased over the same period, emphasizing the difficulty of reducing pollutants with long residence times and diffuse sources [Kolbe *et al.*, 2016]. We organize the discussion of these results around our four initial questions, rephrased here as subsections.

4.1. How General Are C-Q Relationships Across Catchments and How Many of the Nine Possible C-Q Shapes Actually Occur for Each Parameter?

Despite the diversity of catchment characteristics in our study, only two or three C-Q modalities occurred for any given element. This suggests that intrinsic elemental properties (e.g., solubility, reactivity, charge) and extrinsic element-specific differences (e.g., systematic differences in the location of solute sources within and between catchments) exert a strong control on C-Q dynamics. While it is hardly novel to observe that the properties of an element influence how it is sourced and propagated through a system [Basu *et al.*, 2011; Musloff *et al.*, 2015], our study supports the hypothesis that elemental properties set a basic C-Q template that is secondarily influenced by environmental conditions such as land-use, topography, hydrology, and biological activity. The long-term stationarity of C-Q modality for 70% of catchment-element combinations supports this hypothesis, as do previous observations of consistent C-Q relationships across climatic and ecological contexts for major ions (Cl^- , SO_4^{2-} , Mg^{2+} , Ca^{2+}), nutrients, and DOC [Godsey *et al.*, 2009; Meybeck and Moatar, 2012; Thomas *et al.*, 2016]. However, as the title of our paper suggests, we are not implying that elemental properties are solely responsible for C-Q modality. Catchment characteristics strongly influence solute concentrations and may also influence C-Q shape in some circumstances, such as when both hydrology and solute distribution are affected by tile draining and fertilizer application [Musloff *et al.*, 2015]. It is also important to note that our data set consists of low-frequency, long-term measurements. While these data allow powerful description of general patterns, they inevitably obscure short-term C-Q dynamics, when hydrology, biological activity, and antecedent distribution of solute sources likely play a larger role, as evidenced by widespread and diverse observations of hysteresis at the event scale [Evans and Davies, 1998; Hornberger *et al.*, 2001; Chanut *et al.*, 2002; Strohmeier *et al.*, 2013; Thomas *et al.*, 2016].

We found that less than 16% of all C-Q slopes were flat across the hydrograph, contrasting recent work that suggests predominance of chemostatic C-Q relationships for DOC and major ions [Godsey *et al.*, 2009; Creed *et al.*, 2015]. Two reasons for this difference are: 1. the large size of our data set, which increased statistical power to identify significant relationships, and 2. the hydrograph segmentation technique, which avoids confounding patterns for low and high flows [Meybeck and Moatar, 2012]. While C-Q segmentation requires more data than a single analysis across the hydrograph, the fact that approximately two thirds of all catchment-element combinations in our study had a statistically different C-Q slope above and below the median daily flow demonstrates the potential information cost of linear or monotonic analysis of C-Q relationships. While the vast majority of C-Q relationships in our study were chemodynamic, C-Q slopes were relatively low for most parameters, indicating that discharge rather than concentration controls flux for most parameters [Moatar *et al.*, 2013]. In this sense, our study agrees with previous observations that hydrological change varies orders of magnitude more than changes in concentration [Meybeck *et al.*, 2003; Godsey *et al.*, 2009; Creed *et al.*, 2015]. The difference in amplitude of hydrological and biogeochemical fluctuations has been attributed to abundant and homogeneously distributed solute sources, resulting in little change in

concentration with expansion of the terrestrial-aquatic nexus, and covariation between production processes (e.g., weathering, decomposition) and transport processes controlled by hydrology [Godsey et al., 2009; Thompson et al., 2011; Musolff et al., 2015]. This ubiquitous-source hypothesis was supported by the positive relationship between median-flow concentration and flat C-Q behavior, though we point out that the lack of a significant C-Q slope does not indicate temporal chemostasis. Even our low-frequency data set revealed that statistically chemostatic parameters such as TP and PO_4^{3-} still varied dramatically at seasonal and event time-scales, making these parameters more difficult to conceptualize and predict [Johnes, 2007; Cassidy and Jordan, 2011; Jones et al., 2011; Hirsch, 2014; Dupas et al., 2015].

4.2. Biogeochemical and Hydrological Constraints on C-Q Shapes

Several lines of evidence suggest that C-Q response at low flows can be influenced by biogeochemical processes, but that hydrology becomes dominant during high flows (Figure 8). For example, while NO_3^- C-Q relationships are widely reported as chemostatic or negative [Alvarez-Cobelas et al., 2008; Dupas et al., 2016; Thomas et al., 2016], our segmented C-Q curves revealed strong up behavior for the lower half of the hydrograph in 71% of catchments. This could be due to transport limitation (i.e., mobilization of stranded NO_3^- sources as flow increases) or decreases in the relative importance of biogeochemical processes mediating delivery (e.g., denitrification and assimilation). The decrease in $\text{Cl}^-:\text{NO}_3^-$ ratio with discharge, and positive correlation of NO_3^- $b_{50\text{inf}}$ with summer temperature and chlorophyll concentration (both of which are associated with biological processes) support the second hypothesis that biogeochemical processes control low-flow NO_3^- concentration. During low flows, water residence time and the ratio of streambed surface to water volume are high, favoring NO_3^- uptake by attached and sestonic primary producers, and NO_3^- removal in riparian and hyporheic zones by denitrifying microorganisms [Peterson et al., 2001; Mulholland et al., 2008; Zarnetske et al., 2012]. As discharge increases, the capacity of these processes to alter NO_3^- flux decreases rapidly. Similarly, the decoupling of phosphorus and TSS dynamics appears to be partially because of algal effects (based on the correlation with chlorophyll), which decrease in importance as discharge increases. The negative relationship of the $b_{50\text{inf}}$ for both NO_3^- and phosphorus with specific discharge and Q_{mna5} further underlines the importance of base-flow hydrology in modulating nutrient removal capacity. These findings suggest that $b_{50\text{inf}}$ could be used as a proxy of biogeochemical capacity to retain nutrients between streams or for a single stream through time.

Surprisingly, TSS behavior also suggested biogeochemical control of TSS C-Q response during low flows. Though the C-Q of TSS is almost universally reported as strongly positive [Asselman, 2000; Shanley et al., 2011; Raymond et al., 2013], two-thirds of the catchments in our study showed flat TSS C-Q for the bottom




Figure 8. Conceptual diagram of the interplay between biological and hydrological dynamics in determining C-Q relationships for reactive and conservative elements. Hydrological dynamics (dilution effects and changing source area) dominate C-Q slope for all elements above the median flow. At low flows, hydrology is less dominant and biological retention or removal of carbon, nitrogen, and phosphorus become increasingly important, resulting in up behavior for biologically-reactive elements during low-flow conditions (see section 4.2 for detailed discussion).

half of the hydrograph. While erosion and transport processes control TSS delivery during flood events, resulting in strong C-Q slopes, during low flows, algal die-off and biogenic calcite precipitation under high water temperature and pH conditions can influence TSS independent of discharge [Moatar et al., 1999, 2001; Grosbois et al., 2001], as suggested by the positive correlation of Ca^{2+} $b_{50\text{inf}}$ with chlorophyll. This low-flow chemostasis of TSS may have been overlooked in other systems since it would not be detected in short-term studies with a smaller range of hydrological conditions, or if a single C-Q coefficient was calculated. Taken together, the C-Q

patterns of nitrogen, phosphorus, and TSS suggest a model of biological control during low flows with increasing influence from hydrological parameters at high flows (Figure 8). Because the majority of elemental fluxes occur during high flow periods [Meybeck and Moatar, 2012], hydrological controls, rather than biogeochemical controls, modulate fluxes, but biological activity (e.g., denitrification and primary production) influence water quality during the majority of the year (Figure 8). Taking NO_3^- as an example, median concentration is primarily a function of arable land cover (Table 3). When flows are moderate or low, denitrification and uptake by primary producers effectively decrease NO_3^- concentration, improving water quality and habitat throughout the stream network. However, as discharge increases, biological removal capacity becomes insignificant relative to the mass of NO_3^- moving through the system, causing flat C-Q behavior, where NO_3^- flux is a product of median concentration (as determined by land use) and discharge (Figure 8). From a hydrological or “downstream” perspective, one could conclude that biological effects are insignificant, since nutrient delivery to lacustrine and estuarine ecosystems is determined by the interaction between land-use and hydrology. However, from an ecological perspective, freshwater organisms experience concentration, not flux, and biological effects that improve inland water quality even during a part of the year could be important. Because biological mitigation of nutrient pollution is greatest during low flows, when many aquatic organisms pass through critical life stages and are exposed to multiple stressors including low oxygen, high temperature, and decreased habitat [Heugens et al., 2001; Davies and Jackson, 2006; Davis et al., 2015], interventions to improve instream nutrient removal could have substantial benefits in addition to catchment-level efforts to reduce pollutant loading [Pinay et al., 2015; Abbott et al., 2016; Mendoza-Lera and Datry, 2016].

4.3. Does Catchment Size Matter to C-Q Response for Individual Elements?

Though catchment size and associated parameters (e.g., residence time, heterogeneity of source areas, and instream processing) can influence carbon and nutrient dynamics [Ensign and Doyle, 2006; Cory et al., 2014; Catalán et al., 2016], we observed no change in DOC C-Q slopes and only minor changes for nutrients across catchments ranging from 50 to 110,000 km². Concentrations of DOC and NO_3^- did decrease moving downstream, but at the same rate as more conservative elements (Cl^- and Mg^{2+}), suggesting that dilution rather than biological or photochemical removal accounted for the decreases. The predominance of positive DOC C-Q slopes across spatial scales, and particularly during high flows, supports the emerging perspective of hydrological control of DOC flux [Raymond et al., 2016; Moatar et al., 2013]. It also contradicts recent work reporting widespread DOC chemostasis and convergence toward stable DOC concentration with catchment size [Creed et al., 2015]. While residence time, DOC concentration, and biodegradability can be interconnected [Zarnetske et al., 2011; Abbott et al., 2016; Catalán et al., 2016], even the most biodegradable DOC requires weeks to months to be mineralized [Abbott et al., 2014; Larouche et al., 2015; Vonk et al., 2015]. This means that DOC processing timescales are too slow to substantially alter concentration during flood events, and since C-Q modality for DOC is largely determined by high flows, catchment size has little to no influence on C-Q slopes for DOC.

The $b_{50\text{inf}}$ of NO_3^- was the only nutrient C-Q slope to substantially change with catchment size, becoming more positive moving downstream. This shift could be partially due to changes in agricultural activity moving from uplands to lowlands [Musolff et al., 2015; Thomas et al., 2016], but the decrease in Cl^- $b_{50\text{inf}}$ (also associated with agriculture) suggests that the trend is due to the distributed biological effects described in section 4.2. Moving downstream, the increasingly positive C-Q slope during low flows is caused by the cumulative effect of uptake and denitrification in riparian, hyporheic, and instream environments [Burt and Pinay, 2005; Gruber and Galloway, 2008; Mulholland et al., 2008], supporting our conceptual model of biological control during low flows at multiple spatial scales (Figure 8).

4.4. How has Land-Use Change Altered the C-Q Relationships for the Catchments With Long-Term Records?

The comparison of data from 1970–1990 and 1991–2010 revealed that C-Q modalities remained the same for 70% of catchment-element combinations, supporting our hypothesis that both intrinsic and extrinsic elemental properties constrain C-Q shape. This is demonstrated by the contrasting changes observed between the two periods for nitrogen and phosphorus. Phosphorus concentration decreased with time and many catchments experienced changes in C-Q modality, indicating fundamental shifts in phosphorus sources or transport mechanisms. While hydrology has not systematically changed between the time periods, there has been a concerted effort to eliminate phosphorus point sources (e.g., waste water treatment

plants) and this is reflected in the decrease in phosphorus concentration [Minaudo *et al.*, 2015; McDonald *et al.*, 2016], especially for more urban catchments where most human waste is treated (Figure 6 and supporting information Figure S4C). Counterintuitively, the C-Q slope for phosphorus became substantially more positive (increase in b_{50inf}) over the time period, but this was due to decreases in phosphorus concentration at very low flows rather than increases at high flows, emphasizing the importance of considering both C-Q slope and intercept when assessing water quality. NO_3^- concentration on the other hand, increased across the hydrograph between the two time periods (e.g., Figure 7), with few catchments changing C-Q modality. This increase in NO_3^- underlines the difficulty of controlling nonpoint source pollution, particularly since NO_3^- can be stored in groundwater and soil on decadal and centennial timescales [Sebilio *et al.*, 2013; Kolbe *et al.*, 2016], meaning more time is needed for changes in agricultural practice to be reflected in stream concentrations and fluxes. These differences in the effectiveness of phosphorus and nitrogen mitigation raise an unexpected paradox. It is arguably true that our capacity to conceptualize and model nutrient flux is greater for NO_3^- than for phosphorus [Basu *et al.*, 2010; Birgand *et al.*, 2010; Herndon *et al.*, 2015; Pinay *et al.*, 2015; Abbott *et al.*, 2016; Thomas *et al.*, 2016]. However, as shown here, our ability to mitigate nutrient pollution is not necessarily tied to our capacity to predict it.

5. Conclusion

We calculated segmented C-Q slopes and intercepts for four decades of water quality data from 293 catchments in France. We found systematic differences in C-Q behavior for carbon, nutrients, major ions, and suspended sediment. Only two or three of the nine possible C-Q modalities were manifest for each parameter, and these modalities were largely stable through time, suggesting that C-Q dynamics were constrained by intrinsic and extrinsic elemental properties (e.g., solubility, reactivity, and source dynamics). These properties set a basic C-Q template for each parameter, which was secondarily influenced by biological activity during low flows, and the interaction between hydrology and catchment characteristics at high flows. The size of our data set and the segmented C-Q analysis revealed several patterns that challenge current views on C-Q dynamics including predominantly flat (chemostatic) behavior of TSS for the lower half the hydrograph, biological mediation of NO_3^- concentration increasing with catchment size, and positive C-Q behavior of DOC independent of catchment size. Changes over the 40 year record suggest that mitigation of phosphorus pollution reduced concentration and altered C-Q slopes. However, for NO_3^- , concentration continued to rise and C-Q dynamics remained the same despite long-term mitigation efforts. This study demonstrates how large, low-frequency data sets can be analyzed with three conceptually simple indices of ecohydrological behavior (b_{50inf} , b_{50sup} , and C_{Q50}) that can be used as sensitive indicators of interactions between hydrology, biology, and catchment characteristics.

Acknowledgments

Derived data used in this analysis are included in the supporting information. F. Moatar and C. Minaudo were supported by the MAKARA project on the social construction of water quality from the French National Agency (ANR-12-SENV-0009-01). F. Moatar was also supported by a sabbatical grant from the Centre National de la Recherche Scientifique (CNRS). G. Pinay and B. Abbott were supported by the European Union's Seventh Framework Program for research, technological development and demonstration under grant agreement 607150 (FP7-PEOPLE-2013-ITN-INTERFACES - Ecohydrological interfaces as critical hotspots for transformations of ecosystem exchange fluxes and biogeochemical cycling). We thank Loire-Bretagne and Seine Normandie Agencies for long-term water quality acquisition and for giving access to their data set, Michel Meybeck for input on an early version of the manuscript, and Jan Fleckenstein and two anonymous reviewers for thoughtful and detailed criticism. F. Moatar and B.W. Abbott contributed equally to this publication.

References

- Abbott, B. W., J. R. Larouche, J. B. Jones, W. B. Bowden, and A. W. Balsler (2014), Elevated dissolved organic carbon biodegradability from thawing and collapsing permafrost: Permafrost carbon biodegradability, *J. Geophys. Res. Biogeosci.*, *119*, 2049–2063, doi:10.1002/2014JG002678.
- Abbott, B. W., et al. (2016), Using multi-tracer inference to move beyond single-catchment ecohydrology, *Earth Sci. Rev.*, *160*, 19–42, doi:10.1016/j.earscirev.2016.06.014.
- Aitkenhead, J. A., and W. H. McDowell (2000), Soil C:N ratio as a predictor of annual riverine DOC flux at local and global scales, *Global Biogeochem. Cycles*, *14*(1), 127–138, doi:10.1029/1999GB900083.
- Alewell, C., G. Lischeid, U. Hell, and B. Manderscheid (2004), High temporal resolution of ion fluxes in semi-natural ecosystems—Gain of information or waste of resources?, *Biogeochemistry*, *69*(1), 19–35, doi:10.1023/B:BIOG.0000031029.46798.7f.
- Alvarez-Cobelas, M., D. G. Angeler, and S. Sánchez-Carrillo (2008), Export of nitrogen from catchments: A worldwide analysis, *Environ. Pollut.*, *156*(2), 261–269, doi:10.1016/j.envpol.2008.02.016.
- Asselman, N. E. M. (2000), Fitting and interpretation of sediment rating curves, *J. Hydrol.*, *234*(3–4), 228–248, doi:10.1016/S0022-1694(00)00253-5.
- Basu, N. B., et al. (2010), Nutrient loads exported from managed catchments reveal emergent biogeochemical stationarity, *Geophys. Res. Lett.*, *37*, L23404, doi:10.1029/2010GL045168.
- Basu, N. B., S. E. Thompson, and P. S. C. Rao (2011), Hydrologic and biogeochemical functioning of intensively managed catchments: A synthesis of top-down analyses, *Water Resour. Res.*, *47*, W00J15, doi:10.1029/2011WR010800.
- Bierozza, M. Z., and A. L. Heathwaite (2015), Seasonal variation in phosphorus concentration–discharge hysteresis inferred from high-frequency in situ monitoring, *J. Hydrol.*, *524*, 333–347, doi:10.1016/j.jhydrol.2015.02.036.
- Birgand, F., C. Faucheux, G. Gruau, B. Augereau, F. Moatar, and P. Bordenave (2010), Uncertainties in assessing annual nitrate loads and concentration indicators. Part 1: Impact of sampling frequency and load estimation algorithms, *Trans. Am. Soc. Agric. Biol. Eng.*, *53*(2), 437–446.

- Bowes, M. J., W. A. House, R. A. Hodgkinson, and D. V. Leach (2005), Phosphorus-discharge hysteresis during storm events along a river catchment: The River Swale, UK, *Water Res.*, *39*(5), 751–762, doi:10.1016/j.watres.2004.11.027.
- Bowes, M. J., H. P. Jarvie, P. S. Naden, G. H. Old, P. M. Scarlett, C. Roberts, L. K. Armstrong, S. A. Harman, H. D. Wickham, and A. L. Collins (2014), Identifying priorities for nutrient mitigation using river concentration-flow relationships: The Thames basin, UK, *J. Hydrol.*, *517*, 1–12, doi:10.1016/j.jhydrol.2014.03.063.
- Burt, T. P., and G. Pinay (2005), Linking hydrology and biogeochemistry in complex landscapes, *Prog. Phys. Geogr.*, *29*(3), 297–316, doi:10.1191/0309133305pp450ra.
- Butturini, A., M. Alvarez, S. Bernal, E. Vazquez, and F. Sabater (2008), Diversity and temporal sequences of forms of DOC and NO₃-discharge responses in an intermittent stream: Predictable or random succession?, *J. Geophys. Res.*, *113*, G03016, doi:10.1029/2008JG000721.
- Cassidy, R., and P. Jordan (2011), Limitations of instantaneous water quality sampling in surface-water catchments: Comparison with near-continuous phosphorus time-series data, *J. Hydrol.*, *405*(1–2), 182–193, doi:10.1016/j.jhydrol.2011.05.020.
- Catalán, N., R. Marcé, D. N. Kothawala, and L. J. Tranvik (2016), Organic carbon decomposition rates controlled by water retention time across inland waters, *Nat. Geosci.*, *9*(7), 501–504, doi:10.1038/ngeo2720.
- Chanat, J. G., K. C. Rice, and G. M. Hornberger (2002), Consistency of patterns in concentration-discharge plots, *Water Resour. Res.*, *38*(8), 1147, doi:10.1029/2001WR000971.
- Cory, R. M., C. P. Ward, B. C. Crump, and G. W. Kling (2014), Sunlight controls water column processing of carbon in arctic fresh waters, *Science*, *345*(6199), 925–928, doi:10.1126/science.1253119.
- Creed, I. F., et al. (2015), The river as a chemostat: Fresh perspectives on dissolved organic matter flowing down the river continuum, *Can. J. Fish. Aquat. Sci.*, *72*(8), 1272–1285, doi:10.1139/cjfas-2014-0400.
- Davies, S. P., and S. K. Jackson (2006), The biological condition gradient: A descriptive model for interpreting change in aquatic ecosystems, *Ecol. Appl.*, *16*(4), 1251–1266, doi:10.1890/1051-0761(2006)016[1251:TBCGAD]2.0.CO;2.
- Davis, J., et al. (2015), When trends intersect: The challenge of protecting freshwater ecosystems under multiple land use and hydrological intensification scenarios, *Sci. Total Environ.*, *534*, 65–78, doi:10.1016/j.scitotenv.2015.03.127.
- Doyle, M. W., E. H. Stanley, D. L. Strayer, R. B. Jacobson, and J. C. Schmidt (2005), Effective discharge analysis of ecological processes in streams, *Water Resour. Res.*, *41*, W11411, doi:10.1029/2005WR004222.
- Dupas, R., F. Curie, C. Gascuel-Odoux, F. Moatar, M. Delmas, V. Parnaudeau, and P. Durand (2013), Assessing N emissions in surface water at the national level: Comparison of country-wide vs. regionalized models, *Sci. Total Environ.*, *443*, 152–162, doi:10.1016/j.scitotenv.2012.10.011.
- Dupas, R., C. Gascuel-Odoux, N. Gilliet, C. Grimaldi, and G. Gruau (2015), Distinct export dynamics for dissolved and particulate phosphorus reveal independent transport mechanisms in an arable headwater catchment, *Hydrol. Processes*, *29*, 3162–3178, doi:10.1002/hyp.10432.
- Dupas, R., S. Jomaa, A. Musolff, D. Borchardt, and M. Rode (2016), Disentangling the influence of hydroclimatic patterns and agricultural management on river nitrate dynamics from sub-hourly to decadal time scales, *Sci. Total Environ.*, *571*, 791–800, doi:10.1016/j.scitotenv.2016.07.053.
- Edwards, A. M. C. (1973), The variation of dissolved constituents with discharge in some Norfolk rivers, *J. Hydrol.*, *18*(3), 219–242, doi:10.1016/0022-1694(73)90049-8.
- Ensign, S. H., and M. W. Doyle (2006), Nutrient spiraling in streams and river networks, *J. Geophys. Res.*, *111*, G04009, doi:10.1029/2005JG000114.
- European Union (2000), Framework for Community action in the field of water policy. European Union Directive 2000/60/EC, *Off. J. Eur. Commun.*, *372*, 72.
- Evans, C., and T. D. Davies (1998), Causes of concentration/discharge hysteresis and its potential as a tool for analysis of episode hydrochemistry, *Water Resour. Res.*, *34*(1), 129–137, doi:10.1029/97WR01881.
- Gall, H. E., J. Park, C. J. Harman, J. W. Jawitz, and P. S. C. Rao (2013), Landscape filtering of hydrologic and biogeochemical responses in managed catchments, *Landscape Ecol.*, *28*(4), 651–664, doi:10.1007/s10980-012-9829-x.
- Genereux, D. P., H. F. Hemond, and P. J. Mulholland (1993), Use of radon-222 and calcium as tracers in a three-end-member mixing model for streamflow generation on the West Fork of Walker Branch Watershed, *J. Hydrol.*, *142*(1), 167–211, doi:10.1016/0022-1694(93)90010-7.
- Godsey, S. E., J. W. Kirchner, and D. W. Clow (2009), Concentration–discharge relationships reflect chemostatic characteristics of US catchments, *Hydrol. Processes*, *23*(13), 1844–1864, doi:10.1002/hyp.7315.
- Godsey, S. E., et al. (2010), Generality of fractal 1/f scaling in catchment tracer time series, and its implications for catchment travel time distributions, *Hydrol. Processes*, *24*(12), 1660–1671, doi:10.1002/hyp.7677.
- Grosbois, C., P. Négrel, D. Grimaud, and C. Fouillac (2001), An overview of dissolved and suspended matter fluxes in the Loire River Basin: Natural and anthropogenic inputs, *Aquat. Geochem.*, *7*(2), 81–105, doi:10.1023/A:1017518831860.
- Gruber, N., and J. N. Galloway (2008), An Earth-system perspective of the global nitrogen cycle, *Nature*, *451*(7176), 293–296, doi:10.1038/nature06592.
- Guzman, C. D., S. A. Tilahun, A. D. Zegeye, and T. S. Steenhuis (2013), Suspended sediment concentration–discharge relationships in the (sub-) humid Ethiopian highlands, *Hydrol. Earth Syst. Sci.*, *17*(3), 1067–1077, doi:10.5194/hess-17-1067-2013.
- Hartmann, J., N. Jansen, S. Kempe, and H. H. Dürr (2007), Geochemistry of the river Rhine and the Upper Danube: Recent trends and lithological influence on baselines, *J. Environ. Sci. Sustainable Soc.*, *1*, 39–46.
- Hem, J. D. (1948), Fluctuations in concentration of dissolved solids of some southwestern streams, *Eos Trans. AGU*, *29*(1), 80–84, doi:10.1029/TR029i001p00080.
- Herndon, E. M., A. L. Dere, P. L. Sullivan, D. Norris, B. Reynolds, and S. L. Brantley (2015), Landscape heterogeneity drives contrasting concentration–discharge relationships in shale headwater catchments, *Hydrol. Earth Syst. Sci.*, *19*(8), 3333–3347, doi:10.5194/hess-19-3333-2015.
- Heugens, E. H. W., A. J. Hendriks, T. Dekker, N. M. van Straalen, and W. Admiraal (2001), A review of the effects of multiple stressors on aquatic organisms and analysis of uncertainty factors for use in risk assessment, *Crit. Rev. Toxicol.*, *31*(3), 247–284, doi:10.1080/2001409111695.
- Hirsch, R. M. (2014), Large biases in regression-based constituent flux estimates: Causes and diagnostic tools, *J. Am. Water Resour. Assoc.*, *50*(6), 1401–1424, doi:10.1111/jawr.12195.
- Hirsch, R. M., D. L. Moyer, and S. A. Archfield (2010), Weighted Regressions on Time, Discharge, and Season (WRTDS), with an application to Chesapeake Bay river inputs, *J. Am. Water Resour. Assoc.*, *46*(5), 857–880, doi:10.1111/j.1752-1688.2010.00482.x.
- Holmes, R. M., et al. (2012), Seasonal and annual fluxes of nutrients and organic matter from large rivers to the Arctic Ocean and surrounding seas, *Estuaries Coasts*, *35*(2), 369–382, doi:10.1007/s12237-011-9386-6.
- Hooper, R. P., N. Christophersen, and N. E. Peters (1990), Transfer of elements through the hydrological cycle modelling streamwater chemistry as a mixture of soilwater end-members—An application to the Panola Mountain catchment, Georgia, U.S.A., *J. Hydrol.*, *116*(1), 321–343, doi:10.1016/0022-1694(90)90131-G.

- Hornberger, G. M., T. M. Scanlon, and J. P. Raffensperger (2001), Modelling transport of dissolved silica in a forested headwater catchment: The effect of hydrological and chemical time scales on hysteresis in the concentration-discharge relationship, *Hydrol. Processes*, 15(10), 2029–2038, doi:10.1002/hyp.254.
- Jawitz, J. W., and J. Mitchell (2011), Temporal inequality in catchment discharge and solute export: Temporal inequality in catchment discharge and solute export, *Water Resour. Res.*, 47, W00J14, doi:10.1029/2010WR010197.
- Johnes, P. J. (2007), Uncertainties in annual riverine phosphorus load estimation: Impact of load estimation methodology, sampling frequency, baseflow index and catchment population density, *J. Hydrol.*, 332(1–2), 241–258, doi:10.1016/j.jhydrol.2006.07.006.
- Johnson, N. M., G. E. Likens, F. H. Bormann, D. W. Fisher, and R. S. Pierce (1969), A Working Model for the Variation in Stream Water Chemistry at the Hubbard Brook Experimental Forest, New Hampshire, *Water Resour. Res.*, 5(6), 1353–1363, doi:10.1029/WR005i006p01353.
- Jones, A. S., D. K. Stevens, J. S. Horsburgh, and N. O. Mesner (2011), Surrogate measures for providing high frequency estimates of total suspended solids and total phosphorus concentrations, *J. Am. Water Resour. Assoc.*, 47(2), 239–253, doi:10.1111/j.1752-1688.2010.00505.x.
- Kirchner, J. W. (2003), A double paradox in catchment hydrology and geochemistry, *Hydrol. Processes*, 17(4), 871–874, doi:10.1002/hyp.5108.
- Kolbe, T., J. Marçais, Z. Thomas, B. W. Abbott, J.-R. de Dreuzy, P. Rousseau-Gueutin, L. Aquilina, T. Labasque, and G. Pinay (2016), Coupling 3D groundwater modeling with CFC-based age dating to classify local groundwater circulation in an unconfined crystalline aquifer, *J. Hydrol.*, 543, 31–46, doi:10.1016/j.jhydrol.2016.05.020.
- Larouche, J. R., B. W. Abbott, W. B. Bowden, and J. B. Jones (2015), The role of watershed characteristics, permafrost thaw, and wildfire on dissolved organic carbon biodegradability and water chemistry in Arctic headwater streams, *Biogeosciences*, 12(14), 4221–4233, doi:10.5194/bg-12-4221-2015.
- Lehmann, A., and M. Rode (2001), Long-term behaviour and cross-correlation water quality analysis of the River Elbe, Germany, *Water Res.*, 35(9), 2153–60.
- Lischeid, G. (2008), Non-linear visualization and analysis of large water quality data sets: A model-free basis for efficient monitoring and risk assessment, *Stochastic Environ. Res. Risk Assess.*, 23(7), 977–990, doi:10.1007/s00477-008-0266-y.
- McDonald, R. I., K. F. Weber, J. Padowski, T. Boucher, and D. Shemie (2016), Estimating watershed degradation over the last century and its impact on water-treatment costs for the world's large cities, *Proc. Natl. Acad. Sci. U. S. A.*, 113(32), 9117–9122, doi:10.1073/pnas.1605354113.
- McDonnell, J. J., et al. (2007), Moving beyond heterogeneity and process complexity: A new vision for watershed hydrology, *Water Resour. Res.*, 43, W07301, doi:10.1029/2006WR005467.
- Mendoza-Lera, C., and T. Datry (2016), Relating hydraulic conductivity and hyporheic zone biogeochemical processing to conserve and restore river ecosystem services, *Sci. Total Environ.*, doi:10.1016/j.scitotenv.2016.11.166.
- Mendoza-Lera, C., and T. Datry (2017), Relating hydraulic conductivity and hyporheic zone biogeochemical processing to conserve and restore river ecosystem services, *Sci. Total Environ.*, 579, 1815–1821.
- Meybeck, M., and F. Moatar (2012), Daily variability of river concentrations and fluxes: Indicators based on the segmentation of the rating curve, *Hydrol. Processes*, 26, 1188–1207, doi:10.1002/hyp.8211.
- Meybeck, M., L. Laroche, H. H. Dürr, and J. P. M. Syvitski (2003), Global variability of daily total suspended solids and their fluxes in rivers, *Global Planet. Change*, 39(1–2), 65–93, doi:10.1016/S0921-8181(03)00018-3.
- Minaudo, C., M. Meybeck, F. Moatar, N. Gassama, and F. Curie (2015), Eutrophication mitigation in rivers: 30 years of trends in spatial and seasonal patterns of biogeochemistry of the Loire River (1980–2012), *Biogeosciences*, 12(8), 2549–2563, doi:10.5194/bg-12-2549-2015.
- Minaudo, C., F. Moatar, A. Coynel, H. Etcheber, N. Gassama, and F. Curie (2016), Using recent high-frequency surveys to reconstitute 35 years of organic carbon variations in a eutrophic lowland river, *Environ. Monit. Assess.*, 188(1), 41, doi:10.1007/s10661-015-5054-9.
- Moatar, F., and M. Meybeck (2005), Compared performances of different algorithms for estimating annual nutrient loads discharged by the eutrophic River Loire, *Hydrol. Processes*, 19(2), 429–444, doi:10.1002/hyp.5541.
- Moatar, F., F. Fessant, and A. Poirel (1999), pH modelling by neural networks. Application of control and validation data series in the Middle Loire river, *Ecol. Modell.*, 120(2–3), 141–156, doi:10.1016/S0304-3800(99)00098-8.
- Moatar, F., J. Miquel, and A. Poirel (2001), A quality-control method for physical and chemical monitoring data. Application to dissolved oxygen levels in the river Loire (France), *J. Hydrol.*, 252(1–4), 25–36, doi:10.1016/S0022-1694(01)00439-5.
- Moatar, F., M. Meybeck, S. Raymond, F. Birgand, and F. Curie (2013), River flux uncertainties predicted by hydrological variability and riverine material behaviour: Predicting uncertainties when monitoring riverine material fluxes, *Hydrol. Processes*, 27(25), 3535–3546, doi:10.1002/hyp.9464.
- Monteith, D. T., et al. (2007), Dissolved organic carbon trends resulting from changes in atmospheric deposition chemistry, *Nature*, 450(7169), 537–540, doi:10.1038/nature06316.
- Monteith, D. T., P. A. Henrys, C. D. Evans, I. Malcolm, E. M. Shilland, and M. G. Pereira (2015), Spatial controls on dissolved organic carbon in upland waters inferred from a simple statistical model, *Biogeochemistry*, 123(3), 363–377, doi:10.1007/s10533-015-0071-x.
- Mulholland, P. J., et al. (2008), Stream denitrification across biomes and its response to anthropogenic nitrate loading, *Nature*, 452(7184), 202–205, doi:10.1038/nature06686.
- Musolff, A., C. Schmidt, B. Selle, and J. H. Fleckenstein (2015), Catchment controls on solute export, *Adv. Water Resour.*, 86, 133–146, doi:10.1016/j.advwatres.2015.09.026.
- Nicolle, P., V. Andréassian, and E. Sauquet (2013), Blending neighbor-based and climate-based information to obtain robust low-flow estimates from short time series, *Water Resour. Res.*, 49, 8017–8025, doi:10.1002/2012WR012940.
- Oldham, C. E., D. E. Farrow, and S. Peiffer (2013), A generalized Damköhler number for classifying material processing in hydrological systems, *Hydrol. Earth Syst. Sci.*, 17(3), 1133–1148, doi:10.5194/hess-17-1133-2013.
- Onderka, M., S. Wrede, M. Rodný, L. Pfister, L. Hoffmann, and A. Krein (2012), Hydrogeologic and landscape controls of dissolved inorganic nitrogen (DIN) and dissolved silica (DSi) fluxes in heterogeneous catchments, *J. Hydrol.*, 450 451–, 36–47, doi:10.1016/j.jhydrol.2012.05.035.
- Peterson, B. J., et al. (2001), Control of nitrogen export from watersheds by headwater streams, *Science*, 292(5514), 86–90, doi:10.1126/science.1056874.
- Petrone, K. C., L. D. Hinzman, H. Shibata, J. B. Jones, and R. D. Boone (2007), The influence of fire and permafrost on sub-arctic stream chemistry during storms, *Hydrol. Processes*, 21(4), 423–434, doi:10.1002/hyp.6247.
- Pinay, G., S. Peiffer, J.-R. De Dreuzy, S. Krause, D. M. Hannah, J. H. Fleckenstein, M. Sebilo, K. Bishop, and L. Hubert-Moy (2015), Upscaling nitrogen removal capacity from local hotspots to low stream orders' drainage basins, *Ecosystems*, 18(6), 1101–1120, doi:10.1007/s10021-015-9878-5.
- Raymond, P. A., J. E. Saiers, and W. V. Sobczak (2016), Hydrological and biogeochemical controls on watershed dissolved organic matter transport: Pulse-shunt concept, *Ecology*, 97(1), 5–16, doi:10.1890/14-1684.1.
- Raymond, S., F. Moatar, M. Meybeck, and V. Bustillo (2013), Choosing methods for estimating dissolved and particulate riverine fluxes from monthly sampling, *Hydrol. Sci. J.*, 58(6), 1326–1339, doi:10.1080/02626667.2013.814915.

- Schwientek, M., K. Osenbrück, and M. Fleischer (2013), Investigating hydrological drivers of nitrate export dynamics in two agricultural catchments in Germany using high-frequency data series, *Environ. Earth Sci.*, *69*(2), 381–393, doi:10.1007/s12665-013-2322-2.
- Sebilo, M., B. Mayer, B. Nicolardot, G. Pinay, and A. Mariotti (2013), Long-term fate of nitrate fertilizer in agricultural soils, *Proc. Natl. Acad. Sci. U. S. A.*, *110*(45), 18,185–18,189, doi:10.1073/pnas.1305372110.
- Shanley, J. B., W. H. McDowell, and R. F. Stallard (2011), Long-term patterns and short-term dynamics of stream solutes and suspended sediment in a rapidly weathering tropical watershed, *Water Resour. Res.*, *47*, W07515, doi:10.1029/2010WR009788.
- Sloto, R. A., and M. Y. Crouse (1996), HYSEP: A computer program for streamflow hydrograph separation and analysis, *U.S. Geol. Surv. Water Resour. Invest. Rep.*
- Strohmeier, S., K.-H. Knorr, M. Reichert, S. Frei, J. H. Fleckenstein, S. Peiffer, and E. Matzner (2013), Concentrations and fluxes of dissolved organic carbon in runoff from a forested catchment: Insights from high frequency measurements, *Biogeosciences*, *10*(2), 905–916, doi:10.5194/bg-10-905-2013.
- Thomas, Z., B. W. Abbott, O. Troccaz, J. Baudry, and G. Pinay (2016), Proximate and ultimate controls on carbon and nutrient dynamics of small agricultural catchments, *Biogeosciences*, *13*(6), 1863–1875, doi:10.5194/bg-13-1863-2016.
- Thompson, S. E., N. B. Basu, J. Lascurain, A. Aubeneau, and P. S. C. Rao (2011), Relative dominance of hydrologic versus biogeochemical factors on solute export across impact gradients, *Water Resour. Res.*, *47*, W00J05, doi:10.1029/2010WR009605.
- Tian, S., M. A. Youssef, R. P. Richards, J. Liu, D. B. Baker, and Y. Liu (2016), Different seasonality of nitrate export from an agricultural watershed and an urbanized watershed in Midwestern USA, *J. Hydrol.*, *541*, 1375–1384, doi:10.1016/j.jhydrol.2016.08.042.
- Törnqvist, R., J. Jarsjö, J. Thorslund, P. S. C. Rao, N. B. Basu, and G. Destouni (2015), Mechanisms of basin-scale nitrogen load reductions under intensified irrigated agriculture, *Plos One*, *10*(3), e0120015, doi:10.1371/journal.pone.0120015.
- Torres, M. A., A. J. West, and K. E. Clark (2015), Geomorphic regime modulates hydrologic control of chemical weathering in the Andes–Amazon, *Geochim. Cosmochim. Acta*, *166*, 105–128, doi:10.1016/j.gca.2015.06.007.
- Vonk, J. E., S. E. Tank, P. J. Mann, R. G. M. Spencer, C. C. Treat, R. G. Striegl, B. W. Abbott, and K. P. Wickland (2015), Biodegradability of dissolved organic carbon in permafrost soils and aquatic systems: A meta-analysis, *Biogeosciences*, *12*(23), 6915–6930, doi:10.5194/bg-12-6915-2015.
- Walsh, R. P. D., and D. M. Lawler (1981), Rainfall seasonality: Description, spatial patterns and change through time, *Weather*, *36*(7), 201–208.
- Zarnetske, J. P., R. Haggerty, S. M. Wondzell, and M. A. Baker (2011), Labile dissolved organic carbon supply limits hyporheic denitrification, *J. Geophys. Res.*, *116*, G04036, doi:10.1029/2011JG001730.
- Zarnetske, J. P., R. Haggerty, S. M. Wondzell, V. A. Bokil, and R. González-Pinzón (2012), Coupled transport and reaction kinetics control the nitrate source-sink function of hyporheic zones, *Water Resour. Res.*, *48*, W11508, doi:10.1029/2012WR011894.
- Zhang, Q., C. J. Harman, and W. P. Ball (2016), An improved method for interpretation of riverine concentration-discharge relationships indicates long-term shifts in reservoir sediment trapping: An Improved Method on C-Q Relationships, *Geophys. Res. Lett.*, *43*, 10,215–10,224, doi:10.1002/2016GL069945.