



HAL
open science

Effects of prolonged exposure to CO₂ on behaviour, hormone secretion and respiratory muscles in young female rats

J.M. Martrette, Claire Egloff, Céline Clément, Kazutoyo Yasukawa, S.N. Thornton, Marie Trabalon

► **To cite this version:**

J.M. Martrette, Claire Egloff, Céline Clément, Kazutoyo Yasukawa, S.N. Thornton, et al.. Effects of prolonged exposure to CO₂ on behaviour, hormone secretion and respiratory muscles in young female rats. *Physiology & behavior*, 2017, 177, pp.257-262. 10.1016/j.physbeh.2017.05.007 . hal-01526325

HAL Id: hal-01526325

<https://univ-rennes.hal.science/hal-01526325v1>

Submitted on 7 Jul 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effects of prolonged exposure to CO₂ on behavior, hormone secretion and respiratory muscles in young female rats

J.M. Martrette^{a,e}, C. Egloff^{b,e,*}, C. Clément^a, K. Yasukawa^a, S.N. Thornton^c, M. Trabalon^d

^a EA 3450 DevAH, Université de Lorraine, 54505 Vandœuvre-lès-Nancy, France

^b UMR 7039 CNRS, Université de Lorraine, 54505 Vandœuvre-lès-Nancy, France

^c INSERM U_1116, Université de Lorraine, 54505 Vandœuvre-lès-Nancy, France

^d UMR 6552 CNRS, Université de Rennes 1, 35042 Rennes, France

^e Co-first author

* To whom correspondence should be addressed.

Running head: Effects of CO₂ on behaviour and muscles

All the authors declare no conflict of interest.

Dr Claire Egloff

Faculté d'Odontologie

Université de Lorraine

96, Avenue de Lattre de Tassigny

BP 50208, 54004 NANCY, France

Tel : (33)+383 68 29 51

E-mail: claire.juras@univ-lorraine.fr

ABSTRACT

Atmospheric CO₂ concentrations increased significantly over the last century and continuing increases are expected to have significant effects on current ecosystems. This study evaluated the behavioural and physiological (hormone status, muscle structure) effects of prolonged CO₂ exposure in young female Wistar rats exposed at 700 ppm of CO₂ during 6 h a day for 15 days.

Prolonged CO₂ exposure, though not continuous, produced significant disturbances in behaviour with an increase in drinking, grooming and resting, and a reduction in rearing, jumping-play and locomotor activity. Furthermore, CO₂ exposure was accompanied by increased plasma levels of corticosterone, suggesting that prolonged exposure to CO₂ was stressful. The muscular structure can also be modified also when respiratory working conditions change. The expression of myosin heavy chain was significantly affected in the diaphragm and oral respiratory muscles: *Masseter Superficialis* and *Anterior Digastric*. Modified behaviour and hormonal changes both appear to be at the origin of the observed muscular adaptation.

Keywords: behaviour, hormones, CO₂ exposure, rats, respiratory muscle structure, stress

Introduction

Atmospheric carbon dioxide (CO₂) concentrations began increasing with the advent of the industrial revolution and they continue to increase at a rate of approximately 4 parts per million by volume (ppm) annually (20). Over the last two decades, the average CO₂ concentration in the Earth's atmosphere showed a steady rise from 357 ppm in 1992 to 389 ppm in 2011 (53). In 2013, the daily average concentration of carbon dioxide (CO₂) in the atmosphere surpassed 400 ppm (12). Moreover, the mean ambient urban CO₂ concentration (23, 24) is around 400 ppm. At the current rate of increase, CO₂ levels are expected to increase to 750 ppm at the end of this century (20). The increase is expected to have significant effects on physiological changes in all living organisms. Numerous studies have been conducted to determine the effects of elevated CO₂ on plants and their associated communities (behavioural and/or physiological changes in herbivores), but few directly on mammals. On the other hand, work concerning the effects of CO₂ on mammals has used very high doses of CO₂ (20,000 to 60,000 ppm), *i.e.* levels not commonly found in the environment or in the future probable atmosphere. These high doses of CO₂ caused different modifications: hypercapnia, sedation, increased respiratory behaviour (22, 33, 41). Chronic exposure to CO₂ (50,000 ppm), at 6 time points in a 24-h period, increased plasma level of corticosterone in rats (2). These studies showed that an increased CO₂ level in the atmosphere was perceived as a stressful situation by the experimental animals.

However, in addition to CO₂ exposure, the individual's environment also seems to have an impact. Indeed, Clougherty et al. revealed a link between traffic-related air pollution and asthma solely among urban children exposed to familial violence,

suggesting that air pollution was a stressor amplified in the already stressed patient (10). Moreover, Bhat et al. showed that the shape of a rat's housing box could counteract plasma cortisol elevation and oxidative stress caused by chronic restraint. This later study showed that the effect of a stressing agent was fully dependent on the state of the individual and on how he interacted with his environment (4).

Indeed, stress was showed to induce changes of different physiological variables, such as plasma hormone levels and in particular glucocorticoid and thyroid hormones (18, 26, 27, 52). Furthermore, two studies (3, 25) showed that thyroid and glucocorticoid hormones were implicated in the modification of the myosin heavy chain (MHC) isoform expression. Other studies demonstrated that the administration of corticoids affected the diaphragm structure and function (11, 54, 58). This was accompanied by type IIB fibre atrophy in the diaphragm as well as in the peripheral skeletal muscles (43).

The administration of CO₂ by the re-breathing technique produced an increase in the breathing activity amplitude, and the respiratory frequency increased due to a decrease in both the inspiratory and expiratory durations (40). These modifications changed the working conditions of the respiratory muscles, implying a phenotypical adaptation of the muscles (44, 47, 50). These phenotypic modifications to the new environmental requirements generally involved muscle structure, in particular MHC isoform composition (6, 17, 21, 31).

Many studies on CO₂ were carried out in male animals while other results suggested that female rats were more reactive to stress than males (37, 56). Thus for our studies we chose to work on young female rats. Moreover, Cantillon and

Bradford reported that rat gender did not affect upper airway muscle contractile properties (9).

Although many physiological and behavioural effects were described using high doses of CO₂, little is known about the effects of relatively low doses, thus the aim of this present study was to investigate the behaviour, hormonal status (thyroid and glucocorticoid hormones) and respiratory muscle structure of rats chronically exposed to an eventual future condition of atmospheric CO₂ (700 ppm). A non-respiratory muscle related to locomotion, the soleus, was also analysed to determine any peripheral effects of elevated CO₂ exposure.

Materials and methods

Animals and breeding procedures

All experiments were carried out in accordance to the *Guide for the Care and Use of Laboratory Animals* published by the National Research Council (36), the recommendations edited by the European Union (Directive 2010/63/EU) (13), and those of the *Centre National de la Recherche Scientifique* (CNRS, France).

Twenty four female Wistar rats aged 6 weeks, weighing 152 ± 3 g, were purchased from a commercial breeder (IFFA-CREDO, France). On the arrival at the laboratory, the animals were housed in an air-conditioned room (temperature $30 \pm 1^\circ\text{C}$, relative humidity $50 \pm 2\%$) with the lights on from 8:00 pm to 8:00 am (inversion of the natural light cycle to adapt to the natural life conditions of rats which are nocturnal animals). The animals were housed individually in 60x38x20-cm Plexiglas boxes with a metal top and sawdust as bedding. Pellet food and tap water were continuously available. After 1 week of acclimatization, 12 females

were randomly assigned to the CO₂ exposure group, and the other 12 females to the control condition. The animals were kept all the time under filtered air except for the periods of CO₂ exposure in the experimental group. The experiments were performed at the same time of the day and under identical conditions, to insure that all groups were comparable.

Exposure apparatus and experimental procedure

The apparatus included a device for the production of CO₂, a control system for a continuous delivery of CO₂ (with concentration monitoring and recording), and four stainless-steel exposure chambers (100 x 180 x 150 cm) with a hatch glass in the front door. The chambers were located in an air-conditioned room and were equipped with racks which could hold up to 6 Plexiglas housing boxes (60 x 38 x 20 cm).

The control females were treated with charcoal-filtered (CF) air (453.2 ± 13.4 ppm CO₂). All experimental animals were exposed during 15 days to CF air plus 250 ppm CO₂ (685.9 ± 30.2 ppm). These chronic treatments were performed for 6 hours a day during the period of behavioural activity (12:00 AM to 06:00 PM).

Behavioural observations

The behaviour of the rats in their housing boxes was observed on the first day of exposure to CO₂ and after 15 days of exposure. The rats were exposed to CO₂ from 12:00 AM and 06:00 PM and all the observations were carried out during the period of stable gas levels i.e between 03:00 and 06:00 pm.

The behaviour was observed independently and by two experienced and blinded-to-condition observers. The data were recorded using a keyboard event recorder system feeding to a computer for analysis. Separate scores were obtained for each individual. Six types of behaviours displayed by each animal were recorded during a 5 min test with measures being taken at 1s intervals. The six behavioural scores were based on the ethological profile of rodents: inactivity (lying flat or standing still, with eyes open or closed and total absence of movements), jumping-play (jumping up vertically to a wall or to the metal top), rearing (standing on hind legs), grooming (wiping, licking, combing or scratching of any part of the body), drinking and eating.

Physiological analysis

After a 15-day exposure period, the animals were anaesthetized (sodium pentobarbital, i.p., 40mg/kg) and blood samples taken to analyse their physiological state. This was carried out in the neighbourhood around 09:00 am for all animals to minimise diurnal hormonal variations.

Immediately after the anaesthesia (30 s), intra-cardiac blood samplings (1 ml) were performed for corticosterone, thyroxine and triiodothyronine measurements (the duration of blood taking, starting from the departure of the animal from the home box, never exceeded 3 min).

The blood was collected into sterile heparinised syringes fitted with a 26-G needle. For the corticosterone assay, the plasma was immediately separated from the red blood cells by centrifugation at 4°C (15 min at 3,000 rpm) then the extracts were stored at -18°C until the time of the assay. The corticosterone concentration

was measured without an extraction procedure, using a commercially available EIA kit (Assay Designs Inc., USA). The concentration of corticosterone in the plasma samples was calculated from a standard curve and expressed in ng/ml. The intra- and inter-assay coefficients of variation were under 8.4 and 13.1 %, respectively.

Free thyroxine (FT4) and triiodothyronine (FT3) were assayed using commercial RIA kits according to the manufacturer's guidelines (Immunotech SA, France). The concentrations of FT4 and FT3 in the plasma samples were calculated from standard curves and expressed in pg/ml. The intra- and inter-assay coefficients of variation were respectively under 6.7 and 6.5 % for FT4 and under 6.4 and 5.5 % for FT3.

Muscle sampling and myosin extraction

Samples from one locomotor and four respiratory muscles were removed from the animals: the *Soleus* (S, plantarflexion of the foot); *Masseter Superficialis* (MS, propulsive and elevator mandibular muscle, i.e. mouth closing) and *Anterior Digastric* (AD, depressor mandibular muscle, i.e. mouth opening), antagonist muscles related to mouth movements and oral breathing (49, 57); *Levator Nasolabialis* (LN, active sniffing muscle) related to nasal breathing (48); and the Diaphragm (Dia). After dissection, the muscle samples were immediately frozen in liquid nitrogen for protein electrophoretic analyses. Myosin was crudely extracted in a high ionic strength buffer, as described by d'Albis et al. (1).

Electrophoretic analysis of MHC and quantification

The electrophoresis was performed according to the method of Talmadge and Roy (51). Mini-gels were used in the Bio-Rad Mini-protean II Dual Slab Cell. The electrophoresis was performed in a refrigerated room, at a temperature of 6°C for the whole run. To separate all the Myosin Heavy Chains, the duration of the run was of 28 h. The gels were stained with Coomassie blue R-250. The relative amounts of the different MHC were measured using an integration densitometer Bio-Rad GS-800 and analysed with the Quantity one 4.2.1 Program. Only bands representing more than 1% of total MHC were taken into account.

Statistical methods

All results were expressed as mean \pm SEM. The behavioural and physiological data were normally distributed; Student's *t*-test was used to establish the inter-group comparison. Differences were considered significant at $p < 0.05$.

Results

Behaviour

The analysis of the different behaviours is presented in Figs. 1-3. The results in Figure 1 run from the end of the first week of acclimatisation to the experimental setup. It is clear that there were no differences between the two experimental groups. Fig. 2 shows the effect on the behaviour after the first day of exposure to CO₂. During this first day of exposure, there was a significant decrease in rearing ($t = 15.96$, $p = 0.0003$), in association with a significant increase in inactivity ($t = 6.38$, $p = 0.01$), grooming ($t = 4.93$, $p = 0.03$) and drinking ($t = 4.07$, $p = 0.05$).

After 15 days of exposure (Fig. 3), these significant differences were accentuated: rearing ($t = 20.64$, $p < 0.0001$), inactivity ($t = 7.70$, $p = 0.005$), grooming ($t = 9.26$, $p = 0.004$), and drinking ($t = 6.83$, $p = 0.01$). Furthermore, other behaviours were observed to decrease after this long period of exposure: jumping-activity ($t = 5.45$, $p = 0.02$), and locomotor activity ($t = 4.06$, $p = 0.05$).

CO₂ exposure produced a significant increase in the daily amount of water consumed by each female after the first day: mean for the 14 days, 32 ± 1 ml/day vs 27 ± 1 ml/day for control females ($t = 2.85$, $p = 0.05$). The amount of food consumed every day by each animal was similar between the two groups of females: mean for the 15 days, 21 ± 4 g / day for control and 20 ± 2 g / day for ozone group ($t = 0.91$, $p = 0.35$).

Physiological analyses

The plasma levels of corticosterone were increased significantly by CO₂ exposure (Fig. 4): (240 ng/ml in CO₂ vs. 87 ng/ml in Controls, $t = 5.2$, $p = 0.01$).

FT3 and FT4 plasma levels were unchanged under CO₂ versus control group (respectively 3.19 pg/ml vs. 3.18 pg/ml, $t = 0.07$, $p = 0.78$; 16.92 pg/ml vs. 15.67 pg/ml, $t = 1.9$, $p = 0.19$, Fig. 4).

Muscle analyses

The body mass of the rats (185.8 ± 2 g CO₂ group, 194.5 ± 3.6 g control; $t = 4.4$, $p = 0.06$) as well as the relative mass of the five studied muscles was not significantly different between control and CO₂ groups (for the relative mass of the five muscles; $0.1 < t < 2.8$, $0.1 < p < 0.8$). The order of increasing electrophoretic

mobility of adult MHC isoforms was the following: fast 2A, fast 2X, fast 2B and slow 1 type (Fig. 5). Proportions of MHC isoforms were expressed as a relative percentage of the total amount of MHC present in the muscles studied (Table 1).

In the five muscles under control conditions, the 3 fast MHC isoforms could be observed, MHC 2A, 2X and 2B, except for LN in which the MHC 2A isoform was not detected and for S in which only MHC 2A was detected (Table 1). In relative expression, MHC 2B was the most abundant isoform in the AD and LN (44.7 %, 71.2 % respectively), whereas in the MS, MHC 2X was the predominant isoform (47.8 %). MHC 2A had the lowest expression level in S, AD and MS (15.8 %, 22.7 %, 23.9 % respectively). The slow type, MHC 1, was present only in S and Dia (84.2 %, 26.7 % respectively). The distribution between the different MHC isoforms was equivalent in Dia. In the soleus muscle, the slow isoform MHC 1 and only one fast isoform MHC 2A were detected. In relative expression, MHC 1 was the most abundant isoform (84.2 %).

During CO₂ exposure, the relative amounts of MHC protein isoforms were significantly affected in four of the five studied muscles. Indeed, the relative proportion of MHC 2B and MHC 1 significantly decreased to the detriment of MHC 2A in Dia under CO₂ exposure *versus* control. In addition, the relative proportion of MHC 2X increased in AD and decreased in MS. In the soleus muscle, the relative amounts of MHC protein isoforms were also significantly altered with an increase in the relative proportion of MHC 1 and a decrease of MHC 2A.

Discussion

These results specifically show that in sexually immature female rats, during prolonged CO₂ exposure (700 ppm, for 6 hours a day, during 15 days), there was an increase in drinking, grooming and inactivity behaviours, with a decrease in physical activity behaviours (jumping, rearing and locomotor activity).

Previous work concerning the effects of CO₂ on mammals generally used very high doses of CO₂ (20,000 to 60,000 ppm), ie levels not commonly found in the environment or in the future probable atmosphere. Wade et al. used experimental conditions close to the present study, in that they studied rat growth, body composition, and renal function during 30 days of increased ambient CO₂ (3,000, 7,000 and 20,000 ppm) (55).

The increase in drinking could be explained by a specific effect of this exposure to CO₂ which was not continuous. The regulation of breathing relies upon a chemical feedback concerning the levels of CO₂ and O₂. The feedback for CO₂ involves multiple sites including the carotid and receptors in the brain (15, 28). Small increases in CO₂ produce large increases in breathing. For instance, in awake humans at rest, a 1 mmHg increase in PaCO₂ leads to increased ventilation by ~20 - 30% (15). This could easily have increased the respiratory evaporative water loss (46) and thus produced dehydration which stimulated drinking. Wade et al. showed a reduced water turnover in male rats exposed to 3,000, 7,000 and 20,000 ppm (by increased urine output or by reduced water intake) during 30 days (55). They also showed that there were no negative water balances in spite of reduced water turnover.

The amount of food consumed every day by each animal was similar between the two groups of females. Wade et al. showed a slight decrease in food intake, yet the body weight, lean body mass and fat content remained relatively similar (55).

After a chronic exposure to CO₂ there was a significant decrease in rearing, in association with a significant increase in inactivity behaviour, grooming, jumping-activity, and locomotor activity. The observed decrease of rearing can be related to the procedure of behavioural observation in a familiar environment (the housing box) and the particular CO₂ exposure. The reduced physical performance during CO₂ exposure can be linked with another pollutant, O₃ (32), which is supposed to be part of the mechanisms preventing sensory irritation (29).

At CO₂ exposure levels of 10,000-40,000 ppm, the observed deleterious effects on the central nervous system are considered low to mild, and consist in headaches, a reduction of stereoacuity and a decrease in the ability to detect motion (19). A number of studies in humans suggested that CO₂ exposures in the range of 15,000-40,000 ppm did not impair neurobehavioral performances but led to clinical observations of poor attention, erratic behavior, confusion, motor skill impairment and moderate increases in anxiety, apathy, uncooperativeness, desire to leave, and sexual desire (35).

These behavioural changes suggest that chronic exposure to CO₂ could be a stressful situation. Furthermore, the CO₂ exposed group presented increased plasma levels of corticosterone which could be in relation with anxiety behaviour manifested by these females. Chronic exposition to CO₂ (50,000 ppm) increased the plasma level of corticosterone in rats as shown by Altholtz and colleagues (2). Wade et al. showed, in the male rat, an increase of plasma corticosterone only at

20,000 ppm and not at 3,000 and 7,000 ppm (55). To our knowledge, this is the first time that an increased plasma corticosterone level has been shown following chronic exposure to low levels of CO₂.

Chronic episodic hypercapnic hypoxia alters upper airway muscle structure and function in the rat. With respect to maximum velocity of shortening, the type 2B fibre is thought to have the highest velocity, followed by 2X > 2A > 1 (6). Our results have confirmed that the four muscles studied (Dia, LN, AD and MS), which predominantly expressed the type 2 MHC isoforms, were fast-twitch muscles. Among these fast muscles, the LN muscle, containing mainly type 2B (71%), was faster than the AD muscle, containing 45% type 2B, 33% type 2X and 23% type 2A, which in turn was faster than the MS muscle, containing mainly 2X type (48%). Lastly, Dia which contains about the same proportion of the four isoforms is the slowest muscle.

Redline and Strohl showed that the electromyographic activities of the upper airway and thoracic muscles increased at similar rates with an increase in hypercapnia (39). In the CO₂ exposition group, the modifications of MHC distribution in AD and MS could be explained by a modification of the electromyographic activity in these upper air way muscles. Geniohyoid type 1 fibers were reduced and type 2B fibers increased, whereas the sternohyoid muscle had an increase in type 1 and 2A fibers and a decrease in type 2B fibers (34). The primary explanation for these modifications is the working conditions of the muscles, whereas our experimental results bring other possible factors to light such as the hormonal factors or the intracellular pH. CO₂, by diffusing across the

lipid bilayer, can have an effect on intracellular pH. A low pH can alter neuronal function and synaptic transmission (5, 41).

Various hormones are known to cause changes in the expression of MHC isoforms. Thyroid hormones could be an important factor implicated in the muscle structure changes observed after a stressful situation (1, 16) as they are very important in the normal development of vertebrate skeletal muscles (16). All members of the myosin heavy chain multigene family respond to T3. However, the mode of response is not intrinsic to any myosin heavy chain gene, but is determined in a highly muscle type-specific manner (25, 30). Caiozzo et al. showed that combined effects of hyperthyroidism (T3) and a stressful procedure like hindlimb suspension (HS) rapidly converted the soleus into a fast muscle, producing large increases in the relative contents of the fast type 2X and 2B MHC isoforms which were primarily expressed in several populations of hybrid fibres (e.g., types 1/2A/2X, 1/2X/2B, 1/2A/2X/2B) (7). Finally, T3 + HS produced unique populations of hybrid fibers that did not adhere to the 1/2A/2X/2B sequential scheme of MHC plasticity. In a previous publication we showed that under chronic O₃ breathing, the serum level of FT3 was increased, which could have explained the appearance of hybrid fibres responsible for the modification of MHC profile in Dia, AD and MS (32). In contrast to ozone exposure, as shown here, chronic exposure to CO₂ did not modify serum levels of FT3. This could be explained by two mechanisms acting in opposite directions; the glucocorticoid release increased, while the uncontrollable stress inhibited the secretion of thyroid hormones (26, 38).

Chronic exposition to CO₂ increased the plasma level of corticosterone in rats. Falduto et al. showed that glucocorticoid treatment caused an increase in myosin heavy chain 2A expression (14). Glucocorticoids significantly decreased the relative expression of myosin heavy chain isoform MHC 2B in the Dia (38). The level of corticosterone could be one among other hormonal factors acting on muscle structure under CO₂.

In a previous study, Martrette et al. showed that a controllable stressful situation induced a marked increase in the relative expression of MHC 2B in MS and AD muscles, making the studied muscles faster in contraction (31). The authors concluded that in a controllable stress situation, there were probably several factors acting upon muscle MHC distribution. First, behavioural changes could be one of these factors (21). Thus, when the working conditions are changed, marked transitions in the myosin content occur in rat fast- and slow-twitch muscles (50). These modifications generally adapt the muscle to the new environmental requirements. Thus, after a few weeks of synergistic tenotomy, fast muscles become slow, fatigue resistant, and then more adapted to endurance. On the other hand, administration of CO₂ by the rebreathing technique produced an increase in the amplitude of breathing activity, and the respiratory frequency increased due to a decrease in inspiratory and expiratory duration (40). In a human study involving 21 subjects at 15,000 ppm of CO₂ chronic exposure, respiratory acclimatization to CO₂ involved a continuous increase in tidal volume while the respiratory rate declined shortly after an initial increase, associated with an increase in physiological and anatomical dead space (45). Gelhaye et al. showed that orofacial muscles present a profile in MHC adapted to the transition from nasal to oral

breathing, facilitating respiration (17). In the same way, under CO₂, the MHC 2B decreased in the Dia, this muscle adapts its fatigue resistance to the new respiratory frequency.

Conclusion

In conclusion, our work shows that female rats treated with chronic CO₂, an atmospheric pollutant, a situation comparable to the episodes of August 2003 and May 2015 (57) in western European countries, presented modifications of general behaviour, hormonal status (corticosterone), and MHC profile of **diaphragm and oral respiratory muscles**. Modifications of MHC expression could be induced by various factors, particularly behavioural and hormonal factors. These findings show that the control of atmospheric pollutant levels may be of great importance in the long term considering their incidence on organisms.

In future studies, it would be interesting to analyse the long-term effects of these modifications in adult rats and also other atmospheric pollutants alone and/or in combination, and their effects on studied biological parameters.

Acknowledgements

We wish to thank the Institut National de la Recherche Agronomique (INRA) - Université de Lorraine "Ecologie et Ecophysiologie Forestière" for the use of the CO₂ chambers.

References

1. **D'Albis A, Chanoine C, Janmot C, Mira JC, Couteaux R.** Muscle-specific response to thyroid hormone of myosin isoform transitions during rat postnatal development. *Eur J Biochem FEBS* 193: 155–161, 1990.
2. **Altholtz LY, Fowler KA, Badura LL, Kovacs MS.** Comparison of the stress response in rats to repeated isoflurane or CO₂:O₂ anesthesia used for restraint during serial blood collection via the jugular vein. *J Am Assoc Lab Anim Sci JAALAS* 45: 17–22, 2006.
3. **Baldwin KM.** Effects of altered loading states on muscle plasticity: what have we learned from rodents? *Med Sci Sports Exerc* 28: S101–106, 1996.
4. **Bhat MS, Rao G, Murthy KD, Bhat PG.** Housing in pyramid counteracts neuroendocrine and oxidative stress caused by chronic restraint in rats. *Evid-Based Complement Altern Med ECAM* 4: 35–42, 2007.
5. **Bierbower SM, Cooper RL.** The mechanistic action of carbon dioxide on a neural circuit and NMJ communication. *J Exp Zool Part Ecol Genet Physiol* 319: 340–54, 2013.
6. **Bottinelli R, Canepari M, Pellegrino MA, Reggiani C.** Force-velocity properties of human skeletal muscle fibres: myosin heavy chain isoform and temperature dependence. *J Physiol* 495 (Pt 2): 573–586, 1996.
7. **Bottinelli R, Schiaffino S, Reggiani C.** Force-velocity relations and myosin heavy chain isoform compositions of skinned fibres from rat skeletal muscle. *J Physiol* 437: 655–672, 1991.
8. **Caiozzo VJ, Baker MJ, Baldwin KM.** Novel transitions in MHC isoforms: separate and combined effects of thyroid hormone and mechanical unloading. *J Appl Physiol Bethesda Md* 1985 85: 2237–2248, 1998.
9. **Cantillon D, Bradford A.** Effect of gender on rat upper airway muscle contractile properties. *Respir Physiol* 113: 147–156, 1998.
10. **Clougherty JE, Levy JI, Kubzansky LD, Ryan PB, Suglia SF, Canner MJ, Wright RJ.** Synergistic effects of traffic-related air pollution and exposure to violence on urban asthma etiology. *Environ Health Perspect* 115: 1140–1146, 2007.
11. **Dekhuijzen PN, Gayan-Ramirez G, Bisschop A, de Bock V, Dom R, Bouillon R, Decramer M.** Rat diaphragm contractility and histopathology are affected differently by low dose treatment with methylprednisolone and deflazacort. *Eur Respir J* 8: 824–830, 1995.

12. **Dlugokencky EJ, Hall BD, Montzka SA, Dutton G, Muhle J, Elkins JW.** 2014: [Atmospheric composition: Long-lived greenhouse gases] Global Climate [in “State of the Climate in 2013”]. *Bull. Am. Meteorol. Soc. BAMS*.
13. **European Union.** Directive 2010/63/EU of the European Parliament and of the Council of 22 September 2010 on the protection of animals used for scientific purposes [Online]. <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2010:276:0033:0079:en:PDF> [29 Nov. 2015].
14. **Falduto MT, Czerwinski SM, Hickson RC.** Glucocorticoid-induced muscle atrophy prevention by exercise in fast-twitch fibers. *J Appl Physiol Bethesda Md* 1985 69: 1058–1062, 1990.
15. **Feldman JL, Mitchell GS, Nattie EE.** Breathing: rhythmicity, plasticity, chemosensitivity. *Annu Rev Neurosci* 26: 239–266, 2003.
16. **Gambke B, Lyons GE, Haselgrove J, Kelly AM, Rubinstein NA.** Thyroidal and neural control of myosin transitions during development of rat fast and slow muscles. *FEBS Lett* 156: 335–339, 1983.
17. **Gelhaye M, Martrette JM, Legrand-Frossi C, Trabalon M.** Myosin heavy chain expression and muscle adaptation to chronic oral breathing in rat. *Respir Physiol Neurobiol* 154: 443–452, 2006.
18. **Gómez F, Lahmame A, de Kloet ER, Armario A.** Hypothalamic-pituitary-adrenal response to chronic stress in five inbred rat strains: differential responses are mainly located at the adrenocortical level. *Neuroendocrinology* 63: 327–337, 1996.
19. **Guais A, Brand G, Jacquot L, Karrer M, Dukan S, Grévillet G, Molina TJ, Bonte J, Regnier M, Schwartz L.** Toxicity of carbon dioxide: a review. *Chem Res Toxicol* 24: 2061–2070, 2011.
20. **Hall MC, Stiling P, Moon DC, Drake BG, Hunter MD.** Effects of elevated CO₂ on foliar quality and herbivore damage in a scrub oak ecosystem. *J Chem Ecol* 31: 267–286, 2005.
21. **Hartmann N, Martrette JM, Westphal A, Divry M.** Effects of controllable stress on masticatory behaviour and muscle structure: partial protective effect of clomipramine. *Eur J Pharmacol* 366: 19–26, 1999.
22. **Hewett TA, Kovacs MS, Artwohl JE, Bennett BT.** A comparison of euthanasia methods in rats, using carbon dioxide in prefilled and fixed flow rate filled chambers. *Lab Anim Sci* 43: 579–582, 1993.
23. **Idso CD, Idso SB, Idso KE, Brooks T, Balling Jr RC.** Spatial and temporal characteristics of the urban CO₂ dome over Phoenix, Arizona. *Am. Meteorol. Soc.*: 46–48, 1998.

24. **Idso CD, Idso SB, Jr RCB.** The Urban CO₂ Dome of Phoenix, Arizona. *Phys Geogr* 19: 95–108, 1998.
25. **Izumo S, Nadal-Ginard B, Mahdavi V.** All members of the MHC multigene family respond to thyroid hormone in a highly tissue-specific manner. *Science* 231: 597–600, 1986.
26. **Joško J.** Liberation of thyrotropin, thyroxine and triiodothyronine in the controllable and uncontrollable stress and after administration of naloxone in rats. *J Physiol Pharmacol Off J Pol Physiol Soc* 47: 303–310, 1996.
27. **Kant GJ, Bauman RA, Anderson SM, Mougey EH.** Effects of controllable vs. uncontrollable chronic stress on stress-responsive plasma hormones. *Physiol Behav* 51: 1285–1288, 1992.
28. **Lahiri S, Forster RE.** CO₂/H(+) sensing: peripheral and central chemoreception. *Int J Biochem Cell Biol* 35: 1413–1435, 2003.
29. **Lee LY, Dumont C, Djokic TD, Menzel TE, Nadel JA.** Mechanism of rapid, shallow breathing after ozone exposure in conscious dogs. *J Appl Physiol* 46: 1108–1114, 1979.
30. **Li X, Larsson L.** Contractility and myosin isoform compositions of skeletal muscles and muscle cells from rats treated with thyroid hormone for 0, 4 and 8 weeks. *J Muscle Res Cell Motil* 18: 335–344, 1997.
31. **Martrette JM, Hartmann N, Westphal A, Divry M.** Effect of controllable stress on myosin heavy chain expression and muscle-specific protection by clomipramine. *J Muscle Res Cell Motil* 19: 803–810, 1998.
32. **Martrette JM, Thornton SN, Trabalon M.** Prolonged ozone exposure effects behaviour, hormones and respiratory muscles in young female rats. *Physiol Behav* 103: 302–307, 2011.
33. **Maskrey M, Nicol SC.** Responses of conscious rabbits to CO₂ at ambient temperatures of 5, 20, and 35 degrees C. *J Appl Physiol* 47: 522–526, 1979.
34. **McGuire M, MacDermott M, Bradford A.** The effects of chronic episodic hypercapnic hypoxia on rat upper airway muscle contractile properties and fiber-type distribution. *Chest* 122: 1400–1406, 2002.
35. **National Research Council (U.S.), Subcommittee on Emergency and Continuous Exposure Guidance Levels for Selected Submarine Contaminants, National Research Council (U.S.), Committee on Toxicology, National Research Council (U.S.), Board on Environmental Studies and Toxicology.** Emergency and continuous exposure guidance levels for selected submarine contaminants. Volume 1 [Online]. National Academies Press. <http://site.ebrary.com/id/10170925> [29 Nov. 2015].

36. **National Research Council (US) Committee for the Update of the Guide for the Care and Use of Laboratory Animals.** Guide for the Care and Use of Laboratory Animals [Online]. 8th ed. National Academies Press (US). <http://www.ncbi.nlm.nih.gov/books/NBK54050/> [29 Nov. 2015].
37. **Park MK, Hoang TA, Belluzzi JD, Leslie FM.** Gender specific effect of neonatal handling on stress reactivity of adolescent rats. *J Neuroendocrinol* 15: 289–295, 2003.
38. **Prezant DJ, Karwa ML, Richner B, Maggiore D, Gentry EI, Cahill J.** Gender-specific effects of dexamethasone treatment on rat diaphragm structure and function. *J Appl Physiol Bethesda Md* 1985 82: 125–133, 1997.
39. **Redline S, Strohl KP.** Influence of upper airway sensory receptors on respiratory muscle activation in humans. *J Appl Physiol Bethesda Md* 1985 63: 368–374, 1987.
40. **Rigatto H, Lee D, Davi M, Moore M, Rigatto E, Cates D.** Effect of increased arterial CO₂ on fetal breathing and behavior in sheep. *J Appl Physiol Bethesda Md* 1985 64: 982–987, 1988.
41. **Ruffin VA, Salameh AI, Boron WF, Parker MD.** Intracellular pH regulation by acid-base transporters in mammalian neurons. *Front Physiol* 5: 43, 2014.
42. **Saiki C, Mortola JP.** Effect of CO₂ on the metabolic and ventilatory responses to ambient temperature in conscious adult and newborn rats. *J Physiol* 491: 261–269, 1996.
43. **Sasson L, Tarasiuk A, Heimer D, Bark H.** Effect of dexamethasone on diaphragmatic and soleus muscle morphology and fatigability. *Respir Physiol* 85: 15–28, 1991.
44. **Sato K, Tsuji A, Usumoto Y, Kudo K, Yokoyama T, Ikeda N.** Expression of mRNA in the frontal cortex and hypothalamus in a rat model of acute carbon dioxide poisoning. *Leg. Med. Tokyo Jpn.* (July 26, 2015). doi: 10.1016/j.legalmed.2015.07.014.
45. **Schaefer KE, Hastings BJ, Carey CR, Nichols G.** Respiratory acclimatization to carbon dioxide. *J Appl Physiol* 18: 1071–1078, 1963.
46. **Schmidt-Nielsen K, Hainsworth FR, Murrish DE.** Counter-current heat exchange in the respiratory passages: effect on water and heat balance. *Respir Physiol* 9: 263–276, 1970.
47. **Sfondrini G, Reggiani C, Gandini P, Bovenzi R, Pellegrino MA.** Adaptations of masticatory muscles to a hyperpropulsive appliance in the rat. *Am J Orthod Dentofac Orthop* 110: 612–617, 1996.
48. **Soucek G, Breit S, König HE, Liebich HG.** [Functional significance of musculature of the external nose in swine (*Sus scrofa* f. *domestica*)]. *Anat Histol Embryol* 28: 307–314, 1999.

49. **Spyropoulos MN, Tsolakis AI, Alexandridis C, Katsavrias E, Dontas I.** Role of suprahyoid musculature on mandibular morphology and growth orientation in rats. *Am J Orthod Dentofac Orthop Off Publ Am Assoc Orthod Its Const Soc Am Board Orthod* 122: 392–400, 2002.
50. **Swynghedauw B.** Developmental and functional adaptation of contractile proteins in cardiac and skeletal muscles. *Physiol Rev* 66: 710–771, 1986.
51. **Talmadge RJ, Roy RR.** Electrophoretic separation of rat skeletal muscle myosin heavy-chain isoforms. *J Appl Physiol Bethesda Md* 1985 75: 2337–2340, 1993.
52. **Tamasy V, Meisami E, Du JZ, Timiras PS.** Exploratory behavior, learning ability, and thyroid hormonal responses to stress in female rats rehabilitating from postnatal hypothyroidism. *Dev Psychobiol* 19: 537–553, 1986.
53. **United Nations Environment Programme (U.N.E.P.).** Keeping Track of Our Changing Environment: From Rio to Rio+20 (1992-2012) [Online]. Division of Early Warning and Assessment (D.E.W.A.), United Nations Environment Programme (U.N.E.P.), Nairobi: [date unknown]. http://www.unep.org/geo/pdfs/keeping_track.pdf [29 Nov. 2015].
54. **Viires N, Pavlovic D, Pariente R, Aubier M.** Effects of steroids on diaphragmatic function in rats. *Am Rev Respir Dis* 142: 34–38, 1990.
55. **Wade CE, Wang TJ, Lang KC, Corbin BJ, Steele MK.** Rat growth, body composition, and renal function during 30 days increased ambient CO₂ exposure. *Aviat Space Environ Med* 71: 599–609, 2000.
56. **Weinstock M, Razin M, Schorer-Apelbaum D, Men D, McCarty R.** Gender differences in sympathoadrenal activity in rats at rest and in response to footshock stress. *Int J Dev Neurosci* 16: 289–295, 1998.
57. **Van Wessel T, Langenbach GEJ, van Ruijven LJ, Brugman P, van Eijden TMGJ.** Daily number and lengths of activity bursts in rabbit jaw muscles. *Eur J Neurosci* 21: 2209–2216, 2005.
58. **Wilcox PG, Hards JM, Bockhold K, Bressler B, Pardy RL.** Pathologic changes and contractile properties of the diaphragm in corticosteroid myopathy in hamsters: comparison to peripheral muscle. *Am J Respir Cell Mol Biol* 1: 191–199, 1989.
59. Monthly Global CO₂ Average Concentration Passes Symbolic Threshold | Media Centre [Online]. <http://www.wmo.int/media/content/monthly-global-co2-average-concentration-passes-symbolic-threshold> [14 Dec. 2015].

Legends

Table 1. Myosin heavy chain (MHC) distribution in respiratory muscles and in one locomotor muscle in control and CO₂ exposed rats. Values are percentages of total MHC \pm SEM (n=12 per group). Statistical summary: ‡ significantly different from control muscles, $p < 0.05$, ND: not detected, NS: not significant.

Figure 1. Behavioural parameters in control and pre-CO₂ exposed rats. Values are means of behavioural events during the week prior to CO₂ exposure (n=12 per group). NS: not significant.

Figure 2. Behavioural parameters in control and CO₂ exposed rats. Values are means of behavioural events during the first day of CO₂ exposure (n=12 per group). Statistical summary (t-student): * significantly different from control, $p < 0.05$, ** significantly different from control, $p < 0.01$, *** significantly different from control, $p < 0.001$. NS: not significant.

Figure 3. Behavioural parameters in control and CO₂ exposed rats. Values are means of behavioural events during the last day of the 15-day period of CO₂ exposure (n=12 per group). Statistical summary: * significantly different from control, $p < 0.05$, ** significantly different from control, $p < 0.01$, *** significantly different from control, $p < 0.001$. NS: not significant.

Figure 4. Plasma levels of corticosterone, free triiodothyronine (FT3) and free thyroxine (FT4) in control and CO₂ exposed rats. Values are mean \pm SEM (n=12 per group). Statistical summary: ‡ significantly different from control, $p < 0.05$.

Figure 5. Example of Myosin Heavy Chain 2A, 2X, 2B and 1 isoform migration in diaphragm muscle from control and CO₂ exposed rats.

Table 1. Myosin heavy chains (MHC) distribution in respiratory muscles and in one locomotor muscle in control and CO₂ exposed rats. Values are percentages of total MHC \pm SEM (n=12 per group). Statistical summary: ‡ significantly different from control muscles, p < 0.05, ND: not detected, NS: not significant.

Group	MHC type (%)			
	slow 1	2A	fast 2X	2B
<i>Diaphragm (Dia)</i>				
control	26.7 \pm 0.1	22.6 \pm 0.1	31.4 \pm 0.2	19.3 \pm 0.2
CO ₂	25.7 \pm 0.1 ‡	24.3 \pm 0.4 ‡	31.7 \pm 0.1 NS	18.3 \pm 0.2 ‡
<i>Masseter Superficialis (MS)</i>				
control	ND	23.9 \pm 0.5	47.8 \pm 0.2	28.3 \pm 0.5
CO ₂	ND	24.4 \pm 0.2 NS	46.7 \pm 0.1 ‡	28.9 \pm 0.3 NS
<i>Anterior Digastric (AD)</i>				
control	ND	22.7 \pm 0.5	32.6 \pm 0.2	44.7 \pm 0.4
CO ₂	ND	23.0 \pm 0.2 NS	33.3 \pm 0.1 ‡	43.7 \pm 0.1 NS
<i>Levator Nasolabialis (LN)</i>				
control	ND	ND	28.8 \pm 0.3	71.2 \pm 0.3
CO ₂	ND	ND	29.4 \pm 0.3 NS	70.6 \pm 0.3 NS
<i>Soleus (S)</i>				
control	84.2 \pm 0.3	15.8 \pm 0.3	ND	ND
CO ₂	87.6 \pm 0.2 ‡	12.4 \pm 0.2 ‡	ND	ND

Figure 1. Behavioural parameters in control and pre-CO₂ exposed rats. Values are means of behavioural events during the week prior to CO₂ exposure (n=12 per group). NS: not significant.




Figure 2. Behavioural parameters in control and CO₂ exposed rats. Values are means of behavioural events during the first day of CO₂ exposure (n=12 per group). Statistical summary (t-student): * significantly different from control, $p < 0.05$, ** significantly different from control, $p < 0.01$, *** significantly different from control, $p < 0.001$. NS: not significant.




Figure 3. Behavioural parameters in control and CO₂ exposed rats. Values are means of behavioural events during the last day of the 15-day period of CO₂ exposure (n=12 per group). Statistical summary: * significantly different from control, $p < 0.05$, ** significantly different from control, $p < 0.01$, *** significantly different from control, $p < 0.001$. NS: not significant.




Figure 4. Plasma levels of corticosterone, free triiodothyronine (FT3) and free thyroxine (FT4) in control and CO₂ exposed rats. Values are mean \pm SEM (n=12 per group). Statistical summary: ‡ significantly different from control, $p < 0.05$.




Figure 5. Example of Myosin Heavy Chains 2A, 2X, 2B and 1 isoform migration of diaphragm muscle from control and CO₂ exposed rats.

ACCEPTED MANUSCRIPT