



HAL
open science

Redox Control of Skeletal Muscle Regeneration

Emmeran Le Moal, Vincent Pialoux, Gaëtan Juban, Carole Groussard,
Hassane Zouhal, Bénédicte Chazaud, Rémi Mounier

► **To cite this version:**

Emmeran Le Moal, Vincent Pialoux, Gaëtan Juban, Carole Groussard, Hassane Zouhal, et al.. Redox Control of Skeletal Muscle Regeneration. *Antioxidants and Redox Signaling*, 2017, 27 (5), pp.276-310. 10.1089/ars.2016.6782 . hal-01585981

HAL Id: hal-01585981

<https://univ-rennes.hal.science/hal-01585981v1>

Submitted on 28 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comprehensive Invited Review

REDOX CONTROL OF SKELETAL MUSCLE REGENERATION

Emmeran LE MOAL^{1,2,\$}, Vincent PIALOUX^{3,4,\$}, Gaëtan JUBAN¹, Carole GROUSSARD², Hassane ZOUHAL², Bénédicte CHAZAUD^{1,*}, Rémi MOUNIER^{1,*}

1 Institut NeuroMyoGène, Université Claude Bernard Lyon 1, INSERM U1217, CNRS UMR 5310, 69100 Villeurbanne, France

2 Movement, Sport and health Sciences laboratory, M2S, EA1274, University of Rennes 2, 35170 Bruz, France

3 Laboratoire Interuniversitaire de Biologie de la Motricité, EA7424, Université Claude Bernard Lyon 1, Univ Lyon, 69100 Villeurbanne, France

4 Institut Universitaire de France, 75005 Paris, France

\$ equal contribution

* equal contribution

Running title: ROS and skeletal muscle regeneration

Correspondence to:

Rémi Mounier and Bénédicte Chazaud

Institut NeuroMyoGène

UMR CNRS 5310 – INSERM U1217 - Université Claude Bernard Lyon 1

Batiment Gregor Mendel, 16, rue Raphael Dubois, F-69622 Villeurbanne Cedex

Tel: (33) 4 72 43 10 51

remi.mounier@univ-lyon1.fr

benedicte.chazaud@inserm.fr

Word count : 12918

Reference numbers: 364

Number of greyscale illustrations: 13

Number of color illustrations: 2

Reviewing Editors: Charalambos Antoniades, Rafael Lambertucci, Tim Pearson, Vladimir B Ritov, Yuichiro Suzuki and Jose Vina

Abstract

Skeletal muscle shows high plasticity in response to external demand. Moreover, adult skeletal muscle is capable of complete regeneration after injury, due to the properties of Muscle Stem Cells (MuSCs), the satellite cells, which follow a tightly regulated myogenic program to generate both new myofibers and new MuSCs for further needs. Whereas reactive oxygen species (ROS) and reactive nitrogen species (RNS) have long been associated with skeletal muscle physiology, their implication in the cell and molecular processes at work during muscle regeneration is more recent. This review focuses on redox regulation during skeletal muscle regeneration. An overview of the basics of ROS/RNS and antioxidant chemistry and biology occurring in skeletal muscle is first provided. Then, the comprehensive knowledge on redox regulation of MuSCs and their surrounding cell partners (macrophages, endothelial cells) during skeletal muscle regeneration is presented in normal muscle and in specific physiological (exercise-induced muscle damage, aging) and pathological (muscular dystrophies) contexts. Recent advances in the comprehension of these processes has led to the development of therapeutic assays using antioxidant supplementation, which result in inconsistent efficiency, underlying the need for new tools aimed at precisely deciphering and targeting ROS networks. This review should provide an overall insight of the redox regulation of skeletal muscle regeneration while highlighting the limits of the use of non-specific antioxidants to improve muscle function.

1. Introduction

Skeletal muscle is a contractile tissue that accounts for 30-50% of body mass in humans (92). Besides its high plasticity and adaptation to various demands, adult skeletal muscle has the remarkable ability to regenerate *ad integrum* following injury. This regenerative property is mainly due to the Muscle Stem Cells (MuSCs), also called satellite cells, which are activated upon muscle injury and which follow a myogenic differentiation program to fuse and form new myofibers and restore the muscle tissue. Intrinsic and extrinsic cues drive MuSCs through these sequential steps of the regenerative process (15,63,290). Several cell types present in the injured muscle tissue play various roles in the regulation of muscle regeneration (359). Interestingly, functions of MuSCs and their neighboring cells are altered by oxidative stress modulation (44,255,348). For instance, macrophages are a major source of reactive oxygen species (ROS) and reactive nitrogen species (RNS) upon inflammation (16,38). Moreover, ROS and RNS have been recently proposed to be crucial actors of stem cell biology through the modulation of various cellular processes (34).

Concerning skeletal muscle biology, oxidative stress has been mainly investigated in skeletal muscle physiology, myofiber adaptation and contractile properties (22,156,206). It is likely that ROS/RNS-dependent modulation of the various cell types involved in skeletal muscle regeneration play important roles that may be altered in pathological contexts and may be the target of development of therapeutic strategies. The present review focuses on redox regulation of skeletal muscle regeneration. After a presentation of this complex biological process, basics of ROS/RNS and antioxidant chemistry and biology are described in the context of skeletal muscle. Then, the current knowledge related to redox regulation of skeletal

muscle regeneration is presented. Finally, specific alterations of redox regulation are described in various pathophysiological contexts, and therapeutic opportunities to improve skeletal muscle repair are discussed.

2. MuSCs and skeletal muscle regeneration

Skeletal muscle is a highly stable tissue that shows a high capacity to adapt to various physiological demands due to the properties of the myofiber, a multi-nucleated cell extending along the length of the muscle, which is capable of atrophy or hypertrophy. Adaptation is also possible because of the properties of different types of muscle fibers, adapted for different purposes, including slow-twitch type I and fast-twitch type II myofibers. In addition to its adaptive properties, skeletal muscle is capable of full functional recovery after an injury through the process of muscle regeneration, which goes beyond healing as the parenchyma recovers its functionality (**Figure 1**). Skeletal muscle regeneration relies on the properties of MuSCs, which are the adult muscle stem cells that normally reside in a quiescent state between the muscle fiber and its surrounding basal lamina (54). MuSCs are activated upon muscle injury and enter the cell cycle to proliferate. After an expansion phase, a large majority of MuSC progeny enters the myogenic differentiation program. The differentiated myogenic cells fuse to each other to form new myofibers and restore the tissue. A small subset of the expanding MuSCs does not commit into the terminal differentiation program but self-renews to replenish the pool of quiescent MuSCs for further needs. A general presentation of the cellular aspects of skeletal muscle regeneration is provided below since the reader can refer to excellent recent reviews on the topic (92,359). The molecular aspects of skeletal

muscle regeneration will be specifically addressed in the sections dedicated to the redox regulation of muscle regeneration (sections 7 and 8).

2.1. MuSCs and adult myogenesis

While the signaling pathways controlling the activation of MuSCs, *i.e.* their entry into the cell cycle, are still poorly understood and are the subject of intensive investigations, the gene expression program driving activated MuSCs along adult myogenesis is quite well known. MuSCs sequentially express a series of specific transcription factors from their exit from quiescence until their full differentiation and fusion (**Figure 2**) (92,359). At the quiescent state and during the expansion phase, they express Pax7, a transcription factor that is now considered as a canonical marker of MuSCs in various species. Once entered into the cell cycle, MuSCs sequentially express a series of specific muscle transcription factors, the Myogenic Regulatory Factors (MRFs). Structurally, MRFs bind DNA through their basic domain, while a helix–loop–helix motif allows their heterodimerization with E proteins, which mediate the recognition of genomic E-boxes, a motif present in the promoters of muscle-specific genes. While Myf5 is expressed by quiescent/early committed MuSCs, expanding MuSCs express MyoD, which is a strong determinant of their myogenic nature. After the expanding phase, MuSCs exit the cell cycle and downregulate the expression of Pax7 and MyoD while increasing that of Myogenin, which is considered a marker of irreversible terminal myogenic differentiation. Then, differentiated myocytes are capable of fusing together and with pre-existing fibers to reconstitute functional myofibers. At the time of fusion, differentiated MuSCs start to express the sarcomeric machinery of proteins required for contraction. Before exiting the cell cycle to differentiate, a subset of expanding MuSCs self-renews by

decreasing the expression of MyoD and increasing that of Pax7, therefore returning to a G0 quiescent state. The exact time (at the time of activation, during expansion or at the end of the expansion phase before differentiation) at which MuSC self-renew *in vivo* is still debated.

MuSCs are associated with all fiber types albeit with unequal distribution (264). Slow-twitch muscles express two- to three-fold higher MuSC density compared to fast-twitch muscles (284,296). These findings have been recently confirmed *in vivo* in mice using genetically labeled MuSCs (164). The phenotype of MuSCs from adult muscle fibers seems to be similar to their fiber of origin and to the fiber they tend to form in terms of myosin expression and contractile capacity (21,142). Aging is associated with skeletal muscle atrophy and a switch from fast-twitch to slow-twitch fiber type. Concomitantly, the number of MuSCs is reduced in some muscles (gastrocnemius, diaphragm, extraocular muscles) at 20 months from 16 to 25% in mice (164). So far, no study has reported an impact of the type of myofiber – or the corresponding MuSCs – on the efficiency of muscle regeneration.

2.2. MuSC cellular neighborhood

MuSCs are absolutely indispensable for adult skeletal muscle homeostasis, since the specific depletion of Pax7-expressing cells completely prevents post-injury regeneration (188,292). However, numerous studies have recently shown that cells present in the neighborhood of MuSCs are an important support of adult myogenesis (**Figures 3,4**). One can distinguish between resident and non-resident muscle cells.

Among the resident non-myogenic muscle cells that have been shown to participate in muscle regeneration are vascular cells and fibro-adipogenic progenitors (FAPs). Vascular cells comprise endothelial cells (ECs) that specifically interact with

expanding MuSCs in a reciprocal way to promote concomitant myogenesis and angiogenesis (63,177) (**Figure 4**). Vascular cells also include peri-ECs such as pericytes and smooth muscle cells that promote the return of MuSCs to quiescence (2,170). Another cell type that has been recently described in skeletal muscle is the fibro-adipogenic progenitor, which also participates in the regulation of MuSC expansion and differentiation (157,329).

Non-resident muscle cells invade the injured muscle in a tightly organized sequence, as in any other tissue repair process (**Figure 3**). Infiltrating leukocytes exert various actions on FAPs and other immune cells, while macrophages have been shown to exert specific and direct effects on MuSCs (55,56). Invading circulating monocytes become inflammatory macrophages that stimulate the expansion of MuSCs while inhibiting their differentiation. Later on, at the time of resolution of inflammation, macrophages skew their phenotype to an anti-inflammatory status. These restorative macrophages exert specific pro-myogenic effects on MuSCs by stimulating their differentiation (expression of Myogenin) and their fusion (**Figure 4**).

2.3. Time and space orchestration of muscle regeneration – link with muscular diseases

Recent studies have shown that asynchronous injuries (76) or desynchronized delivery of anti-inflammatory cues (257) trigger similar defects in skeletal muscle regeneration, resembling what is observed in some muscle diseases. It is therefore admitted that the MuSC niche - the quiescence niche, as well as the so-called transient niche for a proper implementation of the myogenic program - is tightly regulated in space and time. It is therefore likely that this tight regulation operates

through subtle changes in the cues MuSCs receive all along the regeneration process (56).

Inherited muscular dystrophies are a group of heterogeneous diseases characterized by permanent cycles of necrosis/regeneration that lead to the replacement of muscle fibers by connective and fat tissue, and eventually to the loss of muscle function (57,214). Most of these diseases are caused by a genetic alteration targeting proteins of the dystrophin-glycoprotein complex (DGC), which is located at the sarcolemma and links the myofiber cytoskeleton to the extracellular matrix (6,214) (**Figure 5**). It is believed that a weaker transmission force through the DGC triggers fragility of myofibers, which are therefore more susceptible to damage upon contraction/use (93). This primary event (mutation in a gene encoding for a DGC protein) leads to permanent regeneration processes throughout the tissue, triggering asynchronous delivery of regenerating cues to MuSCs and their close environment. This leads to failure in myogenesis and to secondary events including elevated oxidative stress, altered calcium homeostasis, chronic inflammation and endomysial (between myofibers) fibrosis which eventually replaces the atrophied myofibers (6) (**Figure 6**). Our knowledge of the cellular events in this context is still fragmented. It has been recently shown that in the Duchenne Muscular Dystrophy (DMD) mouse model mdx, FAP number is increased and macrophages exhibit altered inflammatory status while inhibition of the TGF β pathway leads to the apoptosis of FAPs, improving the muscle state (187). The molecular pathways that are altered in such an unbalanced tissue environment are only beginning to be defined for each cell type. For instance, NF- κ B signaling is deregulated in both myogenic cells and macrophages in DMD (3). Notch signaling, which has been shown to regulate MuSC fate, can be targeted to improve muscle homeostasis in this

disease (342). Given that the various cell types in the muscle no longer operate in a synchronized manner, one of the future challenges will be to identify the key pathways controlling the equilibrium of each cell type in degenerative myopathies.

3. Definition of free radicals, oxidative stress and redox signaling

ROS are a family of molecules that are continuously generated and consumed in all living organisms as a consequence of aerobic life (85). In 1985, Helmut Sies proposed oxidative stress as "a disturbance in the pro/antioxidant balance in favor of the former" (306). This definition has been widely associated with the view that ROS exert major biological effects as purely deleterious molecules causing damage to DNA, proteins and lipids, notably during aging (131), chronic diseases (335) or metabolic syndrome (151). During the early nineties, ROS-dependent signaling was discovered (review in (297)), shedding light on the beneficial role of ROS (316). Thus, oxidative stress has been redefined as "a disruption of redox signaling and control" (158). This notion implies that there is an optimal level of ROS/RNS to sustain both cellular homeostasis and adaptive responses, and that both too low and too high levels of ROS/RNS are detrimental to cell functions (108,235). This contemporary definition corresponds to a now well-accepted view that ROS production is crucial for cell functions (75,85,152,230,235). Recent developments in technological tools to detect and quantify ROS and ROS molecular targets, notably *in vivo*, have helped to better understand the role of ROS in numerous cellular processes (30,61,62,230). In the present review, the terms "ROS" and "RNS" include the oxygen and nitrogen radicals as well as some non-radical derivatives of O₂ and N, respectively (129) and are used when the specific entity is not described, as recently recommended by Forman and colleagues (104).

4. Major ROS and RNS in skeletal muscle

From its generation to the terminal product, ROS can undergo a wide range of transformations (**Figure 7**) that elicit specific effects on cell functions.

4.1. Major chemical entities

ROS and RNS can be classified depending on their chemical properties (reactivity, half-life) into two groups. Radical compounds (one electron oxidant) form the first group, whereas non-radical compounds (two electrons oxidant) form the second one (17). These chemical entities can also be classified based on the primary or secondary nature of the radicals.

4.1.1. Superoxide

The radical superoxide ($O_2^{\cdot-}$) is an anion produced by a monoelectronic reduction of an oxygen molecule ($O_2 + 1e^- \rightarrow O_2^{\cdot-}$). $O_2^{\cdot-}$ can diffuse within the cell but, despite a relatively longer half-life as compared with other radicals, $O_2^{\cdot-}$ is neither a strong oxidant, nor a powerful reductant. This is explained by a low standard reduction potential E_h of $O_2^{\cdot-}$ /hydrogen peroxide (H_2O_2) above 0.94V (E_h value *i.e.* measurement of the tendency of chemical species to acquire electrons and to be reduced; the more positive the potential, the higher the tendency to be reduced). By a dismutation reaction, $O_2^{\cdot-}$ can produce H_2O_2 ($O_2^{\cdot-} + O_2^{\cdot-} + (2H^+) \rightarrow 2 H_2O_2 + O_2$). In skeletal muscle as in other tissues, two distinct pools (inner and outer membrane) of mitochondrial $O_2^{\cdot-}$ have been observed (355).

4.1.2. Hydrogen peroxide

H_2O_2 is a two-electron oxidant, non-radical molecule generated from the dismutation of $O_2^{\cdot-}$. This molecule is stable and diffuses within and between cells and

therefore acts as a signaling molecule (338). However, it is a weak oxidizing and reducing agent, as evidenced by its inability to directly oxidize DNA and proteins as well as to induce antioxidant defenses in mammals (83). H_2O_2 is also described as a crucial player in tissue damage and repair (review in (337)). It acts as an alarmin to trigger inflammatory cell recruitment required for appropriate healing (238). However, H_2O_2 may also contribute to chronic tissue injury and subsequent inflammation by inducing irreversible oxidative damage (222). H_2O_2 is now recognized as a key molecule involved in the control of skeletal muscle function such as insulin control and glucose transport (135,338).

4.1.3. Hydroxyl radical

Hydroxyl radical (HO^\bullet) is a powerful oxidant (E_n value $O_2^{\bullet-}/H_2O_2 = 2.34V$) that is generated by reaction of metal ions with H_2O_2 . The high reactivity related to its short half-life (10^{-9} sec) restricts its damaging effects to a local area close to the generation site (129). Additionally, HO^\bullet can be formed from H_2O_2 in the presence of transition metals (iron) through the reactions of Fenton and of Haber-Weiss ($O_2^{\bullet-} + H_2O_2 (+Fe_3/Fe_2) \rightarrow HO^\bullet + O_2 + ^\bullet OH$) to induce oxidative damage (144,165,352).

4.1.4. Nitric oxide

Nitric oxide (NO^\bullet) is a free radical that easily crosses membranes and diffuses between and within cells. NO^\bullet is synthesized from L-arginine by NO synthases (NOS), which are neuronal (nNOS), endothelial (eNOS) and inducible (iNOS) NOS. It acts as a signaling molecule and exerts its effects *via* interactions with heme or non-heme iron and copper or by S-nitrosylation of cysteine redox centers to modulate protein functions (137). Moreover, NO^\bullet reacts with $O_2^{\bullet-}$ to form peroxynitrite (128). nNOS and eNOS are the main NOS found in skeletal muscle (97,311). NO^\bullet is a key

molecule in many physiological functions in skeletal muscle, including muscle regeneration (101,311).

4.1.5. Peroxynitrite

The reaction of $O_2^{\cdot-}$ with NO^{\cdot} forms peroxynitrite, which is a strong oxidant capable of crossing cell membranes. Cell exposure to peroxynitrite produces many effects such as inhibition or depletion of antioxidant enzymes, modification of receptor functions, lipid peroxidation and DNA injury. Recently, peroxynitrite has been shown to induce the activation of mTOR signaling *via* a calcium-dependent signaling leading to skeletal muscle hypertrophy (147).

4.2. Major sources of ROS

In 1954, Commoner and colleagues identified free radicals in skeletal muscle (70). Since then, several sources of ROS production have been identified in various subcellular locations (148) (**Figure 8**).

4.2.1. Mitochondria

Mitochondria are one of the most important sources of ROS in muscle. At least ten sites capable of producing considerable amounts of ROS have been described (123). Early studies suggested that 2-5% of the total oxygen consumed by mitochondria undergo a one-electron reduction and generate $O_2^{\cdot-}$ (39). However, depending on different mitochondrial states of respiration, the percentage of oxygen that forms superoxide ranges from 0.1% in state 3 to 2% in state 4 (266,305). Other studies have shown that both complexes I and III of the electron transport chain are particularly involved in ROS production through a leakage of electrons (20,229). This was demonstrated by the use of specific inhibitors of complex I and III-derived ROS such as rotenone and antimycin, respectively (364).

4.2.2. NADPH oxidases

The NADPH oxidases (NOXs) transfer electrons across biological membranes and form $O_2^{\cdot-}$. NOX family includes 7 NOX isoforms (NOX1, NOX2/gp91^{phox}, NOX3, NOX4, NOX5, DUOX1, DUOX2), 2 organizer subunits (p47^{phox}, NOXO1), 2 activator subunits (p67^{phox}, NOXO1) and 2 DUOX-specific maturation factors (DUOXA1 and DUOXA2). NOX enzymes are primarily known to produce $O_2^{\cdot-}$ in both phagocytic and non-phagocytic cells (26). A small GTPase named Rac1 has been described as an important activator of NOX1 and 2 in various cell types including skeletal muscle (317) and also leads to superoxide production in non-phagocytic cells (140). In skeletal muscle cells, NOX isoforms 2 and 4 are found at the plasma membrane, transverse tubules and sarcoplasmic reticulum where they modulate calcium release (261). They are described as the major source of ROS during skeletal muscle contraction (217). In this context, pharmacological inhibition of NOX-induced ROS production has been widely used, although these inhibitors are mainly unspecific and not isoform selective (review in (8,65)).

4.2.3. Xanthine oxidase

Xanthine oxidoreductase (XOR) is an intracellular enzyme involved in purine catabolism, which includes 2 isoenzymes: xanthine dehydrogenase (XDH) and xanthine oxidase (XO). XOR catalyzes the reduction of hypoxanthine and xanthine in uric acid. The oxidase form is used as an electron acceptor and both hypoxanthine and xanthine are reduced into uric acid and $O_2^{\cdot-}$ (210). The role of XO as a source of ROS generation has been demonstrated using allopurinol, a specific inhibitor. In both animals and humans, allopurinol prevents oxidation of glutathione (GSH) and lipid peroxidation (343). Interestingly, XO is a major source of ROS in skeletal muscle atrophy following unloading and in sarcopenia (82).

4.2.4. Other sources

The endoplasmic reticulum is another organelle involved in ROS production (50,199). In addition, various enzymatic systems such as phospholipase A2 induce ROS production by stimulating NOX. Studies reported both calcium-dependent and calcium-independent forms of phospholipase A2 as key actors in ROS generation in skeletal muscle (124,236).

5. Major antioxidants in skeletal muscle

Enzymatic and non-enzymatic antioxidant systems exist to detoxify excess ROS, prevent and repair oxidative damage, and maintain redox homeostasis in the cell (127). Halliwell and Gutteridge have defined an antioxidant as "any substance that delays, prevents or removes oxidative damage to a target molecule" (129). This definition suggests that this evolutionary conserved system exerts its antioxidant functions in 3 distinct ways: by converting ROS into less deleterious molecules, by reducing pro-oxidant molecules such as metallic ions, and by scavenging ROS. Thus, the complex antioxidant network works synergistically to protect cells from redox disturbances and oxidative damage. Interestingly, there is a growing body of evidence indicating that antioxidants are not only ROS-scavenger molecules, but also act as signaling molecules that are required both for the maintenance of cell homeostasis and for adaptive cell responses (277). Human Atlas Proteome (78,330) and the Redox Atlas of the Mouse (120) provide a global analysis of tissue-specific proteome. Based on these large-scale projects and the available literature, we present hereafter the skeletal muscle tissue-specific antioxidant system.

5.1. Endogenous

Among endogenous antioxidants, GSH, superoxide dismutases (SODs), catalase (CAT) and members of the thioredoxin (Trxs) family contribute to cell protection and regulation of redox signaling.

5.1.1. Superoxide dismutases

SODs are enzymes that catalyze the dismutation of $O_2^{\cdot-}$ into H_2O_2 . SODs are divided into 3 isoenzymes (SOD1, SOD2, SOD3), which all require redox active transition metals in the active site (261). SOD1 (CuZnSOD) is a dimer of 32.5 kDa associated with one copper ion and one zinc ion and is mainly localized in cytosol and the mitochondrial intermembrane space (**Figure 8**). SOD2 (MnSOD) is a tetramer of 24.7 kDa, which is associated with one manganese ion. SOD2 is mainly located in mitochondria. SOD3 is an extracellular tetradimer of 30 kDa, which uses copper-zinc as a cofactor. Deletion of SOD1 in skeletal muscle induces a phenotype of accelerated age-related loss of skeletal muscle mass and function (291).

5.1.2. Catalase

CAT is a homotetramer of 240 kDa composed of 4 subunits, which requires iron as a cofactor. CAT is mainly localized in peroxisomes (**Figure 8**), which were originally named to highlight their content in enzymes either producing (oxidases) or breaking down (catalase) H_2O_2 (24,336). Therefore, it was originally thought that the primary function of these organelles was to control H_2O_2 metabolism (review in (194)). However, peroxisomes also have an established role in lipid metabolism, as well as emerging functions in cellular signaling relevant to metabolism (194). The function of CAT is to catalyze the dismutation of H_2O_2 , which leads to the formation of one molecule of H_2O and of O_2 . It has been shown that a decline of CAT protection, associated with elevated H_2O_2 levels, is observed at the onset of sarcopenia, which is the loss of muscle mass observed during aging (315).

5.1.3. Glutathione

GSH is the most abundant low molecular weight component in the cell. It functions as a buffer of the cellular redox status and as an antioxidant in skeletal muscle (126). This tripeptide synthesized in liver exerts its antioxidant functions through several features: as a substrate for glutathione peroxidases (Gpxs) and as a reducing agent for vitamin C and E radicals. GSH is predominantly found in the cytosol but is also distributed in various organelles such as the nucleus, mitochondria and endoplasmic reticulum (107). GSH is involved in NO[•] metabolism since GSH and peroxynitrite lead to the formation of S-nitroglutathione (GSNO) that is able to generate NO[•] (129). In skeletal muscle fibers, GSH is present at millimolar concentrations. Higher levels of GSH are found in type I muscle fibers, which have an oxidative/aerobic metabolism, as compared with type II fibers, which rely mainly on a glycolytic/anaerobic metabolism (185).

5.1.4. Family of thioredoxins

The thioredoxin family of proteins encompasses Trxs, Gpxs, glutaredoxins (Grx) and peroxiredoxins (Prx). These proteins share common structural sequences and are characterized by their active site motifs, containing either one or two cysteinyl residues. They are essential for the reduction of protein disulfides, protein de/glutathionylation and de/trans/nitrosylation (130,242). The Trx system includes 2 oxidoreductases Trx1 and Trx2 and a thioredoxin reductase (TrxR). Trx1 and TrxR1 isoforms are found in the cytosol, Trx1 is translocated into the nucleus to alter the redox state of transcription factors, while Trx2 and TrxR2 isoforms are located in the mitochondria (283) (**Figure 8**). Trxs serve as donors for Prxs to reduce peroxides. In the Trx system, electrons are transferred during a multiple step process from NADPH to Trxs to ultimately reduce disulfides and generate antioxidants (208). Trx1 and

related redox proteins are constitutively secreted by skeletal muscle cells and have been suggested to belong to myokines (203). Moreover, a decreased expression of Trx1 is observed during muscle atrophy that may be responsible for the enhanced oxidative damage during the late phase of disuse muscle atrophy (207). Grxs also modulate the thiol redox state of several proteins but no study has reported a role in skeletal muscle (130). Gpxs catalyze the reduction of H₂O₂ and organic hydroperoxides to water and alcohol, respectively (42,136). In skeletal muscle, Gpx activity is strongly decreased in cancer cachexia (314) and Gpxs have been involved in the regulation of MuSC function (see section 7). Prxs play multiple roles as peroxidases, sensors and regulators of peroxides. Interestingly, a role for Prx in H₂O₂-mediated cellular signaling has been described (354). The expression of Prx3 is decreased during muscle atrophy (207).

5.2. Exogenous

Exogenous molecules with antioxidant properties are generally classified as vitamins, trace elements or phytochemicals. Vitamin E is a term encompassing 4 tocopherols ($\alpha,\beta,\delta,\gamma$) and 4 tocotrienols ($\alpha,\beta,\delta,\gamma$) molecules. The α -tocopherol is the most abundant form in nature. Vitamin E, considered a master antioxidant, has the ability to prevent lipid peroxidation and other radical-driven events. It is mainly located in the plasma membrane (**Figure 8**) and acts as a chain-breaking antioxidant that prevents the propagation of free radical-reactions (43). Vitamin E reacts with oxidants to form tocopheryl radicals that are further reduced by vitamin C into non-radical α -tocopherols. This process requires GSH that gives an electron, leading to the formation of oxidized GSH. Vitamin E is required for skeletal muscle plasma membrane repair (141,176). However, the underlying cellular and molecular

mechanisms of this phenomenon remain to be determined (166). Ascorbate is the most common form of vitamin C found in the organism. This water-soluble molecule, localized in the extracellular compartment has special functions in redox reactions, as a powerful antioxidant and enzyme cofactor, notably of hydroxylases (204,325). Vitamin C has the ability to protect against lipid peroxidation by acting as a ROS scavenger and by one-electron reduction of lipid hydroperoxyl radicals *via* the vitamin E redox cycle (129). The dehydroascorbic acid (oxidized form of vitamin C) thereby formed can be reduced by TrxR and GSH (189), underlying the synergy within the antioxidant network. Finally, trace elements such as selenium, copper, manganese and zinc play important roles to support antioxidant functions and redox exchanges of many enzymes. Generally, these molecules play a role of substrate or cofactor and their deficiency leads to oxidative stress (116,246). They are likely important in skeletal muscle function, as exemplified by the reduced expression of selenoprotein N (*sepn1*) that triggers a congenital muscle disease (14), highlighting the role of trace elements in skeletal muscle homeostasis.

6. General mechanisms of redox signaling

In skeletal muscle, ROS/RNS have been identified as playing key roles in numerous physiological processes including glucose uptake (163), excitation/contraction coupling (261), mitochondrial biogenesis (145), training adaptations (122,281) and cell death (295). This underlines the need to understand how ROS/RNS act as determinant regulators of biological effectors involved in the control of homeostasis (90,302). It is suggested that oxidation and reduction of the thiol side chain of cysteine amino acids are the major mechanisms by which oxidants integrate signal transduction pathways (85).

6.1. Thiol oxidation as a major mechanism of oxidation-related post-translational modifications

Oxidants mainly exert their effects by targeted modifications of thiol residue (SH) of cysteine amino acids resulting in reversible or irreversible oxidations that depend on the state of oxidation (**Figure 9**). It results in the formation of sulfenic acid (SOH) that can be further oxidized in sulfinic (SO₂H) and in sulfonic (SO₃H) acids. Additional post-translational modifications involving NO[•] or GSH are also observed and identified as S-nitrosylation and S-glutathionylation, respectively (17,35,85). Such chemical reactions are under the control of pKa, pH and local protein environment (127).

6.2. From signal transduction to post-translational modifications

The role of ROS in signaling has emerged with the observation that H₂O₂ production was necessary for normal signaling of growth factors, suggesting that ROS/RNS may act as secondary messengers (316). More recently, Jones highlighted the role of "radical-free" biology of oxidative stress, indicating that the free radical chain reactions are almost completely prevented in biological systems due to the very high concentrations of antioxidant proteins (159). He reported that 10% of unbalanced pro- and antioxidant reactions involve free radical intermediates and that less than 0.1% induce macromolecular damage. Hence, the non-radical H₂O₂ appears to be an ideal candidate for redox-dependent signal transduction (*e.g.* large diffusion, weak oxidant, reversible oxidation inductor) (105,152). Several mechanisms are described for H₂O₂-dependent signal transduction. One proposes H₂O₂-dependent signaling *via* a relay system involving a redox sensor (Prx2), which

oxidizes the target and forms a transient disulfide bond (310). Another mechanism is the floodgate model (354). It is based on the local inactivation of antioxidant defenses, such as Prx1, that allows low H₂O₂ levels to mediate signaling in a specific cell compartment. Although less well-described, two other mechanisms have been proposed. The first includes an intermediate step involving the oxidation of Trx or GSH to the redox relay mechanism, and the second one likely results from the dissociation and subsequent activation of a target protein following H₂O₂ oxidation of a scavenger enzyme (272). In addition, NO[•] and its derivatives (S-nitrosothiols (SNO) and GSNO) are involved in signal transduction (103,152,276,338). These molecules induce modifications of protein structure, thus affecting protein functions as well as the signaling pathways in which these proteins are involved.

Technological advances such as proteomic and redox proteomics (60) have recently revealed that 500 proteins contain reactive and potentially modulatory cysteine residues (162,345). Among the targeted proteins detected in the "redoxome", are transcription factors (NFκB, FOXO), kinases (AMP Kinase, mTOR), protein tyrosine phosphatases, proteases (caspases, metalloproteases), and chaperones proteins. This illustrates the broad impact that oxidation of conserved cysteine residues may have in various physiological processes from signal transduction to gene expression (review in (152)).

In response to oxidative stress, cells modulate their gene expression (7). Several transcription factors have been described as redox-sensitive and are involved in various cellular signaling pathways with distinct outcomes. Nuclear factor kappa B (NF-κB) is the first eukaryotic transcription factor which has been shown to respond directly to oxidative stress in the L6 skeletal muscle myoblasts upon exposure to the non-radical ROS hydrogen peroxide and for which activation was controlled by the

cell glutathione disulfide/glutathione (GSSG/GSH) ratio (301). Additionally, NF- κ B is involved in the regulation of a wide array of cellular processes including immune response, cellular survival, proliferation, and differentiation (review in (18)). Interestingly, NF- κ B activity is impaired as a consequence of S-glutathionylation of IKK- β cysteine 179 and is associated with a decrease of pro-inflammatory signals (275). Redox factor-1 (Ref-1) contains in its N-terminal region a redox regulatory domain, which is characterized by 2 cysteines that are involved in the redox-dependent modification (reduction) of NF- κ B and of other transcription factors such as activator protein-1 (AP-1) (Fos and Jun), p53, and hypoxia-inducible factor (HIF)-1 α . The oxidized form of AP-1 exhibits poor DNA-binding activity, due to the cysteine residues within the DNA-binding domain of both Fos and Jun. Inversely, AP-1 is mainly activated by antioxidants (192,300). Of interest, these redox-sensitive transcription factors NF- κ B and AP1 trigger the synthesis of cytoprotective regulatory proteins such as SODs, catalase and heat shock proteins after the delivery of ROS generated by the skeletal muscle itself upon contractile activity (149).

The transcription factor NF-E2 p45-related factor 2 (Nrf2) is another well-known redox-sensitive master regulator that controls the expression of networks of genes encoding proteins with diverse cytoprotective activities. Nrf2 itself is primarily controlled at the level of protein stability. Under basal conditions, Nrf2 is a short-lived protein that is subjected to continuous ubiquitination and proteasomal degradation. The most common ubiquitin ligases that contribute to the degradation of Nrf2 are Kelch-like ECH-associated protein 1 (Keap1) and a substrate adaptor protein for Cullin 3 (Cul3)/Rbx1 ubiquitin ligase. Keap1 is considered to be a sensor for a wide array of small-molecule activators of Nrf2 including ROS. These latter block the cycle of Keap1-mediated degradation of Nrf2 by modifying specific cysteine residues within

Keap1 (87) or by directly disrupting the Keap1:Nrf2 interaction. As a consequence, Nrf2 is not degraded, and the transcription factor accumulates and translocates to the nucleus, where it forms a heterodimer with a small Maf protein and binds to antioxidant-response elements in the upstream regulatory regions of its target genes, and initiates transcription (133).

In skeletal muscle, which is an active metabolic tissue, insulin signal transduction is crucial. Insulin signaling has been shown to be altered by ROS (such as 4-hydroxy-2-nonenal and peroxynitrate) (13) and to be regulated by the selenoproteins Gpx1 and Sepp1 (through the inhibition of energy metabolic mediators AMPK and Akt) (312).

7. Redox regulation of MuSCs during skeletal muscle regeneration

During skeletal muscle regeneration, MuSCs undergo a series of processes called adult myogenesis, which includes activation and entry in the cycle, proliferation, exit from the cell cycle and commitment into terminal differentiation, followed by fusion into multinucleated cells (myotubes *in vitro* and myofibers *in vivo*) (**Figure 2**). Various aspects of adult myogenesis have been shown to be, or are likely redox-regulated (264,361) (**Figure 10**).

7.1. Viability - apoptosis

ROS and redox reactions have been described to regulate cell survival through the modulation of transcription factors (NF- κ B, AP-1, Nrf2...) and pro-apoptotic proteins (caspases, Bcl-2) (247,270). The role of oxidative stress in cell death pathways is supported by the fact that antioxidants can protect cells against apoptosis (247). In skeletal muscle, long-term exercise has been shown to reduce

ROS generation, to improve antioxidant status and to decrease oxidative stress that should serve to attenuate any ROS-induced activation of apoptosis (265). Elevated apoptotic signaling has been reported in skeletal muscle during aging, stress-induced states and diseases; a phenomenon that plays a role in muscle dysfunction, degradation, and atrophy (265). During fetal myogenesis, analysis of double conditional Pitx2/3 mouse mutants in primary muscle cell cultures reveals excessive upregulation of ROS levels leading to DNA damage and apoptosis of differentiating cells (175). In adult male rats, GSH depletion in skeletal muscle increases caspase-independent signaling and increases mitochondrial-associated apoptotic events to subsequent cell death stimuli (77). Whereas apoptosis in skeletal muscle is low at rest, skeletal muscle injury causes apoptosis and necrosis (307) and is associated with ROS production (341). Therefore, protection of MuSCs from cytotoxic damage is crucial to ensure efficient regeneration. MuSC antioxidant levels have been described as a major determinant of MuSC cell therapy efficiency (91). Several antioxidant pathways have been involved in oxidative stress detoxification of MuSCs. Retinoic acid-Gpx3 pathway (96) and aldehyde dehydrogenase activity (153) improve MuSC survival. Recently, cytoglobin, a heme containing protein has been involved in the resistance to oxidative stress-induced apoptosis that results in incomplete skeletal muscle regeneration (307). Apoptosis being a highly conserved process playing a critical role in tissue homeostasis (**Figure 10**), its putative role in skeletal muscle regeneration remains to be elucidated.

7.2. Quiescence

At rest, MuSCs are maintained in quiescence (G_0 phase) (58). Transcriptomic analysis revealed an antioxidant molecular signature (250) involving the up-

regulation of Gpx3 and TrxR1, suggesting a protective effect of quiescent cells from oxidative insults. During muscle regeneration, while most of MuSCs commit to a myogenic fate and differentiate, a small population self-renews to maintain a stem cell pool for further needs (**Figure 2, 10**). NO[•] was shown to increase MuSC self-renewal in a cGMP-independent pathway. This requires enhanced expression of Vangl2, a member of the Wnt planar cell polarity pathway that controls the homeostatic number of MuSCs (179).

7.3. Activation

Upon muscle injury, MuSCs rapidly enter the cell cycle. Although not fully understood yet, several pathways involved in the control of MuSC activation are redox-regulated (**Figure 10**).

7.3.1. Calmodulin-nitric oxide pathway

Several lines of evidence indicate that NO[•] is involved in MuSC activation. *In vivo* inhibition of NOS activity by the pharmacological inhibitor N(G)-nitro-(L)-arginine methyl ester (L-NAME) prevents the immediate injury-induced MuSC activation and their subsequent proliferation (10,100,319). In the single myofiber model, NO[•] donation (SIN-1) or inhibition (L-NAME) increases and decreases the number of Pax7⁺ quiescent MuSCs, respectively (46). Similar changes of the number of activated MuSCs are observed in NOS deficient mice (nNOS^{-/-} or iNOS^{-/-}) after muscle injury (280). *In vitro*, stretching myofibers induces Hepatocyte Growth Factor (a potent MuSC mitogenic factor) release in a metalloprotease and NO[•]-dependent manner (356). Upstream, calcium-calmodulin, known to associate with NOS in other cell types (347), activates NOS activity in MuSCs; its specific inhibition abrogates the

calcium ionophore-induced MuSC activation (320). Collectively, these data argue for a positive role of NO[•] in MuSC activation following muscle injury.

7.3.2. Sphingolipids

Sphingolipids are a group of lipids playing a crucial role in cell membrane structure and lipid signaling. They may act as second messengers to stimulate oxidant production (11,241). Notably, ROS production from mitochondria has been identified as regulator of sphingosine 1-phosphate (254). Interestingly, sphingolipid signaling is required for the entry of MuSCs in the cell cycle and sphingosine 1-phosphate regulates MuSC proliferation *in vitro* and *in vivo* (233). Consequently, inhibition of sphingolipid signaling impairs skeletal muscle regeneration (119).

7.3.3. p38 MAPK

Mitogen-activated protein kinases (MAPKs) are members of many signaling pathways in eukaryotic cells and are known to be activated in response to oxidative stress *in vitro* and *in vivo* (161). p38 δ/β MAPK are required for MuSC activation, their specific inhibition leading to a reduced expression of MyoD and the acquisition of a quiescent state (327). This may support a role of ROS in p38 δ/β MAPK-dependent MuSC activation.

7.3.4. Notch

The Notch pathway is critical for multiple stages of post-natal myogenesis. Inhibition of Notch signaling abolishes MuSC activation and impairs muscle regeneration (72). In addition to MuSC activation, Notch signaling is also involved in MuSC proliferation, while it must be shut down for their differentiation through antagonism with the Wnt pathway (see below) (40,72). While there is not yet any evidence in muscle, it was recently shown that Notch is modulated by the ROS-producer NOX-1 to control the fate of colonic precursor cells (67), indicating that this

important signaling pathway in myogenesis may also be regulated by ROS during muscle regeneration.

7.3.5. Wnt

The Wnt/ β Catenin pathway plays important various roles in MuSC biology and during regenerative myogenesis (review in (344)). Collectively, several Wnts, through the classical pathway and the planar cell polarity pathway, have been shown to control MuSC fate (179), their motility (31), their proliferation (248) as well as their differentiation (40). In non-muscle systems, several redox molecules regulate Wnt signaling. Trx nucleoredoxin is a putative interactor of Dishevelled along the Wnt/ β Catenin pathway that would suppress the signaling in a redox-dependent manner (115). Another redox molecule, selenium, is associated with the Wnt pathway through Gpx1. Selenium restriction is associated with Wnt activation, whereas high selenium levels inhibits Wnt signaling (41,169). These studies show that ROS regulate the Wnt/ β Catenin pathway. Given its importance in MuSC biology, it is likely that this pathway is also the target of redox regulation in skeletal muscle.

7.4. Proliferation

The cell cycle has emerged as a redox cycle (review in (48,59)). Twenty years ago, Burdon and colleagues stated that low concentrations of oxidants ($O_2^{\cdot-}$, H_2O_2) stimulate mammalian cell proliferation (47). Similarly, $O_2^{\cdot-}$ production by MuSCs depends on NADPH oxidases (Nox2, Nox4, p22(phox), p47(phox), and p67(phox)), and MuSC proliferation is inhibited by antioxidants (N-acetylcysteine (NAC), apocynin) or NOX silencing by controlling cyclin D1 level through Erk1/2, PI-3 kinase/AKT and NF- κ B pathways (223). Furthermore, it has recently been shown that zinc promotes the MuSC proliferation in a dose-dependent manner through the

positive control of PI3K/Akt and ERK signaling pathways, although it is not known whether this occurs through a redox-dependent mechanism (244). Direct evidence for the involvement of ROS comes from genetically deficient mouse strains. Gpx1-deficient MuSCs exhibit lower proliferation rates and decreased differentiation capacity compared with wild-type MuSCs (183). MuSCs from SOD1-G93A mice (a mutation that leads to amyotrophic lateral sclerosis) demonstrate lower proliferation potential, triggering lower differentiation while the expression of cyclin-dependent kinase inhibitor 1A, which is known to promote myoblast differentiation, is reduced (205).

7.5. Myogenic differentiation

Myogenic differentiation is an important step in myogenesis that arises from reprogramming of MuSCs to generate differentiated cells that eventually fuse into plurinucleated myotubes/myofibers. Detoxification and DNA repair appear to be crucial steps before the induction of skeletal muscle differentiation to avoid propagation of genomic abnormalities (263). This suggests that antioxidant and cytoprotective genes could be crucial players involved in the control of efficient myogenic differentiation, and several studies support this notion (**Figure 10**). *In vitro*, MuSC differentiation into myotubes elicits a rapid decline in NOX4 and p67^{phox} expression, suggesting a change in ROS production at the time of myogenic commitment (223). Overexpression and inhibition of DUOX1 in MuSCs reduces and enhances *in vitro* myogenic differentiation, respectively (293). *In vivo*, genetic deletion of the p66^{ShcA} protein adaptor that promotes ROS formation, results in a lower mitochondrial ROS (mtROS) production and a better muscle regeneration (360). In the same line, differentiation defect is observed in a mouse model deleted

for *sepn1* gene (52), confirming previous reports on selenium deprivation of MuSCs (328). A molecular mechanism may be through NF- κ B, which has been shown to regulate myogenic differentiation (18). In addition to NF- κ B, the Nrf2-glutamate-cysteine ligase (GCL)/glutathione reductase (GR)-GSH signaling pathway is involved at the time of myogenic differentiation to regulate the GSH redox status after endogenous H₂O₂ is produced (259). Moreover, it has been shown that the Nrf-2 target gene carbonyl reductase 1 (CRB1), a NADPH-dependent cytosolic enzyme that reduces highly reactive aldehydes and prevents both protein carbonylation and DNA damage, is upregulated during MuSC differentiation. CRB1 inhibition is associated with an increase of ROS production, and with impaired differentiation and skeletal muscle regeneration *in vivo* (190). Consistently, skeletal muscle regeneration is strongly delayed in Nrf2 deficient animals, where alteration in the time course of myogenesis is observed and has been attributed to an alteration of MuSC proliferation/differentiation balance (4,304). Similarly, *in vitro* experiments showed that MuSCs cultured with dithiothreitol (a reductant agent) display a higher expression of the contractile protein myosin heavy chain as compared with those cultured under oxidized conditions (H₂O₂ in medium) (341). Similar results were observed on myogenic differentiation *in vitro* using the antioxidant resveratrol (160,224).

However, other studies support pro-differentiating effects of ROS in skeletal muscle. MuSC differentiation is accompanied by an increase of mtROS production (200), especially H₂O₂ (184,353). As a response, an upregulation of antioxidant enzymes, mainly with peroxidase function such as Prx2, is also observed (109,184,353). Additionally, treatment of MuSCs with mitochondria-targeted antioxidants such as MitoQ, MitoTEMPOL or MCAT alters their differentiation (184),

as does treatment with nordihydroguaiaretic acid (NDGA) (146). Particular attention was given to NO[•] effects on myogenic differentiation. First, experiments using chick MuSCs showed that NOS activity transiently increases at the time of differentiation and requires Ca²⁺, calmodulin, and NADPH (182). This transient activity is associated with the activation of NF-κB sites in the promoter region of the gene encoding iNOS. Interestingly, only differentiating, and not proliferating, cells exhibit these properties (182). Consequently, a NO[•] donor (sodium nitroprusside) induces precocious MuSC fusion, while a NO[•] competitor (NG-monomethyl-L-arginine) delays fusion (181). These results were confirmed in a mouse model in which L-Arginine, the substrate for endogenous synthesis of NO[•], transiently increases NO[•] production and triggers MuSC fusion. L-Arginine effects are prevented by the NOS inhibitor L-NAME and by L-Lysine, a competitive inhibitor of L-Arginine uptake (195). More recently, it has been shown that these NO[•]-mediated effects rely on mitochondria fusion at the time of MuSC fusion since NO[•] inhibition enhances the activity of a pro-fissioning GTPase dynamin-related protein-1 that triggers mitochondrial dysfunction (81).

Collectively, the major concern of all the above studies is the lack or inappropriate characterization and/or quantification of ROS entities. Moreover, opposite conclusions may result from the non-linear dose-response relationship of oxidative stress on biological effects. Indeed, pioneer studies showed that NO[•] exerts positive or negative effects on MuSC proliferation in a dose-dependent way *in vitro* (331). Moreover, several studies showed transient ROS upregulation at the time of differentiation (181,195) suggesting a tight regulation of their release in the MuSC neighborhood. A recent study, using muscle conditional mutants of the homeodomain transcription factors Pitx2/3, highlights the importance of ROS levels in myogenesis (175). Double pitx2/3 mutants show excessive upregulation of ROS levels leading to

DNA damage and apoptosis of differentiating MuSCs, both during fetal and adult myogenesis. Pitx2/3 deficiency triggers the downregulation of antioxidant genes (such as Nrf1) and impairment of mitochondrial function. Interestingly, premature muscle differentiation occurs in single Pitx3 mutants. These various effects are directly linked to dose-dependent changes in ROS levels and can be reversed by lowering ROS with NAC. Thus a controlled increase in ROS is required for MuSC differentiation (175).

Although the impact of oxidative stress on MuSC fate is recognized, only a few studies have explored the regulation of the master genes MyoD and Myogenin by the redox balance. Cellular redox status is reflected in the balance between a number of reduced or oxidized molecular pairs including NADH/NAD. Some transcription factors and DNA modifying enzymes are sensitive to NADH/NAD ratio, implying that cellular redox status regulates gene transcription (192). At the time of MuSC activation, reprogramming of cellular metabolism decreases intracellular NAD⁺ levels, decreasing in turn the activity of Sirtuin 1, a NAD⁺-dependent histone deacetylase. This leads to elevated H4K16 acetylation and activation of MyoD transcription. SIRT1 forms a complex with the acetyltransferase PCAF and MyoD and, when overexpressed, delays muscle differentiation (113,286). These results indicate that the redox status can control the myogenic gene expression program. Later on during myogenesis, exposing differentiating MuSCs to oxidative stress increases the nuclear localization of Focal adhesion kinase (FAK), which interacts with MBD2 (methyl CpG-binding protein 2). This leads to a decreased MBD2 association with HDAC1 (histone deacetylase complex 1) and methyl CpG sites in the myogenin promoter, thus inducing myogenin expression (198). Finally, Heme oxygenase-1 (HMOX1), which is a cytoprotective enzyme and whose expression is induced in response to oxidative

stress, inhibits miRNA processing leading to a decrease of several myomiRs (miR-1, miR-133a/b, miR-206), which triggers a reduced expression of MyoD, myogenin, and myosins. However, it is important to note that this activity is independent of the antioxidative properties of HMOX1 (172).

ROS are clearly involved in the numerous signaling pathways that control the sequential steps of MuSC fate following skeletal muscle injury (**Figure 10**). The recent development of new tools (redox sensors, redox proteomics) will allow for a more accurate identification and quantification of these ROS and for the identification of which entity, at which level, and during which time-period, controls a specific stage of MuSC fate. Finally, MuSCs evolve in a continuously changing environment during muscle regeneration, and receive cues from their immediate microenvironment, which has been shown to be necessary for efficient regeneration, and which is also subjected to redox regulation.

8. Redox regulation of the cells present in the MuSC microenvironment

Skeletal muscle regeneration requires the involvement of many cell types, including vessel cells, interstitial cells and inflammatory cells, such as monocytes/macrophages (55) (**Figures 3,4**).

8.1. Myeloid cells

After skeletal muscle injury, leukocytes rapidly invade the damaged area. While neutrophils rapidly disappear in few days, monocytes convert into macrophages that exert various roles on MuSCs and on the regeneration process (289) (**Figures 4,11**). Numerous data on redox signaling in macrophages are available in the context of infection where they are shown to actively produce ROS (44,106). Inversely, only a

few studies have investigated the redox status and redox-regulation of macrophage biology during tissue injury/repair. In wound healing models in zebrafish and drosophila embryo, leukocyte recruitment depends upon a H₂O₂ signal gradient generated by DUOX activation (237,271). In skeletal muscle, as in other tissues, entering monocytes/macrophages exhibit a damage-associated inflammatory profile and release pro-inflammatory cytokines (TNF- α , IL-1 β , IL-6). In skeletal muscle the most studied redox regulated factor is high-mobility group box 1 (HMGB1), an ubiquitous nuclear protein acting as a Damage Associated Molecular Pattern to signal tissue damage when released extracellularly. When cysteine residues are reduced, HMGB1 promotes the recruitment of inflammatory cells *via* its interaction with the CXCL12 chemokine to bind the CXCR4 receptor. When cysteine residues are oxidized, HGMB1 promotes the release of proinflammatory cytokines (339). After injury, an oxidized form of HMGB1 is detected in the muscle within 6 h (339). Then, an antioxidant response is observed in the regenerating muscle, notably in leukocytes, which express high levels of SOD1 and Trx, suggesting that a biphasic expression of oxidized/reduced forms of HMGB1 is required for skeletal muscle regeneration. In accordance, HMGB1 oxidation abrogates migration of MuSCs and their ability to differentiate into myotubes *in vitro* (341). To date, no study has characterized ROS production in macrophages in the context of skeletal muscle regeneration or tissue repair. iNOS is expressed by inflammatory macrophages that infiltrate a tissue upon infection. It is also expressed by inflammatory macrophages during the first steps of muscle regeneration, its expression declining with time, and notably in restorative anti-inflammatory macrophages (227). It has been shown that mice deficient for iNOS (which is mainly expressed by macrophages infiltrating the injured muscle) fail to regenerate properly because of a defect of MuSC proliferation

and differentiation, associated with an increase of inflammatory markers (280), indicating a role of macrophage-derived iNOS in skeletal muscle regeneration.

The pro-inflammatory phase ends with phagocytosis of muscle debris, at which point macrophages switch their phenotype to give rise to anti-inflammatory/restorative macrophages (15,227). These restorative macrophages produce soluble factors (IL-10, TGF- β , VEGF, IGF-1) to dampen inflammation and stimulate tissue repair, notably triggering myogenic differentiation (**Figure 11**) (15,227,290). The switch (skewing) of macrophage phenotype occurring during skeletal muscle regeneration is a key feature of efficient repair (227). Interestingly, all effectors identified so far as regulators of this process are redox-sensitive. The MKP-1/p38MAPK axis, which belongs to the redox-sensitive MAPK family (7), regulates macrophage phenotype skewing in a timely way to promote the resolution of inflammation (257). Another redox sensitive pathway, CREB-C/EBP β (27) is required for the expression of the anti-inflammatory program in macrophages (285). A third effector is AMPK α 1, which is required in macrophages for the acquisition of the anti-inflammatory phenotype, ensuring efficient muscle regeneration (227). AMPK is linked to redox-based signals, notably it is targeted by pro-oxidant species, suggesting a role for AMPK in the tuning of redox-dependent processes (review in (51)). A fourth molecule has been involved in the skewing of macrophages during skeletal muscle regeneration. Insulin Growth Factor-1 (IGF-1), which is a master regulator of myogenesis and skeletal muscle growth, is expressed mainly by macrophages during muscle regeneration and is required for their skewing in an autocrine way (324). One study reported that ROS suppression (UK-134 and α -lipoic acid) restores IGF-1 sensitivity of diabetic fibroblasts and improves their healing functions *in vitro* and *in vivo* (37).

In vitro studies have shown that ROS/RNS production is needed for macrophage skewing. Inhibition of $O_2^{\cdot-}$ production specifically blocks the differentiation of alternatively activated (G-CSF+IL4) (so-called M2) macrophages. *In vivo*, continuous administration of a ROS inhibitor (butylated hydroxyanisole) efficiently blocks the occurrence of Tumor Associated Macrophages, which exhibit a M2 phenotype in mouse cancer models (362). Similarly, macrophages issued from SOD1^{-/-} mice display a pro-inflammatory profile as compared with WT macrophages (134). Inversely, in SOD1-Tg mice, where SOD1 is overexpressed and H₂O₂ generation is enhanced, macrophages exhibit an anti-inflammatory status, associated with an increased activation of STAT6 (134), which is known to trigger the expression of anti-inflammatory genes in macrophages (178).

8.2. Endothelial cells

Skeletal muscle regeneration is associated with the angiogenic process, in order to rebuild the vascular bed after injury. Privileged interactions have been described between MuSCs and ECs. MuSCs are located close to capillaries and their number increases with the capillarization of myofibers (63). MuSCs express several angiogenic factors, including VEGF, and exhibit angiogenic properties (278). Similarly, ECs release factors supporting MuSC growth. Thus MuSCs and ECs interact to support skeletal muscle regeneration (63,177,226). While the role of ROS in these processes has not yet been investigated in skeletal muscle, there is a considerable amount of evidence coming from other models that ROS modulate angiogenesis through the regulation of redox-sensitive signaling pathways. This results in the control of EC growth, proliferation and migration (review in (95,111)). ECs generate ROS such as $O_2^{\cdot-}$ and H₂O₂. NOX is the main source of ROS in ECs

and is activated by growth factors, cytokines, hormones and hypoxia (125). Exogenous ROS, as well as overexpression of NOX4, increase the expression of VEGF, VEGFR2 and Angiopoietin 1 and stimulate EC proliferation, migration and angiogenesis *in vitro* (1,64,79,334). Furthermore, Angiopoietin 1-induced angiogenesis is enhanced in CAT deficient mice, suggesting an involvement of H₂O₂ in angiogenesis (168). Conversely, in Tie2-driven CAT transgenic mice (where CAT is constitutively expressed in ECs), impairment of neovascularization and reduced vessel sprouting following injury are observed (333). In the same line, silencing the 66^{ShcA} protein adaptor, which promotes ROS formation and by which phosphorylation is enhanced by VEGF, inhibits both VEGF-induced ROS generation and EC proliferation and migration (245).

Few studies have directly investigated the role of ROS in the environment of MuSCs during skeletal muscle regeneration. However, evidence from other models of tissue repair argues for the redox regulation of both immune cells and ECs, which likely takes place in the skeletal muscle milieu.

9. Skeletal muscle regeneration in pathophysiological contexts

9.1. Exercise-induced muscle damage

In a pioneering paper, Davies and colleagues demonstrated an increase in free radical concentrations in rat skeletal muscle following exercise to exhaustion (80). Since then, aerobic and anaerobic exercises have clearly been shown to increase ROS/RNS production in humans (102)). The relationship between ROS and contractile activity has also been widely investigated. NO[•] derivatives and ROS

modulate contractile function of skeletal muscle (273). At rest and during exercise, skeletal muscle produces both species (261) and produces antioxidant defense systems as an adaptive response (154). It is now acknowledged that low and physiological levels of ROS are required for force production, while high levels of ROS promote contractile dysfunction, as a result of oxidative damage of proteins and lipids, triggering muscle weakness and fatigue (261).

Exercise-induced muscle damage (EIMD) is particularly induced by eccentric contractions, which require lengthening of the muscle (112,143,262). Following physical exercise with repeated eccentric contractions, such as long-distance or downhill running, soreness appears within a few hours following the effort, a phenomenon called delayed-onset muscular soreness (DOMS). Despite an unclear etiology, DOMS is characterized by various markers such as histological changes (Z-line disruption), biochemical changes (increased creatine kinase and lactate dehydrogenase in plasma/serum) and altered clinical scores (increased soreness, reduced specific force, decreased range of motion). Several studies showed that damaging exercises are temporally associated with an increase in ROS production (66) and oxidative stress markers (239). Lipid peroxidation markers such as malondialdehyde (MDA), thiobarbituric-acid reactive substances, or protein carbonyls are increased in the blood (99,240) and in the skeletal muscle (218). In parallel, no meaningful alteration of antioxidant enzyme activity is reported in the blood (review in (180)). In whole muscle lysates, the GSH/GSSG ratio is slightly decreased at 2 and 4 days post-exercise, while the levels of antioxidant enzyme activity only slightly increase, most notably when compared with markers of oxidative stress/damage (218,240,313). Moreover as Insulin Receptor Substrate-1 oxidation is induced after a strenuous exercise, it was proposed as a possible explanation for the impairment of

insulin-induced glucose uptake observed in damaged muscle, through the alteration of insulin signaling (12).

A difficulty of such skeletal muscle physiology studies is to evaluate the level of myofiber damage, which is dependent not only on the type of exercise, but also on the status of the subject (healthy/pathologic, trained/sedentary, young/old, etc.) (253). Thus, the link between EIMD and MuSC activation is still unclear. Several levels of EIMD, classified according to the loss of muscle force, have been proposed and linked to the relative extent of inflammation (253). Similar comparative investigations should help to determine the role of ROS in EIMD.

9.2. Aging

Aging is a process characterized by an impairment of functional capacities. Many studies on ROS during aging were inspired by Harman's "free radical theory of aging" (131), which reports that "aging and the degenerative diseases associated are attributed basically to the deleterious side attacks of free radicals on cell constituents and on the connected tissues". Later on, the identification of mitochondria as the main source of ROS and oxidative damage during aging strengthened this free radical theory of aging (221). Interestingly, aging is not only associated with an increase of oxidative damage, but also with an up-regulation of antioxidant enzyme levels in skeletal muscle. Indeed, aging increases skeletal muscle content of Trx1, Trx2, TrxR2, cytosolic and mitochondrial SOD, catalase and Gpxs (86,155).

Aggregation of damaged proteins, mitochondrial dysfunction, epigenetic alterations and accumulation of toxic metabolites (ROS) and of DNA damage are hallmarks of aged stem cells (243). In skeletal muscle, proteomic investigations revealed that aged skeletal muscle is targeted by oxidative protein post-translational

modifications such as carbonylation that alters protein functions (19). Moreover, aging causes a reduction in redox-sensitive proteins involved in energy metabolism, indicating a loss in the flexibility of the redox energy response (211). While numerous studies have investigated the impact of ROS and its regulation on aged skeletal muscle (review in (12,148)), few experimental data establish a link between MuSC aging and ROS (**Figure 12**). In aged muscle, MuSCs exhibit a reduced capacity to proliferate and self-renew while they are prone to differentiation. This is explained by a deregulation of several signaling pathways, including decreased Notch signaling (71), increased Wnt signaling (36), increased JAK-STAT3 (322) and p38 MAPK (32) signaling pathways. All these pathways are redox-sensitive as discussed above (see section 7.3). Only a few studies directly investigate ROS in aged MuSCs. *In vitro*, old MuSCs are refractory to stretch and NO[•] induced activation, a property that is rescued by NO[•] donors (33). Increase in oxidation of DNA and lipids is observed in the elderly MuSCs, together with a reduction in CAT and glutathione transferase activities (114). Similarly, an increase in lipid peroxidation is observed in myotubes derived from aged MuSCs, and is associated with a defective status of the excitation-contraction coupling apparatus (25). Cultured aged MuSCs exhibit increased O₂^{•-}/mitochondrial mass and H₂O₂/mitochondrial mass ratios. The increased ROS production correlates with the corresponding decreased mitochondrial ATP production (220). The antioxidant Nrf2 signaling is impaired in aging muscle (219). Of note, in Nrf2 deficient mice subjected to acute endurance exercise stress, alteration in the expression of the MuSC markers Pax7 and MyoD is observed along with the activation of ubiquitin and apoptotic pathways (234). More recently, studies demonstrated that autophagy has an important role in sustaining the higher bioenergetic demands required for MuSC activation (318) and maintaining MuSC

stemness by preventing the senescence caused by mitochondrial dysfunction and oxidative stress during aging (117). Hence, it is clear that ROS and autophagy participate in skeletal muscle homeostasis (193,268) and are involved in muscle dystrophies (249); however, it is still unclear whether ROS are crucial for the induction of autophagy and whether up or down regulation of these processes induces a detrimental or a beneficial response (282). Another redox sensitive mechanism has recently been involved in the maintenance of aged MuSCs. Reduction of cellular NAD⁺ pool (which is required for mitochondria function) blunts the adaptive mitochondrial unfolded protein response (mtUPR) pathway that leads to the loss of MuSCs and decrease of their regenerative capacities (361). Although the precise mechanisms are still to be deciphered, these studies show that in aged skeletal muscle, higher ROS provide an unfavorable environment, which can adversely affect MuSC survival and functions and limit skeletal muscle repair (9).

9.3. Muscular dystrophies

The increase of oxidative stress in muscular dystrophy, which was acknowledged a long time ago (132,228), has been shown to correlate with the severity of the pathology (274) and is considered to contribute to the pathology of several muscular dystrophies (173). Interestingly, mutations in antioxidant *sepn1* gene cause SEPN1-related myopathy, a different class of muscle disease (14). While oxidative damage is assumed to participate in the pathogenesis of the disease, the exact mechanisms are still poorly understood, particularly at the level of the sarcolemma, where there is no direct experimental evidence of myofiber damage by oxidative stress (review in (321)). Calcium homeostasis is perturbed in muscular dystrophies (6). In this context, it has been shown that S-nitrosylation of the Ryanodine Receptor 1 (RyR1), an

intracellular calcium channel essential for the excitation-contraction coupling of the myofiber, induces a leak of sarcoplasmic reticulum Ca^{2+} , contributing to muscle weakness (28,29,94). Nevertheless, cellular and molecular mechanisms involving ROS signaling in the MuSC defects observed in muscular dystrophies are still poorly understood (**Figure 13**). The first evidence came from cultured MuSCs derived from either normal or dystrophic muscles. In mice, dystrophin deficient MuSC-derived myotubes are more susceptible to free radical-induced injury (269). The relative susceptibility of MuSC-derived myotubes to oxidative stress is associated with the severity of the dystrophy (assessed by the expression of different lengths of the dystrophin protein), and the extent of protein oxidation, suggesting that the DGC complex has important functions in antioxidant defense (88). It was shown that nNOS, which binds to DGC, has no impact on MuSC susceptibility to oxidative stress (363). Inversely, increased susceptibility of dystrophin-deficient-derived myotubes to oxidative stress is associated with altered phosphorylation of MAPKs, including JNK, p38 and ERK1/2 (309), which are both targeted by ROS and involved in myogenesis (see above, section 7). Susceptibility to oxidative stress seems to be a common feature in several muscular dystrophies. Indeed, MuSCs derived from affected muscles in facioscapulohumeral muscular dystrophy (FSHD) patients exhibit higher susceptibility to an induced oxidative stress than MuSCs derived from non-affected muscles of the same patients (350). This is associated with a defect in the size and branching of the differentiated myotubes (23). At the regulatory levels, it was shown that the transcriptional repressor Stra13, which is elevated in dystrophin deficient mdx mice, protects MuSCs from oxidative damage while Stra13 deficiency exacerbates muscular dystrophy (340). Epigenetic regulation has also been reported to control redox signaling and MuSC fate. The Polycomb group protein Bmi1, which

is an essential epigenetic regulator of stem cell function, is upregulated during normal skeletal muscle regeneration, whereas the number of MuSCs expressing high Bmi1 level is reduced in mdx muscle. Upregulation of Bmi-1 in MuSCs in mdx triggers the increase of metallothionein that confers MuSCs the ability to resist oxidative stress and to limit DNA oxidative damage, thus improving the pathology (84). These data confirm that ROS are elevated in muscular dystrophies and that MuSCs are more sensitive to oxidative stress, altering their functions and participating in the skeletal muscle deterioration in these progressive diseases.

Only a few studies have investigated the role of oxidative stress in the non-myogenic cell types in muscular dystrophies. It is assumed that NO deficiency in DMD contributes to the pathogenesis of the disease (69). In transgenic mdx animals expressing normal NO levels (through the constitutive expression of nNOS), an improvement of the muscle state is observed with a decrease of inflammation and of sarcolemma injury (346). In this context, long term treatment of dystrophic mice with HCT 1026, a nitric oxide-releasing derivative of flurbiprofen, ameliorates the morphological, biochemical, and functional phenotype of the diseased muscle (45) through a decrease of the number of inflammatory cells and FAPs, thus reducing fibrosis (74).

10. Therapeutics opportunities

Various strategies have been used to improve recovery of skeletal muscle function. Here, we present data related to exercise and antioxidant supplementation as approaches to improve skeletal muscle homeostasis.

10.1. Exercise as therapy

Skeletal muscle is a secretory organ. Upon contractile activity, it releases myokines that are known to control hypertrophy (LIF, IL-4, IL-6, IL-7, IL-15) (review in (256)). Some myokines have also been identified as crucial molecules involved in skeletal muscle regeneration (290). Few studies have investigated the role of exercise/contractile activity following skeletal muscle injury. It appears that exercise (running) increases skeletal muscle mass and cross sectional area of regenerating muscle as well as recovery of metabolic properties (171,279). These effects can be explained by a higher activation of the mTOR pathway, the master regulator of cell growth and protein synthesis previously described as a redox-regulated molecule (294) and of PGC-1 α (171), the master regulator of oxidative metabolism (191,196).

Exercise, through ROS production, initiates the transcription of antioxidant genes and specific signaling pathways (122,154). Thus, an increased production of ROS may act as an activator of the adaptive response in skeletal muscle, resulting in the increased production of antioxidant enzymes that could facilitate rapid muscle remodeling and provide protection against subsequent periods of damaging exercise (209). However, no study specifically investigated the effect of exercise on ROS in MuSCs. In contrast, endurance exercise of long duration and/or high intensity appears to generate much higher levels of ROS that can overwhelm cellular antioxidant defenses, and may cause tissue damage. This paradoxical effect of physical exercise refers to the hormesis theory (267). Such ROS-induced damage requires effective repair systems, and sufficient recuperation, before exercise adaptation can recommence (288). However, the over generation of ROS observed during acute exercise is blunted in endurance trained subjects, and most studies demonstrate that post-exercise oxidative stress is reduced after endurance exercise training (267). The main mechanism for this reduction in oxidative stress is likely the

upregulation of enzymatic antioxidants in response to an exercise-training program (261). SOD, one of the initial antioxidant enzymes expressed in response to oxidative stress, has been regularly reported to increase in response to exercise training (185) whereas activity of Gpx demonstrates a muscle fiber-specific pattern with type IIa being the most responsive to training (260). On the contrary, measurement of CAT activity in skeletal muscle in response to exercise training gives inconsistent results.

Aged skeletal muscle is more susceptible to damage, particularly following exercise and is unable to rapidly adapt following sequential periods of exercise. However, the elderly who are physically active benefit from exercise-induced adaptation in cellular antioxidant defense systems (326). Improved skeletal muscle mechanics, strength, and endurance make them less vulnerable to acute muscle injury and chronic inflammation (138). Therefore, many critical questions remain regarding the relationship of aging, ROS and exercise. At this time, there is no recommendation for exercising in dystrophic patients. It could be hypothesized that patients receiving pharmacological/gene/cell therapy could practice adapted exercise with exclusive concentric contraction to release beneficial myokines and adaptive responses to free radicals.

10.2. Antioxidant supplementation

10.2.1. In EIMD

The use of antioxidant in the field of physical exercise is still a controversial issue (review in (122,212,258)). Some groups have reported that antioxidant supplementation has a negative effect on training-induced adaptations and health (73,121,201,202,252,281). On the contrary, other groups reported no alterations of

training-induced performance improvements (225,357,358). These conflicting results have to be considered in the light of training status, type of exercise, qualitative and quantitative supplementation. Further studies are needed to elucidate this complex relationship. A wide range of studies has used nutritional antioxidant supplementation as a countermeasure to improve recovery following EIMD (212,231). Indeed, studies have investigated the role of vitamin C, vitamin E, combined antioxidant supplementation or NAC as potent protectors against skeletal muscle damage after eccentric contraction. Antioxidant supplementation before and 7 days after intensive knee maximal eccentric contraction reduces the loss of force following exercise as compared with the control group (303). However, some data indicate that antioxidants may delay muscle recovery (66,218). Indeed, prolonged vitamin C supplementation two days before and fourteen days after downhill running exercise reduces muscle soreness and MDA levels, but is associated with a delayed recovery of muscle function (66). In 2013, a double-blind study suggested that the modulation of redox biology by NAC supplementation for 8 days following quadriceps eccentric contractions at low velocity may interfere with skeletal muscle repair and myogenic program through the modulation of transcription factors involved in inflammation and myogenic program (218). In view of these results, there is little evidence to support a beneficial role of antioxidant supplementation in response to skeletal muscle damage (231).

As for the investigation of the ROS mechanisms in MuSCs, few studies explored the effects of antioxidant treatment on MuSC behavior following EIMD. In fact, to our knowledge, only one group has investigated the effects of antioxidant administration during post-contusion skeletal muscle regeneration at the cellular level. *In vivo* administration of polyphenol in rats 2 weeks before and after the injury induces an

accelerated activation of MuSCs, and consequently, a rapid neofiber formation that is completed earlier than in the control group (232). Furthermore, polyphenol treatment blunts the entry of neutrophils into the injured muscle, while the kinetics of macrophages and of their skewing to restorative macrophages is accelerated, and the production of the anti-inflammatory cytokine IL-10 is increased (174,232).

10.2.2. In aging

So far, most studies performed to counteract oxidative stress in aged muscle have targeted sarcopenia and muscle atrophy. Results consistently report significant improvements of antioxidant biomarkers after antioxidant supplementation while the effects on muscle function are more mitigated (review in (53)) (**Figure 14**). However, increased mitochondrial antioxidant activity, using transgenic mice with targeted overexpression of the human *cat* gene to mitochondria (mtCAT mice), restores age-dependent skeletal muscle dysfunction (332). This suggests that the preservation of muscle function in aging may be specific to the type of antioxidant and their site of action. Studies investigating the functions of MuSCs in aging muscle are rare and relatively new, due to the new development of tools allowing the specific analysis of these cells. *In vitro* treatment of human senescent MuSCs with tocotrienol rich fraction (Vitamin E) reduces their senescent phenotype and increases their proliferation capacity (190). In aged mice, antioxidant treatment (using Trolox, a vitamin E analog) prevents the appearance of senescence markers and restores the expansion capacity of MuSCs, and therefore their regenerative properties, through the recovery of an effective autophagy (117). Similarly, treatment with the NAD precursor nicotinamide riboside restores MuSC concentration of NAD⁺, improves mitochondrial homeostasis and maintains MuSC functions in aging mice (110,361). Moreover, in mice deficient for SIRT1 (an NAD⁺-dependent deacetylase that

increases mitochondrial biogenesis) in MuSCs, nicotinamide riboside beneficial properties are dampened in regenerating muscle, making a link between the NAD⁺-SIRT1 pathway, mitochondrial activity and MuSC function (361) (**Figure 14**).

10.2.3. In muscular dystrophies

More than 30 years ago, thus before the discovery of the dystrophin gene, supplementation with antioxidants (SOD precursors, vitamin E, selenium) was carried out in DMD patients since vitamin E deficiency triggers similar skeletal muscle damage as that observed in DMD patients (6). Since, antioxidants have been shown to reduce necrosis in skeletal muscle of dystrophin deficient mdx mouse (215). NAC administration results in decreased numbers of regenerating myofibers (349). FSHD patients supplemented with vitamin C, vitamin E, zinc gluconate and selenomethionine increase their antioxidant level and improve some physical performance (251). However, it should be acknowledged that most of the results obtained with antioxidant treatments are inconsistent (167), highlighting the complexity of redox biology (**Figure 14**).

Few studies have explored the effects of antioxidant treatment on MuSC behavior (**Figure 14**). *In vitro*, NDGA (inhibitor of Lysyl oxidase) treatment of primary MuSCs issued from a DMD patient inhibits their proliferation while stimulating their differentiation and myotube formation in a dose-dependent manner (118). Green tea polyphenol blend and its major polyphenol (EGCg) efficiently protect mdx myotubes from oxidative damage since mdx MuSCs express 7 fold more 67LR, a receptor for EGCg, than normal MuSCs (89). In desminopathies, mutations trigger the formation of desmin aggregates in MuSCs upon stress conditions, including redox-associated (H₂O₂ and cadmium chloride) stress. Antioxidants (fisetin, NAC, α-tocopherols,

Trolox) prevent desmin-aggregation during stress in MuSCs expressing specific desmin mutants (49,299). Still *in vitro*, it was shown that supplementation of MuSCs with α -tocopherol promotes plasma membrane repair upon oxidant challenge (141). *In vivo*, various antioxidant treatments trigger an improvement of the muscle phenotype of mdx or *sgca* KO (α -sarcoglycan null mice, a severe mouse model of dystrophy) mice, together with an increase in the muscle function. Some studies evaluated the impact of such treatments on MuSCs. Mdx treatment with flavocoxid for 5 weeks increases the number of myogenin-positive cells (216). Similarly, oral administration of the flavonol quercetin during 6 months to adult mdx mice increases the number of new fibers as well as the number of MuSCs, beside a general improvement of the muscle (139). In *sgca* KO mice, NCX 320 (cyclooxygenase inhibiting NO donator) treatment for 8 months leads to an increase in the number of newly formed fibers, as well as an increase of the number of MuSCs. Moreover, MuSCs exhibit an increased ability to differentiate *in vitro* and to express the differentiation markers myogenin and myosin heavy chain, indicative of their good regenerative potential (298). Finally, mdx mice supplemented for 10 weeks with nicotinamide riboside exhibit an increased number of MuSCs while muscle regeneration is improved, indicating that nicotinamide riboside also prevents MuSC senescence in muscular dystrophy (287,361), as it does in aging muscle (see above) **(Figure 14)**.

On the whole, these studies indicate that antioxidant supplementation can benefit muscle health in muscular dystrophies. It is worth mentioning that while the importance of inflammation and inflammatory cells in the establishment and maintenance of the pathology during muscular dystrophies is recognized, the phenotype and functions of these cells have been overlooked. In this context, it is

interesting to mention that antioxidant treatment has been shown to reduce the amount of inflammatory cells in these diseases, and this decrease is associated with the improvement observed in the muscle (298).

11. Perspectives

While redox signaling has long been known to play a role in skeletal muscle homeostasis, it has recently been established that the cellular and molecular processes at work during skeletal muscle regeneration are redox-regulated. Numerous signaling pathways that control MuSC biology as well as their close environment, including immune cells, are targeted by ROS/RNS and are altered in physiological (EIMD, aging) or pathological (muscular dystrophies) situations. Attempts to counteract oxidative stress in such contexts have led to inconsistent results that highlight the complexity of redox biology (6,167). This complexity has begun to be deciphered at the cellular and molecular levels. Measuring ROS concentrations, as well as redox modifications is difficult, notably *in vivo*, given the short half-lives, high specificities and high reactivity of the involved compounds. Very recent years have seen the development of tools to decipher redox complexity. Given the current knowledge, the field should move toward accurate identification of sources of ROS and their molecular targets. Indeed, sensor reporters will help to precisely identify and quantify the ROS/RNS that are produced in a tissue and in a given cell (**Figure 15**). Additionally, techniques of redox proteomics will help to decipher molecular targets of ROS/RNS and the nature of their post-translational modifications (**Figure 15**). Such technological improvements already form the basis of new therapeutic advancements. For example, the development of selective antioxidant probes has led to therapeutic efficiency in preclinical animal studies, and

in some cases, such as with the mitochondria-targeted antioxidant MitoQ, in phase II human trials (308). The design of a specific mitochondria-selective S-nitrosating agent, MitoSNO, has been found to be cardioprotective *in vivo* in mice after ischemia/reperfusion, through the S-nitrosation of mitochondrial complex I, which is the entry point for electrons from NADH into the respiratory chain (62). A genome-wide Cas9-mediated screen enabled the identification of the hypoxia response as protective during mitochondrial respiratory chain inhibition. Genetic or small-molecule activation of the hypoxia response is therefore protective against mitochondrial toxicity (150). NOX are also potent targets for therapeutic developments. New NOX inhibitors, identified through high-throughput screen after structure–activity relationship analysis, showed improved specificity for NOXs and were validated in *in vivo* animal studies although they require further developments to improve isoform selectivity. For instance, the NOX4 inhibitor GKT137831 has been shown to reduce fibrosis in several animal models of chronic diseases and is under clinical development (8). Inversely, after an acute injury, inducing a compartment-specific increase of the NAD^+/NADH ratio (using LbNOX water-forming NOX from *Lactobacillus brevis*) allowed to decipher the NAD^+/NADH ratio driven signaling and to improve proliferative and metabolic defects in mitochondrial deficient cells (323). Several studies recently confirmed that the redox couple NAD^+/NADH is a privileged target to improve cell and tissue homeostasis, including in aged and diseased skeletal muscle (see above, section 10.2) (110,287,361). Therefore, such an integrated approach would be an interesting option to better understand the underlying mechanisms linking redox regulation and skeletal muscle regeneration. Thus, the identification of redox-regulated networks will be useful to design

specifically targeted antioxidant probes in order to improve skeletal muscle phenotype and function.

Author Disclosure Statement

No competing financial interests exist.

Abbreviations

CAT: catalase
DGC: dystrophin-glycoprotein complex
DMD: Duchenne Muscular Dystrophy
DOMS: delayed-onset muscular soreness
ECs: endothelial cells
EIMD: exercise induced muscle damages
eNOS: endothelial NOS
FAPs: FibroAdipogenic Progenitors
FSHD: facioscapulohumeral muscular dystrophy
Gpx: glutathione peroxidase
Grx: glutaredoxin
GSH (GSSG): glutathione
GSNO: S-nitroglutathione
H₂O₂: hydrogen peroxide
HMGB-1: High mobility group box-1
HO•: hydroxyl radical
IGF-1: Insulin Growth Factor-1
iNOS: inducible NOS
L-NAME: N(G)-nitro-(L)-arginine methyl ester
MAPK: mitogen-activated protein kinase
MDA: malondialdehyde
mtROS: mitochondrial ROS
mtUPR: mitochondrial unfolded protein response
MuSC: muscle stem cell
NAC: N-acetylcysteine
NDGA: nordihydroguaiaretic acid
nNOS: neuronal NOS
NO•: Nitric oxide
NOS: NO• synthase
NOX: NADPH oxidase
O₂^{•-}: superoxide
Prx: peroxiredoxin
RNS: reactive nitrogen species
ROS: reactive oxygen species
SEPN1: selenoprotein N
SNO: S-nitrosothiols
SOD: superoxide dismutase
Trx: thioredoxin
TrxR: thioredoxin reductase
XO: xanthine oxidase
XOR: xanthine oxidoreductase

References

1. Abid MR, Tsai JC, Spokes KC, Deshpande SS, Irani K, and Aird WC. Vascular endothelial growth factor induces manganese-superoxide dismutase expression in endothelial cells by a Rac1-regulated NADPH oxidase-dependent mechanism. *FASEB J* 15: 2548-2550, 2001.
2. Abou-Khalil R, Le Grand F, Pallafacchina G, Valable S, Authier FJ, Rudnicki MA, Gherardi RK, Germain S, Chretien F, Sotiropoulos A, Lafuste P, Montarras D, and Chazaud B. Autocrine and paracrine Angiopoietin 1/Tie-2 signalling promotes muscle satellite cell self-renewal. *Cell Stem Cell* 5: 298-309, 2009.
3. Acharyya S, Villalta SA, Bakkar N, Bupha-Intr T, Janssen PM, Carathers M, Li ZW, Beg AA, Ghosh S, Sahenk Z, Weinstein M, Gardner KL, Rafael-Fortney JA, Karin M, Tidball JG, Baldwin AS, and Guttridge DC. Interplay of IKK/NF-kappaB signaling in macrophages and myofibers promotes muscle degeneration in Duchenne muscular dystrophy. *J Clin Invest* 117: 889-901, 2007.
4. Al-Sawaf O, Fragoulis A, Rosen C, Keimes N, Liehn EA, Holzle F, Kan YW, Pufe T, Sonmez TT, and Wruck CJ. Nrf2 augments skeletal muscle regeneration after ischaemia-reperfusion injury. *J Pathol* 234: 538-47, 2014.
5. Albrecht SC, Barata AG, Grosshans J, Teleman AA, and Dick TP. In vivo mapping of hydrogen peroxide and oxidized glutathione reveals chemical and regional specificity of redox homeostasis. *Cell Metab* 14: 819-29, 2011.
6. Allen DG, Whitehead NP, and Froehner SC. Absence of dystrophin disrupts skeletal muscle signaling: roles of Ca²⁺, Reactive Oxygen Species, and Nitric Oxide in the development of muscular dystrophy. *Physiol Rev* 96: 253-305, 2016.
7. Allen RGTresini M. Oxidative stress and gene regulation. *Free Radic Biol Med* 28: 463-99, 2000.
8. Altenhofer S, Radermacher KA, Kleikers PW, Wingler K, and Schmidt HH. Evolution of NADPH Oxidase Inhibitors: Selectivity and Mechanisms for Target Engagement. *Antioxid Redox Signal* 23: 406-27, 2015.
9. Alway SE, Myers MJ, and Mohamed JS. Regulation of satellite cell function in sarcopenia. *Front Aging Neurosci* 6: 246, 2014.
10. Anderson JE. A role for nitric oxide in muscle repair: nitric oxide-mediated activation of muscle satellite cells. *Mol Biol Cell* 11: 1859-1874, 2000.
11. Andrieu-Abadie N, Gouazé V, Salvayre R, and Levade T. Ceramide in apoptosis signaling: relationship with oxidative stress. *Free Radic Biol Med* 31: 717-728, 2001.
12. Aoi W, Naito Y, Tokuda H, Tanimura Y, Oya-Ito T, and Yoshikawa T. Exercise-induced muscle damage impairs insulin signaling pathway associated with IRS-1 oxidative modification. *Physiol Res* 61: 81-8, 2012.
13. Aoi W, Naito Y, and Yoshikawa T. Role of oxidative stress in impaired insulin signaling associated with exercise-induced muscle damage. *Free Radic Biol Med* 65: 1265-72, 2013.
14. Arbogast S Ferreiro A. Selenoproteins and protection against oxidative stress: selenoprotein N as a novel player at the crossroads of redox signaling and calcium homeostasis. *Antioxid Redox Signal* 12: 893-904, 2010.
15. Arnold L, Henry A, Poron F, Baba-Amer Y, van Rooijen N, Plonquet A, Gherardi RK, and Chazaud B. Inflammatory monocytes recruited after skeletal

- muscle injury switch into antiinflammatory macrophages to support myogenesis. *J Exp Med* 204: 1057-1069, 2007.
16. Babior BM. NADPH oxidase: an update. *Blood* 93: 1464-1476, 1999.
 17. Bachi A, Dalle-Donne I, and Scaloni A. Redox proteomics: chemical principles, methodological approaches and biological/biomedical promises. *Chem Rev* 113: 596-698, 2013.
 18. Bakkar NGuttridge DC. NF-kappaB signaling: a tale of two pathways in skeletal myogenesis. *Physiol Rev* 90: 495-511, 2010.
 19. Baraibar MA, Gueugneau M, Duguez S, Butler-Browne G, Bechet D, and Friguet B. Expression and modification proteomics during skeletal muscle ageing. *Biogerontol* 14: 339-352, 2013.
 20. Barja G. Mitochondrial oxygen radical generation and leak: sites of production in states 4 and 3, organ specificity, and relation to aging and longevity. *J Bioenerg Biomembr* 31: 347-366, 1999.
 21. Barjot C, Cotten ML, Goblet C, Whalen RG, and Bacou F. Expression of myosin heavy chain and of myogenic regulatory factor genes in fast or slow rabbit muscle satellite cell cultures. *J Muscle Res Cell Motil* 16: 619-28, 1995.
 22. Barreiro EHussain SN. Protein carbonylation in skeletal muscles: impact on function. *Antioxid Redox Signal* 12: 417-429, 2010.
 23. Barro M, Carnac G, Flavier S, Mercier J, Vassetzky Y, and Laoudj-Chenivesse D. Myoblasts from affected and non-affected FSHD muscles exhibit morphological differentiation defects. *J Cell Mol Med* 14: 275-289, 2010.
 24. Baudhuin P, Beaufay H, and De Duve C. Combined biochemical and morphological study of particulate fractions from rat liver. Analysis of preparations enriched in lysosomes or in particles containing urate oxidase, D-amino acid oxidase, and catalase. *J Cell Biol* 26: 219-43, 1965.
 25. Beccafico S, Puglielli C, Pietrangelo T, Bellomo R, Fano G, and Fulle S. Age-dependent effects on functional aspects in human satellite cells. *Ann N Y Acad Sci* 1100: 345-352, 2007.
 26. Bedard KKrause K-H. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* 87: 245-313, 2007.
 27. Bedogni B, Pani G, Colavitti R, Riccio A, Borrello S, Murphy M, Smith R, Eboli ML, and Galeotti T. Redox regulation of cAMP-responsive element-binding protein and induction of manganous superoxide dismutase in nerve growth factor-dependent cell survival. *J Biol Chem* 278: 16510-16519, 2003.
 28. Bellinger AM, Reiken S, Carlson C, Mongillo M, Liu X, Rothman L, Matecki S, Lacampagne A, and Marks AR. Hypernitrosylated ryanodine receptor calcium release channels are leaky in dystrophic muscle. *Nat Med* 15: 325-30, 2009.
 29. Bellinger AM, Reiken S, Dura M, Murphy PW, Deng SX, Landry DW, Nieman D, Lehnart SE, Samaru M, LaCampagne A, and Marks AR. Remodeling of ryanodine receptor complex causes "leaky" channels: a molecular mechanism for decreased exercise capacity. *Proc Natl Acad Sci U S A* 105: 2198-202, 2008.
 30. Belousov VV, Fradkov AF, Lukyanov KA, Staroverov DB, Shakhbazov KS, Terskikh AV, and Lukyanov S. Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat Method* 3: 281-286, 2006.
 31. Bentzinger CF, von Maltzahn J, Dumont NA, Stark DA, Wang YX, Nhan K, Frenette J, Cornelison DD, and Rudnicki MA. Wnt7a stimulates myogenic stem cell motility and engraftment resulting in improved muscle strength. *J Cell Biol* 205: 97-111, 2014.

32. Bernet JD, Doles JD, Hall JK, Kelly Tanaka K, Carter TA, and Olwin BB. p38 MAPK signaling underlies a cell-autonomous loss of stem cell self-renewal in skeletal muscle of aged mice. *Nat Med* 20: 265-271, 2014.
33. Betters JL, Lira VA, Soltow QA, Drenning JA, and Criswell DS. Supplemental nitric oxide augments satellite cell activity on cultured myofibers from aged mice. *Exp Gerontol* 43: 1094-1101, 2008.
34. Bigarella CL, Liang R, and Ghaffari S. Stem cells and the impact of ROS signaling. *Development* 141: 4206-4218, 2014.
35. Bindoli ARigobello MP. Principles in redox signaling: from chemistry to functional significance. *Antioxid Redox Signal* 18: 1557-1593, 2013.
36. Biressi S, Miyabara EH, Gopinath SD, Carlignani PMM, and Rando TA. A Wnt-TGF β 2 axis induces a fibrogenic program in muscle stem cells from dystrophic mice. *Sci Transl Med* 6: 267ra176, 2014.
37. Bitar MSAI-Mulla F. ROS constitute a convergence nexus in the development of IGF1 resistance and impaired wound healing in a rat model of type 2 diabetes. *Dis Model Mech* 5: 375-88, 2012.
38. Bogdan C, Rölinghoff M, and Diefenbach A. Reactive oxygen and reactive nitrogen intermediates in innate and specific immunity. *Curr Opin Immunol* 12: 64-76, 2000.
39. Boveris AChance B. The mitochondrial generation of hydrogen peroxide. General properties and effect of hyperbaric oxygen. *Biochem J* 134: 707-716, 1973.
40. Brack AS, Conboy IM, Conboy MJ, Shen J, and Rando TA. A temporal switch from notch to Wnt signaling in muscle stem cells is necessary for normal adult myogenesis. *Cell Stem Cell* 2: 50-59, 2008.
41. Brigelius-Flohé RKipp AP. Selenium in the redox regulation of the Nrf2 and the Wnt pathway. *Method Enzymol* 527: 65-86, 2013.
42. Brigelius-Flohé RMaiorino M. Glutathione peroxidases. *Biochim Biophys Acta* 1830: 3289-3303, 2013.
43. Brigelius-Flohé RTraber MG. Vitamin E: function and metabolism. *FASEB J* 13: 1145-1155, 1999.
44. Brüne B, Dehne N, Grossmann N, Jung M, Namgaladze D, Schmid T, von Knethen A, and Weigert A. Redox control of inflammation in macrophages. *Antioxid Redox Signal* 19: 595-637, 2013.
45. Brunelli S, Sciorati C, D'Antona G, Innocenzi A, Covarello D, Galvez BG, Perrotta C, Monopoli A, Sanvito F, Bottinelli R, Ongini E, Cossu G, and Clementi E. Nitric oxide release combined with nonsteroidal antiinflammatory activity prevents muscular dystrophy pathology and enhances stem cell therapy. *Proc Natl Acad Sci U S A* 104: 264-9, 2007.
46. Buono R, Vantaggiato C, Pisa V, Azzoni E, Bassi MT, Brunelli S, Sciorati C, and Clementi E. Nitric oxide sustains long-term skeletal muscle regeneration by regulating fate of satellite cells via signaling pathways requiring Vangl2 and cyclic GMP. *Stem Cells* 30: 197-209, 2012.
47. Burdon RH. Superoxide and hydrogen peroxide in relation to mammalian cell proliferation. *Free Radic Biol Med* 18: 775-794, 1995.
48. Burhans WCHeintz NH. The cell cycle is a redox cycle: linking phase-specific targets to cell fate. *Free Radic Biol Med* 47: 1282-1293, 2009.
49. Cabet E, Batonnet-Pichon S, Delort F, Gausseres B, Vicart P, and Lilienbaum A. Antioxidant treatment and induction of autophagy cooperate to reduce

- desmin aggregation in a cellular model of desminopathy. *PLoS One* 10: e0137009, 2015.
50. Cao SSKaufman RJ. Endoplasmic reticulum stress and oxidative stress in cell fate decision and human disease. *Antioxid Redox Signal* 21: 396-413, 2014.
 51. Cardaci S, Filomeni G, and Ciriolo MR. Redox implications of AMPK-mediated signal transduction beyond energetic clues. *J Cell Sci* 125: 2115-2125, 2012.
 52. Castets P, Bertrand AT, Beuvin M, Ferry A, Le Grand F, Castets M, Chazot G, Rederstorff M, Krol A, Lescure A, Romero NB, Guicheney P, and Allamand V. Satellite cell loss and impaired muscle regeneration in selenoprotein N deficiency. *Hum Mol Genet* 20: 694-704, 2011.
 53. Cerullo F, Gambassi G, and Cesari M. Rationale for antioxidant supplementation in sarcopenia. *J Aging Res* 2012: 316943, 2012.
 54. Chargé SBPRudnicki MA. Cellular and molecular regulation of muscle regeneration. *Physiol Rev* 84: 209-238, 2004.
 55. Chazaud B. Macrophages: supportive cells for tissue repair and regeneration. *Immunobiol* 219: 172-178, 2014.
 56. Chazaud B. Inflammation during skeletal muscle regeneration and tissue remodeling-application to exercise-induced muscle damage management. *Immunol Cell Biol*, 2015.
 57. Chelly JDesguerre I. Progressive muscular dystrophies. *Handb Clin Neurol* 113: 1343-1366, 2013.
 58. Cheung TH, Quach NL, Charville GW, Liu L, Park L, Edalati A, Yoo B, Hoang P, and Rando TA. Maintenance of muscle stem-cell quiescence by microRNA-489. *Nature* 482: 524-528, 2012.
 59. Chiu JDawes IW. Redox control of cell proliferation. *Trends Cell Biol* 22: 592-601, 2012.
 60. Chouchani ET, James AM, Fearnley IM, Lilley KS, and Murphy MP. Proteomic approaches to the characterization of protein thiol modification. *Curr Opin Chem Biol* 15: 120-128, 2011.
 61. Chouchani ET, Kazak L, Jedrychowski MP, Lu GZ, Erickson BK, Szpyt J, Pierce KA, Laznik-Bogoslavski D, Vetrivelan R, Clish CB, Robinson AJ, Gygi SP, and Spiegelman BM. Mitochondrial ROS regulate thermogenic energy expenditure and sulfenylation of UCP1. *Nature* 532: 112-116, 2016.
 62. Chouchani ET, Methner C, Nadtochiy SM, Logan A, Pell VR, Ding S, James AM, Cochemé HM, Reinhold J, Lilley KS, Partridge L, Fearnley IM, Robinson AJ, Hartley RC, Smith RAJ, Krieg T, Brookes PS, and Murphy MP. Cardioprotection by S-nitrosation of a cysteine switch on mitochondrial complex I. *Nat Med* 19: 753-759, 2013.
 63. Christov C, Chrétien F, Abou-Khalil R, Bassez G, Vallet G, Authier F-J, Bassaglia Y, Shinin V, Tajbakhsh S, Chazaud B, and Gherardi RK. Muscle satellite cells and endothelial cells: close neighbors and privileged partners. *Mol Biol Cell* 18: 1397-1409, 2007.
 64. Chua CC, Hamdy RC, and Chua BH. Upregulation of vascular endothelial growth factor by H₂O₂ in rat heart endothelial cells. *Free Radic Biol Med* 25: 891-897, 1998.
 65. Cifuentes-Pagano E, Meijles DN, and Pagano PJ. The quest for selective nox inhibitors and therapeutics: challenges, triumphs and pitfalls. *Antioxid Redox Signal* 20: 2741-54, 2014.
 66. Close GL, Ashton T, Cable T, Doran D, Holloway C, McArdle F, and MacLaren DPM. Ascorbic acid supplementation does not attenuate post-exercise muscle

- soreness following muscle-damaging exercise but may delay the recovery process. *The British Journal of Nutrition* 95: 976-981, 2006.
67. Coant N, Ben Mkaddem S, Pedruzzi E, Guichard C, Tréton X, Ducroc R, Freund J-N, Cazals-Hatem D, Bouhnik Y, Woerther P-L, Skurnik D, Grodet A, Fay M, Biard D, Lesuffleur T, Deffert C, Moreau R, Groyer A, Krause K-H, Daniel F, and Ogier-Denis E. NADPH oxidase 1 modulates WNT and NOTCH1 signaling to control the fate of proliferative progenitor cells in the colon. *Mol Cell Biol* 30: 2636-2650, 2010.
 68. Cocheme HM, Logan A, Prime TA, Abakumova I, Quin C, McQuaker SJ, Patel JV, Fearnley IM, James AM, Porteous CM, Smith RA, Hartley RC, Partridge L, and Murphy MP. Using the mitochondria-targeted ratiometric mass spectrometry probe MitoB to measure H₂O₂ in living *Drosophila*. *Nat Protoc* 7: 946-58, 2012.
 69. Colussi C, Gurtner A, Rosati J, Illi B, Ragone G, Piaggio G, Moggio M, Lamperti C, D'Angelo G, Clementi E, Minetti G, Mozzetta C, Antonini A, Capogrossi MC, Puri PL, and Gaetano C. Nitric oxide deficiency determines global chromatin changes in Duchenne muscular dystrophy. *FASEB Journal* 23: 2131-2141, 2009.
 70. Commoner B, Townsend J, and Pake GE. Free radicals in biological materials. *Nature* 174: 689-691, 1954.
 71. Conboy IM, Conboy MJ, Smythe GM, and Rando TA. Notch-mediated restoration of regenerative potential to aged muscle. *Science* 302: 1575-1577, 2003.
 72. Conboy IM, Rando TA. The regulation of Notch signaling controls satellite cell activation and cell fate determination in postnatal myogenesis. *Dev Cell* 3: 397-409, 2002.
 73. Coombes JS, Powers SK, Rowell B, Hamilton KL, Dodd SL, Shanely RA, Sen CK, and Packer L. Effects of vitamin E and alpha-lipoic acid on skeletal muscle contractile properties. *J Appl Physiol (1985)* 90: 1424-30, 2001.
 74. Cordani N, Pisa V, Pozzi L, Sciorati C, and Clementi E. Nitric oxide controls fat deposition in dystrophic skeletal muscle by regulating fibro-adipogenic precursor differentiation. *Stem Cells* 32: 874-85, 2014.
 75. D'Autréaux B, Toledano MB. ROS as signalling molecules: mechanisms that generate specificity in ROS homeostasis. *Nat Rev Mol Cell Biol* 8: 813-824, 2007.
 76. Dadgar S, Wang Z, Johnston H, Kesari A, Nagaraju K, Chen YW, Hill DA, Partridge TA, Giri M, Freishtat RJ, Nazarian J, Xuan J, Wang Y, and Hoffman EP. Asynchronous remodeling is a driver of failed regeneration in Duchenne muscular dystrophy. *Journal of Cell Biology* 207: 139-158, 2014.
 77. Dam AD, Mitchell AS, Rush JW, and Quadrilatero J. Elevated skeletal muscle apoptotic signaling following glutathione depletion. *Apoptosis* 17: 48-60, 2012.
 78. Dammeyer P, Arnér ESJ. Human Protein Atlas of redox systems - what can be learnt? *Biochim Biophys Acta* 1810: 111-138, 2011.
 79. Datla SR, Peshavariya H, Dusting GJ, Mahadev K, Goldstein BJ, and Jiang F. Important role of Nox4 type NADPH oxidase in angiogenic responses in human microvascular endothelial cells in vitro. *Arterioscler Thromb Vasc Biol* 27: 2319-2324, 2007.
 80. Davies KJ, Quintanilha AT, Brooks GA, and Packer L. Free radicals and tissue damage produced by exercise. *Biochem Biophys Res Commun* 107: 1198-1205, 1982.

81. De Palma C, Falcone S, Pisoni S, Cipolat S, Panzeri C, Pambianco S, Pisconti A, Allevi R, Bassi MT, Cossu G, Pozzan T, Moncada S, Scorrano L, Brunelli S, and Clementi E. Nitric oxide inhibition of Drp1-mediated mitochondrial fission is critical for myogenic differentiation. *Cell Death Differ* 17: 1684-1696, 2010.
82. Derbre F, Ferrando B, Gomez-Cabrera MC, Sanchis-Gomar F, Martinez-Bello VE, Olaso-Gonzalez G, Diaz A, Gratas-Delamarche A, Cerda M, and Viña J. Inhibition of xanthine oxidase by allopurinol prevents skeletal muscle atrophy: role of p38 MAPKinase and E3 ubiquitin ligases. *PLoS One* 7: e46668, 2012.
83. Desaint S, Luriau S, Aude J-C, Rousselet G, and Toledano MB. Mammalian antioxidant defenses are not inducible by H₂O₂. *J Biol Chem* 279: 31157-31163, 2004.
84. Di Foggia V, Zhang X, Licastro D, Gerli MF, Phadke R, Muntoni F, Mourikis P, Tajbakhsh S, Ellis M, Greaves LC, Taylor RW, Cossu G, Robson LG, and Marino S. Bmi1 enhances skeletal muscle regeneration through MT1-mediated oxidative stress protection in a mouse model of dystrophinopathy. *J Exp Med* 211: 2617-2633, 2014.
85. Dickinson BC, Chang CJ. Chemistry and biology of reactive oxygen species in signaling or stress responses. *Nat Chem Biol* 7: 504-511, 2011.
86. Dimauro I, Pearson T, Caporossi D, and Jackson MJ. In vitro susceptibility of thioredoxins and glutathione to redox modification and aging-related changes in skeletal muscle. *Free Radic Biol Med* 53: 2017-27, 2012.
87. Dinkova-Kostova AT, Holtzclaw WD, Cole RN, Itoh K, Wakabayashi N, Katoh Y, Yamamoto M, and Talalay P. Direct evidence that sulfhydryl groups of Keap1 are the sensors regulating induction of phase 2 enzymes that protect against carcinogens and oxidants. *Proc Natl Acad Sci U S A* 99: 11908-13, 2002.
88. Disatnik MH, Chamberlain JS, and Rando TA. Dystrophin mutations predict cellular susceptibility to oxidative stress. *Muscle Nerve* 23: 784-792, 2000.
89. Dorchies OM, Wagner S, Buetler TM, and Ruedg UT. Protection of dystrophic muscle cells with polyphenols from green tea correlates with improved glutathione balance and increased expression of 67LR, a receptor for (-)-epigallocatechin gallate. *Biofactors* 35: 279-294, 2009.
90. Dröge W. Free radicals in the physiological control of cell function. *Physiol Rev* 82: 47-95, 2002.
91. Drowley L, Okada M, Beckman S, Vella J, Keller B, Tobita K, and Huard J. Cellular antioxidant levels influence muscle stem cell therapy. *Mol Ther* 18: 1865-1873, 2010.
92. Dumont NA, Wang YX, and Rudnicki MA. Intrinsic and extrinsic mechanisms regulating satellite cell function. *Development* 142: 1572-1581, 2015.
93. Durbeej M, Campbell KP. Muscular dystrophies involving the dystrophin-glycoprotein complex: an overview of current mouse models. *Curr Opin Genet Dev* 12: 349-61, 2002.
94. Durham WJ, Aracena-Parks P, Long C, Rossi AE, Goonasekera SA, Boncompagni S, Galvan DL, Gilman CP, Baker MR, Shirokova N, Protasi F, Dirksen R, and Hamilton SL. RyR1 S-nitrosylation underlies environmental heat stroke and sudden death in Y522S RyR1 knockin mice. *Cell* 133: 53-65, 2008.

95. Dworakowski R, Alom-Ruiz SP, and Shah AM. NADPH oxidase-derived reactive oxygen species in the regulation of endothelial phenotype. *Pharmacol rep* 60: 21-28, 2008.
96. El Haddad M, Jean E, Turki A, Hugon G, Vernus B, Bonnieu A, Passerieux E, Hamade A, Mercier J, Laoudj-Chenivresse D, and Carnac G. Glutathione peroxidase 3, a new retinoid target gene, is crucial for human skeletal muscle precursor cell survival. *J Cell Sci* 125: 6147-6156, 2012.
97. el-Dwairi Q, Comtois A, Guo Y, and Hussain SN. Endotoxin-induced skeletal muscle contractile dysfunction: contribution of nitric oxide synthases. *Am J Physiol* 274: C770-779, 1998.
98. Ermakova YG, Bilan DS, Matlashov ME, Mishina NM, Markvicheva KN, Subach OM, Subach FV, Bogeski I, Hoth M, Enikolopov G, and Belousov VV. Red fluorescent genetically encoded indicator for intracellular hydrogen peroxide. *Nat Commun* 5: 5222, 2014.
99. Fatouros IG, Chatzinikolaou A, Douroudos, II, Nikolaidis MG, Kyparos A, Margonis K, Michailidis Y, Vantarakis A, Taxildaris K, Katrabasas I, Mandalidis D, Kouretas D, and Jamurtas AZ. Time-course of changes in oxidative stress and antioxidant status responses following a soccer game. *J Strength Cond Res* 24: 3278-3286, 2010.
100. Filippin LI, Cuevas MJ, Lima E, Marroni NP, Gonzalez-Gallego J, and Xavier RM. Nitric oxide regulates the repair of injured skeletal muscle. *Nitric Oxide* 24: 43-49, 2011.
101. Filippin LI, Moreira AJ, Marroni NP, and Xavier RM. Nitric oxide and repair of skeletal muscle injury. *Nitric Oxide* 21: 157-163, 2009.
102. Finaud J, Lac G, and Filaire E. Oxidative stress : relationship with exercise and training. *Sports Med* 36: 327-358, 2006.
103. Finkel T. Signal transduction by reactive oxygen species. *J Cell Biol* 194: 7-15, 2011.
104. Forman HJ, Augusto O, Brigelius-Flohe R, Dennery PA, Kalyanaraman B, Ischiropoulos H, Mann GE, Radi R, Roberts LJ, Vina J, and Davies KJA. Even free radicals should follow some rules: a guide to free radical research terminology and methodology. *Free Radic Biol Med* 78: 233-235, 2015.
105. Forman HJ, Maiorino M, and Ursini F. Signaling functions of reactive oxygen species. *Biochemistry* 49: 835-842, 2010.
106. Forman HJ, Torres M. Redox signaling in macrophages. *Mol Aspects Med* 22: 189-216, 2001.
107. Forman HJ, Zhang H, and Rinna A. Glutathione: overview of its protective roles, measurement, and biosynthesis. *Mol Aspects Med* 30: 1-12, 2009.
108. Forrester MT, Stamler JS. A classification scheme for redox-based modifications of proteins. *Am J Respir Cell Mol Bio* 36: 135-137, 2007.
109. Franco AA, Odom RS, and Rando TA. Regulation of antioxidant enzyme gene expression in response to oxidative stress and during differentiation of mouse skeletal muscle. *Free Radic Biol Med* 27: 1122-1132, 1999.
110. Frederick DW, Loro E, Liu L, Davila A, Jr., Chellappa K, Silverman IM, Quinn WJ, 3rd, Gosai SJ, Tichy ED, Davis JG, Mourkioti F, Gregory BD, Dellinger RW, Redpath P, Migaud ME, Nakamaru-Ogiso E, Rabinowitz JD, Khurana TS, and Baur JA. Loss of NAD Homeostasis Leads to Progressive and Reversible Degeneration of Skeletal Muscle. *Cell Metab* 24: 269-82, 2016.

111. Frey RS, Ushio-Fukai M, and Malik AB. NADPH oxidase-dependent signaling in endothelial cells: role in physiology and pathophysiology. *Antioxid Redox Signal* 11: 791-810, 2009.
112. Fridén J, Seger J, Sjöström M, and Ekblom B. Adaptive response in human skeletal muscle subjected to prolonged eccentric training. *Int J Sports Med* 4: 177-183, 1983.
113. Fulco M, Schiltz RL, Iezzi S, King MT, Zhao P, Kashiwaya Y, Hoffman E, Veech RL, and Sartorelli V. Sir2 regulates skeletal muscle differentiation as a potential sensor of the redox state. *Mol Cell* 12: 51-62, 2003.
114. Fulle S, Belia S, and Di Tano G. Sarcopenia is more than a muscular deficit. *Arch Ital Biol* 143: 229-34, 2005.
115. Funato Y, Michiue T, Asashima M, and Miki H. The thioredoxin-related redox-regulating protein nucleoredoxin inhibits Wnt-beta-catenin signalling through dishevelled. *Nat Cell Biol* 8: 501-508, 2006.
116. Gaetke LM, Chow CK. Copper toxicity, oxidative stress, and antioxidant nutrients. *Toxicology* 189: 147-163, 2003.
117. García-Prat L, Martínez-Vicente M, Perdiguero E, Ortet L, Rodríguez-Ubreva J, Rebollo E, Ruiz-Bonilla V, Gutarra S, Ballestar E, Serrano AL, Sandri M, and Muñoz-Cánoves P. Autophagy maintains stemness by preventing senescence. *Nature* 529: 37-42, 2016.
118. Gati I, Danielsson O, Betmark T, Ernerudh J, Ollinger K, and Dizdar N. Effects of inhibitors of the arachidonic acid cascade on primary muscle culture from a Duchenne muscular dystrophy patient. *Prostaglandins Leukot Essent Fatty Acids* 77: 217-223, 2007.
119. Germinario E, Peron S, Toniolo L, Betto R, Cencetti F, Donati C, Bruni P, and Danieli-Betto D. S1P2 receptor promotes mouse skeletal muscle regeneration. *J Appl Physiol* 113: 707-713, 2012.
120. Godoy JR, Funke M, Ackermann W, Haunhorst P, Oesteritz S, Capani F, Elsässer H-P, and Lillig CH. Redox atlas of the mouse. Immunohistochemical detection of glutaredoxin-, peroxiredoxin-, and thioredoxin-family proteins in various tissues of the laboratory mouse. *Biochim Biophys Acta* 1810: 2-92, 2011.
121. Gomez-Cabrera MC, Domenech E, Romagnoli M, Arduini A, Borrás C, Pallardo FV, Sastre J, and Vina J. Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance. *Am J Clin Nutr* 87: 142-9, 2008.
122. Gomez-Cabrera MC, Salvador-Pascual A, Cabo H, Ferrando B, and Vina J. Redox modulation of mitochondriogenesis in exercise. Does antioxidant supplementation blunt the benefits of exercise training? *Free Radic Biol Med* 86: 37-46, 2015.
123. Goncalves RL, Quinlan CL, Pervoshchikova IV, Hey-Mogensen M, and Brand MD. Sites of superoxide and hydrogen peroxide production by muscle mitochondria assessed ex vivo under conditions mimicking rest and exercise. *J Biol Chem* 290: 209-27, 2015.
124. Gong MC, Arbogast S, Guo Z, Mathenia J, Su W, and Reid MB. Calcium-independent phospholipase A2 modulates cytosolic oxidant activity and contractile function in murine skeletal muscle cells. *J Appl Physiol* 100: 399-406, 2006.
125. Griending KK, Sorescu D, and Ushio-Fukai M. NAD(P)H oxidase: role in cardiovascular biology and disease. *Circ Res* 86: 494-501, 2000.

126. Griffith OW Meister A. Glutathione: interorgan translocation, turnover, and metabolism. *Proc Nat Acad Sci USA* 76: 5606-5610, 1979.
127. Groitl BJ Jakob U. Thiol-based redox switches. *Biochim Biophys Acta* 1844: 1335-1343, 2014.
128. Halliwell B. Free radicals and antioxidants: a personal view. *Nutr Rev* 52: 253-265, 1994.
129. Halliwell B Gutteridge J. *Free Radicals in Biology and Medicine*: Oxford University Press; 2007. 888 p.
130. Hanschmann E-M, Godoy JR, Berndt C, Hudemann C, and Lillig CH. Thioredoxins, glutaredoxins, and peroxiredoxins--molecular mechanisms and health significance: from cofactors to antioxidants to redox signaling. *Antioxid Redox Signal* 19: 1539-1605, 2013.
131. Harman D. Aging: a theory based on free radical and radiation chemistry. *J Gerontol* 11: 298-300, 1956.
132. Haycock JW, MacNeil S, Jones P, Harris JB, and Mantle D. Oxidative damage to muscle protein in Duchenne muscular dystrophy. *Neuroreport* 8: 357-61, 1996.
133. Hayes JDDinkova-Kostova AT. The Nrf2 regulatory network provides an interface between redox and intermediary metabolism. *Trends Biochem Sci* 39: 199-218, 2014.
134. He C, Ryan AJ, Murthy S, and Carter AB. Accelerated development of pulmonary fibrosis via Cu,Zn-superoxide dismutase-induced alternative activation of macrophages. *J Biol Chem* 288: 20745-20757, 2013.
135. Henriksen EJ. Effects of H₂O₂ on insulin signaling the glucose transport system in mammalian skeletal muscle. *Method Enzymol* 528: 269-278, 2013.
136. Herbette S, Roeckel-Drevet P, and Drevet JR. Seleno-independent glutathione peroxidases. More than simple antioxidant scavengers. *FEBS J* 274: 2163-2180, 2007.
137. Hess DT, Matsumoto A, Kim S-O, Marshall HE, and Stamler JS. Protein S-nitrosylation: purview and parameters. *Nat Rev Mol Cell Biol* 6: 150-166, 2005.
138. Hollander J, Fiebig R, Gore M, Ookawara T, Ohno H, and Ji LL. Superoxide dismutase gene expression is activated by a single bout of exercise in rat skeletal muscle. *Pflügers Archiv* 442: 426-434, 2001.
139. Hollinger K, Shanely RA, Quindry JC, and Selsby JT. Long-term quercetin dietary enrichment decreases muscle injury in mdx mice. *Clin Nutr* 34: 515-22, 2015.
140. Hordijk PL. Regulation of NADPH oxidases: the role of Rac proteins. *Circ Res* 98: 453-62, 2006.
141. Howard AC, McNeil AK, and McNeil PL. Promotion of plasma membrane repair by vitamin E. *Nat Commun* 2: 597, 2011.
142. Huang YC, Dennis RG, and Baar K. Cultured slow vs. fast skeletal muscle cells differ in physiology and responsiveness to stimulation. *Am J Physiol Cell Physiol* 291: C11-7, 2006.
143. Hyldahl RD Hubal MJ. Lengthening our perspective: morphological, cellular, and molecular responses to eccentric exercise. *Muscle Nerve* 49: 155-170, 2014.
144. Imlay JA. Pathways of oxidative damage. *Ann Rev Microbiol* 57: 395-418, 2003.

145. Irrcher I, Ljubcic V, and Hood DA. Interactions between ROS and AMP kinase activity in the regulation of PGC-1 α transcription in skeletal muscle cells. *Am J Physiol. Cell Physiol* 296: C116-123, 2009.
146. Ito H, Ueda H, Iwamoto I, Inaguma Y, Takizawa T, Asano T, and Kato K. Nordihydroguaiaretic acid (NDGA) blocks the differentiation of C2C12 myoblast cells. *J Cell Physiol* 202: 874-879, 2005.
147. Ito N, Ruegg UT, Kudo A, Miyagoe-Suzuki Y, and Takeda Si. Activation of calcium signaling through Trpv1 by nNOS and peroxynitrite as a key trigger of skeletal muscle hypertrophy. *Nat Med* 19: 101-106, 2013.
148. Jackson MJ. Control of reactive oxygen species production in contracting skeletal muscle. *Antioxid Redox Signal* 15: 2477-2486, 2011.
149. Jackson MJ, McArdle A. Age-related changes in skeletal muscle reactive oxygen species generation and adaptive responses to reactive oxygen species. *J Physiol* 589: 2139-45, 2011.
150. Jain IH, Zazzeron L, Goli R, Alexa K, Schatzman-Bone S, Dhillon H, Goldberger O, Peng J, Shalem O, Sanjana NE, Zhang F, Goessling W, Zapol WM, and Mootha VK. Hypoxia as a therapy for mitochondrial disease. *Science* 352: 54-61, 2016.
151. James AM, Collins Y, Logan A, and Murphy MP. Mitochondrial oxidative stress and the metabolic syndrome. *Trends Endocrinol Metab* 23: 429-434, 2012.
152. Janssen-Heininger YMW, Mossman BT, Heintz NH, Forman HJ, Kalyanaraman B, Finkel T, Stamler JS, Rhee SG, and van der Vliet A. Redox-based regulation of signal transduction: principles, pitfalls, and promises. *Free Radic Biol Med* 45: 1-17, 2008.
153. Jean E, Laoudj-Chenivresse D, Notarnicola C, Rouger K, Serratrice N, Bonniou A, Gay S, Bacou F, Duret C, and Carnac G. Aldehyde dehydrogenase activity promotes survival of human muscle precursor cells. *J Cell Mol Med* 15: 119-133, 2011.
154. Ji LL. Modulation of skeletal muscle antioxidant defense by exercise: Role of redox signaling. *Free Radic Biol Med* 44: 142-152, 2008.
155. Ji LL, Dillon D, and Wu E. Alteration of antioxidant enzymes with aging in rat skeletal muscle and liver. *Am J Physiol* 258: R918-23, 1990.
156. Ji LL, Kang C, and Zhang Y. Exercise-induced hormesis and skeletal muscle health. *Free Radic Biol Med*: pii: S0891-5849(16)00076-9, 2016.
157. Joe AWB, Yi L, Natarajan A, Le Grand F, So L, Wang J, Rudnicki MA, and Rossi FMV. Muscle injury activates resident fibro/adipogenic progenitors that facilitate myogenesis. *Nat Cell Biol* 12: 153-163, 2010.
158. Jones DP. Redefining oxidative stress. *Antioxid Redox Signal* 8: 1865-1879, 2006.
159. Jones DP. Radical-free biology of oxidative stress. *Am J Physiol. Cell Physiol* 295: C849-868, 2008.
160. Kaminski J, Lancon A, Aires V, Limagne E, Tili E, Michaille JJ, and Latruffe N. Resveratrol initiates differentiation of mouse skeletal muscle-derived C2C12 myoblasts. *Biochem Pharmacol* 84: 1251-1259, 2012.
161. Kang C, O'Moore KM, Dickman JR, and Ji LL. Exercise activation of muscle peroxisome proliferator-activated receptor-gamma coactivator-1 α signaling is redox sensitive. *Free Radic Biol Med* 47: 1394-1400, 2009.
162. Karisch R, Fernandez M, Taylor P, Virtanen C, St-Germain JR, Jin LL, Harris IS, Mori J, Mak TW, Senis YA, Ostman A, Moran MF, and Neel BG. Global

- proteomic assessment of the classical protein-tyrosine phosphatome and "Redoxome". *Cell* 146: 826-840, 2011.
163. Katz A. Modulation of glucose transport in skeletal muscle by reactive oxygen species. *J Appl Physiol* 102: 1671-1676, 2007.
 164. Keefe AC, Lawson JA, Flygare SD, Fox ZD, Colasanto MP, Mathew SJ, Yandell M, and Kardon G. Muscle stem cells contribute to myofibres in sedentary adult mice. *Nat Commun* 6: 7087, 2015.
 165. Kehrer JP. The Haber-Weiss reaction and mechanisms of toxicity. *Toxicology* 149: 43-50, 2000.
 166. Khor SC, Abdul Karim N, Ngah WZ, Yusof YA, and Makpol S. Vitamin E in sarcopenia: current evidences on its role in prevention and treatment. *Oxid Med Cell Longev* 2014: 914853, 2014.
 167. Kim J-H, Kwak H-B, Thompson LV, and Lawler JM. Contribution of oxidative stress to pathology in diaphragm and limb muscles with Duchenne muscular dystrophy. *J Muscle Res Cell Motil* 34: 1-13, 2013.
 168. Kim YM, Kim KE, Koh GY, Ho Y-S, and Lee K-J. Hydrogen peroxide produced by angiopoietin-1 mediates angiogenesis. *Cancer Res* 66: 6167-6174, 2006.
 169. Kipp A, Banning A, van Schothorst EM, Méplan C, Schomburg L, Evelo C, Coort S, Gaj S, Keijer J, Hesketh J, and Brigelius-Flohé R. Four selenoproteins, protein biosynthesis, and Wnt signalling are particularly sensitive to limited selenium intake in mouse colon. *Mol Nutr Food Res* 53: 1561-1572, 2009.
 170. Kostallari E, Baba-Amer Y, Alonso-Martin S, Ngho P, Relaix F, Lafuste P, and Gherardi RK. Pericytes in the myovascular niche promote post-natal myofiber growth and satellite cell quiescence. *Development* 142: 1242-53, 2015.
 171. Koulmann N, Richard-Bulteau H, Crassous B, Serrurier B, Padeloup M, Bigard X, and Banzet S. Physical exercise during muscle regeneration improves recovery of the slow/oxidative phenotype. *Muscle Nerve*: doi: 10.1002/mus.25151, 2016.
 172. Kozakowska M, Ciesla M, Stefanska A, Skrzypek K, Was H, Jazwa A, Grochot-Przeczek A, Kotlinowski J, Szymula A, Bartelik A, Mazan M, Yagensky O, Florczyk U, Lemke K, Zebzda A, Dyduch G, Nowak W, Szade K, Stepniewski J, Majka M, Derlacz R, Loboda A, Dulak J, and Jozkowicz A. Heme oxygenase-1 inhibits myoblast differentiation by targeting myomirs. *Antioxid Redox Signal* 16: 113-27, 2012.
 173. Kozakowska M, Pietraszek-Gremplewicz K, Jozkowicz A, and Dulak J. The role of oxidative stress in skeletal muscle injury and regeneration: focus on antioxidant enzymes. *J Muscle Res Cell Motil* 36: 377-393, 2015.
 174. Kruger MJ, Myburgh KH, and Smith C. Contusion injury with chronic in vivo polyphenol supplementation: leukocyte responses. *Med Sci Sports Exerc* 46: 225-231, 2014.
 175. L'Honoré A, Commère P-H, Ouimette J-F, Montarras D, Drouin J, and Buckingham M. Redox regulation by Pitx2 and Pitx3 is critical for fetal myogenesis. *Dev Cell* 29: 392-405, 2014.
 176. Labazi M, McNeil AK, Kurtz T, Lee TC, Pegg RB, Angeli JPF, Conrad M, and McNeil PL. The antioxidant requirement for plasma membrane repair in skeletal muscle. *Free Radic Biol Med* 84: 246-253, 2015.
 177. Latroche C, Gitiaux C, Chrétien F, Desguerre I, Mounier R, and Chazaud B. Skeletal muscle microvasculature: A highly dynamic lifeline. *Physiology* 30: 417-427, 2015.

178. Lawrence TNatoli G. Transcriptional regulation of macrophage polarization: enabling diversity with identity. *Nat Rev Immunol* 11: 750-761, 2011.
179. Le Grand F, Jones AE, Seale V, Scimè A, and Rudnicki MA. Wnt7a activates the planar cell polarity pathway to drive the symmetric expansion of satellite stem cells. *Cell Stem Cell* 4: 535-547, 2009.
180. Lee J, Goldfarb AH, Rescino MH, Hegde S, Patrick S, and Apperson K. Eccentric exercise effect on blood oxidative-stress markers and delayed onset of muscle soreness. *Med Sci Sport Exerc* 34: 443-448, 2002.
181. Lee KH, Baek MY, Moon KY, Song WK, Chung CH, Ha DB, and Kang MS. Nitric oxide as a messenger molecule for myoblast fusion. *J Biol Chem* 269: 14371-14374, 1994.
182. Lee KH, Kim DG, Shin NY, Song WK, Kwon H, Chung CH, and Kang MS. NF-kappaB-dependent expression of nitric oxide synthase is required for membrane fusion of chick embryonic myoblasts. *Biochem J* 324: 237-242, 1997.
183. Lee S, Shin HS, Shireman PK, Vasilaki A, Van Remmen H, and Csete ME. Glutathione-peroxidase-1 null muscle progenitor cells are globally defective. *Free Radic Biol Med* 41: 1174-1184, 2006.
184. Lee S, Tak E, Lee J, Rashid MA, Murphy MP, Ha J, and Kim SS. Mitochondrial H₂O₂ generated from electron transport chain complex I stimulates muscle differentiation. *Cell Res* 21: 817-834, 2011.
185. Leeuwenburgh C, Hollander J, Leichtweis S, Griffiths M, Gore M, and Ji LL. Adaptations of glutathione antioxidant system to endurance training are tissue and muscle fiber specific. *Am J Physiol* 272: R363-369, 1997.
186. Leichert LI, Gehrke F, Gudiseva HV, Blackwell T, Ilbert M, Walker AK, Strahler JR, Andrews PC, and Jakob U. Quantifying changes in the thiol redox proteome upon oxidative stress in vivo. *Proc Natl Acad Sci U S A* 105: 8197-202, 2008.
187. Lemos DR, Babaeijandaghi F, Low M, Chang CK, Lee ST, Fiore D, Zhang RH, Natarajan A, Nedospasov SA, and Rossi FM. Nilotinib reduces muscle fibrosis in chronic muscle injury by promoting TNF-mediated apoptosis of fibro/adipogenic progenitors. *Nature Medicine*: 10, 2015.
188. Lepper C, Partridge TA, and Fan CM. An absolute requirement for Pax7-positive satellite cells in acute injury-induced skeletal muscle regeneration. *Development* 138: 3639-3646, 2011.
189. Li X, Cobb CE, Hill KE, Burk RF, and May JM. Mitochondrial uptake and recycling of ascorbic acid. *Arch Biochem Biophys* 387: 143-153, 2001.
190. Lim S, Shin JY, Jo A, Jyothi KR, Nguyen MN, Choi TG, Kim J, Park JH, Eun YG, Yoon KS, Ha J, and Kim SS. Carbonyl reductase 1 is an essential regulator of skeletal muscle differentiation and regeneration. *Int J Biochem Cell Biol* 45: 1784-1793, 2013.
191. Lin J, Wu H, Tarr PT, Zhang C-Y, Wu Z, Boss O, Michael LF, Puigserver P, Isotani E, Olson EN, Lowell BB, Bassel-Duby R, and Spiegelman BM. Transcriptional co-activator PGC-1 alpha drives the formation of slow-twitch muscle fibres. *Nature* 418: 797-801, 2002.
192. Liu H, Colavitti R, Rovira, II, and Finkel T. Redox-dependent transcriptional regulation. *Circ Res* 97: 967-74, 2005.
193. Lo Verso F, Carnio S, Vainshtein A, and Sandri M. Autophagy is not required to sustain exercise and PRKAA1/AMPK activity but is important to prevent mitochondrial damage during physical activity. *Autophagy* 10: 1883-94, 2014.

194. Lodhi IJSemenkovich CF. Peroxisomes: a nexus for lipid metabolism and cellular signaling. *Cell Metab* 19: 380-92, 2014.
195. Long JH, Lira VA, Soltow QA, Betters JL, Sellman JE, and Criswell DS. Arginine supplementation induces myoblast fusion via augmentation of nitric oxide production. *J Muscle Res Cell Motil* 27: 577-584, 2006.
196. Lu Z, Xu X, Hu X, Fassett J, Zhu G, Tao Y, Li J, Huang Y, Zhang P, Zhao B, and Chen Y. PGC-1 alpha regulates expression of myocardial mitochondrial antioxidants and myocardial oxidative stress after chronic systolic overload. *Antioxid Redox Signal* 13: 1011-1022, 2010.
197. Lukyanov KABELousov VV. Genetically encoded fluorescent redox sensors. *Biochim Biophys Acta* 1840: 745-756, 2014.
198. Luo SW, Zhang C, Zhang B, Kim CH, Qiu YZ, Du QS, Mei L, and Xiong WC. Regulation of heterochromatin remodelling and myogenin expression during muscle differentiation by FAK interaction with MBD2. *EMBO J* 28: 2568-82, 2009.
199. Malhotra JDKaufman RJ. Endoplasmic reticulum stress and oxidative stress: a vicious cycle or a double-edged sword? *Antioxid Redox Signal* 9: 2277-2293, 2007.
200. Malinska D, Kudin AP, Bejtka M, and Kunz WS. Changes in mitochondrial reactive oxygen species synthesis during differentiation of skeletal muscle cells. *Mitochondrion* 12: 144-148, 2012.
201. Malm C, Svensson M, Ekblom B, and Sjodin B. Effects of ubiquinone-10 supplementation and high intensity training on physical performance in humans. *Acta Physiol Scand* 161: 379-84, 1997.
202. Malm C, Svensson M, Sjoberg B, Ekblom B, and Sjodin B. Supplementation with ubiquinone-10 causes cellular damage during intense exercise. *Acta Physiol Scand* 157: 511-2, 1996.
203. Manabe Y, Takagi M, Nakamura-Yamada M, Goto-Inoue N, Taoka M, Isobe T, and Fujii NL. Redox proteins are constitutively secreted by skeletal muscle. *J Physiol Sci* 64: 401-409, 2014.
204. Mandl J, Szarka A, and Bánhegyi G. Vitamin C: update on physiology and pharmacology. *Br J Pharmacol* 157: 1097-1110, 2009.
205. Manzano R, Toivonen JM, Calvo AC, Oliván S, Zaragoza P, Rodellar C, Montarras D, and Osta R. Altered in vitro proliferation of mouse SOD1-G93A skeletal muscle satellite cells. *Neurodegener Dis* 11: 153-164, 2013.
206. Mason SA, Morrison D, McConell GK, and Wadley GD. Muscle redox signalling pathways in exercise. Role of antioxidants. *Free Radic Biol Med*: pii: S0891-5849(16)00073-3, 2016.
207. Matsushima Y, Nanri H, Nara S, Okufuji T, Ohta M, Hachisuka K, and Ikeda M. Hindlimb unloading decreases thioredoxin-related antioxidant proteins and increases thioredoxin-binding protein-2 in rat skeletal muscle. *Free Radic Res* 40: 715-722, 2006.
208. May JM, Cobb CE, Mendiratta S, Hill KE, and Burk RF. Reduction of the ascorbyl free radical to ascorbate by thioredoxin reductase. *J Biol Chem* 273: 23039-23045, 1998.
209. McArdle A, Vasilaki A, and Jackson M. Exercise and skeletal muscle ageing: cellular and molecular mechanisms. *Ageing Res Rev* 1: 79-93, 2002.
210. McCord JMFridovich I. Superoxide dismutases: you've come a long way, baby. *Antioxid Redox Signal* 20: 1548-1549, 2014.

211. McDonagh B, Sakellariou GK, Smith NT, Brownridge P, and Jackson MJ. Differential cysteine labeling and global label-free proteomics reveals an altered metabolic state in skeletal muscle aging. *J Proteome Res* 13: 5008-5021, 2014.
212. McGinley C, Shafat A, and Donnelly AE. Does antioxidant vitamin supplementation protect against muscle damage? *Sports Med* 39: 1011-1032, 2009.
213. Menger KE, James AM, Cocheme HM, Harbour ME, Chouchani ET, Ding S, Fearnley IM, Partridge L, and Murphy MP. Fasting, but Not Aging, Dramatically Alters the Redox Status of Cysteine Residues on Proteins in *Drosophila melanogaster*. *Cell Rep* 11: 1856-65, 2015.
214. Mercuri EMuntoni F. Muscular dystrophies. *Lancet* 381: 845-860, 2013.
215. Messina S, Altavilla D, Aguenouz Mh, Seminara P, Minutoli L, Monici MC, Bitto A, Mazzeo A, Marini H, Squadrito F, and Vita G. Lipid peroxidation inhibition blunts nuclear factor-kappaB activation, reduces skeletal muscle degeneration, and enhances muscle function in mdx mice. *Am J Pathol* 168: 918-926, 2006.
216. Messina S, Bitto A, Aguenouz M, Mazzeo A, Migliorato A, Polito F, Irrera N, Altavilla D, Vita GL, Russo M, Naro A, De Pasquale MG, Rizzuto E, Musaro A, Squadrito F, and Vita G. Flavocoxid counteracts muscle necrosis and improves functional properties in mdx mice: a comparison study with methylprednisolone. *Exp Neurol* 220: 349-358, 2009.
217. Michaelson LP, Shi G, Ward CW, and Rodney GG. Mitochondrial redox potential during contraction in single intact muscle fibers. *Muscle Nerve* 42: 522-529, 2010.
218. Michailidis Y, Karagounis LG, Terzis G, Jamurtas AZ, Spengos K, Tsoukas D, Chatzinikolaou A, Mandalidis D, Stefanetti RJ, Papassotiriou I, Athanasopoulos S, Hawley JA, Russell AP, and Fatouros IG. Thiol-based antioxidant supplementation alters human skeletal muscle signaling and attenuates its inflammatory response and recovery after intense eccentric exercise. *Am J Clin Nutr* 98: 233-245, 2013.
219. Miller CJ, Gounder SS, Kannan S, Goutam K, Muthusamy VR, Firpo MA, Symons JD, Paine R, 3rd, Hoidal JR, and Rajasekaran NS. Disruption of Nrf2/ARE signaling impairs antioxidant mechanisms and promotes cell degradation pathways in aged skeletal muscle. *Biochim Biophys Acta* 1822: 1038-1050, 2012.
220. Minet ADGaster M. Cultured senescent myoblasts derived from human vastus lateralis exhibit normal mitochondrial ATP synthesis capacities with correlating concomitant ROS production while whole cell ATP production is decreased. *Biogerontol* 13: 277-285, 2012.
221. Miquel J, Economos AC, Fleming J, and Johnson JE, Jr. Mitochondrial role in cell aging. *Exp Gerontol* 15: 575-591, 1980.
222. Mittal M, Siddiqui MR, Tran K, Reddy SP, and Malik AB. Reactive oxygen species in inflammation and tissue injury. *Antioxid Redox Signal* 20: 1126-67, 2014.
223. Mofarrahi M, Brandes RP, Gorlach A, Hanze J, Terada LS, Quinn MT, Mayaki D, Petrof B, and Hussain SNA. Regulation of proliferation of skeletal muscle precursor cells by NADPH oxidase. *Antioxid Redox Signal* 10: 559-574, 2008.

224. Montesano A, Luzi L, Senesi P, Mazzocchi N, and Terruzzi I. Resveratrol promotes myogenesis and hypertrophy in murine myoblasts. *J Transl Med* 11: 310, 2013.
225. Morrison D, Hughes J, Della Gatta PA, Mason S, Lamon S, Russell AP, and Wadley GD. Vitamin C and E supplementation prevents some of the cellular adaptations to endurance-training in humans. *Free Radic Biol Med* 89: 852-62, 2015.
226. Mounier R, Chrétien F, and Chazaud B. Blood vessels and the satellite cell niche. *Curr Top Dev Biol* 96: 121-138, 2011.
227. Mounier R, Théret M, Arnold L, Cuvellier S, Bultot L, Göransson O, Sanz N, Ferry A, Sakamoto K, Foretz M, Viollet B, and Chazaud B. AMPK α 1 regulates macrophage skewing at the time of resolution of inflammation during skeletal muscle regeneration. *Cell Metab* 18: 251-264, 2013.
228. Murphy MEKehrer JP. Free radicals: a potential pathogenic mechanism in inherited muscular dystrophy. *Life Sci* 39: 2271-2278, 1986.
229. Murphy MP. How mitochondria produce reactive oxygen species. *Biochem J* 417: 1-13, 2009.
230. Murphy MP, Holmgren A, Larsson N-G, Halliwell B, Chang CJ, Kalyanaraman B, Rhee SG, Thornalley PJ, Partridge L, Gems D, Nyström T, Belousov V, Schumacker PT, and Winterbourn CC. Unraveling the biological roles of reactive oxygen species. *Cell Metab* 13: 361-366, 2011.
231. Myburgh KH. Polyphenol supplementation: benefits for exercise performance or oxidative stress? *Sports Med* 44: S57-70, 2014.
232. Myburgh KH, Kruger MJ, and Smith C. Accelerated skeletal muscle recovery after in vivo polyphenol administration. *J Nutr Biochem* 23: 1072-1079, 2012.
233. Nagata Y, Partridge TA, Matsuda R, and Zammit PS. Entry of muscle satellite cells into the cell cycle requires sphingolipid signaling. *J Cell Biol* 174: 245-253, 2006.
234. Narasimhan M, Hong J, Atieno N, Muthusamy VR, Davidson CJ, Abu-Rmaileh N, Richardson RS, Gomes AV, Hoidal JR, and Rajasekaran NS. Nrf2 deficiency promotes apoptosis and impairs PAX7/MyoD expression in aging skeletal muscle cells. *Free Radic Biol Med* 71: 402-414, 2014.
235. Nathan CCunningham-Bussell A. Beyond oxidative stress: an immunologist's guide to reactive oxygen species. *Nat Rev Immunol* 13: 349-361, 2013.
236. Nethery D, Stofan D, Callahan L, DiMarco A, and Supinski G. Formation of reactive oxygen species by the contracting diaphragm is PLA(2) dependent. *J Appl Physiol* 87: 792-800, 1999.
237. Niethammer P, Grabher C, Look AT, and Mitchison TJ. A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature* 459: 996-999, 2009.
238. Niethammer P, Grabher C, Look AT, and Mitchison TJ. A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature* 459: 996-9, 2009.
239. Nikolaidis MG, Jamurtas AZ, Paschalis V, Fatouros IG, Koutedakis Y, and Kouretas D. The effect of muscle-damaging exercise on blood and skeletal muscle oxidative stress: magnitude and time-course considerations. *Sports Med* 38: 579-606, 2008.
240. Nikolaidis MG, Paschalis V, Giakas G, Fatouros IG, Koutedakis Y, Kouretas D, and Jamurtas AZ. Decreased blood oxidative stress after repeated muscle-damaging exercise. *Med Sci Sports Exerc* 39: 1080-1089, 2007.

241. Nikolova-Karakashian MN, Reid MB. Sphingolipid metabolism, oxidant signaling, and contractile function of skeletal muscle. *Antioxid Redox Signal* 15: 2501-2517, 2011.
242. Nordberg J, Arnér ES. Reactive oxygen species, antioxidants, and the mammalian thioredoxin system. *Free Radic Biol Med* 31: 1287-1312, 2001.
243. Oh J, Lee YD, and Wagers AJ. Stem cell aging: mechanisms, regulators and therapeutic opportunities. *Nat Med* 20: 870-880, 2014.
244. Ohashi K, Nagata Y, Wada E, Zammit PS, Shiozuka M, and Matsuda R. Zinc promotes proliferation and activation of myogenic cells via the PI3K/Akt and ERK signaling cascade. *Exp Cell Res* 333: 228-237, 2015.
245. Oshikawa J, Kim S-J, Furuta E, Caliceti C, Chen G-F, McKinney RD, Kuhr F, Levitan I, Fukai T, and Ushio-Fukai M. Novel role of p66Shc in ROS-dependent VEGF signaling and angiogenesis in endothelial cells. *Am J Physiol. Heart Circ Physiol* 302: H724-732, 2012.
246. Oteiza PI. Zinc and the modulation of redox homeostasis. *Free Radic Biol Med* 53: 1748-1759, 2012.
247. Ott M, Gogvadze V, Orrenius S, and Zhivotovsky B. Mitochondria, oxidative stress and cell death. *Apoptosis* 12: 913-22, 2007.
248. Otto A, Schmidt C, Luke G, Allen S, Valasek P, Muntoni F, Lawrence-Watt D, and Patel K. Canonical Wnt signalling induces satellite-cell proliferation during adult skeletal muscle regeneration. *J Cell Sci* 121: 2939-2950, 2008.
249. Pal R, Palmieri M, Loehr JA, Li S, Abo-Zahrah R, Monroe TO, Thakur PB, Sardiello M, and Rodney GG. Src-dependent impairment of autophagy by oxidative stress in a mouse model of Duchenne muscular dystrophy. *Nat Commun* 5: 4425, 2014.
250. Pallafacchina G, François S, Regnault B, Czarny B, Dive V, Cumano A, Montarras D, and Buckingham M. An adult tissue-specific stem cell in its niche: a gene profiling analysis of in vivo quiescent and activated muscle satellite cells. *Stem Cell Res* 4: 77-91, 2010.
251. Passerieux E, Hayot M, Jausse A, Carnac G, Gouzi F, Pillard F, Picot M-C, Böcker K, Hugon G, Pincemail J, Defraigne JO, Verrips T, Mercier J, and Laoudj-Chenivresse D. Effects of vitamin C, vitamin E, zinc gluconate, and selenomethionine supplementation on muscle function and oxidative stress biomarkers in patients with facioscapulohumeral dystrophy: a double-blind randomized controlled clinical trial. *Free Radic Biol Med* 81: 158-169, 2015.
252. Paulsen G, Hamarsland H, Cumming KT, Johansen RE, Hulmi JJ, Borsheim E, Wiig H, Garthe I, and Raastad T. Vitamin C and E supplementation alters protein signalling after a strength training session, but not muscle growth during 10 weeks of training. *J Physiol* 592: 5391-408, 2014.
253. Paulsen G, Mikkelsen UR, Raastad T, and Peake JM. Leucocytes, cytokines and satellite cells: what role do they play in muscle damage and regeneration following eccentric exercise? *Exerc Immunol Rev* 18: 42-97, 2012.
254. Pchejetski D, Kunduzova O, Dayon A, Calise D, Seguelas MH, Leducq N, Seif I, Parini A, and Cuvillier O. Oxidative stress-dependent sphingosine kinase-1 inhibition mediates monoamine oxidase A-associated cardiac cell apoptosis. *Circ Res* 100: 41-49, 2007.
255. Pearce EL, Pearce EJ. Metabolic pathways in immune cell activation and quiescence. *Immunity* 38: 633-643, 2013.
256. Pedersen BK, Febbraio MA. Muscles, exercise and obesity: skeletal muscle as a secretory organ. *Nat Rev. Endocrinol* 8: 457-465, 2012.

257. Perdiguero E, Sousa-Victor P, Ruiz-Bonilla V, Jardí M, Caelles C, Serrano AL, and Muñoz-Cánoves P. p38/MKP-1-regulated AKT coordinates macrophage transitions and resolution of inflammation during tissue repair. *J Cell Biol* 195: 307-322, 2011.
258. Peternelj T, Coombes JS. Antioxidant supplementation during exercise training: beneficial or detrimental? *Sports Med* 41: 1043-69, 2011.
259. Piao YJ, Seo YH, Hong F, Kim JH, Kim YJ, Kang MH, Kim BS, Jo SA, Jo I, Jue DM, Kang I, Ha J, and Kim SS. Nox 2 stimulates muscle differentiation via NF-kappaB/iNOS pathway. *Free Radic Biol Med* 38: 989-1001, 2005.
260. Powers SK, Criswell D, Lawler J, Ji LL, Martin D, Herb RA, and Dudley G. Influence of exercise and fiber type on antioxidant enzyme activity in rat skeletal muscle. *Am J Physiol* 266: R375-380, 1994.
261. Powers SK, Jackson MJ. Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiol Rev* 88: 1243-1276, 2008.
262. Proske U, Allen TJ. Damage to skeletal muscle from eccentric exercise. *Exerc Sport Sci Rev* 33: 98-104, 2005.
263. Puri PL, Bhakta K, Wood LD, Costanzo A, Zhu J, and Wang JYJ. A myogenic differentiation checkpoint activated by genotoxic stress. *Nat Genet* 32: 585-593, 2002.
264. Qaisar R, Bhaskaran S, and Van Remmen H. Muscle fiber type diversification during exercise and regeneration. *Free Radic Biol Med*: pii: S0891-5849(16)30004-1, 2016.
265. Quadriatero J, Alway SE, and Dupont-Versteegden EE. Skeletal muscle apoptotic response to physical activity: potential mechanisms for protection. *Appl Physiol Nutr Metab* 36: 608-17, 2011.
266. Quinlan CL, Perevoschikova IV, Goncalves RL, Hey-Mogensen M, and Brand MD. The determination and analysis of site-specific rates of mitochondrial reactive oxygen species production. *Methods Enzymol* 526: 189-217, 2013.
267. Radak Z, Chung HY, Koltai E, Taylor AW, and Goto S. Exercise, oxidative stress and hormesis. *Ageing Res Rev* 7: 34-42, 2008.
268. Rahman M, Mofarrahi M, Kristof AS, Nkengfac B, Harel S, and Hussain SN. Reactive oxygen species regulation of autophagy in skeletal muscles. *Antioxid Redox Signal* 20: 443-59, 2014.
269. Rando TA, Disatnik MH, Yu Y, and Franco A. Muscle cells from mdx mice have an increased susceptibility to oxidative stress. *Neuromuscul Disord* 8: 14-21, 1998.
270. Ray PD, Huang B-W, and Tsuji Y. Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cell Signal* 24: 981-990, 2012.
271. Razzell W, Evans IR, Martin P, and Wood W. Calcium flashes orchestrate the wound inflammatory response through DUOX activation and hydrogen peroxide release. *Curr Biol* 23: 424-429, 2013.
272. Reczek CR, Chandel NS. ROS-dependent signal transduction. *Current Opinion in Cell Biology* 33: 8-13, 2015.
273. Reid MB. Invited Review: redox modulation of skeletal muscle contraction: what we know and what we don't. *J Appl Physiol* 90: 724-731, 2001.
274. Renjini R, Gayathri N, Nalini A, and Srinivas Bharath MM. Oxidative damage in muscular dystrophy correlates with the severity of the pathology: role of glutathione metabolism. *Neurochem Res* 37: 885-898, 2012.

275. Reynaert NL, van der Vliet A, Guala AS, McGovern T, Hristova M, Pantano C, Heintz NH, Heim J, Ho YS, Matthews DE, Wouters EF, and Janssen-Heininger YM. Dynamic redox control of NF-kappaB through glutaredoxin-regulated S-glutathionylation of inhibitory kappaB kinase beta. *Proc Natl Acad Sci U S A* 103: 13086-91, 2006.
276. Rhee SG. Cell signaling. H₂O₂, a necessary evil for cell signaling. *Science* 312: 1882-1883, 2006.
277. Rhee SG, Woo HA. Multiple functions of peroxiredoxins: peroxidases, sensors and regulators of the intracellular messenger H₂O₂, and protein chaperones. *Antioxid Redox Signal* 15: 781-794, 2011.
278. Rhoads RP, Johnson RM, Rathbone CR, Liu X, Temm-Grove C, Sheehan SM, Hoying JB, and Allen RE. Satellite cell-mediated angiogenesis in vitro coincides with a functional hypoxia-inducible factor pathway. *Am J Physiol Cell Physiol* 296: C1321-1328, 2009.
279. Richard-Bulteau H, Serrurier B, Crassous B, Banzet S, Peinnequin A, Bigard X, and Koulmann N. Recovery of skeletal muscle mass after extensive injury: positive effects of increased contractile activity. *Am J Physiol. Cell Physiol* 294: C467-476, 2008.
280. Rigamonti E, Touvier T, Clementi E, Manfredi AA, Brunelli S, and Rovere-Querini P. Requirement of inducible nitric oxide synthase for skeletal muscle regeneration after acute damage. *J Immunol* 190: 1767-1777, 2013.
281. Ristow M, Zarse K, Oberbach A, Klötting N, Birringer M, Kiehntopf M, Stumvoll M, Kahn CR, and Blüher M. Antioxidants prevent health-promoting effects of physical exercise in humans. *Proc Nat Acad Sci USA* 106: 8665-8670, 2009.
282. Rodney GG, Pal R, and Abo-Zahrah R. Redox regulation of autophagy in skeletal muscle. *Free Radic Biol Med* 98: 103-12, 2016.
283. Rohrbach S, Gruenler S, Teschner M, and Holtz J. The thioredoxin system in aging muscle: key role of mitochondrial thioredoxin reductase in the protective effects of caloric restriction? *Am J Physiol Regul Integr Comp Physiol* 291: R927-35, 2006.
284. Rosenblatt JD, Lunt AI, Parry DJ, and Partridge TA. Culturing satellite cells from living single muscle fiber explants. *In Vitro Cell Dev Biol Anim* 31: 773-9, 1995.
285. Ruffell D, Mourkioti F, Gambardella A, Kirstetter P, Lopez RG, Rosenthal N, and Nerlov C. A CREB-C/EBPbeta cascade induces M2 macrophage-specific gene expression and promotes muscle injury repair. *Proc Nat Acad Sci USA* 106: 17475-17480, 2009.
286. Ryall JG, Dell'Orso S, Derfoul A, Juan A, Zare H, Feng X, Clermont D, Koulonis M, Gutierrez-Cruz G, Fulco M, and Sartorelli V. The NAD(+)-dependent SIRT1 deacetylase translates a metabolic switch into regulatory epigenetics in skeletal muscle stem cells. *Cell Stem Cell* 16: 171-83, 2015.
287. Ryu D, Zhang H, Ropelle ER, Sorrentino V, Mazala DA, Mouchiroud L, Marshall PL, Campbell MD, Ali AS, Knowels GM, Bellemin S, Iyer SR, Wang X, Gariani K, Sauve AA, Canto C, Conley KE, Walter L, Lovering RM, Chin ER, Jasmin BJ, Marcinek DJ, Menzies KJ, and Auwerx J. NAD⁺ repletion improves muscle function in muscular dystrophy and counters global PARylation. *Sci Transl Med* 8: 361ra139, 2016.
288. Sachdev SD, Davies KJA. Production, detection, and adaptive responses to free radicals in exercise. *Free Radic Biol Med* 44: 215-223, 2008.

289. Saclier M, Cuvellier S, Magnan M, Mounier R, and Chazaud B. Monocyte/macrophage interactions with myogenic precursor cells during skeletal muscle regeneration. *FEBS J* 280: 4118-4130, 2013.
290. Saclier M, Yacoub-Youssef H, Mackey AL, Arnold L, Ardjoune H, Magnan M, Sailhan F, Chelly J, Pavlath GK, Mounier R, Kjaer M, and Chazaud B. Differentially activated macrophages orchestrate myogenic precursor cell fate during human skeletal muscle regeneration. *Stem Cells* 31: 384-396, 2013.
291. Sakellariou GK, Pye D, Vasilaki A, Zibrik L, Palomero J, Kabayo T, McArdle F, Van Remmen H, Richardson A, Tidball JG, McArdle A, and Jackson MJ. Role of superoxide-nitric oxide interactions in the accelerated age-related loss of muscle mass in mice lacking Cu,Zn superoxide dismutase. *Aging Cell* 10: 749-760, 2011.
292. Sambasivan R, Yao R, Kissenpfennig A, Van Wittenberghe L, Paldi A, Gayraud-Morel B, Guenou H, Malissen B, Tajbakhsh S, and Galy A. Pax7-expressing satellite cells are indispensable for adult skeletal muscle regeneration. *Development* 138: 3647-3656, 2011.
293. Sandiford SD, Kennedy KA, Xie X, Pickering JG, and Li SS. Dual oxidase maturation factor 1 (DUOXA1) overexpression increases reactive oxygen species production and inhibits murine muscle satellite cell differentiation. *Cell Commun Signal* 12: 5, 2014.
294. Sarbassov D, Sabatini DM. Redox regulation of the nutrient-sensitive rapTOR pathway and complex. *J Biol Chem* 280: 39505-39509, 2005.
295. Scherz-Shouval R, Shvets E, Fass E, Shorer H, Gil L, and Elazar Z. Reactive oxygen species are essential for autophagy and specifically regulate the activity of Atg4. *EMBO J* 26: 1749-1760, 2007.
296. Schmalbruch H, Hellhammer U. The number of nuclei in adult rat muscles with special reference to satellite cells. *Anat Rec* 189: 169-75, 1977.
297. Schreck RB, Baeuerle PA. A role for oxygen radicals as second messengers. *Trends Cell Biol* 1: 39-42, 1991.
298. Sciorati C, Miglietta D, Buono R, Pisa V, Cattaneo D, Azzoni E, Brunelli S, and Clementi E. A dual acting compound releasing nitric oxide (NO) and ibuprofen, NCX 320, shows significant therapeutic effects in a mouse model of muscular dystrophy. *Pharmacol Res* 64: 210-217, 2011.
299. Segard BD, Delort F, Bailleux V, Simon S, Leccia E, Gausseres B, Briki F, Vicart P, and Batonnet-Pichon S. N-acetyl-L-cysteine prevents stress-induced desmin aggregation in cellular models of desminopathy. *PLoS One* 8: e76361, 2013.
300. Sen CK, Packer L. Antioxidant and redox regulation of gene transcription. *FASEB J* 10: 709-20, 1996.
301. Sen CK, Rahkila P, and Hanninen O. Glutathione metabolism in skeletal muscle derived cells of the L6 line. *Acta Physiol Scand* 148: 21-6, 1993.
302. Shadel GS, Horvath TL. Mitochondrial ROS signaling in organismal homeostasis. *Cell* 163: 560-569, 2015.
303. Shafat A, Butler P, Jensen RL, and Donnelly AE. Effects of dietary supplementation with vitamins C and E on muscle function during and after eccentric contractions in humans. *Eur J Appl Physiol* 93: 196-202, 2004.
304. Shelar SB, Narasimhan M, Shanmugam G, Litovsky SH, Gounder SS, Karan G, Arulvasu C, Kensler TW, Hoidal JR, Darley-Usmar VM, and Rajasekaran NS. Disruption of nuclear factor (erythroid-derived-2)-like 2 antioxidant

- signaling: a mechanism for impaired activation of stem cells and delayed regeneration of skeletal muscle. *FASEB J* 30: 1865-1879, 2016.
305. Sies H. Role of metabolic H₂O₂ generation: redox signaling and oxidative stress. *J Biol Chem* 289: 8735-41, 2014.
 306. Sies H. Oxidative stress: a concept in redox biology and medicine. *Redox Biol* 4: 180-183, 2015.
 307. Singh S, Canseco DC, Manda SM, Shelton JM, Chirumamilla RR, Goetsch SC, Ye Q, Gerard RD, Schneider JW, Richardson JA, Rothermel BA, and Mammen PPA. Cytoglobin modulates myogenic progenitor cell viability and muscle regeneration. *Proc Nat Acad Sci USA* 111: E129-138, 2014.
 308. Smith RA, Murphy MP. Animal and human studies with the mitochondria-targeted antioxidant MitoQ. *Ann New York Acad Sci* 1201: 96-103, 2010.
 309. Smythe GM, Forwood JK. Altered mitogen-activated protein kinase signaling in dystrophic (mdx) muscle. *Muscle Nerve* 46: 374-383, 2012.
 310. Sobotta MC, Liou W, Stöcker S, Talwar D, Oehler M, Ruppert T, Scharf AND, and Dick TP. Peroxiredoxin-2 and STAT3 form a redox relay for H₂O₂ signaling. *Nat Chem Biol* 11: 64-70, 2015.
 311. Stamler JS, Meissner G. Physiology of nitric oxide in skeletal muscle. *Physiol Rev* 81: 209-237, 2001.
 312. Steinbrenner H. Interference of selenium and selenoproteins with the insulin-regulated carbohydrate and lipid metabolism. *Free Radic Biol Med* 65: 1538-47, 2013.
 313. Sugama K, Suzuki K, Yoshitani K, Shiraishi K, Miura S, Yoshioka H, Mori Y, and Kometani T. Changes of thioredoxin, oxidative stress markers, inflammation and muscle/renal damage following intensive endurance exercise. *Exerc Immunol Rev* 21: 130-142, 2015.
 314. Sullivan-Gunn MJ, Campbell-O'Sullivan SP, Tisdale MJ, and Lewandowski PA. Decreased NADPH oxidase expression and antioxidant activity in cachectic skeletal muscle. *J Cachexia Sarcopenia Muscle* 2: 181-188, 2011.
 315. Sullivan-Gunn MJ, Lewandowski PA. Elevated hydrogen peroxide and decreased catalase and glutathione peroxidase protection are associated with aging sarcopenia. *BMC geriatrics* 13: 104, 2013.
 316. Sundaresan M, Yu ZX, Ferrans VJ, Irani K, and Finkel T. Requirement for generation of H₂O₂ for platelet-derived growth factor signal transduction. *Science* 270: 296-299, 1995.
 317. Sylow L, Moller LL, Kleinert M, Richter EA, and Jensen TE. Rac1--a novel regulator of contraction-stimulated glucose uptake in skeletal muscle. *Exp Physiol* 99: 1574-80, 2014.
 318. Tang AHR, Rando TA. Induction of autophagy supports the bioenergetic demands of quiescent muscle stem cell activation. *EMBO J* 33: 2782-97, 2014.
 319. Tatsumi R, Liu X, Pulido A, Morales M, Sakata T, Dial S, Hattori A, Ikeuchi Y, and Allen RE. Satellite cell activation in stretched skeletal muscle and the role of nitric oxide and hepatocyte growth factor. *Am J Physiol Cell Physiol* 290: C1487-94, 2006.
 320. Tatsumi R, Wuollet AL, Tabata K, Nishimura S, Tabata S, Mizunoya W, Ikeuchi Y, and Allen RE. A role for calcium-calmodulin in regulating nitric oxide production during skeletal muscle satellite cell activation. *Am J Physiol. Cell Physiol* 296: C922-929, 2009.

321. Tidball JG, Wehling-Henricks M. The role of free radicals in the pathophysiology of muscular dystrophy. *J Appl Physiol* (1985) 102: 1677-86, 2007.
322. Tierney MT, Aydogdu T, Sala D, Malecova B, Gatto S, Puri PL, Latella L, and Sacco A. STAT3 signaling controls satellite cell expansion and skeletal muscle repair. *Nat Med* 20: 1182-1186, 2014.
323. Titov DV, Cracan V, Goodman RP, Peng J, Grabarek Z, and Mootha VK. Complementation of mitochondrial electron transport chain by manipulation of the NAD⁺/NADH ratio. *Science* 352: 231-5, 2016.
324. Tonkin J, Temmerman L, Sampson RD, Gallego-Colon E, Barberi L, Bilbao D, Schneider MD, Musaro A, and Rosenthal N. Monocyte/macrophage-derived IGF-1 orchestrates murine skeletal muscle regeneration and modulates autocrine polarization. *Mol Ther* 23: 1189-1200, 2015.
325. Traber MG, Stevens JF. Vitamins C and E: beneficial effects from a mechanistic perspective. *Free Radic Biol Med* 51: 1000-1013, 2011.
326. Tromm CB, Pozzi BG, Paganini CS, Marques SO, Pedrosa GS, Souza PS, Silveira PC, Silva LA, De Souza CT, and Pinho RA. The role of continuous versus fractionated physical training on muscle oxidative stress parameters and calcium-handling proteins in aged rats. *Aging Clin Exp Res*: DOI 10.1007/s40520-015-0501-6, 2015.
327. Troy A, Cadwallader AB, Fedorov Y, Tyner K, Tanaka KK, and Olwin BB. Coordination of satellite cell activation and self-renewal by Par-complex-dependent asymmetric activation of p38 α /beta MAPK. *Cell Stem Cell* 11: 541-53, 2012.
328. Ueda Y, Whanger PD, and Forsberg NE. The effects of selenium deficiency on differentiation, degradation, and cell lysis of L8 rat skeletal muscle cells. *Biol Trace Elem Res* 69: 1-13, 1999.
329. Uezumi A, Fukada S, Yamamoto N, Takeda S, and Tsuchida K. Mesenchymal progenitors distinct from satellite cells contribute to ectopic fat cell formation in skeletal muscle. *Nature Cell Biology* 12: 143-152, 2010.
330. Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C, Sjöstedt E, Asplund A, Olsson I, Edlund K, Lundberg E, Navani S, Szigartyo CA-K, Odeberg J, Djureinovic D, Takanen JO, Hober S, Alm T, Edqvist P-H, Berling H, Tegel H, Mulder J, Rockberg J, Nilsson P, Schwenk JM, Hamsten M, von Feilitzen K, Forsberg M, Persson L, Johansson F, Zwahlen M, von Heijne G, Nielsen J, and Pontén F. Proteomics. Tissue-based map of the human proteome. *Science* 347: 1260419, 2015.
331. Ulibarri JA, Mozdziak PE, Schultz E, Cook C, and Best TM. Nitric oxide donors, sodium nitroprusside and S-nitroso-N-acetylpencillamine, stimulate myoblast proliferation in vitro. *In Vitro Cell Dev Biol Anim* 35: 215-218, 1999.
332. Umanskaya A, Santulli G, Xie W, Andersson DC, Reiken SR, and Marks AR. Genetically enhancing mitochondrial antioxidant activity improves muscle function in aging. *Proc Nat Acad Sci USA* 111: 15250-15255, 2014.
333. Urao N, Sudhakar V, Kim S-J, Chen G-F, McKinney RD, Kojda G, Fukai T, and Ushio-Fukai M. Critical role of endothelial hydrogen peroxide in post-ischemic neovascularization. *PLoS One* 8: e57618, 2013.
334. Ushio-Fukai M, Tang Y, Fukai T, Dikalov SI, Ma Y, Fujimoto M, Quinn MT, Pagano PJ, Johnson C, and Alexander RW. Novel role of gp91(phox)-containing NAD(P)H oxidase in vascular endothelial growth factor-induced signaling and angiogenesis. *Circulation Research* 91: 1160-1167, 2002.

335. Valko M, Leibfritz D, Moncol J, Cronin MTD, Mazur M, and Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 39: 44-84, 2007.
336. Vamecq J, Cherkaoui-Malki M, Andreoletti P, and Latruffe N. The human peroxisome in health and disease: the story of an oddity becoming a vital organelle. *Biochimie* 98: 4-15, 2014.
337. van der Vliet AJanssen-Heininger YM. Hydrogen peroxide as a damage signal in tissue injury and inflammation: murderer, mediator, or messenger? *J Cell Biochem* 115: 427-35, 2014.
338. Veal EA, Day AM, and Morgan BA. Hydrogen peroxide sensing and signaling. *Mol Cell* 26: 1-14, 2007.
339. Venereau E, Casalgrandi M, Schiraldi M, Antoine DJ, Cattaneo A, De Marchis F, Liu J, Antonelli A, Preti A, Raeli L, Shams SS, Yang H, Varani L, Andersson U, Tracey KJ, Bachi A, Uguccioni M, and Bianchi ME. Mutually exclusive redox forms of HMGB1 promote cell recruitment or proinflammatory cytokine release. *J Exp Med* 209: 1519-1528, 2012.
340. Vercherat C, Chung TK, Yalcin S, Gulbagci N, Gopinadhan S, Ghaffari S, and Taneja R. Stra13 regulates oxidative stress mediated skeletal muscle degeneration. *Hum Mol Genet* 18: 4304-4316, 2009.
341. Vezzoli M, Castellani P, Corna G, Castiglioni A, Bosurgi L, Monno A, Brunelli S, Manfredi AA, Rubartelli A, and Rovere-Querini P. High-mobility group box 1 release and redox regulation accompany regeneration and remodeling of skeletal muscle. *Antioxid Redox Signal* 15: 2161-2174, 2011.
342. Vieira NM, Elvers I, Alexander MS, Moreira YB, Eran A, Gomes JP, Marshall JL, Karlsson EK, Verjovski-Almeida S, Lindblad-Toh K, Kunkel LM, and Zatz M. Jagged 1 Rescues the Duchenne Muscular Dystrophy Phenotype. *Cell* 163: 1204-13, 2015.
343. Viña J, Gomez-Cabrera MC, Lloret A, Marquez R, Miñana JB, Pallardó FV, and Sastre J. Free radicals in exhaustive physical exercise: mechanism of production, and protection by antioxidants. *IUBMB life* 50: 271-277, 2000.
344. von Maltzahn J, Chang NC, Bentzinger CF, and Rudnicki MA. Wnt signaling in myogenesis. *Trends Cell Biol* 22: 602-609, 2012.
345. Weerapana E, Wang C, Simon GM, Richter F, Khare S, Dillon MBD, Bachovchin DA, Mowen K, Baker D, and Cravatt BF. Quantitative reactivity profiling predicts functional cysteines in proteomes. *Nature* 468: 790-795, 2010.
346. Wehling M, Spencer MJ, and Tidball JG. A nitric oxide synthase transgene ameliorates muscular dystrophy in mdx mice. *Journal of Cell Biology* 155: 123-131, 2001.
347. Weissman BA, Jones CL, Liu Q, and Gross SS. Activation and inactivation of neuronal nitric oxide synthase: characterization of Ca(2+)-dependent [125I]Calmodulin binding. *Eur J Pharmacol* 435: 9-18, 2002.
348. West AP, Brodsky IE, Rahner C, Woo DK, Erdjument-Bromage H, Tempst P, Walsh MC, Choi Y, Shadel GS, and Ghosh S. TLR signalling augments macrophage bactericidal activity through mitochondrial ROS. *Nature* 472: 476-480, 2011.
349. Whitehead NP, Pham C, Gervasio OL, and Allen DG. N-Acetylcysteine ameliorates skeletal muscle pathophysiology in mdx mice. *J Physiol* 586: 2003-2014, 2008.

350. Winokur ST, Barrett K, Martin JH, Forrester JR, Simon M, Tawil R, Chung SA, Masny PS, and Figlewicz DA. Facioscapulohumeral muscular dystrophy (FSHD) myoblasts demonstrate increased susceptibility to oxidative stress. *Neuromuscul Disord* 13: 322-333, 2003.
351. Winterbourn C. Current methods to study reactive oxygen species--pros and cons. Preface. *Biochim Biophys Acta* 1840: 707, 2014.
352. Winterbourn C, Metodiewa D. Reactivity of biologically important thiol compounds with superoxide and hydrogen peroxide. *Free Radic Biol Med* 27: 322-328, 1999.
353. Won H, Lim S, Jang M, Kim Y, Rashid MA, Jyothi KR, Dashdorj A, Kang I, Ha J, and Kim SS. Peroxiredoxin-2 upregulated by NF- κ B attenuates oxidative stress during the differentiation of muscle-derived C2C12 cells. *Antioxid Redox Signal* 16: 245-261, 2012.
354. Woo HA, Yim SH, Shin DH, Kang D, Yu D-Y, and Rhee SG. Inactivation of peroxiredoxin I by phosphorylation allows localized H₂O₂ accumulation for cell signaling. *Cell* 140: 517-528, 2010.
355. Xu X, Arriaga EA. Qualitative determination of superoxide release at both sides of the mitochondrial inner membrane by capillary electrophoretic analysis of the oxidation products of triphenylphosphonium hydroethidine. *Free Radic Biol Med* 46: 905-913, 2009.
356. Yamada M, Tatsumi R, Kikuri T, Okamoto S, Nonoshita S, Mizunoya W, Ikeuchi Y, Shimokawa H, Sunagawa K, and Allen RE. Matrix metalloproteinases are involved in mechanical stretch-induced activation of skeletal muscle satellite cells. *Muscle Nerve* 34: 313-319, 2006.
357. Yfanti C, Akerstrom T, Nielsen S, Nielsen AR, Mounier R, Mortensen OH, Lykkesfeldt J, Rose AJ, Fischer CP, and Pedersen BK. Antioxidant supplementation does not alter endurance training adaptation. *Med Sci Sports Exerc* 42: 1388-95, 2010.
358. Yfanti C, Nielsen AR, Akerstrom T, Nielsen S, Rose AJ, Richter EA, Lykkesfeldt J, Fischer CP, and Pedersen BK. Effect of antioxidant supplementation on insulin sensitivity in response to endurance exercise training. *Am J Physiol Endocrinol Metab* 300: E761-70, 2011.
359. Yin H, Price F, and Rudnicki MA. Satellite cells and the muscle stem cell niche. *Physiological Reviews* 93: 23-67, 2013.
360. Zaccagnini G, Martelli F, Magenta A, Cencioni C, Fasanaro P, Nicoletti C, Biglioli P, Pelicci PG, and Capogrossi MC. p66(ShcA) and oxidative stress modulate myogenic differentiation and skeletal muscle regeneration after hind limb ischemia. *J Biol Chem* 282: 31453-31459, 2007.
361. Zhang H, Ryu D, Wu Y, Gariani K, Wang X, Luan P, D'Amico D, Ropelle ER, Lutolf MP, Aebbersold R, Schoonjans K, Menzies KJ, and Auwerx J. NAD⁺ repletion improves mitochondrial and stem cell function and enhances life span in mice. *Science*: pii: aaf2693., 2016.
362. Zhang Y, Choksi S, Chen K, Pobezinskaya Y, Linnoila I, and Liu Z-G. ROS play a critical role in the differentiation of alternatively activated macrophages and the occurrence of tumor-associated macrophages. *Cell Res* 23: 898-914, 2013.
363. Zhuang W, Eby JC, Cheong M, Mohapatra PK, Bredt DS, Disatnik MH, and Rando TA. The susceptibility of muscle cells to oxidative stress is independent of nitric oxide synthase expression. *Muscle Nerve* 24: 502-511, 2001.

364. Zorov DB, Juhaszova M, and Sollott SJ. Mitochondrial reactive oxygen species (ROS) and ROS-induced ROS release. *Physiol Rev* 94: 909-50, 2014.

Figure legends

Figure 1. Overview of skeletal muscle regeneration. Left panel presents hematoxylin-eosin (HE) staining of uninjured and regenerating muscle (mouse *Tibialis Anterior* muscle) after an injury triggered by the injection of the venom toxin cardiotoxin. Day 1 after injury, myofibers undergo necrosis (pale pink staining) while the first inflammatory cells (neutrophils and monocytes/macrophages) enter the injured area (purple cells). Day 2 after injury, the number of infiltrating immune cells has increased and is mainly composed of monocytes/macrophages (large purple cells), which phagocytose the necrotic debris and necrotic myofibers. Meanwhile (non visible on HE staining), both MuSCs and FAPs activate and expand. Day 4 after injury, MuSCs differentiate and fuse to form new myofibers (dark pink structures containing one or two large nuclei with a "fried egg" appearance) while angiogenesis takes place (non visible on HE staining). Macrophages are still very numerous and they surround the regenerating myofibers. Day 7 after injury, the number of interstitial cells (FAPs and macrophages) has dropped while the regenerating myofibers grow in size (through both fusion of differentiated MuSCs and hypertrophy). One month after injury, the muscle has come back to its normal appearance, except that the regenerating myofibers are characterized by nuclei in a central - and not peripheral - location. In this model, functionality of the muscle is fully recovered 2 months after injury.

Figure 2. Adult myogenesis. At the quiescent state, most MuSCs express high levels of the transcription factors Pax7 and Myf5. Upon activation, MuSCs enter the cell cycle, still express Pax7 and Myf5 and express MyoD while they are proliferating. Self-renewing MuSCs (grey arrows towards the left) downregulate MyoD expression

while upregulating Pax7 expression to return to quiescence. The majority of expanding MuSCs enter into myogenic differentiation that is sustained by the extinction of Pax7 and Myf5 expression together with the increase of Myogenin and MRF4 expression, which are synthesized once the cells have exited the cell cycle. Later on, contractile proteins are produced in the fully differentiated myogenic cells that fuse to form new myofibers. Maturation (not shown) of the myofibers is achieved by re-innervation through neuromuscular junctions.

Figure 3. Various cell types are present around MuSCs during skeletal muscle regeneration. During the time course of post-injury muscle regeneration, several cell types participate in the MuSC adult myogenesis program. In the model of cardiotoxin injury, MuSCs expand for a couple of days then start to differentiate. At day 4 new regenerating myofibers are visible. FAPs are also activated upon injury and rapidly expand during the first days of regeneration, after which their number rapidly decreases. After injury, the vascular bed is altered and a few days later a reorganization of ECs is observed to form new functional vessels concomitantly to the formation of the new myofibers. Monocytes from the circulation enter into the injured area a few hours after the injury and initially give rise to inflammatory macrophages during the first 1-2 days. Then, upon phagocytosis of the muscle tissue debris, they transform into anti-inflammatory/restorative macrophages that proliferate and stay numerous until the end of the regeneration process, then their number drops back to normal values.

Figure 4. Adult myogenesis during skeletal muscle regeneration is regulated by neighboring cells. The various cell types neighboring MuSCs (see Figure 3)

exert specific functions on MuSCs by delivering specific cues to MuSCs in a timely manner. Proliferation/expansion of activated MuSCs is sustained by FAPs and inflammatory (damage-associated) macrophages. The next step of myogenesis, *i.e.* differentiation, is stimulated by anti-inflammatory (restorative) macrophages as well as by ECs. Finally, the fusion of MuSCs into multinucleated structures is boosted by restorative macrophages. Return to quiescence of a small subset of MuSCs is supported by the action of peri-endothelial cells such as smooth muscle cells and pericytes, which also maintain the stability of vessels/capillaries.

Figure 5. Muscular dystrophies – the DGC complex. Inherited muscular dystrophies originate from mutations in one of the components of the DystroGlycan complex (DGC), which links the actin cytoskeleton of the myofiber to the extracellular matrix. The main proteins are the extracellular α -dystroglycan (α -DG) and laminin, the sarcolemmal β -dystroglycan (β -DG), sarcoglycans (SGCs) and sarcospan (Sar), and the cytosolic proteins dystrophin, syntrophins (SYN), dystrobrevin (DYB), actin and nNOS. The DGC complex provides stability to the membrane structure of skeletal muscle fibers during contraction/relaxation cycles. When mutations target DGC components, the stability is not ensured and fragility of the membrane leads to repeated injuries of the myofibers during contraction.

Figure 6. Permanent regeneration during muscular dystrophies. After an acute injury in normal muscle (top left part of the picture), the process of skeletal muscle regeneration occurs due to synchronous regenerating cues delivered by cells neighboring MuSCs. In muscular dystrophies, repeated injuries, occurring throughout the muscle, lead to the asynchronous delivery of regenerating cues, preventing the

various cell types from being able to carry out the timely orchestration of the process of skeletal muscle regeneration, which is thus profoundly affected. The permanent attempts of regeneration together with the cycles of necrosis lead to failed myogenesis, chronic inflammation, fibrosis and alterations in the vascular bed.

Figure 7. ROS metabolism and biochemistry. Chemical characteristics of ROS reveal their biological properties. Oxygen molecule (O_2) univalent reduction leads to the formation of reactive superoxide anion ($O_2^{\cdot-}$). This primary ROS is a weak oxidant and a weak reductant agent. $O_2^{\cdot-}$ interacts with NO^{\cdot} , transition metals and can naturally dismutate in H_2O_2 . Superoxide dismutases accelerate the rate of this reaction by 10,000. H_2O_2 is a non-radical molecule exhibiting a low reactivity and the ability to cross membrane, suggesting a crucial role in signaling. By contrast, hydroxyl radical (HO^{\cdot}) derived from metals and H_2O_2 interactions is highly reactive and is an unstable molecule that damages unspecified compounds that are located close to its site of production.

Figure 8. Intracellular sources of ROS and RNS and antioxidant network in skeletal muscle cell. Nucleus, mitochondria, endoplasmic reticulum (ER) and peroxisomes are the main sources of ROS in skeletal muscle cell. Additionally, enzymatic complexes generate ROS such as XO and NOX. Antioxidant molecules are mainly localized in the cytosol and in mitochondria. In addition, lipophilic antioxidants (vitamin E and tocopherols) are contained within the lipidic membrane. Dietary antioxidants such as vitamin C and carotenoids enter into cytosol following food ingestion.

Figure 9. Major oxidative thiol-modifications in redox-sensitive cysteine residues. Thiol moiety of redox-sensitive cysteine residues (SH) is vulnerable to oxidation and can lead to various post-translational modifications that are reversible or irreversible depending on the degree of oxidation. The initial reaction product with ROS is sulfenic acid (SOH). Further oxidation of sulfenic acid leads to the formation of irreversible modifications such as sulfinic acid (SO₂H) and subsequently sulfonic acid (SO₃H). Reaction of thiol with ROS can lead to the formation of both intra and intermolecular disulfide bonds. Thiol reaction with GSH and RNS leads to the formation of glutathionylation (-SGSH) and nitrosothiols (-SNO), respectively. Full and dotted arrows represent oxidation and reduction, respectively.

Figure 10. Redox-sensitive targets during adult myogenesis. Each step of adult myogenesis during skeletal muscle regeneration (see Figure 2) is regulated in MuSCs by redox activity. Some redox-related factors have been shown to directly regulate MuSC survival, activation, proliferation, self-renewal and differentiation (upper panel). Moreover, various signaling pathways have been shown to control MuSC fate on the one hand, and are known to be targets of redox regulation on the other hand (lower panel).

Figure 11. Macrophages during skeletal muscle regeneration. Upon skeletal muscle injury, monocytes/macrophages are recruited from blood and invade the tissue as damage-associated macrophages (Ly6C^{pos} cells). They release pro-inflammatory molecules (TNF α , IL6, IL1 β , VEGF) to stimulate MuSC proliferation. Then, upon phagocytosis these macrophages skew their phenotype to acquire a restorative phenotype and secrete compounds that stimulate myogenic differentiation

(IGF-1, TGF β). Macrophage skewing is regulated by a series of effectors, which are regulated by redox mechanisms. Moreover, restorative macrophages express high levels of SOD1 and Trx, which are associated with the reduction of the alarmin HGMB1 that participates to MuSC migration and differentiation. See text for details.

Figure 12. Redox alteration of myogenesis in aged muscle. Fate of MuSCs during post-injury myogenesis is altered in aged muscle, including reduced activation, reduced proliferation, reduced self-renewal and increased differentiation. A series of redox effectors are deregulated in aged MuSCs and are listed in the box. In italic are shown the signaling pathways that are altered in aged MuSCs and known to be redox regulated. \blacktriangleright \blacktriangleleft stand for increase and decrease, respectively.


Figure 13. ROS in MuSCs in muscular dystrophies. The left circle represents a synchronous regeneration process triggered by a single injury. The right circle shows that in muscular dystrophies, repeated injuries trigger the delivery of asynchronous regenerative cues, leading to a failure of the regeneration process, associated with a whole increase of ROS. MuSCs from dystrophic muscle have been shown to be more susceptible to oxidative stress than normal MuSCs. However, only few molecular determinants underlying this susceptibility have been evidenced so far, those are listed in the box. See text for details. \blacktriangleright \blacktriangleleft stand for increase and decrease, respectively.

Figure 14. Therapeutic strategies in aged and dystrophic muscles – effects on MuSCs. Antioxidant supplementation on whole body (upper panel) increases the detection of antioxidant biomarkers, triggers some beneficial effects in muscular



dystrophies, but no improvement of muscle function in both aged and diseased muscle is observed. A variety of antioxidant molecules exert beneficial effects on MuSCs *in vitro* (middle panel), notably they decrease some abnormalities observed in muscular dystrophies. The analysis of MuSCs *in vivo* after treatment in mouse models (lower panel) indicates an overall benefit for MuSC homeostasis via increased energetic metabolism and myogenesis. ↗↘≈ stand for increase, decrease and no effect, respectively.


Figure 15. Future strategies for the study of ROS. The scheme presents interrelated strategies and developing tools that will help to identify precisely which ROS and ROS-dependent signaling are involved in a given biological context. Such knowledge will further allow to define specific targets potentially used in pharmacological trials. Detection and identification of ROS is performed through the development of specific probes (upper left). Several chemical probes have been developed to investigate redox mechanisms, mainly composed of a fluorophore that emits or is liberated upon oxidation (351). More recently, genetically encoded fluorescent redox sensors or injected exomarkers allow the detection and visualization of live variations of ROS or redox regulated molecules (e.g. GSH, NAD) *in vitro* and *in vivo* (upper right) (5,68,98,197). ROS-induced post-translational modifications (PTMs) have recently been described due to the development of redox proteomics, thanks to the setup of cysteine residue labeling (bottom right). Together with crystallography, which describes protein modifications upon redox reaction, these biochemical strategies allowed the identification of new molecular targets of redox reactions (bottom right) (60,61,186,213). The development of this large variety of tools for *in vitro* and *in vivo* investigation of redox biology, together with the

improvement of the probes and of their specificity will enable the development of antioxidant probes to be tested in preclinical and clinical trials (bottom left).




ADULT MYOGENESIS DURING SKELETAL MUSCLE REGENERATION






Myogenic precursor cells




Fibro-Adipogenic Precursors






Endothelial cells




Macrophages











Superoxide
Weak oxidant
Metal interactions
NO· interactions
H₂O₂ source



Hydrogen peroxide
Weak oxidant
Cys/Met oxidation
Signaling

Hydroxyl radical
Highly reactive
Lipids, proteins, and
DNA damages



MITOCHONDRION

SOD1

GPX1/4

GSH

TRX2/TRXR2

SOD2

MYONUCLEUS

CYTOSOL

Enzymatic systems

- Xanthine oxidase
- Cyclooxygenases
- Lipooxygenases
- Cytochrome p450
- Phospholipases
- NO synthases

PEROXISOME

CAT

ER

GSH

SOD1

GRX1

GPX1


PRX1/2


TRX1/TRXR1

Vitamin E


Carotenoids
Vitamin C


NOXs





survival	activation	proliferation	self-renewal	differentiation
Redox related factors				
GPX3 ALDH Cytoglobin	CaM-NOS NO• (HGF, MMP)	NADH-O ₂ ^{•-} GPX1 SOD1	GPX3 TrxR1 NO• (Vangl2)	DUOXA1 p66 ^{ShcA} SEPN Nrf2-GCL/GR-GSH Nrf2-CRB1 NO• DRP1 Prx2
Redox regulated signaling pathways				
NF-κB AP-1 Nrf2 Caspases Bcl-2	S1P MAPKs (p38) Notch Wnt-Se-GPX1	ERK1/2 PI3K Akt/mTOR NF-κB CDKN1 S1P MAPKs (p38) Notch		NF-κB





- ↘ activation upon stretch and NO
- ↘ autophagy


- ↘ catalase
- ↘ glutathione transferase
- ↘ Nrf2 signaling → Pax7 MyoD alteration
- ↘ Nrf2 signaling → ↗ ubiquitin/ apoptotic pathways
- ↘ NAD⁺ → ↘ mtUPR
- ↗ oxidation of DNA and lipid peroxidation
- ↗ superoxide/mitochondrial ratios → ↘ mitochondria ATP production

- ↘ Notch
- ↗ Wnt
- ↗ JAK-STAT3
- ↗ p38 MAPK

Normal muscle



Diseased muscle



MuSCs:

- ↗ susceptibility to oxidative stress
 - altered MAPKs (JNK, p38 and ERK1/2) signaling
 - defect in myotube size
 - defect in branching myotubes/myofibers
- ↗ Stra13
- ↘ Bmi-1 → ↘ MT1

AGING

Antioxidant supplementation:
↗ antioxidant biomarkers
≈ muscle function

MUSCULAR DYSTROPHIES

Antioxidant supplementation:
↗ antioxidant biomarkers
↘ regenerating fibers
↘ necrosis
↗ protein catabolism
↗ muscle phenotype
≈ muscle function

MuSCs- *in vitro* studies

+ Vitamin E
↗ proliferation

+ NDGA
↘ proliferation
↗ differentiation, ↗ myotubes
+ Green tea polyphenol blend
↗ resistance to oxidative damage
+ Antioxidants (fisetin, NAC, vitamin E)
↘ desmin aggregation
+ vitamin E
↗ plasma membrane repair

MuSCs – *in vivo* studies (mouse)

+ Vitamin E
↗ proliferation
↗ autophagy
+ Nicotinamide riboside
↗ mtUPR
↗ mitochondria homeostasis
↗ MuSC regenerative properties

+ Flavocoxid
↗ myogenin-positive cells
+ Quercetin
↗ neofibers, ↗ MuSCs
+ NCX 320
↗ neofibers, ↗ MuSCs
+ Nicotinamide riboside
↗ MuSCs

