



HAL
open science

High-dose thiotepa-based chemotherapy with autologous stem cell support in elderly patients with primary central nervous system lymphoma: a European retrospective study

S. Schorb, C Fox, K Fritsch, L Isbell, N Neubauer, A Tzalavras, R Witherall, C. Choquet, K Kuittinen, D De-Silva, et al.

► To cite this version:

S. Schorb, C Fox, K Fritsch, L Isbell, N Neubauer, et al.. High-dose thiotepa-based chemotherapy with autologous stem cell support in elderly patients with primary central nervous system lymphoma: a European retrospective study. *Bone Marrow Transplantation*, 2017, 52 (8), pp.1113-1119. 10.1038/bmt.2017.23 . hal-01596402

HAL Id: hal-01596402

<https://univ-rennes.hal.science/hal-01596402v1>

Submitted on 27 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

High-dose thiotepa-based chemotherapy with autologous stem cell support in elderly patients with primary central nervous system lymphoma - a European retrospective study

Elisabeth Schorb^{1*}, Christopher P Fox^{2*}, Kristina Fritsch¹, Lisa Isbell¹, Amelie Neubauer³, Asterios Tzalavras³, Ruth Witherall², Sylvain Choquet⁴, Outi Kuittinen⁵, Dunnya De-Silva⁶, Kate Cwynarski⁶, Caroline Houillier⁷, Khê Hoang-Xuan⁷, Valérie Touitou⁸, Nathalie Cassoux⁹, Jean-Pierre Marolleau¹⁰, Jérôme Tamburini¹¹, Roch Houot¹², Vincent Delwail¹³, Gerald Illerhaus³, Carole Soussain^{14*}, and Benjamin Kasenda^{3*}

¹Department of Haematology, Oncology and Stem Cell Transplantation - University of Freiburg, Faculty of Medicine, University of Freiburg, Germany

²Department of Clinical Haematology, Nottingham University Hospitals NHS Trust, Nottingham, UK

³Stuttgart Cancer Centre, Klinikum Stuttgart, Stuttgart, Germany

⁴Department of Haematology, APHP, Sorbonne Universités, Hospital Pitié-Salpêtrière, Paris, France

⁵Department of Oncology, Oulu University Hospital, Oulu, Finland

⁶Department of Haematology, University College Hospital, London, United Kingdom

⁷Department of Neurology, APHP, Sorbonne Universités, UPMC Hospital Pitié-Salpêtrière, LOC network, Paris-France

⁸Department of Ophthalmology, APHP, Sorbonne Universités, UPMC Hospital Pitié-Salpêtrière, LOC network, Paris-France

⁹Department of Ophthalmology, Curie Institute - Rene Huguenin Hospital, LOC network, Saint-Cloud, France

¹⁰Department of Haematology, University Hospital, Amiens, LOC network, France

¹¹Department of Haematology, Cochin University Hospital, LOC network, Paris-France

¹²Department of Haematology, University Hospital, Rennes, LOC network, France

¹³Vincent Delwail, Department of Haematology, University Hospital, Poitiers, LCO network, France

¹⁴Department of Haematology, Curie Institute - Rene Huguenin Hospital, LOC network, Saint-Cloud, France

* authors contributed equally

Running title: High-dose chemotherapy in elderly PCNSL patients

Corresponding author:

Elisabeth Schorb

Department of Haematology, Oncology and Stem Cell Transplantation - University of Freiburg
Hugstetter Str. 55

79106 Freiburg

Germany

Phone: +49-761-270-35360

Fax: +49-761-270-33110

elisabeth.schorb@uniklinik-freiburg.de

Conflict of Interest: The authors declare no conflict of interest.

1 **Abstract**

2 In this retrospective multicentre study, we investigated the outcome of elderly PCNSL
3 patients (≥ 65 years) who underwent HDT-ASCT at eleven centres between 2003
4 and 2016. Endpoints included remission, progression free survival, overall survival
5 and treatment-related mortality.

6 We identified 52 patients (median age 68.5 years, median KPS before HDT-ASCT
7 80%) who all underwent thiotepa-based HDT-ASCT. Fifteen patients (28.8%)
8 received HDT-ASCT as first-line treatment and 37 (71.2%) received it as second or
9 subsequent line. Remission status before HDT-ASCT was: complete remission (CR)
10 34.6%, partial remission (PR) 51.9%, stable disease 3.8%, and progressive disease
11 9.6%. Following completion of HDT-ASCT, 36 patients (69.2%) achieved CR (21.2%
12 first line setting, 48.1% second or subsequent line setting) and 9 (17.3%) PR (5.8%
13 first line setting, 11.5% second or subsequent line setting). With a median follow up
14 of 22 months after HDT-ASCT, median PFS and OS were reached after 51.1 and
15 122.3 months, respectively. 2-year PFS and OS rates were 62.0% and 70.8%,
16 respectively. We observed two HDT-ASCT associated deaths (3.8%).

17 In selected elderly PCNSL patients, HDT-ASCT, using thiotepa-based conditioning
18 regimes, is feasible and effective. Further prospective and comparative studies are
19 warranted to further evaluate the role of HDT-ASCT in elderly PCNSL patients.

20

21

22 **Introduction**

23 Primary central nervous system lymphoma (PCNSL) is an aggressive Non-Hodgkin
24 Lymphoma, typically diffuse large B cell lymphoma (DLBCL), which exclusively involves the
25 central nervous system at diagnosis. It accounts for 3% to 4% of all primary brain tumours
26 and 4% to 6% of extra-nodal lymphomas (1). The incidence of PCNSL in immunocompetent
27 patients has been steadily increasing over the last 30 years (2, 3). However, notwithstanding
28 progress over the last decade with improved immunochemotherapy approaches, PCNSL
29 remains a challenging disease.

30 Patients older than 65 years account for 50% of all PCNSL cases (4). Although some elderly
31 patients may tolerate intensive systemic chemotherapy, they typically experience an inferior
32 prognosis as compared to younger patients with PCNSL. Moreover, elderly patients are
33 vulnerable to iatrogenic toxicity, especially neurotoxicity following whole brain radiation
34 therapy (WBRT) (4); thus they represent a unique treatment subgroup (5, 6). An US registry
35 study of 579 elderly patients diagnosed with PCNSL in the 1990s described a median
36 survival of only 7 months and that WBRT alone was the most common treatment modality
37 (46%) (7). Even with modern conventional chemotherapy protocols, most patients will
38 experience a short progression free survival (PFS) (<12months) and die from disease (8, 9).
39 For younger patients, thiotepa-containing high-dose chemotherapy followed by autologous
40 stem cell transplantation (HDT-ASCT) has been shown to be feasible and effective in both
41 newly diagnosed and relapsed patients with PCNSL (10-15). Due to toxicity and tolerability
42 concerns, this intensive CNS-directed treatment has typically been restricted to patients
43 younger than 65 years of age. However, age alone may not be the appropriate criterion to
44 select patients for this effective treatment approach.

45 We undertook this retrospective, international study to investigate outcome after HDT-ASCT
46 in elderly PCNSL (≥ 65 years) from eleven centres with experienced physicians, who have
47 been treating PCNSL patients and performing clinical trials in PCNSL for many years.

48 **Methods**

49 *Patient selection criteria and data collection*

50 Eligibility criteria for this retrospective multicentre analysis were i) age \geq 65 years at the time
51 of HDT-ASCT; ii) histologically-proven PCNSL (at first diagnosis, no repeat biopsy required
52 at relapse) without systemic lymphoma manifestation at any time; iii) no evidence of
53 immunodeficiency; iv) completed thiotepa-based HDT-ASCT. All centres screened their
54 databases for PCNSL patients \geq 65 years, excluding patients not treated with HDT-ASCT.
55 Data from all study-eligible patients of the eleven cooperating centres were collected using a
56 pre-specified, anonymised case report form, including: patient and tumour characteristics at
57 baseline, treatment, transplantation specific data, main reported toxicities, objective
58 response, site and date of relapse or progression, and survival. Data were checked by the
59 coordinating investigators for consistency and, if necessary, queries resolved with sites
60 before entering data into the central database. All patients provided informed consent for the
61 documentation of anonymised clinical data and the use for scientific publication. The ethics
62 committee of Freiburg University approved the study protocol.

63

64 *Statistical analysis*

65 The principal outcomes of interest were remission status before and after HDT-ASCT
66 (complete remission [CR], partial remission [PR], stable disease [SD], progressive disease
67 [PD]) as reported by the respective centres, PFS (defined as time from HDT-ASCT to
68 progression, relapse or death; whichever occurred first) and overall survival (OS, defined as
69 time from HDT-ASCT to death due to any cause). PFS and OS were estimated using the
70 Kaplan-Meier method including 95% confidence interval (CIs). We additionally stratified
71 response and survival outcomes by line of treatment in which HDT-ASCT was undertaken
72 (first line versus second or subsequent line), and by remission status before HDT-ASCT. The
73 follow-up time was estimated using the inverse Kaplan-Meier method. All analyses are
74 considered exploratory in nature and were conducted using the software package R version
75 3.2.4 (www.r-project.org).

76 **Results**

77 *Patient characteristics*

78 Fifty-two eligible PCNSL patients who were treated with HDT-ASCT between 2004 and 2016
79 were included. Patients' baseline characteristics at time of diagnosis and before HDT-ASCT
80 are summarized in **Table 1**. Forty-eight patients had parenchymal disease manifestation
81 (with or without involvement of deep brain lesions) and four patients had primary vitreoretinal
82 lymphoma without parenchymal manifestations. Prior to HDT-ASCT, most patients had a
83 good clinical performance status (median Karnofsky performance status 80%, range 30% to
84 100%). Eighteen of 52 patients (34.6%) were in CR, 27 of 52 (51.9%) in PR, 2 of 52 (3.8%)
85 had SD, and 5 of 52 (9.6%) had experienced PD following induction treatment.

86 The majority of patients had previously received a HD-MTX-based protocol (98.1%) as first
87 line therapy with 38 of 52 patients (73.1%) having received MTX-AraC-based
88 polychemotherapy, 12 of 52 patients (23.1%) MTX-based polychemotherapy, one patient
89 MTX-monotherapy and another patient MTX-free polychemotherapy. Rituximab was added
90 in 33 of 52 patients (63.5%). Most of the 37 patients with relapsed or refractory disease after
91 first-line therapy received a polychemotherapy of carboplatin, ifosfamide and etoposide with
92 or without rituximab or a AraC - and thiotepa-based salvage regimen. None of the patients
93 had received WBRT before HDT-ASCT; one patient received WBRT as salvage therapy after
94 HDT-ASCT. Five patients received intrathecal therapy with liposomal cytarabine. None of the
95 patients received intraventricular therapy.

96

97 *Conditioning regimen and ASCT-specific data*

98 The majority of patients was conditioned with TT 10-20 mg/kg + carmustine 320-400 mg/m²
99 (61.5%). Remaining patients received TT 10-20 mg/kg + busulfan 3.2-6.4 mg/kg (13.5%), TT
100 250-750 mg/m² + busulfan 2.4–8.0 mg/kg + cyclophosphamide 60-120 mg/kg (13.5%) or TT
101 10 mg/kg as single agent (11.5%). Rituximab was additionally given in 2 patients (3.8%). The
102 median number of reinfused CD34+ hematopoietic stem cells was 5.29 x 10⁶/Kg (range 2.24
103 to 35). Median time to neutrophil engraftment was 10 days (range 6 to 34). We observed two

104 (3.8%) treatment related deaths within 15 days after HDT-ASCT. One patient suffered from
105 sudden death attributed to an acute cardiovascular event whilst the second patient died from
106 infectious complications.

107

108 *Treatment response and survival*

109 **Table 2** summarizes the response status after HDT-ASCT, stratified by line of therapy and
110 remission status prior to HDT-ASCT. Following HDT-ASCT, 45 of 52 patients (86.5%)
111 achieved an objective response (36 CR and 7 PR). One patient (1.9%) experienced SD
112 whilst 5 (9.6%) had PD one month after HDT-ASCT.

113 Of the 15 patients undergoing HDT-ASCT as first-line treatment, 14 patients (93%) achieved
114 an objective response (11 CR and 1 PR). Of the 37 patients undergoing HDT-ASCT as
115 second or subsequent line of treatment 31 patients (83.8%) achieved an objective response.

116 One of five patients with PD before HDT-ASCT achieved ongoing CR (PFS 50 months)
117 without further consolidating treatment. Three of the other four patients achieved partial
118 remission and one patient had stable disease, although all four subsequently experienced
119 PD.

120 After a median follow up of 22.1 months, 36 of 52 patients (69.2%) were still alive with 31
121 free of disease progression after HDT-ASCT. For one patient who experienced PD, the exact
122 date of death was not known, therefore the patient was censored at the date of progression
123 for the OS analysis. Apart from the two patients suffering from treatment related deaths
124 thirteen patients died from progressive disease whereas one patient was lost of follow up
125 after experiencing progressive disease with the exact cause of death being unknown.

126 For the entire cohort, median PFS and OS were reached after 51.1 and 122.3 months,
127 respectively. 2-year PFS and OS probabilities were 62.0% (95% CI 48.4% to 79.6%) and
128 70.8% (95% CI 58.3% to 85.9%), respectively (**Figure 1 and 2**). For the patients undergoing
129 HDT-ASCT as first-line treatment 2-year PFS and OS probabilities were 80% (95% CI 51.6%
130 to 100%) and 80.0% (95% CI 51.6% to 100%), respectively (**Figure 3**). For the patients
131 undergoing HDT-ASCT at second or subsequent line of treatment 2-year PFS and OS

132 probabilities were 54.0% (95% CI 39.1% to 74.4%) and 65.6% (95% CI 51.5% to 83.6%)
133 respectively (**Figure 4**). The PFS and OS rates by remission status before HDT-ASCT are
134 shown in **online figures 1 and 2** suggesting that patients with chemosensitive disease have
135 a better prognosis. Because of the limited patient numbers and events, we did not conduct
136 any statistical testing.

137

138

139 **Discussion**

140 We herein describe outcomes of elderly PCNSL patients who underwent thiotepa-based
141 HDT-ASCT. The overall response rate after HDT-ASCT was 86.5%, with 2-year PFS and OS
142 rates of 62.0% and 70.8%, respectively. Two patients (3.8%) died early of HDT-ASCT-
143 related causes.

144 *Strengths and limitations*

145 To the best of our knowledge, this is the first cohort reporting data on elderly patients who
146 underwent HDT-ASCT for PCNSL. Considering the rarity of the disease, the cohort size is
147 relatively large. Moreover, the dataset has a very low number of missing values and all
148 patients underwent relatively homogenous conditioning with thiotepa-based HDT-ASCT
149 protocols.

150 We recognise that our study has limitations, the first of which is inherent to any transplant
151 analysis; patients were only included if they underwent HDT-ASCT and we are not able to
152 report outcomes on an intent-to-treat basis. Thus, one can only use these outcome data to
153 inform patients about prognosis following HDT-ASCT. Second, we do not have detailed
154 information on relevant co-morbidities that would allow calculation of specific indices that
155 may help to standardise selection of older patients for HDT-ASCT.

156 *Comparison to other studies*

157 WBRT is still employed as a common treatment modality in some countries (7) for elderly
158 PCNSL patients, even though such patients are particularly vulnerable to iatrogenic toxicity,
159 especially neuro-cognitive dysfunction following WBRT (4). Addition of WBRT after
160 methotrexate-based chemotherapy is known to increase the risk of treatment-related
161 neurotoxicity (16). Importantly, in our cohort, none of the patients received consolidating
162 WBRT after HDT-ASCT. Although neurocognitive function was not formally assessed in our
163 cohort, the merits of avoiding WBRT in older PCNSL patients is absolutely clear, particularly
164 given these promising outcomes following HDT-ASCT.

165 Based on a recent systematic review (9), a limited number of prospective multicentre studies
166 focusing on elderly PCNSL patients have been reported (8, 17-21). Most of the studies

167 included HD-MTX in combination with partner chemotherapy agents, but to-date no standard
168 protocol has been defined. Acknowledging the limitations of inter-trial comparison, the best
169 reported response rate was 79% but the corresponding 1-year PFS was only 36% (8). With
170 an overall response rate of 86.5% and a 2-year-PFS of 62% the outcome of our reported
171 population compares favourably with all other trials conducted in this unique subgroup of
172 PCNSL patients. This is even more significant as only less than one third of the present
173 cohort underwent HDT-ASCT in CR. Remarkably, even some patients with chemotherapy-
174 refractory disease achieved sustained objective responses after HDT-ASCT although we
175 acknowledge inherent selection bias within our cohort, likely to be related to favourable
176 performance status and limited co-morbidities.

177 Although experience with HDT-ASCT in PCNSL is limited to prospective non-randomised
178 studies in consolidation of first line therapy or for relapsed patients < 65 years, the results are
179 encouraging; particularly when TT-containing conditioning regimens are used (10-12, 19,
180 22). In a multi-centre retrospective analysis investigating patients with a median age of 52.4
181 years undergoing HDT-ASCT as salvage therapy, 5-year survival rates of patients with
182 chemosensitive relapse was 62% (13). In another large retrospective analysis investigating
183 patients with a median age of 54 years undergoing HDT-ASCT as first line treatment, the
184 reported 2- and 5-year survival rates were 82% and 79%, respectively. Notably, of the
185 reported patients with PD before HDT-ASCT, 7/20 achieved ongoing CR without further
186 treatment suggesting efficacy of HDT-ASCT even in disease refractory to conventionally
187 dosed chemotherapy (23). The thus far published studies on ASCT as first-line treatment or
188 at relapse are summarized in **Table 3**. Acknowledging the limitations of inter-trial
189 comparisons, survival rates of the herein reported elderly population seem to be comparable
190 to those of younger patients.

191 Patients with newly diagnosed PCNSL, aged between 65 and 70 years with ECOG
192 performance status 0-2 have been included in the international, randomised, phase II
193 IELSG32 trial (24), but evidence from prospective clinical trials specifically designed for
194 elderly patients are still lacking.

195 All patients in our cohort received thiotepa-based regimens incorporating a total thiotepa
196 dose of 10-20 mg/kg. In an ongoing German pilot study investigating feasibility of HDT-ASCT
197 in elderly patients, the conditioning regimen comprises busulfan 3.2 mg/kg and thiotepa 10
198 mg/kg (half of the dose routinely administered to younger patients) (DRKS-ID 00008900).
199 Importantly, even for younger patient cohorts, randomised comparisons of different
200 myeloablative combinations and doses have not been conducted; for elderly PCNSL
201 patients, no such data is available.

202 For systemic diffuse large B-cell lymphoma (DLBCL), the current role of HDT-ASCT is
203 restricted to relapsed patients responding to salvage therapy. Notably, the majority of HDT-
204 ASCT studies in this context include younger patients with a median age of 54 years (25).
205 Notwithstanding increasing clinical experience of undertaking HDT-ASCT for older patients
206 with systemic DLBCL, there remains no clear standard for the selection of, or conditioning
207 for, elderly patients undergoing HDT-ASCT.

208 Few data on feasibility and efficacy of HDT-ASCT in elderly patients are available in multiple
209 myeloma (MM), lymphoma and acute leukaemia (26-28). The reported non relapse mortality
210 rates in lymphoma patients older than 70 years who underwent HDT-ASCT differ strongly
211 from 5.2% up to 19% (29-31). In our cohort, only two patients (3.8%) died from treatment
212 related mortality, both within 15 days after HDT-ASCT due to a cardiovascular event and
213 infectious complications.

214 The definition of an elderly patient with regard to therapeutic stratification is unclear;
215 determined by multiple patient- and disease-related parameters. Thus, 'elderly' patients
216 comprise a markedly heterogeneous group and it is unclear how to optimally define the frailty
217 profile in this context. Comorbidity risk scoring, assessment of instrumental activities of daily
218 living, and comprehensive geriatric assessments are likely to be important tools to define
219 treatment-related mortality and overall treatment risk (32-35). To-date, treatment decisions
220 have been largely based on chronological age and performance status. Although
221 standardized assessment scores, especially for cancer patients, are available (36), these are
222 infrequently used because of their complexity. Recently, a score for the quantitation of frailty

223 in designing future clinical MM trials was proposed (37). There is a clear need for a simple
224 and validated tool to inform treatment decisions in elderly PCNSL.

225 *Conclusions*

226 In selected elderly PCNSL patients, HDT-ASCT, using thiotepa-based conditioning regimens,
227 is an effective and safe treatment if conducted at experienced centers, both in first-line and
228 second or subsequent line of treatment. A pilot study investigating feasibility and efficacy of
229 HDT-SCT in PCNSL patients > 65 years is currently recruiting (DRKS-ID 00008900).
230 Prospective trials are needed to better define eligibility for this approach and to further
231 improve therapeutic approaches in this unique and challenging subgroup of patients.

232

233 **Acknowledgements**

234 The authors would like to acknowledge with thanks the contributions of all participating
235 centres.

236

237 **Conflict of Interest**

238 The authors declare no conflict of interest.

239

240

References

1. Panageas KS, Elkin EB, DeAngelis LM, Ben-Porat L, Abrey LE. Trends in survival from primary central nervous system lymphoma, 1975-1999: a population-based analysis. *Cancer*. 2005;104(11):2466-72.
2. Olson JE, Janney CA, Rao RD, Cerhan JR, Kurtin PJ, Schiff D, et al. The continuing increase in the incidence of primary central nervous system non-Hodgkin lymphoma: a surveillance, epidemiology, and end results analysis. *Cancer*. 2002;95(7):1504-10.
3. Makino K, Nakamura H, Kino T, Takeshima H, Kuratsu J. Rising incidence of primary central nervous system lymphoma in Kumamoto, Japan. *Surgical neurology*. 2006;66(5):503-6.
4. Abrey LE, Yahalom J, DeAngelis LM. Treatment for primary CNS lymphoma: the next step. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2000;18(17):3144-50.
5. Sierra del Rio M, Rousseau A, Soussain C, Ricard D, Hoang-Xuan K. Primary CNS lymphoma in immunocompetent patients. *The oncologist*. 2009;14(5):526-39.
6. Jahnke K, Korfel A, Martus P, Weller M, Herrlinger U, Schmitt A, et al. High-dose methotrexate toxicity in elderly patients with primary central nervous system lymphoma. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO*. 2005;16(3):445-9.
7. Panageas KS, Elkin EB, Ben-Porat L, Deangelis LM, Abrey LE. Patterns of treatment in older adults with primary central nervous system lymphoma. *Cancer*. 2007;110(6):1338-44.
8. Omuro A, Chinot O, Taillandier L, Ghesquieres H, Soussain C, Delwail V, et al. Methotrexate and temozolomide versus methotrexate, procarbazine, vincristine, and cytarabine for primary CNS lymphoma in an elderly population: an intergroup ANOCEF-GOELAMS randomised phase 2 trial. *The Lancet Haematology*. 2015;2(6):e251-9.
9. Kasenda B, Ferreri AJ, Marturano E, Forst D, Bromberg J, Ghesquieres H, et al. First-line treatment and outcome of elderly patients with primary central nervous system lymphoma (PCNSL)-a systematic review and individual patient data meta-analysis. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO*. 2015;26(7):1305-13.
10. Illerhaus G, Marks R, Ihorst G, Guttenberger R, Ostertag C, Derigs G, et al. High-dose chemotherapy with autologous stem-cell transplantation and hyperfractionated radiotherapy as first-line treatment of primary CNS lymphoma. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2006;24(24):3865-70.
11. Illerhaus G, Muller F, Feuerhake F, Schafer AO, Ostertag C, Finke J. High-dose chemotherapy and autologous stem-cell transplantation without consolidating radiotherapy as first-line treatment for primary lymphoma of the central nervous system. *Haematologica*. 2008;93(1):147-8.
12. Omuro A, Correa DD, DeAngelis LM, Moskowitz CH, Matasar MJ, Kaley TJ, et al. R-MPV followed by high-dose chemotherapy with TBC and autologous stem-cell transplant for newly diagnosed primary CNS lymphoma. *Blood*. 2015;125(9):1403-10.
13. Soussain C, Choquet S, Fourme E, Delgadillo D, Bouabdallah K, Ghesquieres H, et al. Intensive chemotherapy with thiotepa, busulfan and cyclophosphamide and hematopoietic stem cell rescue in relapsed or refractory primary central nervous system lymphoma and intraocular lymphoma: a retrospective study of 79 cases. *Haematologica*. 2012;97(11):1751-6.

14. Soussain C, Hoang-Xuan K, Taillandier L, Fourme E, Choquet S, Witz F, et al. Intensive chemotherapy followed by hematopoietic stem-cell rescue for refractory and recurrent primary CNS and intraocular lymphoma: Societe Francaise de Greffe de Moelle Osseuse-Therapie Cellulaire. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2008;26(15):2512-8.
15. Illerhaus G KB, Ihorst G, Egerer G, Lamprecht M, Keller U, et al. High-dose chemotherapy with autologous haemopoietic stem cell transplantation for newly diagnosed primary CNS lymphoma: a prospective, single-arm, phase 2 trial. *The Lancet Haematology*. 2016;in press.
16. Doolittle ND, Korfel A, Lubow MA, Schorb E, Schlegel U, Rogowski S, et al. Long-term cognitive function, neuroimaging, and quality of life in primary CNS lymphoma. *Neurology*. 2013;81(1):84-92.
17. Ghesquieres H, Ferlay C, Sebban C, Perol D, Bosly A, Casasnovas O, et al. Long-term follow-up of an age-adapted C5R protocol followed by radiotherapy in 99 newly diagnosed primary CNS lymphomas: a prospective multicentric phase II study of the Groupe d'Etude des Lymphomes de l'Adulte (GELA). *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO*. 2010;21(4):842-50.
18. Hoang-Xuan K, Taillandier L, Chinot O, Soubeyran P, Bogdhan U, Hildebrand J, et al. Chemotherapy alone as initial treatment for primary CNS lymphoma in patients older than 60 years: a multicenter phase II study (26952) of the European Organization for Research and Treatment of Cancer Brain Tumor Group. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2003;21(14):2726-31.
19. Pulczynski EJ, Kuittinen O, Erlanson M, Hagberg H, Fossa A, Eriksson M, et al. Successful change of treatment strategy in elderly patients with primary central nervous system lymphoma by de-escalating induction and introducing temozolomide maintenance: results from a phase II study by the Nordic Lymphoma Group. *Haematologica*. 2015;100(4):534-40.
20. Laack NN, Ballman KV, Brown PB, O'Neill BP, North Central Cancer Treatment G. Whole-brain radiotherapy and high-dose methylprednisolone for elderly patients with primary central nervous system lymphoma: Results of North Central Cancer Treatment Group (NCCTG) 96-73-51. *International journal of radiation oncology, biology, physics*. 2006;65(5):1429-39.
21. Olivier G, Clavert A, Lacotte-Thierry L, Gardembas M, Escoffre-Barbe M, Brion A, et al. A phase 1 dose escalation study of idarubicin combined with methotrexate, vindesine, and prednisolone for untreated elderly patients with primary central nervous system lymphoma. The GOELAMS LCP 99 trial. *American journal of hematology*. 2014;89(11):1024-9.
22. Kasenda B, Schorb E, Fritsch K, Finke J, Illerhaus G. Prognosis after high-dose chemotherapy followed by autologous stem-cell transplantation as first-line treatment in primary CNS lymphoma--a long-term follow-up study. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO*. 2015;26(3):608-11.
23. Schorb E, Kasenda B, Atta J, Kaun S, Morgner A, Hess G, et al. Prognosis of patients with primary central nervous system lymphoma after high-dose chemotherapy followed by autologous stem cell transplantation. *Haematologica*. 2013;98(5):765-70.
24. Ferreri AJ, Cwynarski K, Pulczynski E, Ponzoni M, Deckert M, Politi LS, et al. Chemoimmunotherapy with methotrexate, cytarabine, thiotepa, and rituximab (MATRix regimen) in patients with primary CNS lymphoma: results of the first randomisation of the International Extranodal Lymphoma Study Group-32 (IELSG32) phase 2 trial. *The Lancet Haematology*. 2016;3(5):e217-27.

25. Gisselbrecht C, Glass B, Mounier N, Singh Gill D, Linch DC, Trneny M, et al. Salvage regimens with autologous transplantation for relapsed large B-cell lymphoma in the rituximab era. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2010;28(27):4184-90.
26. Auner HW, Szydlo R, Hoek J, Goldschmidt H, Stoppa AM, Morgan GJ, et al. Trends in autologous hematopoietic cell transplantation for multiple myeloma in Europe: increased use and improved outcomes in elderly patients in recent years. *Bone marrow transplantation*. 2015;50(2):209-15.
27. Jantunen E, Canals C, Rambaldi A, Ossenkoppele G, Allione B, Blaise D, et al. Autologous stem cell transplantation in elderly patients (> or =60 years) with diffuse large B-cell lymphoma: an analysis based on data in the European Blood and Marrow Transplantation registry. *Haematologica*. 2008;93(12):1837-42.
28. Thomas X, Suciú S, Rio B, Leone G, Broccia G, Fillet G, et al. Autologous stem cell transplantation after complete remission and first consolidation in acute myeloid leukemia patients aged 61-70 years: results of the prospective EORTC-GIMEMA AML-13 study. *Haematologica*. 2007;92(3):389-96.
29. Andorsky DJ, Cohen M, Naeim A, Pinter-Brown L. Outcomes of auto-SCT for lymphoma in subjects aged 70 years and over. *Bone marrow transplantation*. 2011;46(9):1219-25.
30. Elstrom RL, Martin P, Hurtado Rua S, Shore TB, Furman RR, Ruan J, et al. Autologous stem cell transplant is feasible in very elderly patients with lymphoma and limited comorbidity. *American journal of hematology*. 2012;87(4):433-5.
31. Chihara D, Izutsu K, Kondo E, Sakai R, Mizuta S, Yokoyama K, et al. High-dose chemotherapy with autologous stem cell transplantation for elderly patients with relapsed/refractory diffuse large B cell lymphoma: a nationwide retrospective study. *Biology of blood and marrow transplantation : journal of the American Society for Blood and Marrow Transplantation*. 2014;20(5):684-9.
32. Morrison VA, Hamlin P, Soubeyran P, Stauder R, Wadhwa P, Aapro M, et al. Approach to therapy of diffuse large B-cell lymphoma in the elderly: the International Society of Geriatric Oncology (SIOG) expert position commentary. *Annals of Oncology*. 2015.
33. Wildes TM, Augustin KM, Sempek D, Zhang QJ, Vij R, Dipersio JF, et al. Comorbidities, not age, impact outcomes in autologous stem cell transplant for relapsed non-Hodgkin lymphoma. *Biology of blood and marrow transplantation : journal of the American Society for Blood and Marrow Transplantation*. 2008;14(7):840-6.
34. Charlson M, Szatrowski TP, Peterson J, Gold J. Validation of a combined comorbidity index. *Journal of clinical epidemiology*. 1994;47(11):1245-51.
35. Hurria A, Togawa K, Mohile SG, Owusu C, Klepin HD, Gross CP, et al. Predicting chemotherapy toxicity in older adults with cancer: a prospective multicenter study. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*. 2011;29(25):3457-65.
36. Hamaker ME, Jonker JM, de Rooij SE, Vos AG, Smorenburg CH, van Munster BC. Frailty screening methods for predicting outcome of a comprehensive geriatric assessment in elderly patients with cancer: a systematic review. *The Lancet Oncology*. 2012;13(10):e437-44.
37. Palumbo A, Brinchen S, Mateos M-V, Larocca A, Facon T, Kumar SK, et al. Geriatric assessment predicts survival and toxicities in elderly myeloma: an International Myeloma Working Group report. *Blood*. 2015.
38. Abrey LE, Moskowitz CH, Mason WP, Crump M, Stewart D, Forsyth P, et al. Intensive methotrexate and cytarabine followed by high-dose chemotherapy with

- autologous stem-cell rescue in patients with newly diagnosed primary CNS lymphoma: An intent-to-treat analysis. *Journal of Clinical Oncology*. 2003;21(22):4151-6.
39. Colombat P, Lemevel A, Bertrand P, Delwail V, Rachieru P, Brion A, et al. High-dose chemotherapy with autologous stem cell transplantation as first-line therapy for primary CNS lymphoma in patients younger than 60 years: a multicenter phase II study of the GOELAMS group. *Bone marrow transplantation*. 2006;38(6):417-20.
 40. Montemurro M, Kiefer T, Schuler F, Al-Ali HK, Wolf HH, Herbst R, et al. Primary central nervous system lymphoma treated with high-dose methotrexate, high-dose busulfan/thiotepa, autologous stem-cell transplantation and response-adapted whole-brain radiotherapy: results of the multicenter Ostdeutsche Studiengruppe Hamato-Onkologie OSHO-53 phase II study. *Annals of Oncology*. 2007;18(4):665-71.
 41. Miyao K, Sakemura R, Imai K, Sakai T, Tsushita N, Kato T, et al. Upfront autologous stem-cell transplantation with melphalan, cyclophosphamide, etoposide, and dexamethasone (LEED) in patients with newly diagnosed primary central nervous system lymphoma. *Int J Hematol*. 2014;100(2):152-8.
 42. Kasenda B, Schorb E, Fritsch K, Finke J, Illerhaus G. Prognosis after high-dose chemotherapy followed by autologous stem-cell transplantation as first-line treatment in primary CNS lymphoma--a long-term follow-up study. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO*. 2012;23(10):2670-5.
 43. Yoon DH, Lee DH, Choi DR, Sohn BS, Kim S, Kim SW, et al. Feasibility of BU, CY and etoposide (BUCYE), and auto-SCT in patients with newly diagnosed primary CNS lymphoma: a single-center experience. *Bone marrow transplantation*. 2011;46(1):105-9.
 44. Brevet M, Garidi R, Gruson B, Royer B, Vaida I, Damaj G. First-line autologous transplantation in stem cell primary CNS lymphoma. *European journal of haematology*. 2005;75(4):288-92.
 45. Alimohamed N, Daly A, Owen C, Duggan P, Stewart DA. Upfront thiotepa, busulfan, cyclophosphamide, and autologous stem cell transplantation for primary CNS lymphoma: a single centre experience. *Leukemia & lymphoma*. 2012;53(5):862-7.
 46. Choi MK, Kang ES, Kim DW, Ko YH, Seok H, Park JH, et al. Treatment outcome of relapsed/refractory primary central nervous system diffuse large B-cell lymphoma: a single-center experience of autologous stem cell transplantation. *Int J Hematol*. 2013;98(3):346-54.
 47. Soussain C, Suzan F, Hoang-Xuan K, Cassoux N, Levy V, Azar N, et al. Results of intensive chemotherapy followed by hematopoietic stem-cell rescue in 22 patients with refractory or recurrent primary CNS lymphoma or intraocular lymphoma. *Journal of Clinical Oncology*. 2001;19(3):742-9.
 48. Welch MR, Sauter CS, Matasar MJ, Faivre G, Weaver SA, Moskowitz CH, et al. Autologous stem cell transplant in recurrent or refractory primary or secondary central nervous system lymphoma using thiotepa, busulfan and cyclophosphamide. *Leukemia & lymphoma*. 2015;56(2):361-7.

Patient characteristics	HDT applied in first line N=15	HDT applied in second/subsequent line N=37	Total N=52
Age at HDT-ASCT median (range)	70 (66, 75)	67 (65, 77)	68.5 (65, 77)
Sex			
female	7 (46.7)	17 (46)	24 (46.2)
male	8 (53.3)	20 (54)	28 (53.8)
KPS at diagnosis (median range)	65 (30, 90)	80 (40, 100)	70 (30, 100)
LDH			
elevated	8 (53.3)	10 (27)	18 (34.6)
not elevated	6 (40)	18 (48.7)	24 (46.2)
unknown	1 (6.7)	9 (24.3)	10 (19.2)
Deep brain structures involved			
yes	10 (66.7)	22 (59.5)	32 (61.5)
no	5 (33.3)	15 (40.5)	20 (38.5)
Ocular involvement			
yes	0 (0)	8 (21.6)	8 (15.4)
no	14 (93.3)	21 (56.8)	35 (67.3)
unknown	1 (6.7)	8 (21.6)	9 (17.3)
Leptomeningeal involvement			
yes		5 (13.5)	5 (9.6)
no	5 (33.3)	20 (54.1)	25 (67.6)
unknown	10 (66.7)	12 (32.4)	22 (42.3)
Histology			
Aggressive B-NHL	15 (100)	34 (91.9)	49 (94.2)
T-NHL	0 (0)	3 (8.1)	3 (5.8)
HDT-ASCT applied in			
1st line	15 (100)		15 (28.8)
2nd line		32 (86.5)	32 (61.5)
3rd line		5 (13.5)	5 (9.6)
KPS before HDT-ASCT median (IQR)	70 (70, 80)	80 (70, 90)	80 (70, 90)
Remission before HDT-ASCT			
CR	4 (26.7)	14 (37.8)	18 (34.6)
PR	11 (73.3)	16 (43.3)	27 (51.9)
SD	0 (0)	2 (5.4)	2 (3.8)
PD	0 (0)	5 (13.5)	5 (9.6)

Table 1: Patients' characteristics. Numbers are frequencies (percentage) unless specified otherwise. Remission status according to IPCG Response criteria

Abbreviations: IQR, interquartile range; KPS, Karnofsky Performance Status; Serum LDH, serum lactate dehydrogenase level, HDT-ASCT, high-dose chemotherapy followed by autologous stem cell transplantation; CR, complete remission, PR, partial remission, SD, stable disease, PD, progressive disease; IPCG, International PCNSL Collaborative Group

		Remission status before HDT-ASCT (1st line setting)				Remission status before HDT-ASCT (2nd or later line setting)			
		CR n=4	PR n=11	SD n=0	PD n=0	CR n=14	PR n=16	SD n=2	PD n=5
Remission status after HDT- ASCT	CR	4	7	0	0	12	11	1	1
	PR	0	3	0	0	0	3	0	3
	SD	0	0	0	0	0	0	0	1
	PD	0	1	0	0	1	2	1	0
	not applicable	0	0	0	0	1	0	0	0

Table 2: Remission status stratified by line of treatment and remission status before HDT-ASCT. In the first line setting, 3/3 patients with PR after HDT-ASCT only had minimal contrast abnormality in brain imaging after HDT-ASCT likely in terms of reactive post-therapy lesions and remained in confirmed complete remission for > 6 months after HDT-ASCT without any additional therapy. In the second or later line setting, only 1/6 patient with PR after HDT-ASCT remained in confirmed remission > 6 months after HDT-ASCT without any additional therapy.

Abbreviations: HDT-ASCT, high dose chemotherapy followed by autologous stem cell transplantation;

CR, complete remission; PR, partial remission; SD, stable disease; PD, progressive disease

<i>N° of patients</i>	<i>Trial design</i>	<i>median age (range)</i>	<i>HD-ASCT setting</i>	<i>Induction regimen</i>	<i>Conditioning regimen</i>	<i>WBRT</i>	<i>Follow-up (median)</i>	<i>OS</i>	<i>TRM</i>	<i>Reference</i>
33	prospective	57 (23-67)	1st line	R-MPV	Bu/TT/Cy	no	45	3 y: 81%	12%	(12)
30	prospective	54 (27-64)	1st line	HD-MTX+AraC/TT	BCNU/TT	yes	63	5 y: 69%	3%	(10)
28	prospective	53 (25-71)	1st line	HD-MTX+AraC	BEAM	no	28	2 y: 55%	4%	(38)
25	prospective	51 (21-60)	1st line	MBVP + IFO/AraC	BEAM	yes	34	4 y: 64%	4%	(39)
23	prospective	55 (18-69)	1st line	HD-MTX	Bu/TT	only if no CR	15	2 y: 48%	13%	(40)
13	prospective	56 (35-65)	1st line	MPV+AraC	LEED	only if no CR	44	3 y: 76%	0%	(41)
13	prospective	54 (38-67)	1st line	HD-MTX + AraC/TT	BCNU/TT	only if no CR	72	5 y: 77%	0%	(11) (42)
11	prospective	52 (33-65)	1st line	HD-MTX + AraC	BUCYE	only if no CR	25	2 y: 89%	0%	(43)
6	prospective	53 (30-66)	1st line	MBVP + IFO/AraC	BEAM	yes	41	2 y: 40%	0%	(44)
105	retrospective	54 (23-70)	1st line	mostly MTX-based polychemotherapy	mostly BCNU/TT	partly	47	5y: 79%	3%	(23)
21	retrospective	56 (34-69)	1st line	MPV + AraC	Bu/TT/Cy	no	60	5 y: 44%	24%	(45)
45	prospective	57 (19-72)	rel/ref	ICE or HD-MTX	Bu/TT	no	53	5 y: 40%	5%	(46)
43	prospective	52 (23-65)	rel/ref	AraC /VP16	Bu/TT/Cy	no	36	2 y: 45%	12%	(14)
22	prospective	53 (27-64)	rel/ref	AraC/VP16	Bu/TT/Cy	no	41	3 y: 64%	4%	(47)
79	retrospective	52 (23-67)	rel/ref	mostly AraC/VP16	TT/Bu/Cy	yes	56	5 y: 51%	8%	(13)
7	retrospective	58 (41-65)	rel/ref	unknown	TT/Bu/Cy	yes	34	not reached	0%	(48)

Table 3: Compendium of publications on HDT-ASCT in PCNSL

Abbreviations: HDT-ASCT, high dose chemotherapy followed by autologous stem cell transplantation; WBRT, whole brain radiation therapy; OS, overall survival; TRM, treatment related mortality; MPV, methotrexate, procarbazine, vincristine; HD-MTX, high-dose methotrexate; AraC, cytarabine; TT, thiotepa; MBVP, methotrexate, carmustine, etoposide, methyprednisolone; ifo, ifosfamide; ICE, ifosfamide, carboplatin, etoposide; VP16, etoposide; Bu, busulfan; BCNU, carmustine; BEAM, carmustine, etoposide, cytarabine, melphalan, LEED, cyclophosphamide, etoposide, melphalan, dexamethasone; BuCy, busulfan, cyclophosphamide; Cy, cyclophosphamide

Figure 1

PFS entire cohort




Figure 2

OS entire cohort





Figure 3

PFS in patients with HCT-ASCT at 1st line



OS in patients with HCT-ASCT at 1st line






Figure 4

PFS in patients with HCT-ASCT at 2nd or later line




OS in patients with HCT-ASCT at 2nd or later line




Online Figure 1

PFS stratified by remission before HCT-ASCT



Online Figure 2

OS stratified by remission before HCT-ASCT



Legend to Figures

Figure 1 Kaplan-Meier plot: Progression Free Survival from time of HDT-ASCT of all evaluable patients

Figure 2 Kaplan-Meier plot: Overall Survival from time of HDT-ASCT of all patients

Figure 3 Kaplan-Meier plot: Progression Free Survival and Overall Survival from time of HDT-ASCT for patients receiving HDT-ASCT during first-line therapy

Figure 4 Kaplan-Meier plot: Progression Free Survival and Overall Survival from time of HDT-ASCT for patients receiving HDT-ASCT at second or later line of treatment

Online Figure 1 Kaplan-Meier plot: Progression Free Survival from time of HDT-ASCT of all patients stratified by remission status before HDT-ASCT

Online Figure 2 Kaplan-Meier plot: Overall Survival from time of HDT-ASCT of all patients stratified by remission status before HDT-ASCT