



HAL
open science

Nuclear Spin Symmetry Conservation in $1\text{H}_2\ 16\text{O}$ Investigated by Direct Absorption FTIR Spectroscopy of Water Vapor Cooled Down in Supersonic Expansion

Robert Georges, X. Michaut, A. Moudens, M. Goubet, O. Pirali, P. Soulard, P. Asselin, T. Huet, Pascal Roy, Martin Fournier, et al.

► **To cite this version:**

Robert Georges, X. Michaut, A. Moudens, M. Goubet, O. Pirali, et al.. Nuclear Spin Symmetry Conservation in $1\text{H}_2\ 16\text{O}$ Investigated by Direct Absorption FTIR Spectroscopy of Water Vapor Cooled Down in Supersonic Expansion. *Journal of Physical Chemistry A*, 2017, 121 (40), pp.7455-7468. 10.1021/acs.jpca.7b06858 . hal-01638856

HAL Id: hal-01638856

<https://univ-rennes.hal.science/hal-01638856>

Submitted on 25 Jan 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Nuclear Spin Symmetry Conservation in $^1\text{H}_2^{16}\text{O}$ Investigated by Direct Absorption FTIR Spectroscopy of Water Vapor Cooled Down in Supersonic Expansion

Robert Georges, *,† Xavier Michaut, ‡ Audrey Moudens, ‡ Manuel Goubet, ¶ Olivier
Pirali, § Pascale Soulard, || Pierre Asselin, || Thérèse Huet, ¶ Pascale Roy, § Martin
Fournier, † and Andrey Vigasin ⊥

†*Institut de Physique de Rennes, UMR6251, CNRS-Université de Rennes 1, Campus de
Beaulieu, 35042 Rennes, France*

‡*Laboratoire d'Etudes du Rayonnement et de la Matière en Astrophysique et Atmosphères,
Sorbonne Universities, UPMC Univ. Paris 06, Université de Cergy-Pontoise, Observatoire
de Paris, PSL Research University, CNRS, F-75252, Paris, France*

¶*Université Lille, CNRS, UMR8523-PhLAM - Physique des Lasers, Atomes et Molécules,
F-59000 Lille, France*

§*Ligne AILES-Synchrotron SOLEIL, L'Orme des Merisiers, 91192 Gif-sur-Yvette, France*

||*Sorbonne Universités, UPMC Univ Paris 06, CNRS, UMR 8233, MONARIS,
F-75005, Paris, France.*

⊥*Obukhov Institute of Atmospheric Physics, Russian Academy of Sciences, 3 Pyzhevsky
per, 119017 Moscow, Russia*

E-mail: robert.georges@univ-rennes1.fr

Abstract

We report the results of an experimental study related to the relaxation of the nuclear spin isomers of the water molecule in a supersonic expansion. Rovibrational lines of both *ortho*- and *para*- spin isomers were recorded in the spectral range of H₂O stretching vibrations around 3700 cm⁻¹ using FTIR direct absorption. Water vapor seeded either in argon, helium or oxygen or in a mixture of oxygen and argon was expanded into vacuum through a slit nozzle. The water vapor partial pressure in the mixture varied over a wide range from 1.5 to 102.7 hPa, corresponding to a water molar fraction varying between 0.2 and 6.5%. Depending on expansion conditions, the effect of water vapor clustering was clearly seen in some of our measured spectra. The Boltzmann-plot of the line intensities allowed the H₂O rotational temperatures in the isentropic core and in the lateral shear layer probed zones of the planar expansion to be determined. The study of the *OPR*, i.e the ratio of the *ortho*- to *para*- absorption line intensities as a function of T_{rot} , did not reveal any signs of the *OPR* being relaxed to the sample temperature. In contrast, the *OPR* was always conserved according to the stagnation reservoir equilibrium temperature. The conservation of the *OPR* was found irrespective of whether water molecule clustering was pronounced or not. Also no effect of the paramagnetic oxygen admixture enhancing *OPR* relaxation was observed.

Introduction

Water is an asymmetric top molecule, and its rovibrational levels are conventionally labelled by $J_{K_a K_c}$, where J is the total rotational angular momentum quantum number and K_a and K_c are quantum numbers of the components of the total angular momentum on the principal axes a and c , respectively. These levels are assigned in the ground vibrational state to either *para* or *ortho* species for even or odd $K_a + K_c$, respectively. The notations are inverted in the antisymmetric vibrational stretch ν_3 state.

The two hydrogen nuclei in a water molecule have either parallel or anti-parallel spins.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Similar to the hydrogen molecule, the conversion between these two forms is strictly forbidden in an isolated water molecule and strongly improbable in the gas phase.¹ Water vapor can therefore, be thought to be a mixture of two stable spin-isomer forms: *ortho* and *para*. *Para*-isomers which have a total nuclear spin statistical weight g_{NS} equal to unity ($g_{\text{NS}} = 2I + 1$, where I is the total nuclear spin: $I_{\text{para}} = 0$, $I_{\text{ortho}} = 1$) are dominant in $^1\text{H}_2^{16}\text{O}$ water vapor at very low temperatures since the water molecule ground state belongs to the *para* sub-system of rotational energy levels. In contrast, thermal population of many rotational levels at elevated temperatures causes *ortho*-isomers to dominate in a mixture because of the higher nuclear spin statistical weight of *ortho* states. The limiting high temperature *ortho*-to-*para* ratio ($OPR = 3$) of the nuclear spin isomer abundances is already reached at temperatures in excess of ca. 50 K and it remains constant at higher temperature, including, of course, room temperature. Some experimental studies (see e.g.²) carried out on the supersonic cooling of water vapor seeded in Ar, relied on the assumption that $OPR = 3$ is preserved in the jet expansion.

An introduction to recent studies of nuclear spin isomer conversion in water vapor can be found in the paper from Manca Tanner *et al.*³ Much attention has been paid in the last few years (see e.g. refs 5-9) to the spectroscopic investigation of whether the normal OPR can be significantly distorted at near room temperature following multiple adsorption and desorption of water molecules onto the surface of various porous media such as adsorbent charcoal or polymer sorbents. Following initial observations dating back almost 30 years,⁴ there were more recent claims⁵ that an increase of OPR up to ten instead of three at near room temperature can be reached using a type of frontal gas-chromatograph column. Moreover, it was claimed in ref.⁵ that the non-equilibrium OPR ratio can be conserved over a long period (months) when a spin-isomer enriched sample is frozen. Once heated and released from the ice, the vapor is still in its non-equilibrium spin-isomer state. These experimental findings were then subject to a contradictory evaluation. There were attempts to reproduce experiments reported in ref.⁵ which yielded either at least partially positive^{6,7} or entirely

1
2
3
4 negative^{1,8} results. To summarize, it can be stated nowadays that at least the idea of very
5
6 slow spin-isomers conversion in the condensed phase has not met any independent confirma-
7
8 tion and has to be ruled out. In contrast, it has been shown that *ortho-para* conversion in
9
10 the condensed phase can occur on a timescale of between milliseconds⁹ and hours.¹⁰⁻¹³ In
11
12 general, the results obtained in ref.⁵ were characterized e.g. in ref.³ as having a “nonstandard
13
14 interpretation”. It was suggested in particular (see e.g.^{8,14}) that microwave radiation-induced
15
16 desorption might be responsible for the effect observed in ref.^{4,5} and that no *ortho-para* en-
17
18 richment occurred at all. This suggestion seems, however, unlikely since the experiments
19
20 with a diode laser spectrometer⁷ apparently have confirmed the phenomenon of *ortho-para*
21
22 enrichment detected previously in ref.⁵ with the use of a Backward Wave Oscillator (BWO).
23

24 The results reported in ref.³ showed a rapid change in the spin-isomer state in a su-
25
26 per-sonic expansion of $^1\text{H}_2^{16}\text{O}$ water vapor seeded in argon at a relatively high water molar
27
28 fraction. Whereas at a low water vapor molar fraction ($0.35\% < x < 0.9\%$) in a mixture
29
30 with the carrier gas, the nuclear spin symmetry was conserved, at higher water molar fracti-
31
32 ons ($0.9\% < x < 1.6\%$) the *OPR* was seen to be completely relaxed, i.e. the measured
33
34 *OPR* responded to a low rotational temperature in expansion. This fast relaxation cannot
35
36 be explained by intramolecular processes in an individual water molecule and was thus in-
37
38 terpreted as a result of water cluster formation which is able presumably to significantly
39
40 facilitate nuclear spin conversion (NSC). The rapid transition from nuclear spin symmetry
41
42 conservation to complete nuclear spin relaxation regimes within the narrow range of water
43
44 mole fractions that was observed in ref.,³ has not found a clear explanation up-to-now. An
45
46 attempt to reproduce the results reported in ref.³ was undertaken recently in ref.¹⁵ using a
47
48 diode laser spectroscopic examination of two water vapor *ortho-* and *para-* absorption lines
49
50 in an H_2O seeded in Ar expansion. Virtually no signs of nuclear spin relaxation was ob-
51
52 served irrespective of whether the clustering in an expansion was definitely present or not.
53
54 No correlation between clustering in the expansion and nuclear-spin isomerization was thus
55
56 observed.
57
58
59
60

1
2
3
4 The idea of water clusters promoting the nuclear spin relaxation process that was put
5 forward in ref.,³ has much in common with an assumption made initially in ref.¹⁶ where
6 preferred clustering of heavy D₂O water molecules from their ground *ortho*-state was first
7 suggested. In their thorough recent study, Vilesov *et al.*¹¹ assumed to have formed small
8 water clusters predominantly made up of *para*-H₂O isomers using water embedded in an
9 Ar matrix at low temperature. By waiting for 1400 minutes for the complete nuclear spin
10 conversion of water molecules in the Ar matrix at 4 K, they produced nearly 100 % of
11 *para* molecules in solid Ar because the mostly populated energetic level at the considered
12 temperature, is the lowest rotational energy level of the *para* state (i.e. $J_{K_a K_c} = 0_{00}$ state
13 in the vibrational ground state). A fast warming up to 50 K of the matrix resulted in a
14 rapid evaporation of Ar atoms, leaving behind water clusters on the cold substrate. The
15 sublimation of such ice showed that the *OPR* of the molecules released into the gas phase
16 was equilibrated at the high temperature limit. If we assume that the ice prior to the
17 sublimation was a pure *para*-ice, the lifetime of *para*-H₂O in ice must be less than ~ 30 min,
18 a value which should be compared to a lifetime of a month, claimed in ref.⁵ The nuclear
19 spin relaxation time in the gas-phase water dimer was estimated in ref.¹¹ to exceed 3 ms.
20 According to ref.,¹¹ rapid nuclear spin relaxation of the water dimer embedded in an Ar
21 matrix is likely to occur on the time-scale of a few minutes. All these observations are
22 invaluable for understanding the nature and the rate of nuclear spin conversion in various
23 media including astrophysical environments. In the paragraphs below we shall address the
24 above estimates while discussing the results of our experiments.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 The present paper aims at further investigation of the water molecule nuclear spin sym-
47 metry conservation and relaxation in a supersonic expansion of ¹H₂¹⁶O water vapor seeded
48 in argon, helium or in oxygen as carrier gases. Mid-infrared spectra were recorded using the
49 Jet-AILES apparatus¹⁷ at the SOLEIL synchrotron facility.
50
51
52
53
54

55 The use of a supersonic jet cross-probed by an FTIR spectrometer has both advantages
56 and disadvantages for the study of relaxation processes. On one hand, rapid adiabatic
57
58
59
60

1
2
3
4 cooling of the core part of the jet gives rise to a significant thermal non-equilibrium which
5
6 can easily be observed in the high-resolution absorption spectra. Low values of the isentropic
7
8 core rotational temperature result in a limited set of rotational levels being populated and
9
10 consequently, only a small number of rotational lines can be seen in the spectrum. Also the
11
12 formation of dimers and larger clusters in the cold part of a jet is clearly seen in the expansion
13
14 of water vapor seeded in Ar as was evidenced e.g. in ref.¹⁸⁻²⁰ The spectral shifts of the small
15
16 cluster stretching bands were monitored in ref.¹⁸⁻²⁰ as a function of successive solvation of
17
18 pure water clusters by argon atoms. On the other hand, only fast relaxation processes can
19
20 be studied in such an experimental geometry where the supersonic jet expanding from the
21
22 slit reaches and crosses the IR beam in few microseconds.

23
24 The effect of lateral shock waves and viscous shear layers was taken into account. This
25
26 effect leads to multi-temperature gas samples being probed at the same time. Thorough
27
28 separation of spectral signatures into low-temperature and high-temperature ones was made
29
30 by means of a Boltzmann plot for the measured absorption line intensities as described below.
31
32 The use of a movable Pitot probe was used to characterize the supersonic expansion which
33
34 was then used for FTIR spectroscopic observations. In spite of the presence of the small water
35
36 cluster signatures in our obtained spectra, no transition to a nuclear spin relaxation regime
37
38 was observed: the *OPR* derived from the spectra was conserved on a level characteristic of
39
40 equilibrium conditions in the stagnation reservoir, that is the *OPR* was always equal to 3
41
42 irrespective of the extremely low rotational temperature. In addition, no change in *OPR*
43
44 was observed when the argon carrier gas was replaced by oxygen. The latter is known to
45
46 enhance nuclear spin relaxation because of its associated magnetic moment.^{10,21,22} Based on
47
48 the results obtained, we are in a position to dismiss the hypothesis of a water clustering
49
50 effect on nuclear spin relaxation and to question the origin of the nuclear spin relaxation
51
52 reported previously for a similar experiment.³
53
54
55
56
57
58
59
60

Experimental set-up and recording of the spectra

Jet-AILES apparatus

The Jet-AILES apparatus was used to record the mid-infrared (MIR) spectrum of water vapor cooled down to a rotational temperature between 10 and 42 K in a supersonic expansion. This experimental set-up has already been described in detail in other studies.¹⁷ Briefly, a planar supersonic jet was formed by expanding water vapor admixed in various carrier gases (argon, helium, oxygen) from a high pressure reservoir (2 bar maximum) to a low pressure chamber through a slit nozzle (60 mm \times 36 μ m or 60 mm \times 18 μ m).

This chamber was evacuated by two Roots pumps (Edwards EH 500 and 2600) backed by a primary pump (Edwards GV80 Drystar), delivering an effective pumping speed of 1800 m³/h. The gas flow was probed with the modulated infrared (IR) light beam issuing from the rear parallel exit of the high-resolution interferometer (Bruker IFS 125HR) equipping the AILES beamline. The IR beam was focused into the gas expansion using a first set of gold coated mirrors with a magnification factor of 1.14. After its absorption by the cold expansion, the IR beam was collected and focused by a second set of mirrors onto a liquid-nitrogen cooled InSb detector connected to the interferometer. A combination of a CaF₂ beamsplitter and a tungsten lamp was chosen to investigate the OH stretching region located around 3700 cm⁻¹.

The spectra were recorded within the spectral range between 3400 cm⁻¹ and 3900 cm⁻¹ where strong absorption by water vapor ν_1 and ν_3 stretching bands can be detected. In high-resolution spectra (Figure 1) the number of intense rotational lines varies significantly as a function of rotational temperature. Despite the significantly good signal-to-noise ratio achieved, the noise level in high resolution spectra did not permit detection of the weak, broad and virtually structureless absorption bands which belong to water clusters. To examine the evolution of cluster content under various expansion conditions, it was necessary to repeat the measurements at a much lower instrumental resolution (1 cm⁻¹). Figure 2 shows that

1
2
3
4 cluster absorption does manifest itself in the region of free-OH vibrations as well as bonded-
5
6 OH vibrations as broad unresolved features accumulating contributions from $(\text{H}_2\text{O})_n$ and
7
8 $(\text{H}_2\text{O})_n\text{Ar}_m$ clusters presumably containing up to dozens of molecules.²⁰

9
10 Each experimental condition (see Table 1) was repeated twice: a first recording was
11
12 performed at high resolution (0.002 cm^{-1} , boxcar apodization, iris aperture: 0.8 mm) to
13
14 determine the rotational temperature and the *OPR*, while a second one was performed at low
15
16 resolution (1 cm^{-1} , triangle apodization, iris aperture: 0.8 mm) to track the presence of water
17
18 clusters. The aperture setting was kept unchanged for high and low resolution recordings
19
20 so that the dimensions of the infrared beam through the flow remain unchanged (see Figure
21
22 3). Note that the spectrometer was maintained under high vacuum ($1.2 \times 10^{-4} - 2.0 \times 10^{-4}$
23
24 hPa) using a turbomolecular pump with active levitated magnetic bearings (Edwards model
25
26 1X455), thus minimizing the contribution of atmospheric water to the absorption spectra.

27
28 The continuous water vapor production was controlled by a vapor generation CEM-
29
30 system (Controlled Evaporation and Mixing, Bronkhorst, model W-303A) consisting of a
31
32 Coriolis liquid mass flow meter (mini Cori-Flow M12 Bronkhorst, $\pm 0.2\%$ of rate), a mass
33
34 flow controller (model F-202AV Bronkhorst, $\pm 0.5\%$ Rd plus $\pm 0.1\%$ FS) for the carrier gas
35
36 and a temperature controlled mixing and evaporation device. This system was used to vary
37
38 the mass flow rate of water vapor between 2 and 100 g/h. The flow rate of the carrier gas
39
40 was varied between 20 and 40 slm (1 standard liter per minute was measured at 0°C and 1
41
42 atm). The water vapor molar fraction was varied between 0.21% and 6.91%. This interval
43
44 was therefore larger than that corresponding to the former study reported in ref.³

45
46 To avoid water condensation, the inlet gas lines were wrapped in heating wires and main-
47
48 tained at the appropriate temperature of 40°C with a PID temperature controller. The gas
49
50 reservoir and the nozzle were made from a block of tempered aluminum. The temperature of
51
52 this element was regulated at 40°C with two heating cartridges and a second PID tempera-
53
54 ture controller. The pressure of the gas reservoir was measured using a bakeable capacitance
55
56 manometer (MKS Baratron 5000 Torr, model 615A) maintained at 100°C to avoid water
57
58
59
60

1
2
3 condensation into the gauge. The pressure of the expansion chamber was measured with a
4
5 100 Torr full scale range Baratron (MKS model 626A).
6
7

8 9 **Structure of the supersonic expansion**

10
11 The supersonic expansion produced by one of the slit nozzles ($60 \text{ mm} \times 18 \text{ }\mu\text{m}$) was ca-
12
13 refully characterized with a Pitot probe. For this purpose, the slit nozzle was placed in a
14
15 different vacuum chamber, at the University of Rennes (France), evacuated by a set of po-
16
17 werful mechanical pumps ($15,000 \text{ m}^3/\text{h}$). A butterfly valve positioned between the vacuum
18
19 chamber and the pumping unit was adjusted in order to match the pumping speed to that
20
21 corresponding to the Jet-AILES apparatus ($1800 \text{ m}^3/\text{h}$). The Pitot probe measured impact
22
23 pressures from which the Mach number of the flow could be extracted. It is constituted of
24
25 a stainless steel capillary (Outer Diameter/Inner Diameter = $1.4/1.1 \text{ mm}$) connected to a
26
27 capacitance manometer (MKS Baratron, 100 Torr) by a flexible tube. Its displacements in
28
29 two perpendicular directions were controlled by two stepper motors with a precision better
30
31 than 0.1 mm.
32
33

34
35 Figure 3 presents maps of various characteristics of the supersonic expansion of pure
36
37 argon (20.1 slm), in a plane perpendicular to the long edge of the slit. The stagnation and
38
39 residual pressures were 1072 and 0.68 hPa, respectively, thus very close to the conditions
40
41 obtained with the Jet-AILES apparatus. Impact pressures measured by the Pitot probe
42
43 are reported in this map, without any further treatment. It allows the diamond shock-cell
44
45 pattern typical of an underexpanded free-jet expansion^{18,19} to be visualized. A succession
46
47 of three supersonic cells each of about 20 mm long can be seen. The first cell (commonly
48
49 known as the “zone of silence”) is terminated by the intersection of the two lateral shocks 20
50
51 mm downstream from the nozzle exit. The lateral extension of this cell is about 8 mm at its
52
53 largest width. The trace of the IR beam is sketched on the figure. The optical axis was set at
54
55 about 5 mm from the nozzle exit. The inner circle of about 1 mm in diameter corresponds to
56
57 the spot focus, positioned at the middle of the slit aperture, while the outer ellipse delimits
58
59
60

1
2
3 the traces of the beam at the two edges of the slit. Part of the infrared beam impinges
4 on the lateral shocks and viscous shear layers which are responsible for a deceleration and
5 compression of the flow. This effect leads to the presence of rotationally “warm” molecules
6 absorbing the IR light along its path that have to be taken into account for the analysis of
7 spectra and to extract the rotational temperature of the water vapor in the cold part of the
8 supersonic expansion.
9

10
11 The hydrodynamic characteristics of the first cell of the jet, delimited by the nozzle exit
12 and the normal shock wave located at about 20 mm downstream, have been extracted from
13 the Pitot probe measurements using the so-called Rayleigh formula.²³ Strictly speaking, this
14 procedure only applies within the assumption of continuous flows, i.e. flows characterized
15 by a Knudsen number (Kn) lower than 0.01. Regarding the present approach, the Knudsen
16 number can be defined as the ratio between the mean free path of the expanding molecules
17 and the outer diameter of the Pitot probe. As displayed in Figure 3 (panel *b*), Kn attains a
18 value of 0.125 just before the shock wave, which denotes a rarefied regime still far from the
19 free molecular flow regime (Kn > 10). It seems appropriate therefore to use this means of
20 characterization for our free jet expansion issuing from a very narrow width. Note that the
21 values returned by the Rayleigh formula are not correct beyond the shock wave which is a
22 source of entropy. The Mach number (Figure 3, panel *c*) attains values up to about 17 in the
23 infrared probed zone, revealing an important enthalpy conversion into kinetic energy, thus
24 yielding very low translational temperatures, as can be seen in panel *d*. The translational
25 temperature is 3.4 K close to the optical axis, which is located at 5 mm downstream from
26 the nozzle. The Pitot probe measurements have been compared to the isentropic model
27 established on the basis of the method of characteristics.²⁴ Calculated and measured tempe-
28 ratures are remarkably close, thus confirming the validity of the measurements returned by
29 the Pitot probe. In the presence of water vapor, the translational temperature is expected
30 to reach slightly higher values. Indeed, adding a polyatomic molecule in the carrier gas
31 modifies the hydrodynamics by decreasing the specific heat ratio of the mixture and conse-
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

quently, the Mach number increase is smaller (some internal energy remains stored in the internal degrees of freedom of the water molecules). The main reason however, originates from thermodynamics: the condensation heat released by the clustering of water molecules is high and causes very significant warming of the flow, as can be seen from Table 1. Another important parameter of the flow field is its molecular density (Figure 3, panel *e*). As for the temperature, the molecular density drop associated with our free jet expansion is extremely abrupt, evolving from $1.7 \times 10^{19} \text{ cm}^{-3}$ at the nozzle exit to $3.4 \times 10^{16} \text{ cm}^{-3}$ at the optical axis location. It is therefore expected that most of the relaxation processes as well as cluster formation that rely on molecular collisions, occur mainly very close to the nozzle exit where the molecular density is still high.

The hydrodynamic time, denoted τ_{hyd} and defined as the time needed by the gas to reach the optical axis from the nozzle exit, can be easily estimated. In a free-jet, the gas reaches its limit velocity v_{∞} very rapidly downstream from the nozzle, i.e. after covering a distance equivalent to only a few slit widths. The limit velocity corresponds to a complete conversion of the enthalpy of the gas into kinetic energy (Mach number $M \rightarrow \infty$; flow temperature $T \rightarrow 0 \text{ K}$) and it is then easily retrieved from the conservation of energy equation:²⁵ $v_{\infty} = (2c_p T_0)^{1/2}$, where c_p is the heat capacity at constant pressure of the carrier gas, and T_0 is the stagnation temperature, fixed at 313 K in the present study. The value obtained for a flow of argon is 570 m/s, which is remarkably close to the value of 558 m/s calculated from the Mach number and the temperature extracted from the Pitot probe measurement ($v = aM$, where $a = \sqrt{\gamma r T}$ is the speed of sound, γ is the specific heat ratio and r is the specific gas constant), thus validating the $T \rightarrow 0 \text{ K}$ approximation. Hence the hydrodynamic times associated with helium, oxygen and argon are 2.8, 6.6 and 8.8 microseconds, respectively. These values are slightly overestimated as the heat of water condensation released in the flow adds a supplement of enthalpy into the flow which can be converted into additional kinetic energy.²⁶ One should keep in mind that these hydrodynamics times are smaller by several orders of magnitude than the timescale of milliseconds relative to *ortho-para* conversion in

1
2
3 the condensed phase.⁹⁻¹¹
4
5
6

7 8 Water clustering 9

10
11 Figure 2 shows that the rotational temperature steadily increases as the water vapor molar
12 fraction increases in the expansion. This can be understood in terms of the progressive
13 clustering of water molecules that results in a release of more latent heat when a more
14 notable portion of a gas in a mixture condenses. The rotational temperature achieved by
15 the water molecules in the expansion is not only governed by the condensation process. The
16 specific heat ratio of the carrier gas is also an important parameter to consider as it defines the
17 Mach number evolution of the expanding flow for a given nozzle geometry. Thus compared
18 to a monoatomic gas expansion, the expansion of a diatomic gas is characterized by a lower
19 cooling rate (i.e. a smaller Mach number increase) and therefore by a higher translational
20 temperature. The rotational temperature of the water molecules is obviously limited by
21 the “translational bath”. As an example, the rotational temperature achieved during our
22 measurements in an expansion of pure oxygen was 31.2 ± 0.6 K whereas no water clustering
23 was detected on the low resolution spectrum. For completeness, it should be noted that
24 the rotational relaxation phenomenon depends on the stagnation pressure and the nature
25 of the collision partner. In this respect, helium is far less efficient than argon for inducing
26 R-T energy transfer. Moreover, as the rotational relaxation is proportional to $P_0 d^*$ (see e.g.
27 ref.²⁷), where P_0 is the stagnation pressure and d^* the slit width, a large stagnation pressure
28 is required for efficient relaxation. However, as helium is lighter than argon by a factor of
29 10, the same molar flow of helium would lead to a stagnation pressure of about $\sqrt{10} \approx 3$
30 times lower than for argon. For these reasons (low R-T energy transfer and low stagnation
31 pressure), the use of helium was rejected in favor of argon. As an example, the maximum
32 flow rate tolerable by our pumping unit (30 slm) was used to inject helium together with a
33 very small amount of water (0.28%), leading to a rotational temperature around 42 K. As
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 for oxygen, no water clustering was observed.

5
6 The extent of water clustering was estimated by comparing the water monomer density
7 *measured* in the cold region of the jet to the monomer density *calculated* assuming no clus-
8 tering. The way the monomer density was extracted from our high resolution absorption
9 spectra is explained in detail in the next section, while the calculation was performed using
10 the standard isentropic model equations. More explicitly, the local molecular density N_{jet} of
11 the flow is expressed as a function of the Mach number:
12
13
14
15
16

$$\frac{N_{\text{jet}}}{N_0} = \left(1 + \frac{\gamma - 1}{\gamma} M^2\right)^{\frac{1}{1-\gamma}}, \quad (1)$$

17
18
19
20
21
22
23 where N_0 is the total molecular density in the reservoir. For a free-jet expansion, the Mach
24 number is modeled using the following analytical expression:
25
26

$$M = \delta^{\frac{\gamma-1}{2}} \sum_{i=0}^{i=3} \frac{A_i}{\delta^i}. \quad (2)$$

27
28
29
30
31
32
33 Here $\delta = x/d^*$ is the dimensionless distance from the nozzle exit. The various A_i coefficients
34 can be found in ref.²⁸ The calculated water monomer density is given by $x_{\text{H}_2\text{O}}N_{\text{jet}}$. As shown
35 in Figure 4, water condensation reaches 80% when argon is used as carrier gas, revealing
36 therefore, an intense water clustering.
37
38
39
40
41
42

43 Extraction of the rotational temperature from spectro- 44 pic data 45 46 47

48
49
50 Our aim was to determine the behaviour of the *OPR* under various experimental conditions.
51 The use of high resolution (0.002 cm^{-1}) makes it possible to accurately fit the observed line
52 profiles and retrieve the fractional populations relative to the rotational energetic levels of
53 the *ortho*- and *para*- molecules. It is then currently possible to determine the rotational tem-
54 perature T_{rot} of the molecules in the line-of-sight of the IR light beam. To take into account
55
56
57
58
59
60

instrument effects inherent to the Fourier Transform spectroscopic technique,²⁹ we analyzed the so-called experimental spectral absorptance (sometime called spectral absorption³⁰) denoted $\alpha(\tilde{\nu})$ as per the recommendation of IUPAC.³¹ Under that in our experimental conditions,²⁰ layers of hot gas surround the cold gas and contribute significantly to the total absorptance of the IR light. In the limit of our experimental accuracy, we extracted the integrated Napierian absorbance associated with the two gaseous zones assuming that the medium can be described by a simplified warm/cold bi-modal model. Analysis of the rotational populations allowed us to estimate a rotational temperature for the warm gas and for the cold gas. The relative integrated absorbances obtained for 4 *ortho* lines and 1 *para* line in the cold gas allow to explore the behaviour of the *OPR* of water within the fast slit-jet expansion. In the following, we describe the procedure used to retrieve the parameters characterizing the cold part of the expansion.

The absorptance $\alpha(\tilde{\nu})$ is obtained by comparing the spectral intensity of light $I(\tilde{\nu})$ at the output of the chamber to the intensity of light at its entrance $I_0(\tilde{\nu})$:

$$\alpha(\tilde{\nu}) = \Delta I(\tilde{\nu})/I_0(\tilde{\nu}) = (I_0(\tilde{\nu}) - I(\tilde{\nu}))/I_0(\tilde{\nu}). \quad (3)$$

$I(\tilde{\nu})$ follows the Beer-Lambert law and is given by :

$$I(\tilde{\nu}) = I_0(\tilde{\nu}) \exp(-A_{fi} \times V(\tilde{\nu})), \quad (4)$$

where A_{fi} is defined as the integrated Napierian absorbance of the line for a transition from a state i to a state f and $V(\tilde{\nu})$ is a normalized line profile (generally a Voigt profile taking into account inhomogeneous Doppler broadening and homogeneous collisional broadening).

A_{fi} is given by the following equation:^{32,33}

$$A_{fi} = \int \sigma(\tilde{\nu}) N L d\tilde{\nu}, \quad (5)$$

where $\sigma(\tilde{\nu})$ is the absorption cross-section at the wavenumbers $\tilde{\nu}$, L the thickness of the sample, N is the number of molecules per unit volume. Then, one has

$$A_{fi} = S_{fi}^N NL, \quad (6)$$

where integrated absorption cross section of the line (also called line intensity in ref^{32,33}) S_{fi}^N ($\text{cm}^{-1}/\text{molecule cm}^{-2}$) is given in the gas phase by:

$$S_{fi}^N = \frac{8\pi^3}{3hc} \tilde{\nu}_{fi} \left[1 - \exp\left(-\frac{hc\tilde{\nu}}{kT}\right) \right] \times \Re_{fi} \times 10^{-36} \times n_i, \quad (7)$$

where T is the temperature, \Re_{fi} (Debye²) the square of the transition moment and n_i represents the fractional population in the initial state. At low temperature (below 296 K), the wavenumber $\tilde{\nu}_{fi}$ of the transition is sufficiently high so that the exponential term vanishes.

Then,

$$S_{fi}^N = \frac{8\pi^3}{3hc} \tilde{\nu}_{fi} \times \Re_{fi} \times 10^{-36} \times n_i. \quad (8)$$

Finally the integrated Napierian absorbance is related to the fractional population through the simple relation:

$$A_{fi} = F_{fi}^N NL \times n_i, \quad (9)$$

where F_{fi}^N is independent of the temperature and afterwards will be called the line strength.

The evolution of the integrated Napierian absorbance is then directly related to the evolution of the population in the initial state. When the fractional populations of the rotational energy levels follow the thermodynamic equilibrium statistics, the population n_i is given by:

$$n_i = \frac{1}{Q(T)} g_i \exp\left(-\frac{hcE_i}{kT}\right) \quad (10)$$

where the energy of the initial level E_i is given in cm^{-1} and T in K, and $Q(T)$ is the partition function, $g_i = g_{\text{NS}} g_J$ with g_{NS} is the degeneracy of the considered Nuclear Spin state and

$g_J = 2J + 1$ is the degeneracy of the rotational state (Zeeman sub-levels).

By extracting the integrated Napierian absorbance of the lines A_{fi} under our experimental conditions, it is then possible to extract the rotational temperature T_{rot} using the following linear equation:

$$\log(A_{fi}/F_{fi}^N/g_J) = \log(NLn_i/(2J + 1)) = -\frac{hc E_i}{k T} + \log\left(\frac{g_{\text{NS}}}{Q(T)}\right) \quad (11)$$

which becomes

$$\log(NLn_i/(2J + 1)) = -\frac{E_i}{T} + \log\left(\frac{g_{\text{NS}}}{Q(T)}\right) \quad (12)$$

when the initial energy E_i is expressed in Kelvins.

In the spectral region around 3700 cm^{-1} , the Doppler Half Width at Half Maximum (HWHM) ranges between $1 \cdot 10^{-3}$ and $5 \cdot 10^{-3} \text{ cm}^{-1}$ for temperatures between 10 and 250 K. It is then necessary to take into account the width of the apparatus function to accurately model the line shape. The width of the apparatus function ($\approx 2 \cdot 10^{-3} \text{ cm}^{-1}$) has to be taken into account to perform accurate analysis of the line shape. Experimentally, the intensity of the light is given by the convolution product:

$$I_{\text{exp}}(\tilde{\nu}) = I(\tilde{\nu}) * f_{\text{app}}(\tilde{\nu}) \quad (13)$$

where the apparatus function $f_{\text{app}}(\tilde{\nu})$ is modelled by the Fourier transform of the Boxcar function:

$$f_{\text{app}}(\tilde{\nu}) = \frac{2}{R} \times \text{sinc}\left(\frac{2\tilde{\nu}}{R}\right). \quad (14)$$

Here $\text{sinc}(x) = \frac{\sin(\pi \cdot x)}{\pi \cdot x}$. The resolution R is given by $R = 1/\text{MOPD}$ where MOPD is the Maximum Optical Path Difference. In our case, the resolution corresponds to the resolution defined by the Bruker company divided by 0.9, then $R = 2.22 \cdot 10^{-3} \text{ cm}^{-1}$. The experimental absorption becomes:

$$\alpha_{\text{exp}}(\tilde{\nu}) = \frac{(I_0(\tilde{\nu}) - I(\tilde{\nu})) * f_{\text{app}}(\tilde{\nu})}{I_0(\tilde{\nu}) * f_{\text{app}}(\tilde{\nu})} \quad (15)$$

1
2
3 For FTIR spectroscopy, the entrance intensity $I_0(\tilde{\nu})$ is spectrally broad compared to the
4 width of the apparatus function. Then, $I_0(\tilde{\nu}) * f_{\text{app}}(\tilde{\nu}) \approx I_0(\tilde{\nu})$ and the absorptance is given
5 by:
6
7

$$\alpha_{\text{exp}}(\tilde{\nu}) = ((I_0(\tilde{\nu}) - I(\tilde{\nu})) * f_{\text{app}}(\tilde{\nu})) / I_0(\tilde{\nu}) \approx \alpha(\tilde{\nu}) * f_{\text{app}}(\tilde{\nu}) \quad (16)$$

8
9
10
11 then

$$\alpha_{\text{exp}}(\tilde{\nu}) = (1 - \exp(-A_{fi} \times V(\tilde{\nu}))) * f_{\text{app}}(\tilde{\nu}) \quad (17)$$

12
13
14
15
16
17 The variations of absorption lines intensities have been plotted as a function of T_{rot} for a
18 series of reference lines listed in Table 2.
19

20
21 Since the gas temperature in the shear layers is significantly higher than in the central
22 part of the jet, more observed rotational lines are due to absorption by the warm gas than
23 by the cold one. We performed the analysis of the individual lineshapes based on a two-steps
24 procedure which uses a homemade computational code written in MATLAB language:
25
26
27

28
29
30 (i) In the first step, the absorption $\alpha_{\text{exp}}(\tilde{\nu})$ of each individual line was fitted using a
31 Gaussian profile $V(\tilde{\nu})$. The only fitted parameters were the integrated Napierian absorbance
32 A_{fi} and the HWHM of the Gaussian profile. For experimental conditions relative to higher
33 water concentrations, contribution from the low temperature gas could be neglected for
34 transitions originating from an initial energetic level higher than 190 K. Indeed for these
35 transitions, the lineshape was perfectly fitted using Eq. 15 and the rotational diagram shows
36 a linear behaviour for energy E_i above 190 K (see Figure 5). Application of Eq. 9 yielded
37 the averaged column density NL^{hot} listed in Table 1 for each line. It should be noted that for
38 the calculation of the population factor n_i appearing in Eq. 9, the OPR value was assumed
39 to be close to 3. Because of the high temperature considered here, this value is justified by
40 the fact that calculations give values close to 3 irrespective of the nuclear spin equilibration
41 dynamics (fast or slow nuclear spin conversion; see text below and Eqs. 17–21). Application
42 of Eqs. 11-12 to the same integrated absorbances, led to the rotational temperatures T_{hot} ,
43 also listed in Table 1. For the case of a flow of 100 g/h of water in 30 slm of Ar, the
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 rotational diagram gives a high rotational temperature $T_{\text{hot}} = 251 \pm 11$ K very close to the
4 translational temperature $T = 260 \pm 40$ K extracted from the HWHM Gaussian lineshape.
5 For the “low temperature” lines, the difference between the calculated and experimental
6 spectral absorptance shows a typical residue presented in Figure 6.a (case of the $1_{11} \leftarrow 1_{10}$
7 transition). For the case of low water concentration, our signal-to-noise ratio was too low to
8 permit detection of high- J rotational lines. Therefore the only observable transitions were
9 those originating from initial energy states E_i lower than 190 K. The rotational diagrams
10 for *ortho* and *para* were linear for energy lower than 190 K. This means that the populations
11 followed a Boltzmann distribution allowing us to extract rotational temperatures directly
12 using Eq. 12. The rotational temperatures extracted for the *ortho* and *para* molecules are
13 listed in Table 1.
14
15
16
17
18
19
20
21
22
23
24

25
26 (ii) In a second step, we focused our analysis on spectra recorded with higher water
27 concentration. We fitted the lines corresponding to transitions related to initial rotational
28 states below 190 K, including the contribution of the warm gas. The absorptance was then
29 fitted using:
30
31
32
33
34
35

$$\alpha_{\text{exp}}(\tilde{\nu}) = (1 - \exp(-A_{fi}^{\text{cold}} \times V^{\text{cold}}(\tilde{\nu}) - A_{fi}^{\text{hot}} \times V^{\text{hot}}(\tilde{\nu}))) * f_{\text{app}}(\tilde{\nu}) \quad (18)$$

36
37
38
39 The Figure 6.b shows that the discrepancies between the calculated and the experimental
40 absorptances are drastically reduced for a flow of 100 g/h of water in 30 slm of Argon. The
41 only parameters in this second step were A_{fi}^{cold} and the HWHM of the Gaussian profile
42 $V^{\text{cold}}(\tilde{\nu})$. In this extreme case, the contribution of the hot molecules counts for a third of the
43 total absorbance. We were then able to remove the hot gas contribution from the absorbance
44 of these low- J rotational lines. Figure 5 shows that the populations reported for the cold
45 molecules in the rotational diagram follow a straight line.
46
47
48
49
50
51
52

53
54 In all cases, rotational temperatures for *ortho* and *para* subsystems were determined in-
55 dependently and were found to be nearly equal (see Table 1). The integrated absorbances
56 obtained for the “cold” transitions and the average rotational temperatures in the various
57
58
59
60

1
2
3 experimental conditions are listed in Table 3. The Figures 7, 8, 9 and 10 show the evolution
4 of the *ortho/para* relative absorbances for several transitions as a function of rotational tempe-
5 rature. As these ratios indicate that the *OPR* has not been relaxed during the cooling down,
6 we calculated the n_i fractional population using a conversion factor $(CF)_{\text{slow}}$ as described
7 by Eq. 23. The NL^{cold} column densities estimated for the cold gas in various conditions are
8 reported in Table 3 and can be compared to the warm gas contribution NL^{hot} . They have
9 been used to estimate the fraction of condensed water.
10
11
12
13
14
15
16

17
18 Finally, we evaluated a possible contribution of homogeneous collisional broadening under
19 the conditions of pressure and temperature of our experiment using the broadening coeffi-
20 cients tabulated in the HITRAN database.³² Under the various experimental conditions
21 used in this work, the collisional width was always less than 10^{-7} cm^{-1} (HWHM), justifying
22 *a posteriori* the use of a Gaussian profile for the lineshape analysis.
23
24
25
26
27

28 Although the infrared light beam diameter is rather large compare to the flow field
29 dimension, two well defined low- and high-rotational temperatures could be extracted from
30 our spectra using the above two-steps procedure, which seems contradictory regarding the
31 density and temperature gradients highlighted by the Pitot probe measurements (see Figure
32 3). In the zone of silence, it appears that beyond a certain distance from the nozzle, the gas
33 absorption becomes too weak to contribute significantly. Another assumption is that the
34 rotational temperature no longer relaxes beyond a certain distance from the nozzle, due to
35 a lack of collisions.
36
37
38
39
40
41
42
43
44
45

46 Discussion

47
48
49 To investigate the behaviour of the nuclear spin isomers spin species after the fast expansion,
50 we followed the procedure used by Manca Tanner et al.³ As the experimental data showed
51 that rotational populations of *ortho* and *para* molecules were equilibrated at the same ro-
52 tational temperature T we conducted our analysis of the *OPR* using the following method.
53
54
55
56
57
58
59
60

The ratio of integrated absorbance of one *ortho* fi transition to one *para* $f'i'$ transition is proportional to the ratio of the fractional population of molecules in the two considered rotational states:

$$\frac{A_{fi}}{A_{f'i'}} = \frac{F_{fi}^N}{F_{f'i'}^N} \times \frac{n_i}{n_{i'}} = \frac{F_{fi}^N}{F_{f'i'}^N} \times \frac{g_i}{g_{i'}} \exp\left(-\frac{hc(E_i - E_{i'})}{kT}\right) \times CF$$

$$\frac{A_{fi}}{A_{f'i'}} = \frac{F_{fi}^N}{F_{f'i'}^N} \times \frac{g_{NS_i} g_{J_i}}{g_{NS_{i'}} g_{J_{i'}}} \exp\left(-\frac{hc(E_i - E_{i'})}{kT}\right) \times CF \quad (19)$$

where $g_{NS} = 2I + 1$ is the nuclear spin states degeneracy given for a value of the total nuclear spin momentum number I and CF is a Conversion Factor.

OPR is a key parameter for our analysis. The OPR at a temperature T can be defined assuming that the *ortho* and *para* molecules behave as independent species (however both in contact with the same thermal bath of the expanding gas).³⁴ It can be calculated from separate partition functions evaluated for each nuclear spin conformers $Q_{ortho}(T)$ and $Q_{para}(T)$ expected at thermal equilibrium. When the $OPR(T)$ is out of equilibrium but the rotational *ortho* and *para* sub-systems are equilibrated, the coefficient of proportionality between the total *ortho* fractional populations n_{ortho} and $Q_{ortho}(T)$ is different of the one between n_{para} and $Q_{para}(T)$. The CF conversion factor in Eq. 19 is then given by :

$$OPR(T) = \frac{Q_{ortho}(T)}{Q_{para}(T)} \times CF = \frac{\sum g_{NS}(ortho) g_J(ortho) \exp(-\frac{hcE_{ortho}}{kT})}{\sum g_{NS}(para) g_J(para) \exp(-\frac{hcE_{para}}{kT})} \times CF \quad (20)$$

Here and below the summation extends to all the *ortho* or *para* lines, respectively. Two limiting cases can be distinguished.

(i) In the case of Fast Nuclear Spin Conversion (designated by Fast in the following), the equipartition of the nuclear spin states gives a $(CF)_{Fast}$ factor equal to 1. As a result, equation (20) becomes:

$$OPR(T) = 3 \frac{\sum g_J(ortho) \exp(-\frac{hcE_{ortho}}{kT})}{\sum g_J(para) \exp(-\frac{hcE_{para}}{kT})} \quad (21)$$

(ii) In the case of Slow Nuclear Spin Conversion (Slow in the following), we assume that the total number of molecules in each nuclear spin configuration remains constant during the cooling. Starting from a gaseous mixture with an OPR of 3, the ratio of the absorbances has to be calculated with a temperature dependent $(CF)_{\text{Slow}}$ parameter imposing an $OPR(T) = 3$ irrespective of the temperature:

$$OPR(T) = 3(CF)_{\text{Slow}} \times \left(\frac{\sum g_J(\text{ortho}) \exp\left(-\frac{hcE_{\text{ortho}}}{kT}\right)}{\sum g_J(\text{para}) \exp\left(-\frac{hcE_{\text{para}}}{kT}\right)} \right) = 3 \quad (22)$$

thus giving

$$(CF)_{\text{Slow}} = 1/ \left(\frac{\sum g_J(\text{ortho}) \exp\left(-\frac{hcE_{\text{ortho}}}{kT}\right)}{\sum g_J(\text{para}) \exp\left(-\frac{hcE_{\text{para}}}{kT}\right)} \right) \quad (23)$$

$(CF)_{\text{Slow}}$ is calculated using equation (23) to determine the theoretical ratio $A_{fi}/A_{f'i'}$ between absorbances of two transitions in the case of Slow NSC. In the section below, we shall compare the measured ratios of integrated absorption for selected pairs of *ortho*- and *para*-transitions with what can be expected for the limiting Fast or Slow nuclear spin conversion cases.

The theoretical ratio of integrated absorbances $A_{fi}/A_{f'i'}$ was calculated using the relationship (19) into which the CF parameter was substituted either assuming it equals 1 in the case of Fast conversion or equation (23) in the case of Slow conversion. The values of J quantum number, the energy of the low lying level of a selected transition and the line intensities $S_{fi}^N(296K)$ were taken from HITRAN database.³² Note that the line strengths F_{fi} that enter formula (19) were calculated from line intensities $S_{fi}^N(296K)$ reported in Table 2. The error is between 5 and 10 % which gives an uncertainty of 10-20 % on the relative absorbances.

Then the calculated values of $A_{fi}/A_{f'i'}$ were compared with the ratio of absorbances retrieved from the fit of experimentally recorded line profiles and given in Table 3. Comparison between experimental measurements and expected values are presented in Figures 7, 8, 9 and 10. Confidence domains presented in these figures for the theoretical values in case of Fast

1
2
3 or Slow NSC are calculated using 10 % error given by the uncertainties on line intensities
4
5 $S_{fi}^N(296K)$ reported in the HITRAN Database.³²
6
7

8 Figures 7, 8, 9, and 10 show that in all cases, our measured ratios of integrated absorban-
9 ces are in agreement with an assumption of Slow NSC. It is seen that even the use of oxygen
10 instead of argon as a carrier gas, does not result in the change of the nuclear spin conversion
11 rate. We do not observe any effect of cluster formation on the change in the nuclear spin
12 conversion rate as was suggested in ref.,³ despite clear evidence of cluster signatures in our
13 recorded low resolution spectra and an important fraction of condensed water molecules,
14 as estimated from our high resolution spectra. The rate of water condensation in an ex-
15 pansion of argon was estimated by comparing the monomer concentration extracted from
16 the line absorbances to the value given by the isentropic model for which a total absence
17 of condensation is assumed. As shown in Figure 4, water condensation reaches 80% when
18 argon is used as the carrier gas, revealing therefore, intense water clustering. Under our
19 experimental conditions, a fast nuclear spin conversion would then be expected even at low
20 temperature when the two models give very different *ortho/para* intensity ratios. Figures
21 7, 8, 9, and 10 show that such an effect does not occur. One explanation could arise from
22 the expansion dynamics differing between our study and that performed by Manca Tanner
23 et al.³ At first glance, the differences are quite subtle. Indeed, the flow characteristics in the
24 isentropic part of a supersonic expansion are governed by only a few parameters linked to
25 the nozzle geometry and the stagnation conditions (namely temperature, pressure, specific
26 heat ratio of the expanding mixture, water molar fraction). It appears that in both studies,
27 a slit nozzle has been used, argon was mainly used and very similar stagnation pressures
28 and temperatures have been set, so that very similar flow expansions are therefore expected.
29 One has to notice however, that the wider 100 μm width nozzle used in ref. 3 (versus 18
30 and 36 μm in the present work) certainly leads to a slightly lower cooling rate. However, as
31 in this former study the flow was probed farther from the nozzle exit (10 mm *vs* 5 mm), the
32 temperature and molecular density reached by the flow for equivalent gas compositions are
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 expected to be very similar to those obtained in the present work. However, it is interesting
4
5 to notice that the extracted rotational temperature is systematically higher by about 5 K
6
7 for equivalent partial pressures of water (see table 6 of ref. 3). One explanation could be
8
9 attributed to a more pronounced water condensation induced by the wider slit nozzle used
10
11 in ref. 3. Finally, to be complete, one has to point out, a hydrodynamic time (as defined
12
13 previously) two times larger in ref. 3 ($\sim 18 \mu s$ vs $\sim 9 \mu s$) because of the larger probing
14
15 distance. We are considering using a Laval nozzle to better control water clustering and to
16
17 increase the hydrodynamic time (see e.g. ref.³⁵). It would provide us with new insights on
18
19 the effects of the hydrodynamic time and the rate of water condensation in the expansion.
20
21

22 23 24 Conclusion

25
26
27 The present paper was aimed at a further examination of water vapor nuclear spin-isomerization.
28
29 In recent years the mechanism and the effectiveness of nuclear spin conversion among wa-
30
31 ter vapor *ortho*- and *para*- spin isomers have been the subject of a number of studies that
32
33 often gave rise to contradictory conclusions. We have conducted a Jet-FTIR examination
34
35 of several water vapor absorption lines seen in H₂O seeded expansions using pure argon or
36
37 helium as carrier gases. Some experiments were carried out with dioxygen as carrier gas or
38
39 co-added to expanding mixture of gases in order to accelerate *ortho-para* conversion due to
40
41 collisions with paramagnetic oxygen molecules. To summarize, we did not observe any signs
42
43 of nuclear spin relaxation in the wide range of expansion conditions which were used:
44
45

- 46 • H₂O molar fraction: from 0.2% to 6.5%;
- 47
- 48 • rotational temperature range: from 10 K to 42 K;
- 49
- 50 • transitions explored: one *para* line ($1_{01} \leftarrow 0_{00}$) and four *ortho* lines ($2_{02} \leftarrow 1_{01}$; $0_{00} \leftarrow$
51
52 1_{01} ; $1_{11} \leftarrow 1_{10}$; $2_{11} \leftarrow 1_{10}$).
- 53
54
55

56
57 In any event the measured *OPR* corresponded to the temperature in the stagnation
58
59
60

1
2
3 reservoir and was not relaxed to very low rotational temperatures measured at the sampled
4 volume of the jet. Moreover, in spite of clear signatures of water clusters which were formed
5 at some experimental conditions, we did not observe any effect of clustering on the rate of
6 nuclear spin isomer conversion. This result agrees with a conclusion that has been drawn from
7 a similar experiment in ref.¹⁵ and differs significantly from what has been reported previously
8 by ref.³ Based on the results obtained, we are in a position to state that at least the suggestion
9 of ref.³ that the transition from nuclear spin conservation to nuclear spin relaxation regimes
10 is due to clustering in an expansion, is not supported by our measurements, also in agreement
11 with the previous work.¹⁵
12
13
14
15
16
17
18
19
20
21
22
23

24 Acknowledgement

25
26
27 The authors are grateful to SOLEIL and the AILES staff for providing technical support
28 under the proposal 20120156. X. M. and A. M. are grateful to the French National Research
29 Agency (Project ANR GASOSPIN No. 09-BLAN-0066-01) for the financial support of the
30 calculation code development. A.V. acknowledges partial support from the RFBR Grant
31 15-05-00736 and the Russian Acad. Sci. Program 9. This work was supported by the
32 Programme National “Physique et Chimie du Milieu Interstellaire” (PCMI) of CNRS/INSU
33 with INC/INP co-funded by CEA and CNES. The authors warmly thank the reviewers for
34 their valuable comments and suggestions which have helped to improve the quality of our
35 manuscript.
36
37
38
39
40
41
42
43
44
45
46
47

48 References

- 49
50
51 (1) Cacciani, P.; Cosléou, J.; Khelkhal, M. Nuclear Spin Conversion in H₂O. *Phys. Rev. A*
52 **2012**, *A85*, 012521-1-8.
53
54
55
56 (2) Chapman, W.B.; Kulcke, A.; Blackmon, B.W.; Nesbitt, D.J. Rotationally Inelastic
57
58
59
60

- 1
2
3
4 Scattering of Jet Cooled H₂O with Ar: State-to-State Cross Sections and Rotational
5 Alignment Effects. *J. Chem. Phys.* **1999**, *110*, 8543–8554.
6
7
8
9 (3) Manca Tanner, C.; Quack, M.; Schmidiger, D. Nuclear Spin Symmetry Conservation
10 and Relaxation in Water (¹H₂¹⁶O) Studied by Cavity Ring-Down (CRD) Spectroscopy
11 of Supersonic Jets. *J. Phys. Chem. A* **2013**, *117*, 10105–10118.
12
13
14
15 (4) Konyukhov, V.K.; Tikhonov, V.I.; Tikhonova, T.L.; Faizulaev, V.N. Separation of Spin-
16 Modifications of Water and Heavy-Water Molecules. *Pis'ma v Zhurnal Tekhnicheskoi*
17 *Fiziki* **1986**, *12*, 1438–1441, 1986 [in Russian].
18
19
20
21
22 (5) Tikhonov, V.; Volkov, A. Separation of Water into its *ortho* and *para* Isomers. *Science*
23 **2002**, *296*, 2363.
24
25
26
27 (6) Stepanov, E.V.; Tikhonov, V. I.; Milyaev, V.A. Diagnostics of *ortho* and *para* Water
28 Isomers with Tunable Diode Lasers. *Quantum Electron.* **2005**, *35*, 205 [in Russian].
29
30
31
32 (7) Stepanov, E. V.; Zyryanov, P. V.; Milyaev, V. A. Laser Analysis of the Relative Content
33 of *ortho*- and *para*-Water Molecules for the Diagnostics of Spin-Selective Processes in
34 Gaseous Media. *Phys. Wave Phenom.* **2010**, *18*, 33–43.
35
36
37
38
39 (8) Veber, S. L.; Bagryanskaya, E. G.; Chapovsky, P. L. On the Possibility of Enrichment
40 of H₂O Nuclear Spin Isomers by Adsorption. *J. Exp. Theor. Phys.* **2006**, *102*, 76–83.
41
42
43
44 (9) Buntkowsky¹, G.; Limbach, H.-H.; Walaszek, B.; Adamczyk, A.; Xu, Y.; Breitzke, H.;
45 Schweitzer, A.; Gutmann, T.; Wächtler, M.; Frydel, J. et al. Mechanisms of Dipolar
46 Ortho/Para-H₂O Conversion in Ice. *Z. Phys. Chem.* **2008**, *222*, 1049–1063 . DOI
47 10.1524.zpch.2008.5359
48
49
50
51
52
53 (10) Abouaf-Marguin, L.; Vasserot, A.-M.; Pardanaud, C.; Michaut, X. Nuclear Spin Con-
54 version of Water Diluted in Solid Argon at 4.2 K: Environment and Atmospheric Im-
55 purities Effects. *Chem. Phys. Lett.* **2007**, *447*, 232–235.
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- (11) Sliter, R.; Gish, M.; Vilesov, A.F. Fast Nuclear Spin Conversion in Water Clusters and Ices: A Matrix Isolation Study. *J. Phys. Chem. A* **2011**, *115*, 9682–9688.
- (12) Turgeon, P.-A.; Ayotte, P.; Lisitsin, E.; Meir, Y.; Kravchuk, T.; Alexandrowicz G. Preparation, Isolation, Storage, and Spectroscopic Characterization of Water Vapor Enriched in the *ortho*-H₂O Nuclear Spin Isomer. *Phys. Rev. A* **2012** *86*, 062710.
- (13) Turgeon, P.-A.; Vermette, J.; Alexandrowicz, G.; Peperstraete, Y.; Philippe, L.; Bertin, M.; Fillion, J.-H.; Michaut, X.; Ayotte, P. Confinement Effects on the Nuclear Spin Isomer Conversion of H₂O. *J. Chem. Phys. A* **2017** *121*, 1572–1577.
- (14) Bunker, Ph.; Jensen, P. In *Frontiers of Molecular Spectroscopy*; Ed. by Jaan Laane; ELSEVIER, 2008, p. 331.
- (15) Votava, O.; Horká-Zelenková, V.; Rakovský, J.; Svoboda, V.; Fárnik, M. H₂O Nuclear-Spin Conservation/Relaxation in Supersonic Jet Expansion Probed by Quantitative Direct Absorption Spectroscopic Detection. *The 23-rd International Conference on High Resolution Molecular Spectroscopy, 2 – 6 september 2014* Book of abstracts, Bologna, P. 134.
- (16) Konyukhov, V.K.; Prokhorov, A.M.; Tikhonov, V.I.; Faizulaev, V.N. Rotationally Selective Condensation of Heavy Water in a Supersonic Carbon Dioxide Jet. *JETP Lett.* **1986**, *43*, 85.
- (17) Cirtog, M.; Asselin, P.; Soulard, P.; Tremblay, B.; Madebène, B.; Alikhani, M.E.; Georges, R.; Moudens, A.; Goubet, M.; Huet, T.R. et al. The (CH₂)₂O-H₂O Hydrogen Bonded Dimer. *Ab initio* Calculations and FTIR Spectroscopy from Neon Matrix, and a New Supersonic Jet Experiment Coupled to the Infrared AILES Beamline of Synchrotron SOLEIL. *J. Phys. Chem. A* **2011**, *115*, 2523–2532.
- (18) Häber, T.; Schmitt, U.; Emmeluth, C.; Suhm, M. A. Ragout-jet FTIR Spectroscopy

- of Cluster Isomerism and Cluster Dynamics: from Carboxylic Acid Dimers to N₂O Nanoparticles. *Faraday Discuss.* **2001**, *118*, 331–359.
- (19) Zischang, J.; Suhm, M.A. Infrared Absorption Imaging of 2D Supersonic Jet Expansions: Free Expansion, Cluster Formation, and Shock Wave Patterns. *J. Chem. Phys.* **2013**, *139*, 024201.
- (20) Moudens, A.; Georges, R.; Goubet, M.; Makarewicz, J.; Lokshtanov, S. E.; Vigasin, A. A. Direct Absorption Spectroscopy of Water Clusters Formed in a Continuous Slit Nozzle Expansion. *J. Chem. Phys.* **2009**, *131*, 204312.
- (21) Chehrouri, M.; Fillion, J.-H.; Chaabouni, H.; Mokrane, H.; Congiu, E.; Dulieu, F.; Matar, E.; Michaut, X.; Lemaire, J.L. Nuclear Spin Conversion of Molecular Hydrogen on Amorphous Solid Water in the presence of O₂ traces. *Phys. Chem. Chem. Phys.* **2011**, *13*, 2172–2178.
- (22) Redington, R.L.; Milligan, D.E. Molecular Rotation and *ortho-para* Nuclear Spin Conversion of Water Suspended in Solid Ar, Kr, and Xe. *J. Chem. Phys.* **1963**, *39*, 1276.
- (23) Georges, R.; Bonnamy, A.; Benidar, A.; Decroi, M.; Boissoles, J. FTIR Free-Jet Set-up for the High Resolution Spectroscopic Investigation of Condensable Species. *Mol. Phys.* **2002**, *100*, 1551–1558.
- (24) Miller, D. R. In *Atomic and Molecular Beam Methods*; Ed. by G. Scoles; Oxford University Press, New York, 1988, Vol. 1.
- (25) Pauly, H. *Atom, Molecule, and Cluster Beams I: Theory, Production and Detection of Thermal Energy Beams*; Springer-Verlag: Berlin Heidelberg, Germany, 2000.
- (26) Bonnamy, A.; Georges, R.; Benidar, A.; Boissoles, J.; Canosa, A.; Rowe, B.R. Infrared Spectroscopy of (CO₂)_N Nanoparticles (30 < N < 14500) Flowing in a Uniform Supersonic Expansion. *J. Chem. Phys.* **2003**, *118*, 3612–3621.

- 1
2
3
4 (27) Belikov, A.E.; Sharafutdinov, R.G. Rotational Relaxation Time in Free Jets of He + N₂
5 Mixtures. *Chem. Phys. Lett.* **1995**, *241*, 209–214.
6
7
8
9 (28) Herman M.; Georges R.; Hepp M.; Hurtmans D. High resolution Fourier transform
10 spectroscopy of jet-cooled molecules. *Int. Rev. Phys. Chem.* **2000**, *19*, 277–325.
11
12
13 (29) *Handbook of High Resolution Spectroscopy*; Ed. by M. Quack and F. Merkt; John Wiley
14 and Sons, **2011**, *vol. II*, 970–977.
15
16
17
18 (30) Mandin, J.-Y.; Flaud, J.-M.; Camy-Peyret C.; Guelachvili, G. Measurements and Cal-
19 culations of Self-Broadening Coefficients of Lines Belonging to the ν_2 Band of H₂¹⁶O. *J.*
20 *Quant. Spectrosc. Radiat. Transfer* **1979**, *23*, 351–370.
21
22
23
24
25 (31) PAC, Glossary of Terms Used in Photochemistry (IUPAC Recommendations 1996).
26 **1996**, *68*, 2223 doi:10.1351/pac199668122223
27
28
29
30 (32) Rothman, L.S.; Jacquemart, D.; Barbe, A.; Chris Benner, D.; Birk, M.; Brown, L.R.;
31 Carleer, M.R.; Chackerian Jr., Chance, K.; Coudert, L.H. et al. The HITRAN 2004
32 Molecular Spectroscopic Database *J. Quant. Spectrosc. Radiat. Transfer* **2005**, *96*, 139–
33 204.
34
35
36
37
38
39 (33) Simeckova M.; Jacquemart D.; Rothmana L.S.; Gamache R. R.; Goldman A. Ein-
40 stein A-Coefficients and Statistical Weights for Molecular Absorption Transitions in
41 the HITRAN Database. *J. Quant. Spectrosc. Radiat. Transfer* **2006**, *98*, 130–155.
42
43
44
45
46 (34) Votava O.; Masat M.; Pracna P.; Kassi S.; Campargue A. Accurate Determination of
47 Low State Rotational Quantum Numbers ($J < 4$) from Planar-Jet and Liquid Nitrogen
48 Cell Absorption Spectra of Methane Near 1.4 micron. *Phys. Chem. Chem. Phys* **2010**,
49 *12*, 3145–3155.
50
51
52
53
54
55 (35) Bourgalais, J.; Roussel, V.; Capron, M.; Benidar, A.; Jasper, A. W.; Kloppenstein, S.
56
57
58
59
60

1
2
3 J.; Biennier, L.; Le Picard, S. D. Low Temperature Kinetics of the First Steps of Water
4 Cluster Formation. *Phys. Rev. Lett.* **2016**, *116*, 113401.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Experimental conditions

Carrier gas			H ₂ O			P _{H₂O}	P ₀ ^b	P _{ch} ^c	d ^{*d}	T _{rot} (O/P) ^e	NL _{cold} ^f	T _{hot} ^g	NL _{hot} ^f	τ _{hyd} ^h	x _{cond} ⁱ
slm															
Ar	He	O ₂	g/h	slm	x ^a %	hPa	hPa	hPa	μm	K	molec/cm ²	K	molec/cm ²	μs	%
20			2	0.041	0.21	1.73	843	0.71	36	10.7±1.6 / -	1.1±0.3	-	-	8.8	75
20			4	0.083	0.41	3.46	843	0.71	36	9.8±0.6 / -	3.0±0.5	-	-	8.8	65
20			8	0.166	0.82	6.94	843	0.71	36	11.6±0.3 / 13.4±2.7	4.3±0.3	288±50	5.2±0.1	8.8	75
20			10	0.238	1.04	8.68	843	0.71	36	12.0±0.6 / 12.7±0.5	4.8±0.2	167±13	5.6±0.1	8.8	78
20			12	0.249	1.23	10.37	843	0.71	36	13.0±0.6 / 13.3±0.1	6.0±0.6	176±30	5.6±1.0	8.8	77
20			20	0.415	2.03	17.15	845	0.71	36	20.2±0.5 / 18.9±0.4	11.8±1.3	225±25	10.2±1.4	8.8	72
20			30	0.622	3.01	25.68	853	0.72	36	22.4±0.3 / 24.1±1	15.3±0.7	257±15	17.6±2.4	8.8	76
30			20	0.415	1.36	16.35	1202	1.04	36	18.9±1.3 / 18.7±0.4	8.8±0.8	-	-	8.8	78
30			50	1.037	3.34	51.20	1533	1.09	18	25.3±0.6 / 25.7±1.3	13.5±1.5	237±13	28.4±2.2	8.8	79
30			100	2.074	6.47	102.74	1588	1.11	18	32.3±0.4 / 31.9±0.4	22.2±1.7	251±11	53.5±1.7	8.8	83
		20	8	0.166	0.82	8.88	1083	0.69	18	31.3±0.5 / 30.4±4	10.4±0.1	-	-	6.6	0
20		20	8	0.166	0.41	8.25	2013	1.33	18	21.8±2 / 18.0±4	7.8±1.2	-	-	7.6	0
	30		4	0.083	0.28	1.54	549	1.11	36	41.8±3 / 39.7±4	3.8±0.2	-	-	2.8	0

^awater vapor/carrier gas molar ratio; ^bstagnation pressure; ^cresidual pressure in the expansion chamber; ^dnozzle width (the slit length is 60 mm); ^erotational temperature extracted from *ortho* (O) or *para* rovibrational lines (P); ^fcolumn densities for hot and cold gases (see text for details); ^g rotational temperature of the warm residual gas in shear layers; ^hhydrodynamic time; ⁱfraction of condensed water vapor.

Table 2: Assignments and some parameters of the H₂O absorption lines of the ν_3 band used in this work

Transition	Spin-isomer	Line position, ^a cm ⁻¹	Lower energy level, ^a cm ⁻¹ / K	Line Intensity 10 ⁻¹⁹ S _{fi} ^N , ^a cm/molec	Line Strength, 10 ¹⁸ F _{fi} ^N , cm/molec
1 ₀₁ ←2 ₀₂	<i>para</i>	3709.40227	70.0908/100.8457	0.702	3.44721
1 ₁₀ ←1 ₁₁	<i>para</i>	3759.84448	37.1371/53.4324	0.5809	4.05056
1 ₀₁ ←0 ₀₀	<i>para</i>	3779.49307	0	0.4559	7.96176
2 ₁₂ ←1 ₁₁	<i>para</i>	3796.43951	37.1371/53.4324	0.5527	3.85393
3 ₀₃ ←2 ₀₂	<i>para</i>	3820.73852	70.0908/100.8457	0.8754	4.29869
1 ₁₁ ←2 ₁₂	<i>ortho</i>	3712.20451	79.4964/114.3783	1.501	2.57184
0 ₀₀ ←1 ₀₁	<i>ortho</i>	3732.13435	23.7944/34.2351	1.303	2.83838
1 ₁₁ ←1 ₁₀	<i>ortho</i>	3749.32915	42.3717/60.9638	1.743	4.15566
2 ₁₁ ←2 ₁₂	<i>ortho</i>	3769.88893	79.4964/114.3783	0.7786	1.33407
2 ₀₂ ←1 ₀₁	<i>ortho</i>	3801.41863	23.7944/34.2351	2.312	5.03633
2 ₁₁ ←1 ₁₀	<i>ortho</i>	3807.01357	42.3717/60.9638	1.562	3.72412
3 ₁₃ ←2 ₁₂	<i>ortho</i>	3816.09164	79.4964/114.3783	2.312	3.96143
2 ₂₀ ←1 ₀₁	<i>ortho</i>	3863.31982	23.7944/34.2351	0.0456	0.09933

^a – From ref. 32

Table 3: Rotational temperatures T_{rot} and integrated Napierian absorbances A associated with the molecules present in the cold part of supersonic expansion

Carrier gas / slm			H ₂ O	T_{rot}^a	$AR(0_{00})^b$	$AR(1_{01})^c$	$AP(1_{01})^d$	$AQ(1_{10})^e$	$AR(1_{10})^f$
Ar	He	O ₂	g/h	K	<i>para</i>	<i>ortho</i>	<i>ortho</i>	<i>ortho</i>	<i>ortho</i>
					10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³	10 ⁻³
20			2	10.7 ±1.8	0.754 ±0.014	0.792 ±0.012	1.36 ±0.03	0.084 ±0.004	0.067 ±0.008
20			4	9.8 ±1.5	1.56 ±0.02	1.65 ±0.02	2.78 ±0.03	0.169 ±0.006	0.178 ±0.007
20			8	11.8 ±1.2	2.21 ±0.02	2.32 ±0.02	3.86 ±0.04	0.336 ±0.007	0.316 ±0.009
20			10	12.2 ±0.4	2.58 ±0.02	2.64 ±0.02	4.36 ±0.04	0.397 ±0.007	0.374 ±0.008
20			12	13.3 ±0.2	2.74 ±0.02	2.87 ±0.03	4.72 ±0.05	0.499 ±0.010	0.478 ±0.006
20			20	19.5 ±0.7	3.31 ±0.02	3.53 ±0.03	5.82 ±0.06	1.10 ±0.01	1.010 ±0.008
20			30	22.8 ±1.0	3.36 ±0.02	3.69 ±0.03	6.31 ±0.06	1.56 ±0.01	1.45 ±0.01
30			20	18.8 ±0.1	2.33 ±0.02	2.53 ±0.03	4.37 ±0.04	0.825 ±0.018	0.749 ±0.016
30			50	25.4 ±0.2	2.69 ±0.02	2.94 ±0.02	5.07 ±0.04	1.29 ±0.01	1.20 ±0.01
30			100	32.1 ±0.4	2.77 ±0.02	3.49 ±0.03	6.09 ±0.06	2.16 ±0.02	2.01 ±0.02
		20	8	31.2 ±0.6	1.25 ±0.02	1.59 ±0.02	2.75 ±0.02	0.978 ±0.016	0.884 ±0.016
20		20	8	21.6 ±2.6	1.90 ±0.02	1.92 ±0.03	3.48 ±0.03	0.690 ±0.014	0.703 ±0.018
	30		4	41.3 ±1.2	0.28 ±0.01	0.451 ±0.006	0.786 ±0.008	0.330 ±0.007	0.328 ±0.008

^a average rotational temperature;

^b $1_{01} \leftarrow 0_{00}$; ^c $2_{02} \leftarrow 1_{01}$; ^d $0_{00} \leftarrow 1_{01}$; ^e $1_{11} \leftarrow 1_{10}$; ^f $2_{11} \leftarrow 1_{10}$ transitions.




Figure 1: Selection of jet-cooled transmittance spectra recorded at 0.002 cm^{-1} resolution to avoid line saturation effects. Water clustering, driven by the partial pressure of water vapor present in the flow, is one of the means used to control the temperature of the flow (argon in these examples). Rotational temperature increases from the lower to the upper panel as the water molar fraction increases. In our study, relative intensities were calculated using four *ortho* transitions (given using the notation “band $\nu_3 \leftarrow$ ground state” by: $2_{02} \leftarrow 1_{01}$; $0_{00} \leftarrow 1_{01}$; $1_{11} \leftarrow 1_{10}$; $2_{11} \leftarrow 1_{10}$) and one *para* transition ($1_{01} \leftarrow 0_{00}$).




Figure 2: Series of low resolution absorbances (offset for clarity) of water vapor cooled in argon (400 co-added scans at 1 cm^{-1} resolution). Water clustering is driven by the water vapor partial pressure (as indicated in the rightmost part of the figure) in the stagnation reservoir. In addition to the partially resolved water monomer lines, both the signatures of free OH (around 3710 cm^{-1}) and bound OH vibrations are observed.²⁰ The signatures of the water dimer and trimer can be seen at 3601 and 3533 cm^{-1} , respectively, for conditions leading to weak clustering. The broad absorption observed below 3600 cm^{-1} , corresponds to much larger water clusters that form at enhanced water vapor partial pressure.




Figure 3: Impact pressure measured by a Pitot probe (a). This 2D map was used to visualize the flow structure in a section perpendicular to the long edge of the slit, resulting from an expansion of argon under conditions very similar to those used for the spectral recordings ($P_0 = 1072 \text{ hPa}$, $P_{\text{ch}} = 0.683 \text{ hPa}$, $T_0 = 295 \text{ K}$, slit width: 18 microns, slit length: 60 mm). The path of the infrared beam is indicated by the dashed lines (circle = focal point; ellipse = edges of the slit). Various flow parameters have been extracted from measurements relative to the plane of symmetry: Knudsen number (b); Mach number (c); translational temperature (d) compared to a calculation using the method of characteristics (solid line); molecular density (e) compared to a calculation using the method of characteristics (solid line).




Figure 4: Concentration of condensed water vapor in an expansion of argon. These values, estimated from the measured monomer absorption, reveal a strong water condensation (between 65 and 83 %) even for small partial pressures of water vapor.




Figure 5: Example of rotational diagram used to extract the rotational temperature from absorbance spectra in the case of the presence of hot and cold molecules in the path of the infrared beam (here 100 g/h of H₂O in 30 slm of Argon). The quantity $(NLn_i/(2J+1))$ (see Eq. 10) extracted from the fitting of the different lines, is represented as a function of the energy level E_i of the initial state of the transition. The open circles and open squares represent the data obtained with a single Gaussian profile for *ortho* and *para* lines respectively. The data for energies E_i higher than 190 K follow a straight line and the slope gives the rotational temperature of the hot molecules (251 ± 11 K). The warm gas contribution is then removed from the low- J rotational line absorption by a fitting procedure taking into account the hot and cold contributions (see text for details), leading to a corrected value of the cold gas rotational temperature for the *ortho* molecules (32.3 ± 0.4 K) very close to the rotational temperature extracted from the lines associated with the *para* molecules (31.9 ± 0.4 K). A total of 8 low- J rotational lines have been used to estimate the cold rotational temperature.




Figure 6: Absorbance of the $1_{11} \leftarrow 1_{10}$ transition observed with a flow of 100 g/h of water in 30 slm of Ar. The experimental data (line) are compared to the calculated data (dashed line) after a fitting procedure (a) with a single Gaussian profile and (b) with two Gaussian profiles associated with one contribution coming from “hot” gas (red dash lines) and one contribution from “cold” gas (blue short dash-dot line) convoluted by the appropriate apodization function (Boxcar). The obs-calc residue is depicted in the bottom part of each panel.





Figure 7: Relative integrated absorbances $A_{1_{01} \leftarrow 0_{00}} / A_{2_{02} \leftarrow 1_{01}}$ measured as a function of rotational temperature. The expected behaviour in case of Slow NSC (solid line) and in case of Fast NSC (dashed line) is also represented. The uncertainty of 10% on the theoretical values are shown in gray are derived from HITRAN³² data. Triangle symbols represent data obtained in pure oxygen or in a mixture with equal Ar and O₂ flow rates as a carrier gas, as shown in Table 3, whereas the star refers to the data obtained in an expansion of He instead of Ar as a carrier gas. The x-axis error bars reflect the uncertainty of the rotational temperature retrieved from the rotational diagram. The y-axis error bars originate from the most severe estimate of uncertainty in the line intensities measurements following subtraction, when possible, of the hot gas contribution to the low-temperature line absorption.



45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 8: Relative integrated absorbances $A_{101 \leftarrow 000} / A_{000 \leftarrow 101}$ measured as a function of rotational temperature. See Fig. 7 caption for more details.




Figure 9: Relative integrated absorbances $A_{1_{01} \leftarrow 0_{00}} / A_{1_{11} \leftarrow 1_{10}}$ measured as a function of rotational temperature. See Fig. 7 caption for more details.





Figure 10: Relative integrated absorbances $A_{1_{01} \leftarrow 0_{00}} / A_{2_{11} \leftarrow 1_{10}}$ measured as a function of rotational temperature. See Fig. 7 caption for more details.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


TOC Graphic




Para


Ortho






?






2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60






1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56



100
90
80
70
60
50





ACS Paragon Plus Environment
Partial pressure of water vapor in the reservoir (hPa)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46








$AR(0_{00}) / AQ(1_{10})$

Relative Absorbance



$AR(0_{00}) / AR(1_{10})$

Relative Absorbance

