



# **A Review of Physical Activity and Circulating miRNA Expression: Implications in Cancer Risk and Progression**

Suzanne Dufresne, Amélie Rébillard, Paola Muti, Christine M. Friedenreich,  
Darren R. Brenner

## **► To cite this version:**

Suzanne Dufresne, Amélie Rébillard, Paola Muti, Christine M. Friedenreich, Darren R. Brenner. A Review of Physical Activity and Circulating miRNA Expression: Implications in Cancer Risk and Progression. *Cancer Epidemiology, Biomarkers and Prevention*, 2018, 27 (1), pp.11-24. 10.1158/1055-9965.EPI-16-0969 . hal-01713229

**HAL Id: hal-01713229**

**<https://univ-rennes.hal.science/hal-01713229>**

Submitted on 28 Feb 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**A review of physical activity and circulating-miRNA expression: Implications in cancer risk  
and progression**

Suzanne Dufresne<sup>1</sup>, Amélie Rébillard<sup>1</sup>, Paola Muti<sup>2</sup>, Christine M. Friedenreich<sup>3,4,5</sup>, Darren R.  
Brenner<sup>3,4,5</sup>

1. Movement, Sport and Health Sciences Laboratory, EA 1274, University Rennes 2, ENS  
Rennes, Bruz, France.

2. Department of Oncology, McMaster University, Hamilton Ontario, Canada

3. Department of Cancer Epidemiology and Prevention Research, CancerControl Alberta,  
Alberta Health Services, Calgary, Alberta, Canada

4. Department of Oncology, Cumming School of Medicine, University of Calgary, Calgary,  
Alberta, Canada

5. Department of Community Health Sciences, Cumming School of Medicine, University of  
Calgary, Calgary, Alberta, Canada

**Running title:** Physical activity, MiRNA expression and cancer

**Key words:** microRNAs; c-miRNA; physical activity; exercise; cancer

**Abbreviations list:**

AGO2: Argonaute2

CCL2: Chemokine ligand 2

c-miRNA: circulating miRNA

CPK: Phosphocreatine Kinase

DNA: DeoxyriboNucleic Acid

EGFR: Epidermal Growth Factor Receptor

ERK: Extracellular signal-Regulated Kinases

- 1 FIIT: Frequency, Intensity, Timing and Type
- 2 HDL: High Density Lipoprotein
- 3 IGF1R: Insulin-like Growth Factor 1 Receptor
- 4 LDL: Low Density Lipoprotein
- 5 MAPK: Mitogen-Activated Protein Kinases
- 6 miRNA: microRNA
- 7 mRNA: messenger RNA
- 8 MVB: Multi-Vesicular Bodies
- 9 PA: Physical Activity
- 10 PBMC: Peripheral Blood Mononuclear Cell
- 11 PDCD4: Programmed cell death 4
- 12 PI3K: PhosphoInositide 3-Kinase
- 13 PLC: PhosphoLipase C
- 14 PTEN: Phosphatase and tensin homolog
- 15 RT-qPCR: real-time quantitative reverse transcription polymerase chain reaction
- 16 RNA: RiboNucleic Acid
- 17 RNase: ribonuclease
- 18 RTK: Receptor Tyrosine-Kinase
- 19 STAT3: Signal Transducer and Activator of Transcription 3
- 20 TIMP3: Tissue Inhibitor of MetalloProteinases-3
- 21  $VO_{2max}$ : maximal oxygen consumption
- 22

**Financial support:**

Suzanne Dufresne was supported by the Ministère de l'Enseignement Supérieur et de la Recherche (France) while writing this review paper.

Dr Darren Brenner is supported by a Capacity Development Award Cancer Prevention Development Career Award from the Canadian Cancer Society (#703917).

Dr Christine Friedenreich is supported by an Alberta Innovates-Health Solutions Health Senior Scholar Award and an Alberta Cancer Foundation Weekend to End Women's Cancers Breast Cancer Chair.

Dr Paola Muti is supported by the ArcelorMittal Dofasco Chair.

Dr Amélie Rebillard is financed by Ministère de l'Enseignement Supérieur et de la Recherche (France).

**Corresponding author**

Dr. Darren Brenner  
Department of Cancer Epidemiology and Prevention Research  
CancerControl Alberta, Alberta Health Services  
Holy Cross Centre, Room 513  
Box ACB, 2210 2<sup>nd</sup> Street S.W.  
Calgary, Alberta, CANADA T2S 3C3  
Tel: (403) 698-8178  
Fax: (403) 476-2654  
Email: darren.brenner@albertahealthservices.ca

## 1   **ABSTRACT**

2           The role of circulating microRNAs (c-miRNAs) in carcinogenesis has garnered  
3   considerable scientific interest. MiRNAs may contribute actively to cancer development and  
4   progression, making them potential targets for cancer prevention and therapy. Lifestyle factors  
5   such as physical activity (PA) have been shown to alter c-miRNA expression but the subsequent  
6   impact on cancer risk and prognosis is unknown. To provide a better understanding of how PA  
7   reduces the risk of cancer incidence and improves patient outcomes, we conducted a review of  
8   the impact of PA on c-miRNA expression, which includes a comprehensive synthesis of studies  
9   examining the impacts of acute and chronic exercise on expression of c-miRNAs. While the  
10   variability in methods used to assess miRNA expression creates challenges in comparing and/or  
11   synthesizing the literature, results to date suggest that the circulating form of several miRNAs  
12   known for playing a role in cancer (c-miR-133, c-miR-221/22, c-miR126 and c-let-7) are altered  
13   by both acute and chronic PA. Additional research should develop standardized procedures for  
14   assessing both c-miRNA and PA measurement to improve the comparability of research results  
15   regarding the direction and amplitude of changes in c-miRNAs in response to PA.

## 1 INTRODUCTION

2 MicroRNAs (miRNAs) are implicated in the etiology of various diseases, including  
3 cancer (1), and understanding their biologic characteristics has led to novel insights for cancer  
4 diagnosis, treatment and prognosis (2). MiRNAs are considered to be stable in healthy  
5 individuals, but external factors including lifestyle can impact their expression. Several studies  
6 have reported that physical activity (PA) can modulate miRNAs' expression (3–5). The impact of  
7 PA on reducing cancer risk and improving cancer progression has been well documented, but the  
8 exact mechanisms underlying these relations are not yet completely understood (6,7). A better  
9 understanding of altered expression and function of miRNAs in response to various forms of PA  
10 could assist in clarifying the role of PA in cancer prevention and prognosis.

## 12 MiRNAs and cancer

13 MiRNAs are a class of small noncoding RNAs that regulate gene expression post-  
14 transcriptionally, either by mRNA cleavage, mRNA destabilization, or inhibition of translation  
15 (8). Since the first miRNA was discovered by Lee *et. al.* in 1993, more than 2500 human miRNA  
16 sequences have been identified (miRBase) (9). This rapid expansion of discovery can be partially  
17 explained by the broad importance of miRNAs in regular bodily functions. MiRNAs are  
18 predicted to modulate more than 60% of protein-coding genes (10) and are involved in numerous  
19 integral cellular processes, including development, proliferation, metabolism, and signal  
20 transduction (11,12). Given their regulatory importance, it is not surprising that miRNAs are  
21 widely implicated in carcinogenesis and progression. The mechanistic role of miRNAs in cancer  
22 was first discussed by Johnson *et. al.* in 2005 (13) and the number of studies has been rapidly  
23 expanding ever since. In both solid and hematological tumors, numerous miRNAs have been  
24 found to be altered, and their expression are associated with the severity and stage at diagnosis

(14). In breast cancer, for example, several miRNAs have been observed as very early biomarkers of this disease (15).

The role of miRNAs in carcinogenesis and cancer progression is twofold. Both *in vivo* and *in vitro* studies have demonstrated they target either tumor suppressors or oncogenes (16). Expression of particular miRNAs can downregulate various oncogenes or re-express tumor suppressor genes, leading to tumor suppression, whereas upregulation of other miRNAs, also referred to as oncomiRs, inhibits tumor suppressor genes or over-express oncogenes, thereby promoting tumor cell proliferation and metastasis (16).

MiRNAs are found in tissues and organs and are also released into the circulation in a remarkably stable form (17). The expression of those c-miRNAs has been found to be modified in numerous cancers including, but not limited to: colorectal, prostate, breast, hepatocellular, gastric and head and neck cancers (1,18–24). Altered c-miRNA expression was first observed in B-cell lymphoma patients by Lawrie *et. al.* in 2008 (25) and it is now established that, on average, the expression of 100 c-miRNAs are altered in each cancer type. More importantly, the discovery of c-miRNAs in body fluids resulted in the implication of new possible pathways through which miRNAs may impact tumor development and progression, and led to novel insights for their use as therapeutic tools for cancer (27).

### **MiRNAs as potential mediators of the association between physical activity and cancer**

PA has been shown to reduce cancer risk and progression (6,28–31). Multiple hypothesized biologic mechanisms have been proposed to explain these benefits (6,29,32) including reductions in chronic inflammation, regulation of metabolic factors, changes in insulin resistance, enhanced immune system function, and levels of circulating sex hormones, myokines or adipokines (6,29,32,33). The exact mechanisms, however, whereby PA alters cancer risk and

progression at various cancer sites are not fully understood and additional pathways are likely. Since miRNAs are now well recognized for playing important roles in carcinogenesis and cancer outcomes, they could be involved in the PA-related benefits towards cancer.

The impact of PA on miRNAs has been investigated by several research groups, but mainly in the context of physical performance, muscle pathogenesis and muscular dysfunction related to aging, as well as in muscle disorders associated with diabetes, cancer and inflammation (3). The majority of the miRNAs shown to be altered by either acute or chronic exercise in these studies are muscle-specific miRNAs, also called myomiRs. MiRNAs are differentially expressed within various organs, and if a miRNA is expressed more than 20-fold in a specific tissue compared with the mean of its expression in other tissues, it is thus considered as a tissue-specific miRNA (34). The expression of a variety of miRNAs, and particularly myomiRs, is modulated by PA within muscle tissues (4,35) as well as in plasma (5). Zacharewicz *et. al.* (4) have published a detailed review on miRNA expression within muscles in response to PA. Here we restrict our review to miRNA expression in blood following acute or chronic PA.

C-miRNAs are currently being extensively studied due to several advantages from a clinical and etiological perspective including a remarkable stability in the circulation, resistance to RNase activity (36), and preservation through multiple freeze-thaw cycles (37). Consequently, c-miRNAs can be easily evaluated by standard and relatively inexpensive qRT-PCR techniques, even in low concentrations (38), and their measurement is a non-invasive procedure (39). This is in stark contrast to the analysis of miRNA expression within specific tissues, which require biopsies. Based on these differences, c-miRNAs measured from stored blood samples will prove advantageous in the conduct of large epidemiological studies of PA-related cancer etiology. For these reasons, in this review we focused on the role of c-miRNAs as an underlying mechanism



mediating PA's benefits against cancer through the exploration of potential links between PA-induced c-miRNA modulation and cancer-related c-miRNA expression.

The majority of c-miRNAs are derived from leukocytes and endothelial cells (40) but they can also originate from organs exposed to high blood flow (3). When exercising, the blood flow is significantly elevated, with up to a 80-fold blood flow increase in times of high physical output (41,42), leading to expression changes in miRNAs originating from various organs (43–45), and in particular from skeletal muscles. The release of miRNAs in the blood circulation can originate from destroyed tissues (46,47), especially if the exercise load is great (48). However, increases in the expression of c-miRNAs have also been reported in the absence of cell damage markers in plasma (49). MiRNAs can, in fact, be exported by active transport systems, either encapsulated in extracellular vesicles (such as exosomes) (46,50,51), or associated with protein or lipid-based complexes (e.g., Argonaute2 (*AGO2*), high density lipoproteins (*HDL*) and low density lipoproteins (*LDL*)) (36,52–54). Importantly, increasing evidence shows that miRNAs, through their circulating form, can be transported from donor cells to recipient cells, where they exert their functions (18,54). C-miRNAs are thus being recognized as important components in intercellular communication (53,55,56), and have notably been demonstrated to play key roles in crucial cellular processes, such as apoptosis, proliferation, metastasis, and immunity (57,56). For example, exosomal-miR-105 secreted by breast cancer cells and transferred to endothelial cells have been shown to promote metastasis (58).

## MATERIALS AND METHODS

We conducted a literature search up to March of 2016 in PubMed using the following search strategy: « miRNAs AND exercise » or « miRNAs AND physical activity » or « microRNAs AND exercise » or « microRNAs AND physical activity ». Only studies conducted in humans containing data measuring circulating miRNAs were included. We did not employ a minimum sample size threshold and studies with small numbers of participants were also included, as the total number of studies meeting our inclusion criteria was relatively small. Studies examining populations with disease were excluded, and all studies included in the review presented measurements among healthy subjects as underlying diseases may impact c-miRNAs. To our knowledge, there are no data on the effect of PA on c-miRNA expression in cancer patients. One study has investigated the impact of PA on miRNA expression within tumor tissue based on an animal model, and is included in the discussion section. We also analyzed reference lists of the identified studies for additional relevant articles. The title and abstract were examined, and full text was obtained if the paper seemed eligible. No period or language restrictions were applied. We compiled the results from studies investigating c-miRNA expression in response to a single bout of PA (acute PA) in either normally active individuals (Table 1) or trained subjects (Table 2) as well as the impact of a training period and/or regular PA (chronic PA) on basal c-miRNA expression (Table 3).

Tables 1 and 2 include details on study (design, sample size, timing of blood draws), participant (age, gender, PA level) and protocol characteristics (training program if any, type of exercise (resistance or endurance exercise). Included in these tables are descriptions of the number of miRNAs screened, number of miRNAs altered by PA, miRNAs measured but not altered, and undetectable or unreliable miRNAs.

With regards to the type of exercise, we considered predominantly endurance exercises as exercises involving the whole body and increasing oxidative capacity and aerobic endurance (e.g. a marathon run), while resistance exercises were defined as exercises using machines, weights, or even individuals' body weight to stimulate muscle hypertrophy and increase muscular strength or power (e.g. leg press, squats, pull-ups). We highlight the distinctions between normally active and "trained" participants, based on information provided by the studies. We considered the subjects trained if: 1) the subjects underwent a specific training program described in the study; 2) a large bout of exercise such as a marathon run was included in the protocol as undergoing such an effort implies previous training (even though not always specified in the reviewed studies); 3) the authors stated that the participants were "trained individuals" and/or part of a competitive program. We considered the subjects "untrained" or "normally active" if the participants engaged in recreational activity  $\leq 4$  hours per week and/or if specified in the study. In total, we included 5 studies examining normally active individuals and 10 studies investigating trained individuals, according to our criteria.

To facilitate comparisons across studies, we present the data for c-miRNAs that were examined in multiple studies. We first synthesize and discuss the results obtained for c-miRNA expression following acute PA, followed by a discussion and synthesis of results examining basal miRNA expression in response to chronic PA.

PA has also been shown to modulate miRNA expression within peripheral blood mononuclear cells (PBMCs). PBMCs are leukocytes with a round nucleus, and comprise lymphocytes (T cells, B cells, and natural killer cells), monocytes, and dendritic cells (59). Similar to c-miRNAs, PBMC miRNAs can easily be detected by standard RT-qPCR techniques, but it remains controversial whether their expression is similar or not to whole blood miRNAs (60,61). Thus, even though two different research groups found an impact of acute PA in

1 circulating leukocytes (62–65), those results are not discussed in this review.

2

3

## RESULTS

We have identified a total number of 16 studies in our review, which are displayed in tables 1, 2 and 3. Among these studies, 14 investigated c-miRNA expression in response to a single bout of exercise (66–68,48,69,49,65,70–76), and six (49,66,71,75,77,78) evaluated the impact of chronic exercise on c-miRNA expression.

### **Impact of acute physical activity on c-miRNA expression**

Acute PA refers to a single isolated PA session. It is clear that a single bout of PA alters expression of several c-miRNAs shortly after its completion (Table 1 and 2). However, the modalities of PA (frequency, intensity, timing and type (FITT)) and the individual's physical fitness levels (trained or untrained) may impact those changes. Modifications in several c-miRNAs are not always observed immediately after PA, but are seen only after 2-3 hours of rest. For example, Banzet *et. al.* reported no changes in miRNA expression immediately after the completion of a single bout of eccentric exercise (30 minutes downhill walking) but observed an increase in miR-1, miR-133a, miR-133b, miR-499-5p and miR-208b six hours post bouts of PA (68). These c-miRNAs then returned to baseline expression after periods of inactivity, which has been reported to be relatively slow in some studies (66–68,70,72). For example, decreased expression of c-miR-221 was still observed by Sawada *et. al.* three days post-completion of a resistance exercise session (67).

### ***Muscle- or cardiac-specific miRNAs***

In healthy individuals, muscle- or cardiac specific miR-1, miR-133, miR-206, miR-499, miR-208 are expressed at low levels in the circulation, whereas miR-486 is generally found in higher concentrations (79,74). In response to PA, several studies have shown an increase in the circulating forms of miR-1, miR-133, miR-206, miR-208 and miR-499 immediately after the

completion of various PA modalities (FITT) (48,68,70,72,73,75,76), and a decrease in miR-486 (66). It can be noted that specific exercise modalities appear to impact changes in miRNA levels. Banzet *et. al.* found different c-miRNA expression patterns in normally active participants who underwent a downhill walk versus others who completed an uphill walk (68). Downhill walking is considered an eccentric exercise as the muscles actively lengthen during this effort, whereas uphill walking induces active shortening of the muscles and is thus considered a concentric exercise. Interestingly, Banzet *et. al.* found significantly higher muscle/cardiac specific c-miRNA expression for participants who walked downhill compared to subjects walking uphill (68). Thus, these data suggest that the type of muscle contraction involved in the exercises impacts c-miRNA expression. Furthermore, data from various studies also suggests that muscle/cardiac c-miRNAs can be differentially expressed depending on PA's intensity. For example, when looking at c-miR-133, Uhlemann *et. al.* showed that modifications in c-miR-133 expression in response to PA were correlated with phosphocreatine kinase (*CPK*) activity, a marker for muscle damages: the more damaging the exercise was, the most altered c-miR-133 expression was (48).

Modulations in muscle/cardiac c-miRNA expression post-exercise appear to be temporary, yet the exact delay before a return to baseline values is unclear. When examining measures taken one day post-exercise, most studies showed that muscle and cardiac specific c-miRNAs had returned to their basal value. Others, however, reported a decrease in expression of those c-miRNAs compared to immediately after exercise, but still significantly elevated compared to baseline values. Focusing on c-miR-1 for example, four studies found similar expression of this c-miRNA between pre-exercise and 24 hours post exercise (49,68,75,76), while one other observed significantly higher expression compared to baseline value (but attenuated from expression measured immediately after exercise) (70).

Interestingly, high inter-individual variation in c-miR-499 expression has been reported in

response to an acute bout of endurance exercise in two different studies (68,72) which suggests that c-miR-499 is highly dependent on participants' characteristics, notably their training status. MiR-499 is a cardiac-specific miRNA and studies have shown its circulating form may represent a marker of myocardial injury (80–84). Strenuous exercises such as marathon running can induce transient myocardial injuries (85–88), this risk being higher in less trained individuals (87,89). Therefore, the inter-individual variation observed in c-miR-499 expression following a marathon run may reflect variation in training among participants, and/or cardiac muscle exhaustion.

### ***Other circulating miRNAs***

Non muscle- or cardiac-specific c-miRNAs have also been reported to be impacted by acute PA, but the exact direction and magnitude of those changes remain unclear. Two different research groups observed an up-regulation of c-miR-126, an endothelial-specific miRNA involved in angiogenesis (90), immediately after a single bout of PA (48,72), while no statistical differences in c-miR-126 expression pre- and post-PA have also been reported (48,69). Interestingly, Uhlemann *et. al.* measured c-miR-126 expression before and after an acute bout of PA: subjects were either trained or untrained and the exercises which had to be completed varied among protocols in type, duration and intensity (48). They observed an upregulation in c-miR-126 in response to a single maximal symptom-limited test performed by healthy individuals, as well as in trained men who underwent four hours of bicycling at 70% of their anaerobic threshold, and in trained runners who completed a marathon. However, Uhlemann *et. al.* also reported no significant difference in c-miR-126 expression in trained subjects who performed a resistance training session compared to basal values (48). Since miR-126 is an endothelial-specific miRNA, the authors explained the increase in c-miR-126 expression following acute exercise observed in three of their protocols by exercise-induced endothelial damages, while they

suggested that the resistance exercise session, which resulted in unaltered c-miR-126 expression, did not cause such damages (48).

C-miR-146a, a miRNA known for playing a role in inflammation and immunity (91,92) has also been found to be altered by acute PA, however, the effects appear to be highly dependent on the type of activity intervention. Two studies have reported an up-regulation of this c-miRNA in response to a single bout of endurance exercise in trained individuals (49,72), but Nielsen *et al.* observed opposite results with a decrease in c-miR-146a in response to a similar type of PA in normally active subjects (71). In contrast, Van Craenenbroeck *et al.* reported no change in c-miR-146a expression in response to a single maximal symptom-limited test, while Sawada *et al.* measured no alteration in c-miR-146a expression in healthy men immediately after they performed resistance exercise session but found a decrease in its expression three days post-exercise (67). Interestingly, in trained athletes, Baggish *et al.* found an upregulation miR-146a in response to an acute bout of PA both before and after a training program, the magnitude of the elevation being higher after completion of the training program (93).

Similarly, changes in the circulating expression of miR-221 and miR-222, important players in vascular biology (94), have been reported, but high variations can be found between the studies (49,67,71). In response to a 60 min cycle ergometer exercise bout below anaerobic threshold, c-miR-221 has been found to be down-regulated in normally active participants by Nielsen *et al.* (71), as opposed to Baggish *et al.* who showed an up-regulation of c-miR-221 expression in trained individuals following an acute exhaustive cycling exercise (49). When assessing the impact of a short exhaustive bout of PA on those c-miRNA expression, Sawada *et al.* did not report any changes immediately after PA completion, but found a decrease three days later (67). C-miR-222 expression in response to an acute bout of PA has only been investigated by two different studies: one of them showed an upregulation of this c-miRNA in trained men



undergoing an exhaustive bout of exercise (49), while the other one reported no change in its expression between before and after a resistance exercise session (67). The differences observed between studies are likely attributable to differences in protocols, suggesting that alterations of c-miRNAs in response to acute PA are dependent on the dose and type of exercise (FIIT) as well as participants' characteristics (e.g. age, gender, fitness level, health status, personal history, diet, smoking habits, etc).

### **Impact of chronic physical activity on c-miRNA expression**

Chronic PA is defined as regular PA done over an extended time period (a minimum of several weeks). Interestingly, contrary to what is observed for c-miRNA expression following an acute bout of PA (Table 1 and 2), there is a general trend for a down-regulation of c-miRNAs expression in response to chronic PA (Table 3). Among the six different studies investigating the impact of chronic PA on resting expression of c-miRNAs we included in our review, four were intervention trials (49,66,71,75) and two were observational studies (77,78). The blood samples analyzed in those research projects were taken at rest, at least 12 hours after the last training session.

#### ***Muscle- or cardiac-specific miRNAs***

Cardiac- or muscle specific c-miRNA expression changes in response to chronic PA are hard to measure because of their low concentrations in humans at rest, consequently, there are limited results thus far. For c-miR-1, one study reported no change between its resting expression before and after a 10-week marathon training period in either trained or untrained individuals (75), whereas two other research groups found its expression too low to interpret its measurement (66,78). A trend towards a decrease in c-miR-133 expression in response to a supervised training on a cycle ergometer was found by Nielsen *et. al.* (71), while other studies reported no changes

(49,75) or non-detectable amounts for the quantification of c-miR-133 (66,78). Several studies measured the resting expression of c-miR-206 (66,78), c-miR-208 (66) and c-miR-499 (66,78) before and after a training period, but their expressions were too low to be interpreted. Finally, Aoi *et. al.* found that a four week cycling program down-regulates c-miR-486 basal expression (66).

### ***Other circulating miRNAs***

When evaluating c-let-7d expression, a miRNA that has a crucial role in cell division and differentiation (95,96), Nielsen *et. al.* showed a down-regulation of its baseline expression after a 12 week training period (71) while Bye *et. al.*'s observational study (77) showed that subjects with high maximal oxygen uptake ( $VO_{2max}$ ) ( $145.2 \pm 20.7$  mL/kg<sup>0.75</sup>/min) had a lower c-let-7d expression compared to individuals engaging in similar activity levels but with lower  $VO_{2max}$  ( $101.1 \pm 18.0$  mL/kg<sup>0.75</sup>/min).

For c-miR-21, Nielsen *et. al.* reported that baseline expression was decreased after chronic PA (71), while the results obtained from Bye *et. al.*'s observational study shows that independently of activity levels, individuals with high  $VO_{2max}$  have decreased c-miR-21 basal expression compared to subjects with lower  $VO_{2max}$  (77). However, Baggish *et. al.* found an up-regulation of c-miR-21 after 13 weeks of rowing training, while Wardle *et. al.* reported no alteration in expression between athletes (endurance or resistance trained) and controls, but showed a significant increase in resistance trained individuals when compared to endurance athletes (72,78).

C-miR-221 and c-miR-222 expression in response to chronic PA remains unclear. Baggish *et. al.* reported increased expression of c-miR-222 after a 13-week training program (72), in line with Wardle *et. al.*'s findings when comparing endurance trained athletes with control individuals (78). However, Wardle *et. al.* found a down-regulation of c-miR-222 in

1 resistance trained athletes, and Bye *et. al.*'s reported a decrease in c-miR-222 expression in  
2 individuals with high  $VO_{2max}$  when compared to subjects with a lower  $VO_{2max}$  (77,78). Similarly,  
3 various results have been found for the changes of c-miR-221 expression in response to chronic  
4 PA: Baggish *et. al.* reported an increase in its baseline expression after a training period (72),  
5 whereas Wardle *et. al.* found no statistical differences between trained athletes (endurance or  
6 resistance) and controls, but a down-regulation in resistance trained athletes when compared with  
7 endurance trained athletes (78).

## DISCUSSION

### Potential impact of PA-induced c-miRNA changes on cancer risk, progression, and treatment

Overall, the current results available from the scientific literature suggest that several c-miRNAs are impacted by acute PA (miR-1, miR-133, miR-206, miR-208, miR-499, miR-486, miR-126, miR-146, miR-221 and miR-222) and/or by chronic PA (miR-133, miR-486, let-7d, miR-21, miR-222, and miR-221). Importantly both acute PA and chronic PA appear to induce changes in c-miRNA expression depending on exercise modalities and individuals' fitness status. While results to date are promising and suggest a clear impact of PA on miRNA expression, comparisons between studies are challenging and efforts should be devoted towards standardized protocols and procedures for miRNA collection, analysis and reporting.

The studies included in this review suggest that PA modulates c-miRNA expression in healthy individuals. Several studies have also observed that PA can alter c-miRNA expression within disease populations such as patients with chronic kidney diseases (69) and pre-diabetic individuals (97). To our knowledge, however, no data exists regarding the impact of PA on miRNA in cancer patients. PA also appears to be able to influence miRNA expression within organs other than skeletal muscles (98,99). Taken together, these elements suggest that PA may influence c-miRNA expression in cancer patients, perhaps via c-miRNA intercellular communication. This hypothesized mechanism is illustrated in Figure 1.

Throughout this section, we discuss how modulation of specific c-miRNA expression by PA might impact cancer risk, progression and treatments, and provide some examples of promising early findings. Figure 2 presents hypothesized examples of how PA could impact cancer via c-miRNA modulation using c-miR-133, c-miR-221/222, c-miR-126 and c-let-7.

#### *Impact of miRNAs on cancer risk and development*

Multiple miRNAs are involved in DNA repair, checkpoint functions, tumor suppression, etc. (100), and their modulation by PA might play an important role in cancer risk and progression. For example, PA can alter c-miR-133 expression (48,68,70–73,75,76,101), a well known myoMiR participating in myoblast differentiation which has also been identified as a tumor suppressor (102) in several cancers, including ovarian, colorectal, bladder, breast, prostate, and gastric cancers (101,103–108). MiR-133 has also been shown to be modified within muscle tissues in response to acute and chronic PA. More specifically, acute bouts of exercise seem to increase muscular miR-133 expression (109,110), while training tends to decrease expression (109,26). These data suggest that miR-133 can translocate from muscle tissue to blood vessels, and that this miRNA can impact cancer progression as depicted in Figure 1. Several studies suggest that miR-133 targets several oncogenes, such as the epidermal growth factor receptor (*EGFR*) (101) and the insulin-like growth factor 1 receptor (*IGF1R*) (108,111). When activated, those oncogenes stimulate various pathways causing deregulation in several cell processes, eventually leading to carcinogenesis. For example, activation of *EGFR* pathway leads to the stimulation of intracellular signaling cascades such as the mitogen-activated protein kinases/extracellular signal-regulated kinases (*MAPK/ERK*) pathway, which plays a role in cell cycle progression, differentiation, proliferation, and apoptosis (112,113). When activated, the *EGFR* pathways also stimulate the phosphoinositide 3-kinase - protein kinase B (*PI3K/AKT*) cascade, known for its crucial role in regulation of apoptosis and protein synthesis (114,115). An increase in c-miR-133 by PA originating from muscle tissue, may therefore impact tumor cells and regulate target oncogenes and associated pathways.

Similarly, let-7 is a tumor suppressor and pro-apoptotic miRNA (116) whose expression is decreased in numerous cancers (117), and reported to be modulated in plasma by exercise (71,77). While no direct evidence was identified in humans, in an animal model, breast tumor-

bearing mice who underwent five weeks of interval exercise training (treadmill running) had increased let-7 expression within the tumor itself when compared to breast-tumor sedentary mice (118). Let-7 is known for inhibiting mRNA translation of well-known oncogenes, including the *RAS* family (*HRAS*, *KRAS*, and *NRAS*) (13) and *c-MYC* (119) (120). When activated, *Ras* stimulates several signaling pathways, including *MAPK* cascade and *PI3K/AKT* (121), thereby influencing many cellular functions.

We must note the complex and often paradoxical role of miRNAs, with several miRNAs exerting stimulatory as well as inhibitory effects depending on cancer type or stage of the disease (122–124). MiR-221 and miR-222 for example have a dual role: they can either act as tumor suppressors as oncogenes. On one hand, miR-221 and miR-222 have been found to relente cancer progression in several cancer types, and their expression in the circulation have been found to be altered by PA in several studies (49,71,72,77,78). In gastrointestinal stromal tumors for example, miR-221 and miR-222 are thought to have prophylactic effects by negatively regulating the stem cell factor receptor *KIT*, and are found under-expressed in tumors when compared to healthy tissues (125–127). *KIT* is a receptor tyrosine-kinase (*RTK*), and its activation leads to the stimulation of several intracellular signaling pathways, including the signal transducer and activator of transcription 3 (*STAT3*), *PI3K*, phospholipase C (*PLC*), and the *MAPK* cascade (128–130). *KIT* promotes cell survival, proliferation and motility (131), and its inhibition by miR-221 and miR-222 therefore contributes to carcinogenesis suppression. Modulation of those miRNAs by PA could thereby inhibit *KIT* activation, consequently lowering the risk of developing cancer.

On the other hand, miR-222 and miR-221 have been shown to act as oncomiRs in other cancer types. In lung and liver cancer for example, these miRNAs have been shown to inhibit the action of the phosphatase and tensin homolog (*PTEN*), known to inhibit *MAPK/ERK* pathway, and the tissue inhibitor of metalloproteinases-3 (*TIMP3*), thereby enhancing cell proliferation and

migration through *PI3K/AKT* pathway (132). The dual role played by miR-222 and miR-221 suggest that clarifying the influence of exercise modalities (endurance vs resistance, short vs long duration, regular vs irregular training, etc.), tissue and/or cancer site of interest will be extremely important in subsequent research.

### ***Impact of miRNAs on cancer invasion and metastasis***

PA also modulates c-miRNAs involved in cell proliferation, invasion and metastasis (Figure 2). For example miR-21 is altered by resistance and endurance training (71,72,78) and is also known for participating in tumor invasion. In fact, *in vitro* studies have shown that in several cancer types, miR-21 knock-down mice displayed suppression of cell proliferation and tumor growth (133), as well as reduced invasion and metastasis (133–135). Furthermore miR-21 negatively regulates tumor suppressor programmed cell death 4 (*PDCD4*) (136) and downstream signaling targets (137,138) in colorectal cell lines. The results in this review (Tables 1, 2 and 3) suggest that acute exercise can transiently upregulate c-miR-21 expression (65,93), while chronic exercise (which reflects physiological adaptations) can also lead to alterations in miR-21 expression within circulation (71,77), however, the influence of chronic PA is less clear. By lowering miR-21 expression within cancer cells, PA may restore *PDCD4* and *PTEN* activation and limit cancer proliferation. This is supported by literature that suggests PA is able to modulate miR-21 within multiple tissues/fluids, for example muscle tissue in mice (139) as well as in plasma in healthy human subjects (93,65,71,77,78), as well as in tumors of breast tumor-bearing mice (118) (Figure 1).

The up-regulation of c-miR-126 by acute PA (48,72) might also represent a pathway through which PA impacts cancer progression. In breast cancer, miR-126 regulates the tumor microenvironment composition by directly inhibiting stromal cell-derived factor-1 alpha (*CXCL12*) expression and indirectly suppressing chemokine ligand 2 (*CCL2*) expression in

cancer cells (140). These two chemokines play a role in recruiting stromal cells to the primary tumor microenvironment (141,142), thereby leading to cancer cell invasion and metastasis. Therefore, the increase in c-miR-126 expression in response to acute PA followed by active transport into tumor cells could lead to inhibition of *CXCL12* and *CCL2*, consequently suppressing cancer expansion by modifying the tumor microenvironment composition (48).

Overall, PA-induced c-miRNA expression changes could have the potential to impact tumorigenesis and cancer development. Importantly, several studies suggest that re-expression of tumor-suppressor miRNAs within tumor tissues can inhibit tumor growth, thereby representing a promising therapeutic tool against cancer (18,143–145). For example, re-expression of let-7 within various tumor types have been proven to slow cancer growth (95,146–148) and is thus considered a promising tool against cancers under-expressing let-7 family members (146,149). Additional research, particularly large intervention studies in populations of cancer patients are needed to determine the exact impact of PA on miRNA expression and potential roles in cancer therapy.

### **Limitations of the research to date**

Investigating c-miRNAs in response to PA presents several challenges. First, to date, there are few studies that have compared similar c-miRNAs using comparable methods to enable direct comparisons across studies. Although some consistency has been observed for several c-miRNAs such as miR-133 (48,70,72,73,75,76), contradictory results have often been observed. This inconsistency can largely be attributed to discrepancy in methods across studies, including: 1) differences in sample collection; 2) post-processing of samples and; 3) the use of serum or plasma for miRNAs' extraction.

Different normalization strategies have been used between the reviewed studies. The



current results available on c-miRNA expression changes in response to PA should therefore be carefully interpreted. For example, Nielsen *et. al.* used a stable expressed c-miRNA to account for the biological variation between samples, whereas Baggish *et. al.* used a synthetic spike in approach (71,72). Some studies also included a hemolysis control phase in the study design, since it has been shown that hemolysis occurring during blood collection has substantial impact on the miRNA content in plasma/serum (150). This issue occurs because erythrocytes contain numerous miRNAs that unavoidably will contaminate a plasma sample if the erythrocyte bursts during sampling (151).

Another methodologic factor likely to impact the reliability of the results obtained from the various studies reviewed here is the fact that miRNA concentrations vary between serum and corresponding plasma samples (152). Analyzing and comparing miRNA expression from serum and plasma within or between studies should be done carefully.

To date, miRNA expression in response to PA has mainly been measured in the circulation and in muscles, but it would also be interesting to study the changes in miRNA expression in other tissues. More specifically, investigating miRNA expression after PA in target tissues, and more specifically tumor tissues, would provide novel data of how PA can impact cancer through miRNA modulation. Tissue-specific approaches would also aid in a better understanding of the role of c-miRNAs in specific cancer sites that may result in the development of targeted novel therapies.

## CONCLUSIONS AND FUTURE DIRECTIONS

PA represents a lifestyle behaviour that influences the expression of several c-miRNAs, some of which have been associated with carcinogenesis and cancer progression. Our results suggest that alteration of miRNAs within the circulation is dependent on the type/modality/frequency of exercise as well as on the participants characteristics. Furthermore, the evidence to date on c-miRNA expression in response to PA is limited by small sample sizes without standardized measures of miRNA or physical activity. miRNAs appear to have a meaningful impact on cancer risk and progression, however, the effect vary between cancer types. Therefore, it appears essential to provide a better understanding of how various types of PA in a specific population could impact c-miRNA expression: it could represent a useful tool for healthcare practionners in establishing and monitoring PA programs for cancer patients.

Future research should focus on large epidemiologic studies with standardized blood storage and collection as well as standardized measures of c-miRNA expression and objective measures of PA. Additional consistency is needed to provide more meaningful conclusions as well as standarization of PA measurement. Further investigation into the effects of PA on miRNAs is necessary and could have implication both for cancer prevention, treatment and survival outcomes.



## REFERENCES

1. Ha T-Y. MicroRNAs in Human Diseases: From Cancer to Cardiovascular Disease. *Immune Netw.* 2011;11:135–54.
2. Schoof CRG, Botelho EL da S, Izzotti A, Vasques L dos R. MicroRNAs in cancer treatment and prognosis. *Am J Cancer Res.* 2012;2:414–33.
3. Aoi W. Frontier impact of microRNAs in skeletal muscle research: a future perspective. *Front Physiol.* 2014;5:495.
4. Zacharewicz E, Lamon S, Russell AP. MicroRNAs in skeletal muscle and their regulation with exercise, ageing, and disease. *Front Physiol.* 2013;4:266.
5. Xu T, Liu Q, Yao J, Dai Y, Wang H, Xiao J. Circulating microRNAs in response to exercise. *Scand J Med Sci Sports.* 2015;25:e149–54.
6. Friedenreich CM, Orenstein MR. Physical Activity and Cancer Prevention: Etiologic Evidence and Biological Mechanisms. *J Nutr.* 2002;132:3456S–3464S.
7. Ballard-Barbash R, Friedenreich CM, Courneya KS, Siddiqi SM, McTiernan A, Alfano CM. Physical Activity, Biomarkers, and Disease Outcomes in Cancer Survivors: A Systematic Review. *JNCI J Natl Cancer Inst.* 2012;104:815–40.
8. He L, Hannon GJ. MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev Genet.* 2004;5:522–31.
9. miRBase [Internet]. [cited 2016 Mar 17]. Available from: [www.mirbase.org](http://www.mirbase.org)
10. Friedman RC, Farh KK-H, Burge CB, Bartel DP. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res.* 2009;19:92–105.
11. Bartel DP. MicroRNAs: Genomics, Biogenesis, Mechanism, and Function. *Cell.* 2004;116:281–97.
12. Aoi W, Sakuma K. Does regulation of skeletal muscle function involve circulating microRNAs? *Front Physiol.* 2014;5:39.
13. Johnson SM, Grosshans H, Shingara J, Byrom M, Jarvis R, Cheng A, et al. RAS Is Regulated by the let-7 MicroRNA Family. *Cell.* 2005;120:635–47.
14. Garzon R, Marcucci G, Croce CM. Targeting microRNAs in cancer: rationale, strategies and challenges. *Nat Rev Drug Discov.* 2010;9:775–89.
15. Muti P, Sacconi A, Hossain A, Donzelli S, Ben Moshe NB, Ganci F, et al. Downregulation of microRNAs 145-3p and 145-5p Is a Long-term Predictor of Postmenopausal Breast Cancer Risk: The ORDET Prospective Study. *Cancer Epidemiol Biomarkers Prev.* 2014;23:2471–81.
16. Zhang B, Pan X, Cobb GP, Anderson TA. microRNAs as oncogenes and tumor suppressors. *Dev Biol.* 2007;302:1–12.
17. Mitchell PS, Parkin RK, Kroh EM, Fritz BR, Wyman SK, Pogosova-Agadjanyan EL, et al. Circulating microRNAs as stable blood-based markers for cancer detection. *Proc Natl Acad Sci U S A.* 2008;105:10513–8.
18. Wang W-T, Chen Y-Q. Circulating miRNAs in cancer: from detection to therapy. *J Hematol Oncol J Hematol Oncol.* 2014;7:86.
19. Ng EK, Chong WW, Jin Hongchuan, Lam EK, Shin VY, Yu J, et al. Differential expression of microRNAs in plasma of colorectal cancer patients: A potential marker for colorectal cancer screening. *Gut* [Internet]. 2009; Available from: <http://gut.bmj.com/content/early/2009/02/06/gut.2008.167817.abstract>

20. Ozen M, Creighton CJ, Ozdemir M, Ittmann M. Widespread deregulation of microRNA expression in human prostate cancer. *Oncogene*. 2007;27:1788–93.
21. Iorio MV, Ferracin M, Liu CG, Veronese A, Spizzo R, Sabbioni S, et al. MicroRNA gene expression deregulation in human breast cancer. *Cancer Res*. 2005;65.
22. Hung C-H, Chiu Y-C, Chen C-H, Hu T-H. MicroRNAs in Hepatocellular Carcinoma: Carcinogenesis, Progression, and Therapeutic Target. *BioMed Res Int*. 2014;2014:486407.
23. Liu H-S, Xiao H-S. MicroRNAs as potential biomarkers for gastric cancer. *World J Gastroenterol WJG*. 2014;20:12007–17.
24. Chang SS, Jiang WW, Smith I, Poeta LM, Begum S, Glazer C, et al. MicroRNA alterations in Head and Neck Squamous Cell Carcinoma. *Int J Cancer J Int Cancer*. 2008;123:2791–7.
25. Lawrie CH, Gal S, Dunlop HM, Pushkaran B, Liggins AP, Pulford K, et al. Detection of elevated levels of tumour-associated microRNAs in serum of patients with diffuse large B-cell lymphoma. *Br J Haematol*. 2008;141:672–5.
26. Keller A, Leidinger P, Gislefoss R, Haugen A, Langseth H, Staehler P, et al. Stable serum miRNA profiles as potential tool for non-invasive lung cancer diagnosis. *RNA Biol*. 2011;8:506–16.
27. Kosaka N, Iguchi H, Ochiya T. Circulating microRNA in body fluid: a new potential biomarker for cancer diagnosis and prognosis. *Cancer Sci*. 2010;101:2087–92.
28. Jones LW, Eves ND, Peppercorn J. Pre-exercise screening and prescription guidelines for cancer patients. *Lancet Oncol*. 2010;11:914–6.
29. Rogers C, Colbert L, Greiner J, Perkins S, Hursting S. Physical Activity and Cancer Prevention. *Sports Med*. 2008;38:271–96.
30. Na H-K, Oliynyk S. Effects of physical activity on cancer prevention. *Ann N Y Acad Sci*. 2011;1229:176–83.
31. Betof AS, Lascola CD, Weitzel DH, Landon CD, Scarbrough PM, Devi GR, et al. Modulation of Murine Breast Tumor Vascularity, Hypoxia, and Chemotherapeutic Response by Exercise. *J Natl Cancer Inst [Internet]*. 2015;107. Available from: <http://jnci.oxfordjournals.org/content/107/5/djv040.abstract>
32. McTiernan A. Mechanisms linking physical activity with cancer. *Nat Rev Cancer*. 2008;8:205–11.
33. Brenner DR, Neilson HK, Courneya KS, Friedenreich CM. Physical Activity After Breast Cancer: Effect on Survival and Patient-Reported Outcomes. *Curr Breast Cancer Rep*. 2014;6:193–204.
34. Lee EJ, Baek M, Gusev Y, Brackett DJ, Nuovo GJ, Schmittgen TD. Systematic evaluation of microRNA processing patterns in tissues, cell lines, and tumors. *RNA*. 2008;14:35–42.
35. Pasiakos SM, McClung JP. miRNA Analysis for the Assessment of Exercise and Amino Acid Effects on Human Skeletal Muscle. *Adv Nutr Int Rev J*. 2013;4:412–7.
36. Turchinovich A, Burwinkel B. Distinct AGO1 and AGO2 associated miRNA profiles in human cells and blood plasma. *RNA Biol*. 2012;9:1066–75.
37. Chen X, Ba Y, Ma L, Cai X, Yin Y, Wang K, et al. Characterization of microRNAs in serum: a novel class of biomarkers for diagnosis of cancer and other diseases. *Cell Res*. 2008;18:997–1006.
38. Chen C, Ridzon DA, Broomer AJ, Zhou Z, Lee DH, Nguyen JT, et al. Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Res*. 2005;33:e179–e179.
39. Brase JC, Wuttig D, Kuner R, Sultmann H. Serum microRNAs as non-invasive

biomarkers for cancer. *Mol Cancer*. 2010;9:1–9.

40. Pritchard CC, Kroh E, Wood B, Arroyo JD, Dougherty KJ, Miyaji MM, et al. Blood cell origin of circulating microRNAs: a cautionary note for cancer biomarker studies. *Cancer Prev Res Phila Pa*. 2012;5:492–7.
41. Laughlin MH, Korthuis RJ, Duncker DJ, Bache RJ. Control of Blood Flow to Cardiac and Skeletal Muscle During Exercise. *Compr Physiol* [Internet]. John Wiley & Sons, Inc.; 2010. Available from: <http://dx.doi.org/10.1002/cphy.cp120116>
42. Boushel R, Langberg H, Olesen J, Nowak M, Simonsen L, Bülow J, et al. Regional blood flow during exercise in humans measured by near-infrared spectroscopy and indocyanine green. *J Appl Physiol*. 2000;89:1868–78.
43. Cortez MA, Calin GA. MicroRNA identification in plasma and serum: a new tool to diagnose and monitor diseases. *Expert Opin Biol Ther*. 2009;9:703–11.
44. Hanke M, Hoefig K, Merz H, Feller AC, Kausch I, Jocham D, et al. A robust methodology to study urine microRNA as tumor marker: microRNA-126 and microRNA-182 are related to urinary bladder cancer. *Urol Oncol Semin Orig Investig*. 28:655–61.
45. Zubakov D, Boersma AWM, Choi Y, van Kuijk PF, Wiemer EAC, Kayser M. MicroRNA markers for forensic body fluid identification obtained from microarray screening and quantitative RT-PCR confirmation. *Int J Legal Med*. 2010;124:217–26.
46. Creemers EE, Tijssen AJ, Pinto YM. Circulating MicroRNAs: Novel Biomarkers and Extracellular Communicators in Cardiovascular Disease? *Circ Res*. 2012;110:483–95.
47. Xu L, Yang B, Ai J. MicroRNA transport: A new way in cell communication. *J Cell Physiol*. 2013;228:1713–9.
48. Uhlemann M, Möbius-Winkler S, Fikenzer S, Adam J, Redlich M, Möhlenkamp S, et al. Circulating microRNA-126 increases after different forms of endurance exercise in healthy adults. *Eur J Prev Cardiol*. 2014;21:484–91.
49. Baggish AL, Hale A, Weiner RB, Lewis GD, Systrom D, Wang F, et al. Dynamic regulation of circulating microRNA during acute exhaustive exercise and sustained aerobic exercise training. *J Physiol*. 2011;589:3983–94.
50. Chen X, Liang H, Zhang J, Zen K, Zhang C-Y. Secreted microRNAs: a new form of intercellular communication. *Trends Cell Biol*. 2012;22:125–32.
51. Zhang J, Li S, Li L, Li M, Guo C, Yao J, et al. Exosome and Exosomal MicroRNA: Trafficking, Sorting, and Function. *Genomics Proteomics Bioinformatics*. 2015;13:17–24.
52. Vickers KC, Palmisano BT, Shoucri BM, Shamburek RD, Remaley AT. MicroRNAs are Transported in Plasma and Delivered to Recipient Cells by High-Density Lipoproteins. *Nat Cell Biol*. 2011;13:423–33.
53. Boon RA, Vickers KC. Intercellular Transport of MicroRNAs. *Arterioscler Thromb Vasc Biol*. 2013;33:186–92.
54. Turchinovich A, Samatov TR, Tonevitsky AG, Burwinkel B. Circulating miRNAs: cell–cell communication function? *Front Genet*. 2013;4:119.
55. Katsuda T, Kosaka N, Ochiya T. The roles of extracellular vesicles in cancer biology: Toward the development of novel cancer biomarkers. *PROTEOMICS*. 2014;14:412–25.
56. Yang Q, Diamond MP, Al-Hendy A. The emerging role of extracellular vesicle-derived miRNAs: implication in cancer progression and stem cell related diseases. *J Clin Epigenetics*. 2016;2:13.
57. Fabbri M, Paone A, Calore F, Galli R, Croce CM. A new role for microRNAs, as ligands of Toll-like receptors. *RNA Biol*. 2013;10:169–74.
58. Zhou W, Fong MY, Min Y, Somlo G, Liu L, Palomares MR, et al. Cancer-secreted miR-

- 105 destroys vascular endothelial barriers to promote metastasis. *Cancer Cell*. 2014;25:501–15.
59. Twine NC, Stover JA, Marshall B, Dukart G, Hidalgo M, Stadler W, et al. Disease-associated Expression Profiles in Peripheral Blood Mononuclear Cells from Patients with Advanced Renal Cell Carcinoma. *Cancer Res*. 2003;63:6069–75.
60. Mookherjee N, El-Gabalawy HS. High degree of correlation between whole blood and PBMC expression levels of miR-155 and miR-146a in healthy controls and rheumatoid arthritis patients. *J Immunol Methods*. 2013;400–401:106–10.
61. Atarod S, Smith H, Dickinson A, Wang X-N. MicroRNA levels quantified in whole blood varies from PBMCs. *F1000Research*. 2014;3:183.
62. Radom-Aizik S, Zaldivar F, Oliver S, Galassetti P, Cooper DM. Evidence for microRNA involvement in exercise-associated neutrophil gene expression changes. *J Appl Physiol*. 2010;109:252–61.
63. Radom-Aizik S, Zaldivar F, Leu S-Y, Adams GR, Oliver S, Cooper DM. Effects of Exercise on microRNA Expression in Young Males Peripheral Blood Mononuclear Cells. *Clin Transl Sci*. 2012;5:32–8.
64. Radom-Aizik S, Zaldivar F, Haddad F, Cooper DM. Impact of brief exercise on peripheral blood NK cell gene and microRNA expression in young adults. *J Appl Physiol*. 2013;114:628–36.
65. Tonevitsky AG, Maltseva DV, Abbasi A, Samatov TR, Sakharov DA, Shkurnikov MU, et al. Dynamically regulated miRNA-mRNA networks revealed by exercise. *BMC Physiol*. 2013;13:9–9.
66. Aoi W, Ichikawa H, Mune K, Tanimura Y, Mizushima K, Naito Y, et al. Muscle-enriched microRNA miR-486 decreases in circulation in response to exercise in young men. *Front Physiol*. 2013;4:80.
67. Sawada S, Kon M, Wada S, Ushida T, Suzuki K, Akimoto T. Profiling of Circulating MicroRNAs after a Bout of Acute Resistance Exercise in Humans. Zuo Z, editor. *PLoS ONE*. 2013;8:e70823.
68. Banzet S, Chennaoui M, Girard O, Racinais S, Drogou C, Chalabi H, et al. Changes in circulating microRNAs levels with exercise modality. *J Appl Physiol*. 2013;115:1237–44.
69. Van Craenenbroeck AH, Ledeganck KJ, Van Ackeren K, Jürgens A, Hoymans VY, Fransen E, et al. Plasma levels of microRNA in chronic kidney disease: patterns in acute and chronic exercise. *Am J Physiol - Heart Circ Physiol*. 2015;309:H2008–16.
70. Mooren FC, Viereck J, Krüger K, Thum T. Circulating micrornas as potential biomarkers of aerobic exercise capacity. *Am J Physiol - Heart Circ Physiol*. 2014;306:H557–63.
71. Nielsen S, Åkerström T, Rinnov A, Yfanti C, Scheele C, Pedersen BK, et al. The miRNA Plasma Signature in Response to Acute Aerobic Exercise and Endurance Training. Eckel J, editor. *PLoS ONE*. 2014;9:e87308.
72. Baggish AL, Park J, Min P-K, Isaacs S, Parker BA, Thompson PD, et al. Rapid upregulation and clearance of distinct circulating microRNAs after prolonged aerobic exercise. *J Appl Physiol*. 2014;116:522–31.
73. Gomes CPC, Oliveira-Jr GP, Madrid B, Almeida JA, Franco OL, Pereira RW. Circulating miR-1, miR-133a, and miR-206 levels are increased after a half-marathon run. *Biomarkers*. 2014;19:585–9.
74. Cui SF, Li W, Niu J, Zhang CY, Chen X, Ma JZ. Acute responses of circulating microRNAs to low-volume sprint interval cycling. *Front Physiol*. 2015;6:311.
75. Clauss S, Wakili R, Hildebrand B, Kääb S, Hoster E, Klier I, et al. MicroRNAs as Biomarkers for Acute Atrial Remodeling in Marathon Runners (The miRathon Study ? A Sub-

Study of the Munich Marathon Study). PLoS ONE. 2016;11:e0148599.

76. Min P-K, Park J, Isaacs S, Taylor BA, Thompson PD, Troyanos C, et al. Influence of statins on distinct circulating microRNAs during prolonged aerobic exercise. *J Appl Physiol*. 2016;120:711–20.
77. Bye A, Røsjø H, Aspenes ST, Condorelli G, Omland T, Wisløff U. Circulating MicroRNAs and Aerobic Fitness – The HUNT-Study. Fehlings M, editor. PLoS ONE. 2013;8:e57496.
78. Wardle SL, Bailey MES, Kilikevicius A, Malkova D, Wilson RH, Venckunas T, et al. Plasma MicroRNA Levels Differ between Endurance and Strength Athletes. van Wijnen A, editor. PLoS ONE. 2015;10:e0122107.
79. Aoi W, Naito Y, Takagi T, Tanimura Y, Takanami Y, Kawai Y, et al. A novel myokine, secreted protein acidic and rich in cysteine (SPARC), suppresses colon tumorigenesis via regular exercise. *Gut*. 2013;62:882–9.
80. Corsten MF, Miranda R, Kasmieh R, Krichevsky AM, Weissleder R, Shah K. MicroRNA-21 knockdown disrupts glioma growth in vivo and displays synergistic cytotoxicity with neural precursor cell delivered S-TRAIL in human gliomas. *Cancer Res*. 2007;67.
81. Wang GK, Zhu JQ, Zhang JT, Li Q, Li Y, He J, et al. Circulating microRNA: a novel potential biomarker for early diagnosis of acute myocardial infarction in humans. *Eur Heart J* [Internet]. 2010;31. Available from: <http://dx.doi.org/10.1093/eurheartj/ehq013>
82. Adachi T, Nakanishi M, Otsuka Y, Nishimura K, Hirokawa G, Goto Y, et al. Plasma MicroRNA 499 as a Biomarker of Acute Myocardial Infarction. *Clin Chem* [Internet]. 2010;56. Available from: <http://dx.doi.org/10.1373/clinchem.2010.144121>
83. Zhang L, Chen X, Su T, Li H, Huang Q, Wu D, et al. Circulating miR-499 are novel and sensitive biomarker of acute myocardial infarction. *J Thorac Dis*. 2015;7:303–8.
84. Xin Y, Yang C, Han Z. Circulating miR-499 as a potential biomarker for acute myocardial infarction. *Ann Transl Med*. 2016;4:135.
85. Neumayr G, Gaenger H, Pfister R, Sturm W, Schwarzacher SP, Eibl G, et al. Plasma levels of cardiac troponin I after prolonged strenuous endurance exercise. *Am J Cardiol*. 2001;87:369–71.
86. Urhausen A, Scharhag J ürge., Herrmann M, Kindermann W. Clinical significance of increased cardiac troponins T and I in participants of ultra-endurance events. *Am J Cardiol*. 2004;94:696–8.
87. Neilan TG, Januzzi JL, Lee-Lewandrowski E, Ton-Nu T-T, Yoerger DM, Jassal DS, et al. Myocardial Injury and Ventricular Dysfunction Related to Training Levels Among Nonelite Participants in the Boston Marathon. *Circulation*. 2006;114:2325.
88. Jassal DS, Moffat D, Krahn J, Ahmadie R, Fang T, Eschun G, et al. Cardiac Injury Markers in Non-elite Marathon Runners. *Int J Sports Med*. 2009;30:75–9.
89. Fortescue EB, Shin AY, Greenes DS, Mannix RC, Agarwal S, Feldman BJ, et al. Cardiac Troponin Increases Among Runners in the Boston Marathon. *Ann Emerg Med*. 2007;49:137–143.e1.
90. Wang S, Aurora AB, Johnson BA, Qi X, McAnally J, Hill JA, et al. An Endothelial-specific microRNA Governs Vascular Integrity and Angiogenesis. *Dev Cell*. 2008;15:261–71.
91. Baldeon R. L, Weigelt K, de Wit H, Ozcan B, van Oudenaren A, Sempértegui F, et al. Decreased Serum Level of miR-146a as Sign of Chronic Inflammation in Type 2 Diabetic Patients. PLoS ONE. 2014;9:e115209.
92. Saba R, Sorensen DL, Booth SA. MicroRNA-146a: A Dominant, Negative Regulator of the Innate Immune Response. *Front Immunol*. 2014;5:578.



93. Baggish AL, Hale A, Weiner RB, Lewis GD, Systrom D, Wang F, et al. Dynamic regulation of circulating microRNA during acute exhaustive exercise and sustained aerobic exercise training. *J Physiol [Internet]*. 2011;589. Available from: <http://dx.doi.org/10.1113/jphysiol.2011.213363>
94. Chistiakov DA, Sobenin IA, Orekhov AN, Bobryshev YV. Human miR-221/222 in Physiological and Atherosclerotic Vascular Remodeling. *BioMed Res Int*. 2015;2015:354517.
95. Boyerinas B, Park S-M, Hau A, Murmann AE, Peter ME. The role of let-7 in cell differentiation and cancer. *Endocr Relat Cancer*. 2010;17:F19–36.
96. Roush S, Slack FJ. The let-7 family of microRNAs. *Trends Cell Biol*. 2008;18:505–16.
97. Párrizas M, Brugnara L, Esteban Y, González-Franquesa A, Canivell S, Murillo S, et al. Circulating miR-192 and miR-193b Are Markers of Prediabetes and Are Modulated by an Exercise Intervention. *J Clin Endocrinol Metab*. 2014;100:E407–15.
98. Melo SFS, Barauna VG, Neves VJ, Fernandes T, Lara L da S, Mazzotti DR, et al. Exercise training restores the cardiac microRNA-1 and -214 levels regulating Ca(2+) handling after myocardial infarction. *BMC Cardiovasc Disord*. 2015;15:166.
99. Oghbaei H, Ahmadi Asl N, Sheikhzadeh F, Alipour MR, khamaneh AM. The Effect of Regular Moderate Exercise on miRNA-192 Expression Changes in Kidney of Streptozotocin-Induced Diabetic Male Rats. *Adv Pharm Bull*. 2015;5:127–32.
100. Payne CM, Crowley-Skillicorn C, Bernstein C, Holubec H, Bernstein H. Molecular and cellular pathways associated with chromosome 1p deletions during colon carcinogenesis. *Clin Exp Gastroenterol*. 2011;4:75–119.
101. Cui W, Zhang S, Shan C, Zhou L, Zhou Z. microRNA-133a regulates the cell cycle and proliferation of breast cancer cells by targeting epidermal growth factor receptor through the EGFR/Akt signaling pathway. *FEBS J*. 2013;280:3962–74.
102. Rao X, Di Leva G, Li M, Fang F, Devlin C, Hartman-Frey C, et al. MicroRNA-221/222 confers breast cancer fulvestrant resistance by regulating multiple signaling pathways. *Oncogene*. 2011;30:1082–97.
103. Guo J, Xia B, Meng F, Lou G. miR-133a suppresses ovarian cancer cell proliferation by directly targeting insulin-like growth factor 1 receptor. *Tumor Biol*. 2013;35:1557–64.
104. Dong Y, Zhao J, Wu C-W, Zhang L, Liu X, Kang W, et al. Tumor Suppressor Functions of miR-133a in Colorectal Cancer. *Mol Cancer Res*. 2013;11:1051–60.
105. Yoshino H, Chiyomaru T, Enokida H, Kawakami K, Tatarano S, Nishiyama K, et al. The tumour-suppressive function of miR-1 and miR-133a targeting TAGLN2 in bladder cancer. *Br J Cancer*. 2011;104:808–18.
106. Chiyomaru T, Enokida H, Tatarano S, Kawahara K, Uchida Y, Nishiyama K, et al. miR-145 and miR-133a function as tumour suppressors and directly regulate FSCN1 expression in bladder cancer. *Br J Cancer*. 2010;102:883–91.
107. Kojima S, Chiyomaru T, Kawakami K, Yoshino H, Enokida H, Nohata N, et al. Tumour suppressors miR-1 and miR-133a target the oncogenic function of purine nucleoside phosphorylase (PNP) in prostate cancer. *Br J Cancer*. 2012;106:405–13.
108. Gong Y, Ren J, Liu K, Tang L-M. Tumor suppressor role of miR-133a in gastric cancer by repressing IGF1R. *World J Gastroenterol WJG*. 2015;21:2949–58.
109. Nielsen S, Scheele C, Yfanti C, Åkerström T, Nielsen AR, Pedersen BK, et al. Muscle specific microRNAs are regulated by endurance exercise in human skeletal muscle. *J Physiol*. 2010;588:4029–37.
110. Russell AP, Lamon S, Boon H, Wada S, Güller I, Brown EL, et al. Regulation of miRNAs in human skeletal muscle following acute endurance exercise and short-term endurance training.

J Physiol. 2013;591:4637–53.

111. Huang M-B, Xu H, Xie S-J, Zhou H, Qu L-H. Insulin-Like Growth Factor-1 Receptor Is Regulated by microRNA-133 during Skeletal Myogenesis. de Windt LJ, editor. PLoS ONE. 2011;6:e29173.
112. Neuzillet C, Tijeras-Raballand A, de Mestier L, Cros J, Faivre S, Raymond E. MEK in cancer and cancer therapy. *Pharmacol Ther.* 2014;141:160–71.
113. McCubrey JA, Steelman LS, Chappell WH, Abrams SL, Wong EW, Chang F, et al. Roles of the Raf/MEK/ERK pathway in cell growth, malignant transformation and drug resistance. *Biochim Biophys Acta.* 2007;1773:1263–84.
114. Jorissen RN, Walker F, Pouliot N, Garrett TP., Ward CW, Burgess AW. Epidermal growth factor receptor: mechanisms of activation and signalling. *Exp Cell Res.* 2003;284:31–53.
115. Franke TF, Hornik CP, Segev L, Shostak GA, Sugimoto C. PI3K//Akt and apoptosis: size matters. *Oncogene.* 22:8983–98.
116. Long D, Chan CY, Ding Y. Analysis of microRNA-target interactions by a target structure based hybridization model. *Pac Symp Biocomput Pac Symp Biocomput.* 2008;64–74.
117. Tong AW, Nemunaitis J. Modulation of miRNA activity in human cancer: a new paradigm for cancer gene therapy? *Cancer Gene Ther.* 2008;15:341–55.
118. Isanejad A, Alizadeh AM, Amani Shalamzari S, Khodayari H, Khodayari S, Khor V, et al. MicroRNA-206, let-7a and microRNA-21 pathways involved in the anti-angiogenesis effects of the interval exercise training and hormone therapy in breast cancer. *Life Sci.* 2016;151:30–40.
119. Sampson VB, Rong NH, Han J, Yang Q, Aris V, Soteropoulos P, et al. MicroRNA let-7a down-regulates MYC and reverts MYC-induced growth in Burkitt lymphoma cells. *Cancer Res.* 2007;67.
120. Dang CV. MYC, Metabolism, Cell Growth, and Tumorigenesis. *Cold Spring Harb Perspect Med* [Internet]. 2013;3. Available from: <http://perspectivesinmedicine.cshlp.org/content/3/8/a014217.abstract>
121. Castellano E, Downward J. RAS Interaction with PI3K: More Than Just Another Effector Pathway. Santos E, editor. *Genes Cancer.* 2011;2:261–74.
122. Shenouda SK, Alahari SK. MicroRNA function in cancer: oncogene or a tumor suppressor? *Cancer Metastasis Rev.* 2009;28:369.
123. Di Leva G, Piovan C, Gasparini P, Nguan A, Taccioli C, Briskin D, et al. Estrogen Mediated-Activation of miR-191/425 Cluster Modulates Tumorigenicity of Breast Cancer Cells Depending on Estrogen Receptor Status. Clurman BE, editor. *PLoS Genet.* 2013;9:e1003311.
124. Costa-Pinheiro P, Ramalho-Carvalho J, Vieira FQ, Torres-Ferreira J, Oliveira J, Gonçalves CS, et al. MicroRNA-375 plays a dual role in prostate carcinogenesis. *Clin Epigenetics.* 2015;7:42.
125. Koelz. Down-regulation of miR-221 and miR-222 correlates with pronounced Kit expression in gastrointestinal stromal tumors. *Int J Oncol* [Internet]. 2011 [cited 2016 Jan 28];38. Available from: <http://www.spandidos-publications.com/ijo/38/2/503>
126. Ihle MA, Trautmann M, Kuentstlinger H, Huss S, Heydt C, Fassunke J, et al. miRNA-221 and miRNA-222 induce apoptosis via the KIT/AKT signalling pathway in gastrointestinal stromal tumours. *Mol Oncol.* 2015;9:1421–33.
127. Gits CMM, van Kuijk PF, Jonkers MBE, Boersma AWM, van IJcken WF, Wozniak A, et al. MiR-17-92 and miR-221/222 cluster members target KIT and ETV1 in human gastrointestinal stromal tumours. *Br J Cancer.* 2013;109:1625–35.
128. Hemesath TJ, Price ER, Takemoto C, Badalian T, Fisher DE. MAP kinase links the transcription factor Microphthalmia to c-Kit signalling in melanocytes. *Nature.* 1998;391:298–

301.

129. Chian R, Young S, Danilkovitch-Miagkova A, Rönstrand L, Leonard E, Ferrao P, et al. Phosphatidylinositol 3 kinase contributes to the transformation of hematopoietic cells by the D816V c-Kit mutant. *Blood*. 2001;98:1365–73.

130. Ning Z-Q, Li J, Arceci RJ. Activating Mutations of c-Kit at Codon 816 Confer Drug Resistance in Human Leukemia Cells. *Leuk Lymphoma*. 2001;41:513–22.

131. Linnekin D. Early signaling pathways activated by c-Kit in hematopoietic cells. *Int J Biochem Cell Biol*. 1999;31:1053–74.

132. Garofalo M, Leva GD, Romano G, Nuovo G, Suh S-S, Ngankea A, et al. MiR-221&222 regulate TRAIL-resistance and enhance tumorigenicity through PTEN and TIMP3 down-regulation. *Cancer Cell*. 2009;16:498–509.

133. Si M-L, Zhu S, Wu H, Lu Z, Wu F, Mo Y-Y. miR-21-mediated tumor growth. *Oncogene*. 2006;26:2799–803.

134. Wang N, Zhang C, He J, Duan X, Wang Y, Ji X, et al. miR-21 Down-Regulation Suppresses Cell Growth, Invasion and Induces Cell Apoptosis by Targeting FASL, TIMP3, and RECK Genes in Esophageal Carcinoma. *Dig Dis Sci*. 2013;58:1863–70.

135. Zhu S, Wu H, Wu F, Nie D, Sheng S, Mo Y-Y. MicroRNA-21 targets tumor suppressor genes in invasion and metastasis. *Cell Res*. 2008;18:350–9.

136. Asangani IA, Rasheed SAK, Nikolova DA, Leupold JH, Colburn NH, Post S, et al. MicroRNA-21 (miR-21) post-transcriptionally downregulates tumor suppressor Pcd4 and stimulates invasion, intravasation and metastasis in colorectal cancer. *Oncogene*. 2007;27:2128–36.

137. Yang EV, Sood AK, Chen M, Li Y, Eubank TD, Marsh CB, et al. Norepinephrine Up-regulates the Expression of Vascular Endothelial Growth Factor, Matrix Metalloproteinase (MMP)-2, and MMP-9 in Nasopharyngeal Carcinoma Tumor Cells. *Cancer Res*. 2006;66:10357–64.

138. Leupold JH, Asangani I, Maurer GD, Lengyel E, Post S, Allgayer H. Src Induces Urokinase Receptor Gene Expression and Invasion/Intravasation via Activator Protein-1/p-c-Jun in Colorectal Cancer. *Mol Cancer Res*. 2007;5:485–96.

139. Aoi W, Naito Y, Mizushima K, Takanami Y, Kawai Y, Ichikawa H, et al. The microRNA miR-696 regulates PGC-1 $\alpha$  in mouse skeletal muscle in response to physical activity. *Am J Physiol - Endocrinol Metab*. 2010;298:E799.

140. Zhang Y, Yang P, Sun T, Li D, Xu X, Rui Y, et al. miR-126 and miR-126\* repress recruitment of mesenchymal stem cells and inflammatory monocytes to inhibit breast cancer metastasis. *Nat Cell Biol*. 2013;15:284–94.

141. Teicher BA, Fricker SP. CXCL12 (SDF-1)/CXCR4 Pathway in Cancer. *Clin Cancer Res*. 2010;16:2927–31.

142. Qian B-Z, Li J, Zhang H, Kitamura T, Zhang J, Campion LR, et al. CCL2 recruits inflammatory monocytes to facilitate breast tumor metastasis. *Nature*. 2011;475:222–5.

143. Costa PM, Pedrosa de Lima MC. MicroRNAs as Molecular Targets for Cancer Therapy: On the Modulation of MicroRNA Expression. *Pharmaceuticals*. 2013;6:1195–220.

144. Mishra PJ, Merlino G. MicroRNA reexpression as differentiation therapy in cancer. *J Clin Invest*. 2009;119:2119–23.

145. Rothschild SI. microRNA therapies in cancer. *Mol Cell Ther*. 2014;2:1–8.

146. Barh D, Malhotra R, Ravi B, Sindhurani P. MicroRNA let-7: an emerging next-generation cancer therapeutic. *Curr Oncol*. 2010;17.

147. Wang X, Cao L, Wang Y, Wang X, Liu N, You Y. Regulation of let-7 and its target

oncogenes (Review). *Oncol Lett.* 2012;3:955–60.

148. Kolenda T, Przybyła W, Teresiak A, Mackiewicz A, Lamperska KM. The mystery of let-7d – a small RNA with great power. *Contemp Oncol.* 2014;18:293–301.

149. Chiu S-C, Chung H-Y, Cho D-Y, Chan T-M, Liu M-C, Huang H-M, et al. Therapeutic Potential of MicroRNA Let-7: Tumor Suppression or Impeding Normal Stemness. *Cell Transplant.* 2014;23:459–69.

150. Kirschner MB, Edelman JJB, Kao SC-H, Vallety MP, van Zandwijk N, Reid G. The Impact of Hemolysis on Cell-Free microRNA Biomarkers. *Front Genet.* 2013;4:94.

151. Chen S-Y, Wang Y, Telen MJ, Chi J-T. The Genomic Analysis of Erythrocyte microRNA Expression in Sickle Cell Diseases. Zhang B, editor. *PLoS ONE.* 2008;3:e2360.

152. Wang K, Yuan Y, Cho J-H, McClarty S, Baxter D, Galas DJ. Comparing the MicroRNA Spectrum between Serum and Plasma. Ahuja SK, editor. *PLoS ONE.* 2012;7:e41561.

**Table 1: Effect of acute physical activity on the expression of c-miRNAs in healthy and normal activity subjects**

Reference	Subjects	Type of acute PA	Plasma/serum base	No of miRNAs screened	MiRNAs altered by PA (compared to basal values)		Unaltered miRNAs among the measured ones	miRNAs not detectable or unreliable	
					Immediately post-ex	During recovery			
Acute endurance exercise									
Aoi <i>et. al.</i> , 2013 (66)	10 males 21.5±4.5 yrs	60min cycling exercise (70% VO <sub>2</sub> max)	Serum	-	miR-486↓	Post 3h: miR-486↓	NONE	miR-1 miR-133a miR-133b	miR-206 miR-208b miR-499
Banzet <i>et. al.</i> , 2013(68)	9 males 27 to 36 yrs	30min uphill walking exercise (concentric)	Plasma	-	miR-181b↑ miR-214↑ miR-499-5p↑	Post 2h and 6h: miR-133a↑ miR-133b↑  Post 2h, 6h, and 1 day: miR-499-5p↑	miR-181a	miR-206 miR-208a	
Banzet <i>et. al.</i> , 2013 (68)	9 males 27 to 36 yrs	30min downhill walking exercise (eccentric)	Plasma	-	NONE	Post 2h and 6h: miR-133a↑ miR-133b↑  Post 2h, 6h, and 1 day: miR-499-5p↑  Post 6h: miR-208b↑	miR-499-5p miR-214	miR-181a miR-181b	miR-206 miR-208a
Uhlemann <i>et. al.</i> , 2014 (48)	7 males and 6 females 30.4±2.0 yrs	Maximal cycle ergometry test	Plasma	-	miRNA-126↑	No recovery blood draw time points	miR-133		
Van Craenenbroeck <i>et. al.</i> , 2015 (69)	7 males and 5 females 43.4±4.7 yrs	Maximal cycle ergometry test	Plasma	-	miR-150↑	No recovery blood draw time points	miR-146a miR-21	miR-126 miR-210	
Acute resistance exercise									
Sawada <i>et. al.</i> , 2013 (67)	3 males 29.9±1.2 yrs	Bench press and leg press (5 sets of 10 reps with 1min rest between sets)	Serum	1458	NONE	Post 1 day: miR-149*↑  Post 3 days: miR-146a↓ miR-221↓	miR-20a miR-21 miR-133a miR-210	miR-222 miR-328 miR-1908	miR-299-5p

\*Results of a systematic review of studies investigating the impact of acute PA on c-miRNA expression in healthy and “normal” activity levels humans subjects either immediately after acute exercise completion and/or several hours up to a couple days after.

Abbreviations: ex: exercise; wk: week; yrs: years.

**Table 2: Effect of acute physical activity on the expression of c-miRNAs at various time points in healthy trained subjects**

Reference	Subjects	Training program	Type of acute PA	Plasma/ Serum base	No of miRNAs screened	MiRNAs altered by PA (compared to basal values)			Unaltered miRNAs among the measured ones	miRNAs not detectable or unreliable	
						Immediately post-ex		During recovery			
Acute endurance exercise											
Baggish <i>et. al.</i> , 2011 (49)	10 males 19.1±0.6 yrs	13 weeks rowing training	Maximal cycle ergometry test	Plasma	-	miR-146a↑ miR-222↑	miR-21↑ miR-221↑	No recovery blood draw time points		miR-20a miR-328	miR-210 miR-133a
Tonevitsky <i>et. al.</i> , 2013 (65)	8 males 21.7±2.6 yrs	Regular ski training	30min run on a treadmill (80% VO <sub>2</sub> max)	Serum	200	miR-24-2-5p↑ miR-27a-5p↑ miR-181a-5p↑		<u>Post 1/2h and post 1h:</u> miR-24-2p↓ miR-27a-5p↓ miR-181-5p↓ <u>Post 1h:</u> miR-21-5p↑			
Mooren <i>et. al.</i> , 2014 (70)	14 males 42.8±6.0 yrs	Regular endurance training	Marathon run	Plasma	-	miR-1↑ miR-133a↑ miR-206↑	miR-208b↑ miR-499↑	<u>Post 1 day:</u> miR-1↑ miR-133a↑	miR-206↑	miR-21 miR-155	
Nielsen <i>et. al.</i> , 2014 (71)	13 males 28±8 yrs	12 weeks endurance training (cycle ergometry)	60min cycle ergometry exercise (65% Pmax)	Plasma	188	miR-221↓ miR-30b↓ let7i↓ miR-652↓	miR-106a↓ miR-146a↓ miR-151-3p↓ miR-151-5p↓	<u>Post 1h:</u> miR-338-3p↑ miR-330-3p↑ miR-223↑ miR-139-5p↑ <u>Post 3h:</u> miR-1↑ miR-133a↑‡	miR-143↑ miR-145↑‡ miR-424↑‡		
Baggish <i>et. al.</i> , 2014 (72)	21 males 51.8±1.4 yrs	Regular endurance training	Marathon run	Plasma	-	miR-126↑ miR-1↑ miR-133a↑ miR-208a↑	miR-499-5p↑ miR-208a↑ miR-134↑ miR-146a↑	<u>Post 1 day:</u> miR-499-5p↑			
Uhlemann <i>et. al.</i> , 2014 (48)	13 males 32.4±2.3 yrs	Regular endurance training	4h cycle ergometry exercise (70% anaerobic threshold)	Plasma	-	miR-126↑		<u>Post 1h:</u> NONE <u>Post 1 day:</u> NONE		miR-133	
Uhlemann <i>et. al.</i> , 2014	22 males 56.8±5.2 yrs	Regular endurance	Marathon run	Plasma	-	miR-126↑ miR-133↑		No recovery blood draw time points			

(48)	yrs	training							
Gomes <i>et. al.</i> , 2014 (73)	5 males 31.6±4.39 yrs	Regular endurance training	Half-marathon run	Plasma	-	miR-1↑ miR-133a↑ miR-206↑	No recovery blood draw time points		
Cui <i>et. al.</i> , 2015 (74)	18 males 20.23±0.97 yrs	Regular exercise training (activity type not specified)	Cycle ergometry sprint intervals (2 sets of 30s all out sprints with 4min rest between sets)	Plasma	-	miR-1↓ miR-133a↓ miR-133b↓ miR-122↓ miR-16↓	No recovery blood draw time points	miR-206 miR-499	
Clauss <i>et. al.</i> , 2016 (75)	15 males 40.1±1.4 yrs	Marathon training ≤40km/wk for 10 wk	Marathon run	Plasma	-	miR-1↑ miR-133a↑	<u>Post 1 day:</u> miR-133a↑	miR-30a miR-26a miR-29b	
Clauss <i>et. al.</i> , 2016 (75)	15 males 40.0±1.7 yrs	Marathon training ≥55km/wk for 10 wk	Marathon run	Plasma	-	miR-1↑ miR-133a↑ miR-30a↑	<u>Post 1 day:</u> NONE	miR-26a miR-29b	
Min <i>et. al.</i> , 2016 (76)	20 males and 8 females 53.0±6.5 yrs	Regular endurance training	Marathon run	Plasma	-	miR-1↑ miR-133a↑ miR-206↑	<u>Post 1 day:</u> miR-499-5p↑		
<b>Acute resistance exercise</b>									
Uhlemann <i>et. al.</i> , 2014 (48)	4 males and 7 females 37±2 yrs	Non-supervised regular training (activity type not specified)	Lat pulldown, leg press and butterfly (3 sets of 15 reps with 1min rest between sets)	Plasma	-	miR-133↑	<u>Post 1h:</u> NONE	miR-126	

Legend: ‡ is placed after borderline significant miRNAs

\*Results of a systematic review of studies investigating the impact of acute PA on c-miRNA expression in healthy trained human subjects either immediately after acute exercise completion and/or several hours up to a couple days after.

Abbreviations: No: number; ex: exercise; reps: repetitions; wk: week; yrs: years.



**Table 3: Effect of chronic/regular/long-term physical activity on c-miRNA levels in healthy subjects**

Reference	Type of study	Participants characteristics	Training program	Blood draw timing & plasma /serum base	No of miRNAs screened	MiRNAs altered by chronic PA		miRNAs measured but not altered by chronic PA		miRNAs not detectable/ unreliable	
Chronic endurance exercise											
Baggish <i>et. al.</i> , 2011 (49)	Intervention	10 males 19.1±0.6 yrs Very trained	13 weeks rowing training	At least 12h post exercise -plasma	-	miR-146a↑ miR-222↑ miR-21↑	miR-221↑ miR-20a↑	miR-328 miR-210	miR-133a		
Aoi <i>et. al.</i> , 2013 (66)	Intervention	10 males 21.5±4.5 yrs Normally active	4 weeks cycling training	2 days post training period -serum	-	miR-486↓				miR-1 miR-133a miR-133b miR-206	miR-208b miR-486 miR-499
Nielsen <i>et. al.</i> , 2014 (71)	Intervention	7 males 28±5 yrs Normally active	12 weeks of supervised training on cycle ergometer	3 days and 5 days post training period -plasma	188	miR-342-3p↓ let-7d↓ miR-103↑ miR-107↑ miR-766↓ miR-25↓ miR-148a↓	miR-185↓ miR-21↓ miR-148b↓ ‡ miR-133a↓ ‡ miR-92a↓ ‡ miR-29b↓ ‡				
Wardle <i>et. al.</i> , 2015 (78)	Observational	20 males 22.6±3.7 yrs Trained (endurance athletes) vs Control (not active)	Trained: 13h of training per week	At least 12h post PA -plasma	-	miR-222↑		miR-16-2 miR-20a-1 miR-103a miR-192	miR-451 miR-21 miR-146a miR-221	miR-1 miR-133a miR-206 miR-499	
Clauss <i>et. al.</i> , 2016 (75)	Intervention	15 males 40.1±1.4 yrs Normally active	Marathon training: Running of ≤ 40km/week for 10 weeks	After the training period -plasma	-	NONE		miR-1 miR-133a miR-30a	miR-26a miR-29b		
Clauss <i>et. al.</i> , 2016 (75)	Intervention	15 males 40.0±1.7 yrs Trained	Marathon training: Running of ≥ 55km/week for 10 weeks	After the training period -plasma	-	NONE		miR-1 miR-133a miR-30a	miR-26a miR-29b		
Chronic resistance exercise											

Wardle <i>et al.</i> , 2015 (78)	Observational	20 males 22.2±2.1 yrs Trained (resistance athletes) vs Control (not active)	Trained: 13h of training per week	At least 12h post PA -plasma	-	miR-222↓	miR-16-2 miR-20a-1 miR-103a miR-192	miR-451 miR-21 miR-146a miR-221	miR-1 miR-133a miR-206 miR-499
Unspecified chronic exercise (endurance and/or resistance exercise)									
						Screening:			
						Males and females:			
						miR-210↓			
						miR-125a ↓			
						miR-652 ↑			
						Males only: Women only:			
						miR-151↑ miR-210↓			
						miR-29a↓ miR-125a↓			
						let-7d↓ miR-21↓			
						miR-125 in the validation cohort			
						Validation cohort:			
						Males and females:			
						miR-222↓			
						miR-210↓			
						Males only:			
						miR-21↓			

Legend: ‡ is placed after borderline significant miRNAs

\*Results of a literature review of intervention and observational studies investigating the impact of chronic PA on c-miRNA expression in humans.

Abbreviations: No: number; yrs: years.

## **Figure Legend**

**Figure 1:** Hypothesized pathways of miRNA transport from muscles to tumor cells via the circulation.

MiRNAs can be exported from one cell to another via plasma through various active transport systems including: exosomes developed within multivesicular bodies (MVBs); released into the blood circulation after fusion of the MVBs with the plasma membrane; transport through protein or lipid-based complexes, such as *HDL*, and delivered to recipient cells.

Abbreviations: MVB: Multi-Vesicular Body.

**Figure 2:** Hypothesized mechanisms of action of physical activity on cancer risk and progression through c-miRNA modulation. Multiple c-miRNAs (MiR-126, MiR-221/222, Mir-133 and Let-7) which impact pathways involved in cancer risk and progression have been shown to be altered by physical activity.

# Muscle

# Plasma

# Tumor



