



Genetic characterization of a VanG-type vancomycin-resistant *Enterococcus faecium* clinical isolate

Mohamed Sassi, Francois Guerin, Léonie Leseq, Christophe Isnard, Marguerite Fines-Guyon, Vincent Cattoir, Jean-Christophe Giard

► To cite this version:

Mohamed Sassi, Francois Guerin, Léonie Leseq, Christophe Isnard, Marguerite Fines-Guyon, et al.. Genetic characterization of a VanG-type vancomycin-resistant *Enterococcus faecium* clinical isolate. *Journal of Antimicrobial Chemotherapy*, 2018, 73 (4), pp.852-855. 10.1093/jac/dkx510 . hal-01743400

HAL Id: hal-01743400

<https://univ-rennes.hal.science/hal-01743400>

Submitted on 23 Aug 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Genetic characterization of a VanG-type vancomycin-resistant *Enterococcus faecium*
clinical isolate**

Mohamed SASSI¹, François GUERIN^{2,3}, Léonie LESEC², Christophe ISNARD^{2,3}, Marguerite
FINES-GUYON^{2,3}, Vincent CATTOIR^{4,5}, Jean-Christophe GIARD^{2*}

Running title: VanG-type vancomycin-resistant in *E. faecium*

¹Université Rennes 1, Laboratoire de Biochimie Pharmaceutique, Inserm U1230 - UPRES EA
2311, Rennes, France; ²Université de Caen Normandie, EA4655 U2RM (équipe «Antibio-
résistance»), Caen, France; ³CHU de Caen, Service de Microbiologie, Caen, France; ⁴CHU de
Rennes – Hôpital Ponchaillou, Service de Bactériologie-Hygiène hospitalière, Rennes,
France; ⁵CNR de la Résistance aux Antibiotiques (laboratoire associé «Entérocoques»),
Rennes, France

*Corresponding author. Tel: +(33) (0) 2 31 06 33 28; Fax: +(33) (0) 2 31 06 45 73; E-mail:
jean-christophe.giard@unicaen.fr

(Word count: Synopsis = 210; Text = 1418; 1 Table; 1 Figure; 16 References)

Synopsis

Objectives: to characterize phenotypically and genotypically the first *Enterococcus faecium* clinical isolate harbouring a *vanG* operon.

Methods: The antibiotic-resistance profile of *E. faecium* 16-346 was determined and its whole genome sequenced using the PacBio technology. Attempts to transfer vancomycin resistance by filter mating were performed and the inducibility of expression of the *vanG* operon was studied by reverse-transcription quantitative PCR (RT-qPCR) in the presence or absence of subinhibitory concentrations of vancomycin.

Results: *E. faecium* 16-346 was resistant to rifampicin (Minimum Inhibitory Concentration (MIC) > 4 mg/L) erythromycin (MIC > 4 mg/L), tetracycline (MIC > 16 mg/L) and vancomycin (MIC of 8 mg/L) but susceptible to teicoplanin (MIC of 0.5 mg/L). The strain harboured the *vanG* operon into its chromosome, integrated into a 45.5-kb putative mobile genetic element, similar to that of *E. faecalis* BM4518. We were unable to transfer vancomycin resistance from *E. faecium* 16-346 to *E. faecium* BM4107 and *E. faecalis* JH2-2. Lastly, transcription of the *vanG* gene was inducible by vancomycin.

Conclusions: This is to the best of our knowledge the first report of a VanG-type vancomycin-resistant strain of *E. faecium*. Despite the alarm pulled because of the therapeutic problems caused by vancomycin-resistant enterococci, our work shows that new resistant loci can still be found in *E. faecium*.

Introduction

Commensal bacteria of the human gut microbiota, enterococci have become major opportunistic pathogens.¹ In addition, vancomycin-resistant *Enterococcus faecium* (VREF) is currently a member of the list of bacteria recently edited by the World Health Organization (WHO), for which new antibiotics are urgently needed (<http://www.who.int/mediacentre/news/releases/2017/bacteria-antibiotics-needed/en/>). This points to the worldwide emergence of this species, especially hospital-adapted strains belonging to clonal complex (CC) 17, that leads to major concerns in clinical settings.² Glycopeptide resistance is due to the presence of *van* operons that encode enzymes catalyzing the production of modified peptidoglycan precursors ending in D-Ala-D-Lac (VanA, B, D and M) or D-Ala-D-Ser (VanC, E, G, L and N).² These precursors have much less affinity for vancomycin than the D-Ala-D-Ala motif, leading to a resistant phenotype (MIC of vancomycin > 4 mg/L).² Moreover, these operons are tightly regulated by VanS/VanR-type two-component system, where VanS is a membrane-bound histidine kinase and VanR a transcriptional regulator.³ Because some *van* operons are part of mobile genetic elements, the glycopeptide resistance has spread among enterococci.

The *vanG* operon, in which the VanG ligase allows synthesis of precursor ending in D-Ala-D-Ser, has only been detected in a few *Enterococcus faecalis* clinical isolates and confers moderate levels of resistance to vancomycin (MIC of 16 mg/L) but not to teicoplanin.^{4,5} It is characterized by a “three component” regulatory system comprising an additional repressor (VanU_G) also involved in the control of the *vanG* operon transcription.⁶

We report, to the best of our knowledge, the first clinical isolate of *E. faecium* harbouring a *vanG* operon. The genome of this strain (*E. faecium* 16-346) was completely sequenced and we identified in the chromosome an acquired putative mobile genetic element containing the *vanG* locus.

69

70 **Materials and methods**

71 **Bacterial strains and MIC determination**

72 *E. faecium* 16-346 was obtained in 2016 from a fecal sample of a non-infected 40-year-old
73 female patient hospitalized in the medical center of Sainte-Foy-l'Argentière, France.

74 *E. faecium* BM4107 and *E. faecalis* JH2-2 were used for filter mating experiments.

75 MICs of 20 antibiotics (chloramphenicol, daptomycin, gentamicin, linezolid, rifampicin,
76 trimethoprim/sulfamethoxazole, quinupristin/dalfopristin, tetracycline, erythromycin,
77 oxacillin, ampicillin, penicillin G, vancomycin, levofloxacin, tigecycline, moxifloxacin,
78 clindamycin, streptomycin, ciprofloxacin and nitrofurantoin) were determined using a
79 SensititreTM automated system (Thermo Fisher Scientific, Waltham, MA, USA) according to
80 the manufacturer's instructions.

81

82 **Whole genome sequencing and accession number**

83 The whole genome sequence of the *E. faecium* 16-346 strain was determined with the PacBio
84 technology (GATC Biotech, Konstanz, Germany). The sequence was annotated using the
85 NCBI Prokaryotic Genome Annotation Pipeline.⁷ The sequence was submitted to GenBank
86 and assigned accession no. CP021849.

87

88 **Filter mating**

89 Transfer of vancomycin resistance from *E. faecium* 16-346 to *E. faecium* BM4107 and *E.*
90 *faecalis* JH2-2 was attempted by filter mating. Transconjugants were selected on brain heart
91 infusion (BHI) agar containing rifampicin (60 mg/L), fusidic acid (50 mg/L) and vancomycin
92 (4 mg/L).

93

Reverse-transcription quantitative PCR (RT-qPCR) experiments

The expression of the *vanG* gene was assessed by RT-qPCR using RNAs extracted from bacterial cells (OD₆₀₀ of 0.5) grown in the absence or in the presence of subinhibitory concentrations of vancomycin (i.e., 1 or 2 mg/L). PCR amplification was carried out with primers specific for *vanG* (*vanG*-F: (5'-TTCGTGCAGGCTCTTCCTTT-3'; *vanG*-R: 5'-CACAACCGACTTCAAAGCCG-3') or *adk* (housekeeping reference-gene) as described.⁸

Results and discussion

Characterization of *E. faecium* 16-346

E. faecium 16-346 was resistant to vancomycin (MIC of 8 mg/L). It was also resistant to rifampicin (MIC > 4 mg/L), erythromycin (MIC > 4 mg/L), and tetracycline (MIC > 16 mg/L) but remained susceptible to teicoplanin (MIC of 0.5 mg/L) and to the other antibiotics tested. Screening by specific PCR for the presence of the *vanA*, *vanB*, *vanD*, *vanM* and *vanN* genes, known to confer glycopeptide resistance in this species was negative.² Primers specific for the other *van* operons, not yet described in *E. faecium* were tested, and an amplification of the *vanG* gene was obtained. *vanG*-type operons have only been found in *E. faecalis* clinical isolates from Australia and Canada as well as in various *Clostridium* sp.^{4,5,9,10} In the latter, *vanG* did not confer vancomycin resistance but they may however be considered as a possible reservoir. It has been shown that part of the microflora of the human gastrointestinal tract harboured the *vanG* gene.¹¹ Indeed, from 248 rectal swabs samples, Domingo and coworkers showed that 9.3% were positive for *vanG*, but none was associated with enterococci.¹² Since, to our knowledge, no VanG-type *E. faecium* have been identified so far, we further analyzed this strain by whole genome sequencing. With a size of 2,736,595 bp, the genome harboured 2,824 putative genes (2,730 CDS) (Figure S1). This strain was plasmid free and belonged to sequence type ST121. Search for resistance

genes revealed the presence of the *vanG* operon, but also that of *erm*(Y) and *msr*(C), both involved in macrolide resistance, as well as that of *tet*(M) and *tet*(L) that confer tetracycline resistance (Table 1). These data were in agreement with the resistance phenotype of *E. faecium* 16-346.

Analysis of the *vanG* operon

The sequence of the *vanG* operon was very similar to those from *E. faecalis* BM4518, WCH9 and G1-0247 (Figure 1).^{4,5,13} Indeed, the *van* genes (*U_G*, *R_G*, *S_G*, *Y_G*, *W_G*, *G_G*, *XY_G*, and *T_G*) were between 98.6% and 100% identical (Table 1). Thus, after VanN, VanG is the second ligase able to synthesize D-Ala-D-Ser-ending precursors in *E. faecium*.¹⁴ This is consistent with low level vancomycin resistance (MICs of 8-16 mg/L), since the precursors ending with D-Ala-D-Ser present only a seven-fold less affinity for the antibiotic.¹⁵ In *vanY_G* of *E. faecalis* BM4518, a frameshift mutation is present.¹³ In *E. faecium*, *vanY_G* has no mutations as in the *vanY_G* from *E. faecalis* G1-0247 strain.⁵

Twenty two bp imperfect inverted repeats (IR) framed the mobile genetic element (Figure 1). In *E. faecalis* BM4518, the element is 240 kb long, whereas, in *E. faecium*, its size was of 45.5 kb comprising 43 genes (Table S1). Of note, the sequence of the left IR was identical to that of *E. faecalis* G1-0247 (CGGTAGTACTTCTTTCCCACAA) and diverged from that of BM4518-by 2 bases (CGGTGGTACTGCTTTCCCACAA).^{5,13} Surprisingly, the sequence of the left direct repeat (DR) (TGGA) was not the same as the TTGA sequence of the right DR (Figure 1). Based on the sequence of *E. faecium* Aus0004, TGGA appears as the motif present into the gene in which the element has been inserted¹⁶. It may be hypothesized that the target site of integration could be only the GA 2-bp sequence. To date, in *E. faecalis*, two types of *vanG* operon have been characterized, one from strain BM4518, WCH9 and G1-0247 (*vanG*), and another from N03-0233 (*vanG2*), lacking the *vanY_G* gene and showing also a 2-bp (CA)

DR.⁵ Interestingly, in all the VanG-type strains (clinical isolates or transconjugants) insertion of the *vanG* cluster systematically occurred into the gene encoding a RNA methyltransferase, whereas the *vanG2* is located into the *dctP* gene (encoding a subunits of the TRAP dicarboxylate transporter).¹³ This was not observed in *E. faecium* where the element was integrated into a gene encoding a hypothetical protein (annotated EFAU004_00391) (Figure 1).

Attempts to transfer and inductibility of expression of the *vanG* operon

We were unable to transfer vancomycin resistance from *E. faecium* 16-346 to *E. faecium* BM4107 and *E. faecalis* JH2-2. Similarly, McKessar and co-workers failed to transfer *vanG* operons from *E. faecalis* to *E. faecium*.⁴ In *E. faecalis* BM4518, *erm*(B) was co-transferred with *vanG* and transconjugants could only be obtained on erythromycin-containing plates.¹³ In *E. faecium* 16-346, the mobile genetic element did not harbour *erm*(B) and macrolide resistance was likely due to the presence of the *erm*(Y) and *msr*(C) genes located 231 kb and 522 kb downstream, respectively (Table 1). As in *E. faecalis* G1-0247 and N03-0233, the functions necessary for the transfer should not efficient or the experimental conditions were not adapted.

In order to test if *vanG* expression was inducible, RT-qPCR experiments were carried out from cells grown with 0, 1 or 2 mg/L of vancomycin (corresponding to $\frac{1}{8}$ and $\frac{1}{4}$ of MIC, respectively). As expected, *vanG* transcription was increased 3 and 5 fold ($p<0.05$) in the presence of 1 and 2 mg/L of antibiotic, respectively. This strongly suggests that transcriptional regulation may be similar to that of *E. faecalis* where the repressor VanU_G and the activator VanR_G (both present in *E. faecium*) compete to the same *P_{YG}* promoter leading to a rheostatic control of *vanG* operon expression.⁶

169 **Conclusion**

170 To our knowledge, this is the first report of a *vanG*-type locus acquired by a strain of *E.*
171 *faecium*. This points out that the surveillance of VREF remains crucial to overcome the spread
172 of this major opportunistic pathogen.

173

174 **Acknowledgements**

175 The authors thank M. Auzou for excellent technical assistance.

176 **Funding**

177 This work was supported by internal funding.

178 **Transparency declarations**

179 None to declare.

180

References

- 1 Arias CA, Murray BE. The rise of the *Enterococcus*: beyond vancomycin resistance. *Nat Rev Microbiol* 2012; **10**: 266-78.
- 2 Cattoir V, Giard JC. Antibiotic resistance in *Enterococcus faecium* clinical isolates. *Expert Rev Anti Infect Ther* 2014; **12**: 239-48.
- 3 Depardieu F, Podglajen I, Leclercq R *et al.* Modes and modulations of antibiotic resistance gene expression. *Clin Microbiol Rev* 2007; **20**: 79-114.
- 4 McKessar SJ, Berry AM, Bell JM *et al.* Genetic characterization of *vanG*, a novel vancomycin resistance locus of *Enterococcus faecalis*. *Antimicrob Agents Chemother* 2000; **44**: 3224-8.
- 5 Boyd DA, Du T, Hizon R *et al.* VanG-type vancomycin-resistant *Enterococcus faecalis* strains isolated in Canada. *Antimicrob Agents Chemother* 2006; **50**: 2217-21.
- 6 Depardieu F, Mejean V, Courvalin P. Competition between VanU(G) repressor and VanR(G) activator leads to rheostatic control of *vanG* vancomycin resistance operon expression. *PLoS Genet* 2015; **11**: e1005170.
- 7 Angiuoli SV, Gussman A, Klimke W *et al.* Toward an online repository of standard operating procedures (SOPs) for (meta)genomic annotation. *Omics* 2008; **12**: 137-41.
- 8 Sinel C, Cacaci M, Meignen P *et al.* Subinhibitory concentrations of ciprofloxacin enhance antimicrobial resistance and pathogenicity of *Enterococcus faecium*. *Antimicrob Agents Chemother* 2017; **61**: 2763-16.
- 9 Berthet N, Périchon B, Mazuet C *et al.* A *vanG*-type locus in *Clostridium argentinense*. *J Antimicrob Chemother* 2015; **70**: 1942-5.
- 10 Peltier J, Courtin P, El Meouche I *et al.* Genomic and expression analysis of the *vanG*-like gene cluster of *Clostridium difficile*. *Microbiology* 2013; **159**: 1510-20.

- 206 **11** Domingo MC, Huletsky A, Giroux R *et al.* *vanD* and *vanG*-like gene clusters in a
207 *Ruminococcus* species isolated from human bowel flora. *Antimicrob Agents Chemother* 2007;
208 **51**: 4111-7.
- 209 **12** Domingo MC, Huletsky A, Giroux R *et al.* High prevalence of glycopeptide resistance
210 genes *vanB*, *vanD*, and *vanG* not associated with enterococci in human fecal flora. *Antimicrob*
211 *Agents Chemother* 2005; **49**: 4784-6.
- 212 **13** Depardieu F, Bonora MG, Reynolds PE *et al.* The *vanG* glycopeptide resistance operon
213 from *Enterococcus faecalis* revisited. *Mol Microbiol* 2003; **50**: 931-48.
- 214 **14** Lebreton F, Depardieu F, Bourdon N *et al.* D-Ala-D-Ser VanN-type transferable
215 vancomycin resistance in *Enterococcus faecium*. *Antimicrob Agents Chemother* 2011; **55**:
216 4606-12.
- 217 **15** Courvalin P. Vancomycin resistance in Gram-positive cocci. *Clin Infect Dis* 2006; **42**
218 Suppl 1: S25-34.
- 219 **16** Lam MM, Seemann T, Bulach DM *et al.* Comparative analysis of the first complete
220 *Enterococcus faecium* genome. *J Bacteriol* 2012; **194**: 2334-41.

221

Table 1: Antimicrobial-resistance genes identified in the genome sequence of *E. faecium* 16-246.

| Gene | % Identity | Position in contig | Phenotype | Accession no. |
|--------------------------|---------------------|--------------------|--|---------------|
| <i>vanT_G</i> | 100.00 ^a | 1442739..1444877 | Vancomycin resistance | AY271782 |
| <i>vanXY_G</i> | 100.00 ^a | 1444870..1445634 | Vancomycin resistance | AY271782 |
| <i>vanG</i> | 100.00 ^a | 1445631..1446680 | Vancomycin resistance | AY271782 |
| <i>vanW_G</i> | 99.88 ^a | 1446682..1447527 | Vancomycin resistance | AY271782 |
| <i>vanY_G</i> | 100.00 ^b | 1447603..1448442 | Vancomycin resistance | DQ212986 |
| <i>vanS_G</i> | 99.82 ^a | 1448579..1449683 | Vancomycin resistance | AY271782 |
| <i>vanR_G</i> | 100.00 ^a | 1449697..1450404 | Vancomycin resistance | AY271782 |
| <i>vanU_G</i> | 100.00 ^a | 1450406..1450633 | Vancomycin resistance | AY271782 |
| <i>tet</i> (M) | 97.45 ^c | 1707444..1709362 | Tetracycline resistance | EU182585 |
| <i>tet</i> (L) | 99.85 ^d | 1709491..1710832 | Tetracycline resistance | M29725 |
| <i>erm</i> (Y) | 81.51 ^e | 1715249..1715977 | Macrolide resistance | AB014481 |
| <i>msr</i> (C) | 98.99 ^f | 2006190..2007668 | Macrolide, Lincosamide Streptogramin resistance | AY004350 |

^aIdentity with the homologous gene from *E. faecalis* BM4518

^bIdentity with the homologous gene from *E. faecalis* G1-0247

^cIdentity with the homologous gene from *Streptococcus suis* T2S3

^dIdentity with the homologous gene from pLS1 plasmid of *Streptococcus agalactiae*

^eIdentity with the homologous gene from the pMS97 plasmid of *Staphylococcus aureus* RN4220

^fIdentity with the homologous gene from *E. faecium* TX2465

Legends

Figure 1: Schematic of the chromosome region harbouring the *vanG* operon of *E. faecium* 16-346, compared to the corresponding region of *E. faecalis* BM4518.

DR-L, direct repeat left; DR-R, direct repeat right; IR-L, inverted repeat left; IR-R, inverted repeat right.

Table S1: Genes identified in the mobile genetic element containing the *vanG* operon in *E. faecium* 16-346.

Figure S1: Circle representation of the genome of *E. faecium* 16-346 compared to four other *E. faecium* genomes.

Figure 1