

Synthesis of (Poly)halo-substituted Diarylsulfones through Palladium Catalyzed C-H Bond Sulfonylation using (Poly)Halobenzenesulfonyl Chlorides

Arpan Sasmal, Jitendra K. Bera, Henri Doucet, Jean-François Soulé

▶ To cite this version:

Arpan Sasmal, Jitendra K. Bera, Henri Doucet, Jean-François Soulé. Synthesis of (Poly)halo-substituted Diarylsulfones through Palladium Catalyzed C-H Bond Sulfonylation using (Poly)Halobenzenesulfonyl Chlorides. European Journal of Organic Chemistry, 2018, 2018 (44), pp.6114-6120. 10.1002/ejoc.201800819. hal-01915420

HAL Id: hal-01915420 https://univ-rennes.hal.science/hal-01915420

Submitted on 7 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Synthesis of (Poly)halo-substituted Diarylsulfones through Palladium Catalyzed C–H Bond Sulfonylation using (Poly)Halobenzenesulfonyl Chlorides

Arpan Sasmal,^[a] Jitendra K. Bera,^[b] Henri Doucet^{*[a]} and Jean-François Soulé^{*[a]}

Abstract: The reactivity of (poly)halo-substituted benzenesulfonyl chlorides in palladium-catalyzed *ortho*-directed C–H bond sulfonylation of 2-arylpyridines was investigated. The use of Ag₂CO₃ in concert with Cu(OAc)₂ was found to be critical to promote the chemoselective sulfonylation reaction. Fluoro-, chloro-, bromo-, and even iodo-benzenesulfonyl chlorides react nicely to afford halo-substituted diarylsulfones in good to high yields without cleavage of the C–halo bonds, allowing further transformations.

Introduction

Unsymmetrical diarylsulfones are important motifs, which are embedded in many drugs and pharmacologically active compounds. For examples, Dapsone, which exhibits antibacterial properties, is administrated for the treatment of leprosy and for the treatment of pneumocystis pneumonia (Figure 1 left).^[1] L-737, 126 is a non-nucleoside inhibitor of HIV-1 reverse transcriptase (Figure 1 middle).^[2] Its structure contains a chlorine atom, which is very important for its activity. CX157 is a fluorinated fused diarylsulfone, which has been in phase II clinical trials for the treatment of depression (Figure 1 right). There is a high demand for the preparation of diarylsulfones using mild reaction conditions with high functional group tolerance, including C–halo bonds.



Figure 1. Examples of Relevant Diarylsulfones

[a] Mr. A. Sasmal, Dr. H. Doucet, and Dr. J.-F. Soulé Univ Rennes, CNRS, ISCR UMR 6226, F-35000 Rennes, France email : henri.doucet@univ-rennes1.fr, jean-francois.soule@univrennes1.fr
[b] Prof. J. K. Bera Department of Chemistry and Center for Environmental Science and Engineering Indian Institute of Technology Kanpur Kanpur 208016. India

Supporting information for this article is given via a link at the end of the document.((Please delete this text if not appropriate))

Although there are different reported methods for the synthesis of diarylsulfones, most of them suffer from limitations. For example, the oxidation of sulfides could be only achieved with substrates which are compatible with oxidizing agents.^[3] Friedel–Crafts-type sulfonylation of arenes are often carried out under harsh reaction conditions and are not regiospecific, leading to mixtures of compounds.^[4] Nucleophilic substitution of sulfonate esters or coupling of sulfinates required the use of organometallic reagents.^[5] Alternatively, transition-metal catalyzed C-H bond activation/functionalization allows the preparation of organic molecules in more efficient and shorter ways with higher atom economy compared to the traditional cross-coupling reactions.^[6] Most of the reported protocols for ortho-regioselective aryl C-H bond sulfonylation reaction required a chelate directing group^[7] or occurred at a remote C-H bond.^[8] In 2009, V. Dong and coworkers reported Pd-catalyzed ortho-directed C-H bond sulfonylation of 2-arylpridines from benzenesulfonyl chlorides (Figure 2a).^[9] However, no examples using halo-substituted benzenesulfonyl chlorides were described, although it could allow the synthesis of halo-substituted diarylsulfones, which are useful synthetic intermediates.^[10] Sodium arylsulfinates have been also employed for sulfonylation, although scarce examples involving chemoselective C-H bond functionalization tolerant to C-halo bond were reported.^[11] In 2012, formal C-H bond sulfonylations of 2-arylpyridines were achieved through sequential metal-free ortho-directed borylation followed by copper-catalyzed sulfonylation using sodium arylsulfinates. These conditions tolerated a chloro substituent on the phenyl unit of 2phenylpyridine derivatives. However, there was no example tolerant to C-Br or C-I bonds (Figure 2b).^[12] In 2016, Li and coworkers reported the rhodium-catalyzed sulfonylation of 2arylpyridines (Figure 2c).^[13] The reaction proceeded by initial Rh(III)-catalyzed C-H hyperiodination of arene at room temperature followed by uncatalyzed nucleophilic functionalization using sodium arylsulfinates. The procedure required the use of 1.6 equivalent of hypervalent iodine, making this protocol not very user- and eco-friendly for the preparation of (poly)halo-substituted diarylsulfones. The authors shown that the reaction was tolerant to C-CI and C-Br bonds. However, no example with substrate bearing a C-I bond was reported.

In 2015, our group shown that (poly)halo-benzenesulfonyl chlorides, which are commercially available at an affordable cost and are easy to handle, undergo palladium-catalyzed chemoand regio-selective C–H bond arylation of heteroarenes allowing the formation of (poly)halo-substituted bi(hetero)aryls, including iodo-substituted ones.^[14] In our continuous efforts to employ (poly)halo-benzenesulfonyl chlorides in chemoselective C–H

bond functionalizations, we wish to report a practical and costeffective method for the synthesis of unsymmetrical halodiarylsulfones through a palladium catalyzed C–H bond sulfonylation using (poly)halo-benzenesulfonyl chlorides (Figure 2d).

a) Preparation of Diarylsulfones through Palladium-Catalyzed C–H Bond Sulfonylation with Arylsulfonyl Chlorides (V. Dong)^{[9]}



b) Two Steps Procedure for the Preparation of Halo-Substituted Diarylsulfones *via* Sequential Borylation and Copper Catalysis (H. Fu)^[12]



11 examples including 2 containing a C-CI bond

c) Two Steps Procedure for the Preparation of Halo-Substituted Diarylsulfones via Combination of Rhodium-Catalyzed C–H Bond Activation and Hypervalent



Figure 2. C-H Bond Functionalization using Benzenesulfonyl chloride or Sodium Sulfinates

Results and Discussion

2-Phenylpyridine and 4-bromobenzenesulfonyl chloride were selected as model substrates to study the Pd-catalyzed chemoselective C-H bond sulfonylation. Using our previous conditions -namely 5 mol% PdCl₂(CH₃CN)₂ in the presence of Li₂CO₃ as base in 1,4-dioxane- which are tolerant to C-halo bonds in Pd-catalyzed desulfitative arylation of heteroarenes,[14] no coupling product was obtained (Table 1, entry 1). The conditions reported by Dong et al. for the sulfonylation of 2phenylpyridine with benzenesulfonyl chloride (i.e. 10 mol% PdCl₂(CH₃CN)₂ associated to K₂CO₃ as base in 1,4-dioxane in the presence of 4Å molecular sieves), [9a] also failed, and only degradation products from 4-bromobenzenesulfonyl chloride were observed (Table 1, entry 2). When the reaction was carried out without molecular sieves, again no formation of 1 was observed (Table 1, entry 3). However, the use of Aq₂CO₃ as base allowed the formation of 2-(2-((4bromophenyl)sulfonyl)phenyl)pyridine 1 in 27% vield (Table 1. entry 4). Noteworthy, the reaction is highly chemoselective toward S-CI bond activation and no coupling product derived from the activation of the C-Br bond was detected by ¹H NMR or GC/MS analysis of the crude mixture. This chemoselectivity might be explained by the fast oxidation addition of the sulforvl chloride function to a Pd^{II} species as reported by Dong.^[9] In the presence of benzenesulfonyl chlorides using Ag₂CO₃ as base, the formation of Pd⁰ species is suppressed and no oxidative addition of the C-Br bond to palladium is observed. Pd(OAc)₂ displayed a higher catalytic activity for this sulfonylation reaction than PdCl₂(CH₃CN)₂ and allowed the formation of the brominated sulfone 1 in 50% yield (Table 1, entry 5). Previously, we had shown that copper salts can affect the outcome of coupling reactions with benzenesulfonyl chlorides, especially for substrates containing a nitrogen atom.[15] Therefore, we investigated the effect of set of copper salts as oxidant on the reactivity (Table 1, entries 6-9). The reaction was completely inhibited in the presence of 1 equivalent of Cul; whereas, better yields of 65% and 75% in 1 were obtained in the presence of 1 equivalent of CuOTf or CuBr as additives. Finally, Cu(OAc)2 nH2O was found the be the most efficient additive for this reaction. In the presence of $Cu(OAc)_2 \cdot nH_2O$, the amount of Ag_2CO_3 can be decreased to 1 equivalent without loss of activity, and the bromosubstituted diarylsulfone 1 was isolated in 78% yield (Table 1, entry 10). However, the use of only 0.5 equivalent of $Cu(OAc)_2 nH_2O$ resulted in a lower yield of **1** (Table 1, entry 11). Other solvent such as CPME, DMF, or 1,2-dichloroethane (CICH₂CH₂CI) did not allow any improvement (Table 1, entries 12-14). Notably, in all cases, we did not observe the formation of arylated products resulting from a desulfitative coupling,[16] probably due to the conformation of the cyclometalated palladium(IV) sulfinate, which allows quite easily the formation of the carbon-sulfur bond.[9b]



[a] Determined by ¹H NMR spectroscopy using 1,3,5-trimethoxybenzene as internal standard. The number in parentheses shows the isolated yield. [b] Reaction performed in CPME instead of 1,4-dioxane. [c] Reaction performed in DMF instead of 1,4-dioxane. [d] Reaction performed in CICH₂CH₂Cl instead of 1,4-dioxane.

Having the optimized reaction conditions in hands, we investigated the reactivity of other benzenesulfonyl chlorides bearing C-halo bonds (Scheme 1). The reaction was also tolerant toward the very reactive C-I bonds, as 4iodobenzenesulfonyl chloride chemoselectively reacted with 2phenylpyridine to afford the diarylsulfone 2 in 75% yield, without C-I bond cleavage. The synthesis of diarylsulfones bearing C-CI or C-F bonds, such as compounds 3-5, can also be achieved in good yields using this protocol. From 3-bromo-, 3-chloro- or 3fluorobenzenesulfonyl chlorides, the halo-substituted diarylsulfones 6-8 were obtained in 71%, 69% and 72% yields, respectively; whereas, no reaction occurred using 2bromobenzenesulfonyl chloride.



Scheme 1. Reactivity of Halo-Substituted Benzenesulfonyl Chlorides in Pd-Catalyzed C–H Bond Sulfonylation of 2-Phenylpyridine

We next evaluated the reaction scope with respect to the 2arylpyridine derivatives in reactions with halo-substituted benzenesulfonyl chlorides (Scheme 2). 2-(Naphthalen-1yl)pyridine also underwent palladium-catalyzed C-H bond sulfonylation with 4-bromo- or 4-iodo-benzenesulfonyl chlorides to afford the diarylsulfones 10 and 11 in 72% and 68% yields, respectively. An electron-withdrawing group such as trifluoromethyl on the phenyl ring of the 2-phenylpyridine slightly decreased the efficiency of the C-H bond sulfonylation reaction, as the desired sulfonated product 12 was obtained in 70% yield. In contrast, when the phenyl ring bears an electron-donating group such as a methyl, the reaction with 4-iodo- or 4-bromobenzenesulfonyl chlorides delivered the halo-substituted diarylsulfones 13 and 14 in 77% and 78% yields, respectively. The substituent on the pyridine ring seems to have a minor influence on the reactivity of 2-phenylpyridine derivatives in Pdcatalyzed C-H bond sulfonylation with halo-benzenesulfonyl chlorides. Indeed, from 2-phenyl-5-(trifluoromethyl)pyridine and 4-bromo- or 4-iodo-benzenesulfonyl chlorides, the optimized reaction condition allowed the exclusive formation of the halosubstituted diarylsulfones 15 and 16 in good yields. 2-Phenylnicotinonitrile also reacted nicely with 4-bromo-, or 4-iodo-, benzenesulfonyl chlorides to deliver the halo-substituted diarylsulfones 17 and 18 in 68 and 61% yields, respectively.



Scheme 2. Scope of 2-Arylpyridine Derivatives in Pd-Catalyzed C–H Bond Sulfonylation with Halo-Benzenesulfonyl Chlorides

So far, all the reported protocols for the ortho-sulfonylation of halo-substituted 2-arylpyridine derivatives required two steps procedures (Figure 2b and 2c).^[12-13] Therefore, we investigated the reactivity of halo-substituted 2-arylpyridine derivatives in such Pd-catalyzed direct sulfonylation reaction (Scheme 3). Under optimized reaction conditions, ortho-C-H bond sulfonylation of 2-(4-bromophenyl)pyridine was selectively achieved with 4-nitro or 4-(trifluroromethyl)benzenesulfonyl chlorides to give the diarylsulfones 19 and 20 in 78% and 82% yields, respectively without cleavage of the C-Br bonds. A similar reactivity's trend was observed with 3-chloro-2-phenylpyridine, which allowed the formation of 21 and 22 in high yields from with 4-nitro- and 4-(trifluroromethyl)benzenesulfonyl chlorides, respectively. From 3chloro-2-phenylpyridine and halo-benzenesulfonyl chlorides, the polyhalogenated diarylsulfones 23 and 24 were synthetized in good yields. Noteworthy, no cleavage of both C-halo bond -i.e., on 2-phenylpyridine derivatives and on halo-benzenesulfonyl chlorides- was observed.



Scheme 3. Reactivity of Halo-substituted 2-Arylpyridine Derivatives in Pd-Catalyzed C–H Bond Sulfonylation with (Halo)-Benzenesulfonyl Chlorides

We further demonstrated the potential of chemoselective C–H bond sulfonylation of 2-arylpyridines with 4-halobenzene sulfonyl chlorides by performing successive C–H bond functionalizations (Scheme 4). The direct arylation of 2-isopropyl-4-methylthiazole at C5 position using 2-(2-((4-bromophenyl)sulfonyl)phenyl)pyridine (1) as aryl source was performed using the our previously optimized catalytic system [i.e., PdCI(C₃H₅)(dppb) (1 mol%) in the presence of KOAc (2 equiv.) as base in DMA at 150 °C],^[17] to afford the product **25** in 85% yield.



Scheme 4. Reactivity of Halo-substituted 2-Arylpyridine Derivatives in Pd-Catalyzed C–H Bond Arylation using a Thiazole Derivative

Conclusions

We have developed a one step synthetic scheme for access to (poly)halosubstituted diarylsulfones from commercially available substrates. The reaction proceeds *via* Pd-catalyzed C–H sulfonylation of from 2-arylpyridine derivatives using poly(halo)-substituted benzenesulfonyl chlorides. The use of Ag₂CO₃ in concert with Cu(OAc)₂ in the presence of a catalytic amount of phosphine-free Pd(OAc)₂ was found to be critical to promote these chemoselective sulfonylation reactions. These reaction conditions were found to be tolerant to C–Br and even C–I bonds, allowing further functionalizations. These reaction conditions also tolerated C–Br and C-Cl bonds on the 2-arylpyridines.

Experimental Section

General: All reactions were carried out under argon atmosphere with standard Schlenk-tube techniques. HPLC grade 1,4-dioxane was stored under argon and used without further purification. ¹H NMR spectra were recorded on Bruker GPX (400 MHz or 300 MHz) spectrometers. Chemical shifts (d) were reported in parts per million relative to residual chloroform (7.26 ppm for ¹H ; 77.0 ppm for ¹³C), constants were reported in Hertz. ¹H NMR assignment abbreviations were the following: singlet (s), doublet (d), triplet (t), quartet (q), doublet of doublets (dd), doublet of triplets (dt), and multiplet (m). ¹³C NMR spectra were recorded at 100 MHz on the same spectrometer and reported in ppm. All reagents were weighed and handled in air.

General procedure for synthesis of (poly)halo-substituted diarylsulfones: To a 25 mL oven dried Schlenk tube, arylsulfonyl chloride (1.5 mmol), 2-phenylpyridine derivative (1 mmol), Ag_2CO_3 (0.275 g, 1 mmol), $Cu(OAc)_2$ nH₂O (181 mg, 1mmol), 1,4-dioxane (2 mL) and Pd(OAc)₂ (11.1 mg, 0.05 mmol) were successively added. The reaction mixture was evacuated by vacuum-argon cycles (5 times) and stirred at 140 °C (oil bath temperature) for 16 h (see tables and schemes). After cooling the reaction at room temperature and concentration, the crude mixture was purified by silica column chromatography to afford the (poly)halo-substituted diarylsulfones.

2-(2-((4-Bromophenyl)sulfonyl)phenyl)pyridine (1): 2-phenylpyridine (0.155 g, 1 mmol) and 4-bromobenzenesulfonyl chloride (0.383 g, 1.5 mmol) affords **1** in 78% (0.292 g) yield. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.43 (s, 1H), 8.35 (d, *J* = 7.7 Hz, 1H), 7.80 (t, *J* = 7.7 Hz, 1H), 7.73 – 7.63 (m, 2H), 7.57 (d, *J* = 7.7 Hz, 1H), 7.48 (d, *J* = 8.6 Hz, 2H), 7.41 (d, *J* = 7.2 Hz, 1H), 7.38 – 7.33 (m, 3H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 156.6, 148.5, 140.9, 140.5, 138.9, 135.7, 133.5, 132.1, 131.9, 129.2, 129.2, 128.8, 128.0, 125.8, 122.7. This is a known compound and the spectral data are identical to those reported in literature.^[13]

2-(2-((4-lodophenyl)sulfonyl)phenyl)pyridine (2): 2-phenylpyridine (0.155 g, 1 mmol) and 4-iodobenzenesulfonyl chloride (0.454 g, 1.5 mmol) affords **2** in 75% (0.316 g) yield. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.39 (ddd, *J* = 1.0, 1.8, 4.9 Hz, 1H), 8.34 (dd, *J* = 1.2, 7.6 Hz, 1H), 7.74 (td, *J* = 1.8, 7.7 Hz, 1H), 7.70 – 7.59 (m, 4H), 7.52 (dt, *J* = 1.1, 7.9 Hz, 1H), 7.40 – 7.35 (m, 1H), 7.29 (ddd, *J* = 1.2, 4.9, 7.6 Hz, 1H), 7.16 (d, *J* = 8.6 Hz, 2H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 159.0, 149.5, 140.9, 139.4, 139.4, 137.2, 135.5, 131.6, 130.6, 130.3, 128.7, 128.0, 125.3, 121.5, 92.6. Elemental analysis: calcd (%) for C₁₇H₁₂INO₂S (421.25): C 48.47, H 2.97; found: C 48.51, H 3.09.

2-(2-((4-Chlorophenyl)sulfonyl)phenyl)pyridine (3): 2-phenylpyridine (0.155 g, 1 mmol) and 4-chlorobenzenesulfonyl chloride (0.317 g, 1.5 mmol) affords **3** in 68% (0.224 g) yield. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.39 (d, J = 4.8 Hz, 1H), 8.34 (dd, J = 1.6, 7.7 Hz, 1H), 7.75 (td, J = 1.8, 7.7 Hz, 1H), 7.70 – 7.60 (m, 2H), 7.53 (d, J = 8.0 Hz, 1H), 7.43 – 7.36 (m, 3H), 7.29 – 7.27 (m, 3H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 156.5, 148.4, 140.8, 139.8, 139.3, 138.8, 135.6, 133.4, 132.0, 129.1, 129.0, 128.8, 128.7, 125.7, 122.6. This is a known compound and the spectral data are identical to those reported in literature.^[13]

2-(2-((4-Fluorophenyl)sulfonyl)phenyl)pyridine (4): 2-phenylpyridine (0.155 g, 1 mmol) and 4-fluorobenzenesulfonyl chloride (0.291 g, 1.5 mmol) affords 4 in 71% (0.222 g) yield. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.42 (d, J = 5.0 Hz, 1H), 8.36 (dd, J = 1.1, 7.8 Hz, 1H), 7.77 (td, J = 1.8, 7.7 Hz, 1H), 7.70 – 7.61 (m, 2H), 7.56 (dt, J = 1.1, 7.9 Hz, 1H), 7.52 – 7.45 (m, 2H), 7.39 (dd, J = 1.6, 7.3 Hz, 1H), 7.30 (dd, J = 2.1, 7.3 Hz, 1H), 7.00 (dd, J = 8.0, 9.2 Hz, 2H). ¹⁹F{¹H} NMR (376.5 MHz, CDCl₃) δ (ppm) -104.9.

 ^{13}C NMR (100 MHz, CDCl₃) δ (ppm) 165.1 (d, J = 255.5 Hz), 156.7, 148.5, 140.8, 139.2, 137.4, 135.7, 133.3, 132.1, 130.5 (d, J = 9.6 Hz), 129.1, 128.8, 125.9, 122.7. 115.9 (d, J = 22.7 Hz). Elemental analysis: calcd (%) for C1₁H1₂FNO₂S (313.34): C 65.16, H 3.86; found: C 65.28, H 3.79.

2-(2-((3,4-Difluorophenyl)sulfonyl)phenyl)pyridine (5): 2-phenylpyridine (0.155 g, 1 mmol) and 3,4-difluorobenzenesulfonyl chloride (0.319 g, 1.5 mmol) affords **5** in 64% (0.212 g) yield. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.43 (ddd, J = 1.0, 1.8, 4.9 Hz, 1H), 8.33 (dd, J = 1.6, 7.8 Hz, 1H), 7.78 (td, J = 1.8, 7.7 Hz, 1H), 7.73 – 7.60 (m, 2H), 7.54 (dt, J = 1.1, 7.8 Hz, 1H), 7.39 (dd, J = 1.6, 7.4 Hz, 1H), 7.31 (dddd, J = 1.4, 4.8, 7.9, 9.1 Hz, 3H), 7.17 – 7.08 (m, 1H). ¹⁹F{¹H} NMR (376.5 MHz, CDCl₃) δ (ppm) -129.1 (d, J = 20.8 Hz), -134.2 (d, J = 20.8 Hz). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 156.2, 153.2 (dd, J = 12.9, 253.7 Hz), 149.8 (dd, J = 12.9, 253.7 Hz), 148.5, 140.9, 138.8, 138.3 (t, J = 4.9 Hz), 135.8, 133.6, 132.2, 129.4, 128.9, 125.8, 125.0 (dd, J = 4.0, 7.7 Hz), 122.9, 117.9 (dd, J = 1.9, 20.1 Hz), 117.8 (d, J = 18.6 Hz). Elemental analysis: calcd (%) for C₁₇H₁₁F₂NO₂S (331.33): C 61.63, H 3.35; found: C 61.79, H 3.59.

2-(2-((3-Bromophenyl)sulfonyl)phenyl)pyridine (6): 2-phenylpyridine (0.155 g, 1 mmol) and 3-bromobenzenesulfonyl chloride (0.383 g, 1.5 mmol) affords **6** in 71% (0.266 g) yield. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.43 (d, J = 4.9 Hz, 1H), 8.37 (dd, J = 2.0, 7.4 Hz, 1H), 7.80 (dt, J = 1.6, 7.7 Hz, 1H), 7.68 (dt, J = 1.7, 6.5 Hz, 2H), 7.63 – 7.58 (m, 1H), 7.58 – 7.52 (m, 2H), 7.50 – 7.44 (m, 1H), 7.43 – 7.39 (m, 1H), 7.38 – 7.32 (m, 1H), 7.26 – 7.21 (m, 1H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 156.3, 148.6, 143.2, 140.9, 138.8, 135.8, 135.7, 133.6, 132.1, 130.8, 130.3, 129.3, 128.9, 126.4, 125.7, 123.0, 122.4. Elemental analysis: calcd (%) for C₁₇H₁₂BrNO₂S (374.25): C 54.56, H 3.23; found: C 54.59, H 3.41.

2-(2-((3-Fluorophenyl)sulfonyl)phenyl)pyridine (8): 2-phenylpyridine (0.155 g, 1 mmol) and 3-fluorobenzenesulfonyl chloride (0.291 g, 1.5 mmol) affords **8** in 72% (0.226 g) yield. ¹H NMR (400 MHz, CDCl₃) 8.42 (d, J = 4.9 Hz, 1H), 8.39 – 8.35 (m, 1H), 7.82 – 7.75 (m, 1H), 7.74 – 7.62 (m, 2H), 7.55 (d, J = 7.8 Hz, 1H), 7.41 (d, J = 7.1 Hz, 1H), 7.39 – 7.28 (m, 3H), 7.22 – 7.12 (m, 2H). ¹⁹F{¹H} NMR (376.5 MHz, CDCl₃) $\overline{0}$ (ppm) -110.3. ¹³C NMR (100 MHz, CDCl₃) $\overline{0}$ (ppm) 162.0 (d, J = 251.3 Hz), 156.6, 148.5, 143.6 (d, J = 6.7 Hz), 141.0, 138.9, 135.7, 133.5, 132.1, 130.5 (d, J = 7.7 Hz), 129.4, 128.8, 125.7, 123.5 (d, J = 3.3 Hz), 122.8, 120.0 (d, J = 21.4 Hz), 115.2 (d, J = 24.8 Hz). Elemental analysis: calcd (%) for C₁₇H₁₂FNO₂S (313.36): C 65.16, H 3.86; found: C 65.18, H 3.91.

2-(2-((4-lodophenyl)sulfonyl)naphthalen-1-yl)pyridine (10): 2-(naphthalen-1-yl)pyridine (0.205 g, 1 mmol) and 4-iodobenzenesulfonyl chloride (0.454 g, 1.5 mmol) affords **10** in 72% (0.339 g) yield. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.54 (dt, J = 1.2, 5.0 Hz, 1H), 8.33 (d, J = 8.8 Hz, 1H), 8.10 (d, J = 8.9 Hz, 1H), 7.95 (d, J = 8.2 Hz, 1H), 7.88 (td, J = 1.9, 7.7 Hz, 1H), 7.75 – 7.69 (m, 2H), 7.62 (ddd, J = 1.2, 6.8, 8.2 Hz, 1H), 7.59 – 7.55 (m, 1H), 7.46 – 7.39 (m, 2H), 7.26 (d, J = 7.9 Hz, 2H), 7.18 (d, J = 8.6 Hz, 1H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 154.9, 148.8, 141.4, 139.5, 138.0, 136.1, 135.7, 135.5, 132.8, 129.5, 129.0, 128.9, 128.1, 127.7, 127.6, 127.3, 123.6, 122.9, 100.6. Elemental analysis: calcd (%) for C₂₁H₁₄INO₂S (471.31): C 53.52, H 2.99; found: C 53.51, H 2.76.

2-(2-((4-Bromophenyl)sulfonyl)naphthalen-1-yl)pyridine (11): 2-(naphthalen-1-yl)pyridine (0.205 g, 1 mmol) and 4-bromobenzenesulfonyl chloride (0.383 g, 1.5 mmol) affords **11** in 68% (0.288 g) yield. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.56 (d, J = 5.0 Hz, 1H), 8.31 (d, J = 8.9 Hz, 1H), 8.10 (d, J = 8.8 Hz, 1H), 7.96 (d, J = 8.2 Hz, 1H), 7.91 (t, J = 7.8 Hz, 1H), 7.64 (d, J = 7.4 Hz, 1H), 7.62 – 7.57 (m, 1H), 7.51 (d, J = 8.6 Hz, 2H), 7.45 (t, J = 7.3 Hz, 4H), 7.18 (d, J = 8.6 Hz, 1H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 154.9, 148.8, 140.7, 139.4, 136.1, 135.7, 135.5, 132.8, 132.0, 129.5, 129.2, 128.9, 128.1, 128.0, 127.7, 127.6, 127.3, 123.6, 123.0. Elemental analysis: calcd (%) for C₂₁H₁₄BrNO₂S (422.31): C 59.44, H 3.33 found: C 59.67, H 3.49.

2-(2-((4-Bromophenyl)sulfonyl)-4-(trifluoromethyl)phenyl)pyridine

(12): 2-(4-(trifluoromethyl)phenyl)pyridine (0.223 g, 1 mmol) and 4-bromobenzenesulfonyl chloride (0.283 g, 1.5 mmol) affords 12 in 70% (0.309 g) yield. ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.63 (s, 1H), 8.40 (d, J = 4.8 Hz, 1H), 7.92 (d, J = 8.0 Hz, 1H), 7.78 (dt, J = 4.8, 7.8 Hz, 1H), 7.53 (dd, J = 7.9, 10.8 Hz, 2H), 7.47 (d, J = 8.3 Hz, 2H), 7.38 – 7.23 (m, 3H). ¹⁹F{¹H} NMR (376.5 MHz, CDCl₃) δ (ppm) -62.7. ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 155.3, 148.7, 144.2, 140.3, 139.5, 136.0, 133.0, 132.1, 131.4 (q, J = 33.6 Hz), 130.0 (m), 129.3, 128.6, 126.4 (q, J = 4.1 Hz), 125.8, 123.3 123.0 (q, J = 272.1 Hz). Elemental analysis: calcd (%) for C₁₈H₁₁F₃BrNO₂S (442.25): C 48.89, H 2.51; found: C 48.96, H 2.69.

2-(2-((4-lodophenyl)sulfonyl)-4-methylphenyl)pyridine (13): 2-(*p*-tolyl)pyridine (0.169 g, 1 mmol) and 4-iodobenzenesulfonyl chloride (0.454 g, 1.5 mmol) affords **13** in 77% (0.335 g) yield. ¹H NMR (400 MHz, CDCl₃) 8.42 – 8.38 (m, 1H), 8.16 (s, 1H), 7.75 (td, J = 2.9, 5.7 Hz, 1H), 7.69 (d, J = 8.6 Hz, 1H), 7.55 – 7.46 (m, 2H), 7.42 (d, J = 9.2 Hz, 1H), 7.33 – 7.25 (m, 3H), 7.19 (d, J = 7.9 Hz, 1H), 2.54 (s, 3H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 148.4, 141.3, 139.2, 138.5, 137.8, 135.6, 134.1, 132.0, 129.5, 129.1, 129.0, 128.9, 126.0, 122.6, 100.5, 21.2. Elemental analysis: calcd (%) for C₁₈H₁₄INO₂S (435.28): C 49.67, H 3.24; found: C 49.41, H 3.55.

2-(2-((4-Bromophenyl)sulfonyl)-4-methylphenyl)pyridine (14): 2-(*p*-tolyl)pyridine (0.169 g, 1 mmol) and 4-bromobenzenesulfonyl chloride (0.383 g, 1.5 mmol) affords **14** in 78% (0.303 g) yield. ¹H NMR (400 MHz, CDCl₃) δ (ppm) ¹H NMR (400 MHz, CDCl₃) δ 8.41 (d, *J* = 4.3 Hz, 1H), 8.16 (s, 1H), 7.76 (t, *J* = 7.7 Hz, 1H), 7.56 – 7.53 (m, 1H), 7.51 – 7.49 (m, 1H), 7.50 – 7.46 (m, 2H), 7.36 (d, *J* = 8.7 Hz, 2H), 7.32 – 7.29 (m, 2H), 2.54 (s, 3H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 156.6, 148.4, 140.6, 139.2, 138.5, 138.0, 135.7, 134.1, 132.0, 131.8, 129.5, 129.1, 127.9, 126.0, 122.6, 21.3. Elemental analysis: calcd (%) for C₁₈H₁₄BrNO₂S (388.28): C 55.68, H 3.63; found: C 55.81, H 3.70.

2-(2-((4-Iodophenyl)sulfonyl)phenyl)-5-(trifluoromethyl)pyridine (15): 2-phenyl-5-(trifluoromethyl)pyridine (0.223 g, 1 mmol) and 4-iodobenzenesulfonyl chloride (0.454 g, 1.5 mmol) affords **15** in 75% (0.367 g) yield. ¹H NMR (400 MHz, CDCI₃) 8.68 (s, 1H), 8.34 (dd, *J* = 1.6, 7.7 Hz, 1H), 8.01 (dd, *J* = 2.0, 8.2 Hz, 1H), 7.74 (d, *J* = 8.5 Hz, 2H), 7.71 – 7.66 (m, 3H), 7.39 (dd, *J* = 2.3, 7.4 Hz, 1H), 7.26 – 7.21 (m, 2H). ¹⁹F{¹H} NMR (376.5 MHz, CDCI₃) δ (ppm) -62.3. ¹³C NMR (100 MHz, CDCI₃) δ (ppm) 160.3, 145.3 (q, *J* = 4.0 Hz), 141.0, 139.5, 139.1, 138.1, 133.6, 132.9 (q, *J* = 3.5 Hz), 131.8, 129.5, 128.9, 125.7 (q, *J* = 33.5 Hz), 125.5, 123.4 (q, *J* = 272.1 Hz), 100.9. Elemental analysis: calcd (%) for C₁₈H₁₁F₃INO₂S (489.25): C 44.19, H 2.27; found: C 44.28, H 2.56.

2-(2-((4-Bromophenyl)sulfonyl)phenyl)-5-(trifluoromethyl)pyridine

(16): 2-phenyl-5-(trifluoromethyl)pyridine (0.223 g, 1 mmol) and 4bromobenzenesulfonyl chloride (0.383 g, 1.5 mmol) affords 16 in 74% (0.327 g) yield. ¹H NMR (400 MHz, CDCl₃) 8.67 (s, 1H), 8.32 (dd, J = 1.3, 7.7 Hz, 1H), 8.00 (dd, J = 2.2, 9.0 Hz, 1H), 7.74 – 7.64 (m, 3H), 7.50 (d, J = 8.6 Hz, 2H), 7.40-7.35 (m, 3H). ¹⁹F{¹H} NMR (376.5 MHz, CDCl₃) δ (ppm) -62.3. ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 160.3, 145.3 (q, J = 4.0 Hz), 140.3, 139.5, 139.1, 133.6, 132.9 (q, J = 3.5 Hz), 132.1, 131.8, 129.6, 129.1, 128.4, 127.0, 125.8 (q, J = 33.1 Hz), 125.5, 123.1 (q, J = 272.5 Hz). Elemental analysis: calcd (%) for C₁₈H₁₁BrF₃NO₂S (442.25): C 48.89, H 2.51; found: C 49.05, H 2.80.

2-(2-((4-Bromophenyl)sulfonyl)phenyl)nicotinonitrile (18): 2-phenylnicotinonitrile (0.180 g, 1 mmol) and 4-bromobenzenesulfonyl chloride (0.383 g, 1.5 mmol) affords **18** in 61% (0.244 g) yield. ¹H NMR (400 MHz, CDCl₃) 8.75 (d, *J* = 3.8 Hz, 1H), 8.26 (dd, *J* = 2.0, 7.2 Hz, 1H), 8.09 (dd, *J* = 1.7, 7.8 Hz, 1H), 7.79 – 7.71 (m, 2H), 7.63 – 7.57 (m, 4H), 7.51 (dd, *J* = 4.8, 8.1 Hz, 1H), 7.45 (dd, *J* = 2.1, 6.7 Hz, 1H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 160.3, 151.2, 140.7, 140.2, 139.2, 137.2, 133.7, 132.3, 131.9, 131.3, 130.5, 130.4, 129.3, 129.3, 128.5, 122.8. Elemental analysis: calcd (%) for C₁₈H₁₁BrN₂O₂S (399.26): C 54.15, H 2.78; found: C 53.87, H 3.04.

2-(4-Bromo-2-((4-nitrophenyl)sulfonyl)phenyl)pyridine (19): 2-(4-bromophenyl)pyridine (0.234 g, 1 mmol) and 4-nitrobenzenesulfonyl chloride (0.332 g, 1.5 mmol) affords **19** in 78% (0.327 g) yield. ¹H NMR (400 MHz, CDCl₃) 8.56 (d, J = 2.0 Hz, 1H), 8.34 (d, J = 4.5 Hz, 1H), 8.22 (d, J = 8.9 Hz, 2H), 7.87 (dd, J = 2.1, 8.1 Hz, 1H), 7.81 (td, J = 1.8, 7.8 Hz, 1H), 7.73 (d, J = 8.9 Hz, 2H), 7.55 (d, J = 7.8 Hz, 1H), 7.37 – 7.34 (m, 1H), 7.32 (d, J = 8.2 Hz, 1H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 155.4, 150.1, 148.4, 146.9, 139.8, 139.6, 137.0, 136.3, 133.7, 132.5, 129.0, 125.6, 123.8, 123.3, 123.1. Elemental analysis: calcd (%) for C₁₇H₁₁BrN₂O4S (419.24): C 48.70, H 2.64; found: C 48.79, H 2.91.

2-(4-Bromo-2-((4-(trifluoromethyl)phenyl)sulfonyl)phenyl)pyridine

(20): 2-(4-bromophenyl)pyridine (0.234 g, 1 mmol) and 4-(trifluoromethyl)benzenesulfonyl chloride (0.367 g, 1.5 mmol) affords 20 in 82% (0.362 g) yield. ¹H NMR (400 MHz, *d*₆-DMSO) 8.50 (d, J = 2.0 Hz, 1H), 8.33 (ddd, J = 0.9, 1.8, 4.8 Hz, 1H), 8.07 (dd, J = 2.0, 8.2 Hz, 1H), 7.95 – 7.84 (m, 5H), 7.51 – 7.46 (m, 2H), 7.41 (ddd, J = 1.1, 4.8, 7.6 Hz, 1H). ¹⁹F{¹H} NMR (376.5 MHz, *d*₆-DMSO) δ (ppm) -61.6. ¹³C NMR (100 MHz, *d*₆-DMSO) δ (ppm) 171.9, 155.7, 148.6, 140.4, 140.2, 137.6, 137.0, 134.6, 133.3 (q, J = 33.1 Hz), 132.7, 129.0, 126.6 (q, J = 5.1 Hz), 124.8, 123.8 (q, J = 273.5 Hz), 123.7, 122.6. Elemental analysis: calcd (%) for C₁₈H₁₁BrF₃NO₂S (442.25): C 48.89, H 2.51; found: C 49.04, H 2.11.

3-Chloro-2-(2-((4-nitrophenyl)sulfonyl)phenyl)pyridine (21): 3-chloro-2-phenylpyridine (0.189 g, 1 mmol) and 4-nitrobenzenesulfonyl chloride (0.332 g, 1.5 mmol) affords **21** in 80% (0.299 g) yield. ¹H NMR (400 MHz, CDCl₃) 8.41 (dd, J = 1.4, 4.7 Hz, 1H), 8.32 (dd, J = 1.5, 8.4 Hz, 1H), 8.28 (d, J = 8.8 Hz, 2H), 7.88 (d, J = 8.8 Hz, 2H), 7.85 (dd, J = 1.4, 8.1 Hz, 1H), 7.77 (td, J = 1.5, 7.5 Hz, 1H), 7.71 (td, J = 1.5, 7.6 Hz, 1H), 7.42 – 7.39 (m, 1H), 7.39 – 7.35 (m, 1H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 155.1, 150.2, 147.5, 146.2, 138.5, 137.8, 137.3, 134.1, 132.1, 131.7, 130.4, 129.8, 129.1, 124.5, 124.0. Elemental analysis: calcd (%) for C₁₇H₁₁ClN₂O₄S (374.80): C 54.48, H 2.96; found: C 54.30, H 2.87.

3-Chloro-2-(2-((4-(trifluoromethyl)phenyl)sulfonyl)phenyl)pyridine (22): 3-chloro-2-phenylpyridine (0.189 g, 1 mmol) and 4-(trifluoromethyl)benzenesulfonyl chloride (0.367 g, 1.5 mmol) affords 22 in

79% (0.314 g) yield. ¹H NMR (400 MHz, CDCl₃) 8.41 (d, J = 4.6 Hz, 1H), 8.31 (dd, J = 2.0, 7.8 Hz, 1H), 7.86 – 7.79 (m, 3H), 7.77 – 7.67 (m, 4H), 7.41 – 7.32 (m, 2H). ¹⁹F{¹H} NMR (376.5 MHz, CDCl₃) δ (ppm) -63.2. ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 155.2, 146.3, 145.3, 138.4, 138.3, 137.1, 134.6 (q, J = 34.1 Hz), 133.8, 132.1, 131.5, 130.2, 129.7, 128.4, 126.0 (q, J = 3.5 Hz), 124.3, 123.2 (q, J = 272.5 Hz). Elemental analysis: calcd (%) for C₁₈H₁₁ClF₃NO₂S (397.80): C 54.35, H 2.79; found: C 54.58, H 2.51.

3-Chloro-2-(2-((4-iodophenyl)sulfonyl)phenyl)pyridine (23): 3-chloro-2-phenylpyridine (0.189 g, 1 mmol) and 4-iodobenzenesulfonyl chloride (0.454 g, 1.5 mmol) affords **23** in 58% (0.264 g) yield. ¹H NMR (400 MHz, CDCl₃) 8.43 (dd, *J* = 1.5, 4.7 Hz, 1H), 8.26 (dd, *J* = 1.6, 7.7 Hz, 1H), 7.83 – 7.78 (m, 2H), 7.74 – 7.63 (m, 3H), 7.40 – 7.32 (m, 4H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 155.3, 146.3, 141.4, 138.8, 138.1, 138.1, 137.1, 133.5, 132.1, 131.4, 130.0, 129.6, 129.2, 124.2, 100.9. Elemental analysis: calcd (%) for C₁₇H₁₁ClINO₂S (455.69): C 44.81, H 2.43; found: C 45.03, H 2.89.

2-(2-((4-Bromophenyl)sulfonyl)phenyl)-3-chloropyridine (24): 3-chloro-2-phenylpyridine (0.189 g, 1 mmol) and 4-bromobenzenesulfonyl chloride (0.383 g, 1.5 mmol) affords **24** in 61% (0.249 g) yield. ¹H NMR (400 MHz, CDCl₃) 8.44 (dd, J = 1.5, 4.7 Hz, 1H), 8.27 (dd, J = 1.3, 7.4 Hz, 1H), 7.81 (dd, J = 1.5, 8.2 Hz, 1H), 7.71 (td, J = 1.6, 7.5 Hz, 1H), 7.66 (td, J = 1.7, 7.6 Hz, 1H), 7.60 – 7.49 (m, 4H), 7.39 – 7.31 (m, 2H). ¹³C NMR (100 MHz, CDCl₃) δ (ppm) 155.4, 146.3, 140.8, 138.8, 138.2, 137.1, 133.5, 132.1, 132.1, 131.4, 130.0, 129.6, 129.4, 128.3, 124.2. Elemental analysis: calcd (%) for C₁₇H₁₁BrCINO₂S (408.69): C 49.96, H 2.71; found: C 50.21, H 2.52.

Preparation of the PdCI(C₃H₅)(dppb) catalyst:^[18] An oven-dried 40 mL Schlenk tube equipped with a magnetic stirring bar under argon atmosphere, was charged with [Pd(C₃H₅)Cl]₂ (182 mg, 0.5 mmol) and dppb (426 mg, 1 mmol). 10 mL of anhydrous dichloromethane were added, then, the solution was stirred at room temperature for twenty minutes. The solvent was removed in vacuum. The powder was used without purification. (³¹P NMR 381 MHz, CDCl₃) δ = 19.3 (s).

2-Isopropyl-4-methyl-5-(4-((2-(pyridin-2-

yl)phenyl)sulfonyl)phenyl)thiazole (25): To a 25 mL oven dried Schlenk tube, 2-(2-((4-bromophenyl)sulfonyl)phenyl)pyridine (1) (0.186 g, 0.5 mmol) and 2-isopropyl-4-methylthiazole (0. 106 g, 0.75 mmol) KOAc (0.196 g, 2 mmol), DMA (4 mL) and PdCl(C₃H₅)(dppb) (12.2 mg, 0.02 mmol) were successively added. The reaction mixture was evacuated by vacuum-argon cycles (5 times) and stirred at 150 °C (oil bath temperature) for 16 h. After cooling the reaction at room temperature and concentration. the crude mixture was purified by silica column chromatography to afford **25** in 87% (0.189 g). ¹H NMR (400 MHz, CDCl₃) δ (ppm) 8.38 (ddd, J = 2.0, 3.2, 4.2 Hz, 2H), 7.75 (td, J = 1.8, 7.7 Hz, 1H), 7.70 - 7.61 (m, 2H), 7.57 (d, J = 7.8 Hz, 1H), 7.48 (d, J = 8.6 Hz, 2H), 7.41 - 7.38 (m, 1H), 7.36 - 7.32 (m, 2H), 7.30 - 7.26 (m, 1H), 3.41 - 3.19 (m, 1H), 2.45 (s, 3H), 1.41 (d, J = 6.9 Hz, 6H). ¹³C NMR (100 MHz, CDCl₃) \overline{o} (ppm) 176.7, 156.7, 148.5, 148.4, 140.9, 140.0, 139.3, 137.4, 135.7, 133.4, 132.1, 129.2, 129.0, 128.9, 128.8, 128.1, 125.9, 122.7, 33.5, 23.2, 16.4. Elemental analysis: calcd (%) for C24H22N2O2S2 (434.57): C 66.33, H 5.10; found: C 66.38, H 5.21.

Acknowledgements

A.S. acknowledges the Indo-French Centre for the Promotion of Advanced Research (IFCPAR, N $^{\circ}$ 5705-1) for a PhD grant and research support.

Keywords: C–H bond • Sulfonylation • Palladium • Halo Bonds • Sulfones

- R. L. Lopez de Compadre, R. A. Pearlstein, A. J. Hopfinger, J. K. Seydel, J. Med. Chem. 1987, 30, 900-906.
- [2] T. M. Williams, T. M. Ciccarone, S. C. MacTough, C. S. Rooney, S. K. Balani, J. H. Condra, E. A. Emini, M. E. Goldman, W. J. Greenlee, J. Med. Chem. 1993, 36, 1291-1294.
- [3] R. S. Varma, R. K. Saini, H. M. Meshram, *Tetrahedron Lett.* 1997, 38, 6525-6528.
- [4] a) B. M. Graybill, *J. Org. Chem.* **1967**, *32*, 2931-2933; b) M. Ueda, K. Uchiyama, T. Kano, Synthesis **1984**, 1984, 323-325; c) S. J. Nara, J. R. Harjani, M. M. Salunkhe, *J. Org. Chem.* **2001**, *66*, 8616-8620.
 [5] a) H. Gilman, N. J. Beaber, C. H. Myers, *J. Am. Chem. Soc.* **1925**, *47*,
- a) H. Gilman, N. J. Beaber, C. H. Myers, J. Am. Chem. Soc. 1925, 47, 2047-2052; b) J. M. Baskin, Z. Wang, Org. Lett. 2002, 4, 4423-4425; c) B. P. Bandgar, S. V. Bettigeri, J. Phopase, Org. Lett. 2004, 6, 2105-2108; d) W. Zhu, D. Ma, J. Org. Chem. 2005, 70, 2696-2700; e) A. Kar, I. A. Sayyed, W. F. Lo, H. M. Kaiser, M. Beller, M. K. Tse, Org. Lett. 2007, 9, 3405-3408; f) H. Yang, Y. Li, J. Min, W. Junmei, H. Fu, Chem. Eur. J. 2011, 17, 5652-5660; g) E. J. Emmett, B. R. Hayter, M. C. Willis Angew. Chem. Int. Ed. 2013, 52, 12679-12683.
- a) F. Kakiuchi, T. Kochi, Synthesis 2008, 3013-3039; b) L. Ackermann, [6] R. Vicente, A. R. Kapdi, Angew. Chem. Int. Ed. 2009, 48, 9792-9826; c) F. Bellina, R. Rossi, Tetrahedron 2009, 65, 10269-10310; d) X. Chen, K. M. Engle, D.-H. Wang, J.-Q. Yu, Angew. Chem. Int. Ed. 2009, 48, 5094-5115; e) T. W. Lyons, M. S. Sanford, Chem. Rev. 2010, 110, 1147-1169; f) E. M. Beck, M. J. Gaunt, *Top. Curr. Chem.* **2010**, *292*, 85-121; g) T. Satoh, M. Miura, Synthesis **2010**, 3395-3409; h) C.-L. Sun, B.-J. Li, Z.-J. Shi, Chem. Commun. 2010, 46, 677-685; i) S. H. Cho, J. Y. Kim, J. Kwak, S. Chang, Chem. Soc. Rev. 2011, 40, 5068-5083; j) B. Liu, J. Li, . Song, J. You, Chem. Eur. J. 2012, 18, 10830-10833; k) M. C. White, Synlett 2012, 23, 2746-2748; I) J. Yamaguchi, A. D. Yamaguchi, K. Itami, Angew. Chem. Int. Ed. 2012, 51, 8960-9009; m) S. I. Kozhushkov, L. Ackermann, *Chem. Sci.* **2013**, *4*, 886-896; n) R. Rossi, F. Bellina, M. Lessi, C. Manzini, *Adv. Synth. Catal.* **2014**, 356, 17-117; o) M. Zhang, Y. Zhang, X. Jie, H. Zhao, G. Li, W. Su, Org. Chem. Front. 2014, 1, 843-895; p) M. R. Yadav, R. K. Rit, M. Shankar, A. K. Sahoo, Asian J. Org. Chem. 2015, 4, 846-864; q) K. Hirano, M. Miura, Chem. Lett. 2015, 44, 878-873; r) K. Yuan, J.-F. Soulé, H. Doucet, ACS Catal. 2015, 5, 978-991.
- [7] For examples of *ortho* C–H bond sulfonylations of aromatic amides utilizing an *N*,*N*-bidentate directing group and nickel(II) catalysis, see: A. Yokota, N. Chatani, *Chem. Lett.* 2015, *44*, 902-904.
 [8] a) For examples of selective remote C–H bond sulfonylations with
- [8] a) For examples of selective remote C–H bond sulfonylations with benzenesulfonyl chlorides, see: O. Saidi, J. Marafie, A. E. W. Ledger, P. M. Liu, M. F. Mahon, G. Kociok-Köhn, M. K. Whittlesey, C. G. Frost, J. Am. Chem. Soc. 2011, 133, 19298-19301; b) G. Li, X. Lv, K. Guo, Y. Wang, S. Yang, L. Yu, Y. Yu, J. Wang, Org. Chem. Front. 2017, 4, 1145-1148; c) H.-W. Liang, K. Jiang, W. Ding, Y. Yuan, L. Shuai, Y.-C. Chen, Y. Wei, Chem. Commun. 2015, 51, 16928-16931; d) J.-M. Li, J. Weng, G. Lu, A. S. C. Chan, Tetrahedron Lett. 2016, 57, 2121-2124.
- [9] a) X. Zhao, E. Dimitrijević, V. M. Dong, J. Am. Chem. Soc. 2009, 131, 3466-3467; b) X. Zhao, V. M. Dong, Angew. Chem. Int. Ed. 2011, 50, 932-934.
- [10] H. Li, A. Sasmal, X. Shi, J.-F. Soulé, H. Doucet, Org. Biomol. Chem. 2018, (DOI: 10.1039/C1038OB00632F).
- [11] a) For examples of ortho C–H bond sulfonylations with sodium sulfinates utilizing an *N*,*N*-bidentate directing group and copper(II) catalysis, see: W.-H. Rao, B.-F. Shi, Org. Lett. 2015, 17, 2784-2787; b) S. Liang, N. W. Liu, G. Manolikakes, *Adv. Synth. Catal.* 2016, 358, 159-163; c) For examples of ortho C–H bond sulfonylations with sodium sulfinates utilizing an *N*,*N*-bidentate directing group and nickel(II) catalysis, see: S. L. Liu, X. H. Li, S. S. Zhang, S. K. Hou, G. C. Yang, J. F. Gong, M. P. Song, *Adv. Synth. Catal.* 2017, 359, 2241-2246; d) For examples of the synthesis of halo-substituted diarylsulfones *via* insertion of sulfur dioxide, see: H. Xia, Y. An, X. Zeng, J. Wu, *Chem. Commun.* 2017, 53, 12548-12551.
- [12] L. Niu, H. Yang, D. Yang, H. Fu, Adv. Synth. Catal. 2012, 354, 2211-2217.
- [13] F. Wang, X. Yu, Z. Qi, X. Li, Chem. Eur. J. 2016, 22, 511-516.
- [14] A. Skhiri, A. Beladhria, K. Yuan, J.-F. Soulé, R. Ben Salem, H. Doucet, Eur. J. Org. Chem. 2015, 4428-4436.
- [15] a) I. Idris, F. Derridj, S. Djebbar, J.-F. Soulé, H. Doucet, *Tetrahedron* 2015, 71, 9617-9625; b) K. Yuan, J.-F. Soulé, V. Dorcet, H. Doucet, ACS *Catal.* 2016, 8121-8126.
- [16] a) S. R. Dubbaka, P. Vogel, Org. Lett. 2003, 6, 95-98; b) S. R. Dubbaka, P. Vogel, Chem. Eur. J. 2005, 11, 2633-2641.
- [17] X. Shi, J.-F. Soulé, H. Doucet, J. Org. Chem. 2017, 82, 3886-3894.

[18] T. Cantat, E. Génin, C. Giroud, G. Meyer, A. Jutand, J. Organomet. Chem. 2003, 687, 365-376.

Entry for the Table of Contents (Please choose one layout)

Layout 1:

FULL PAPER

Layout 2:

FULL PAPER



A one step procedure for the synthesis of (poly)halo-substituted diarylsulfones was developed. The reaction involves *ortho* C–H bond sulfonylation of 2-arylpyridines with (poly)halo-substituted benzenesulfonyl chlorides. These reaction conditions tolerated C–F, C–CI, C–Br and even C–I bonds.

C-H bond Functionalization*

A. Sasmal, J. K. Bera, H. Doucet* and J.-F. Soulé*

Page No. – Page No.

Synthesis of (Poly)halo-substituted Diarylsulfones through Palladium Catalyzed C–H Bond Sulfonylation using (Poly)Halobenzenesulfonyl Chlorides