



HAL
open science

Comment on Couzi et al. (2018): a phenomenological model for structural transitions in incommensurate alkane/urea inclusion compounds

B. Toudic, Laurent Guérin, Céline Mariette, I. Frantsuzov, P. Rabiller, Claude Ecolivet, Mark D. Hollingsworth

► To cite this version:

B. Toudic, Laurent Guérin, Céline Mariette, I. Frantsuzov, P. Rabiller, et al.. Comment on Couzi et al. (2018): a phenomenological model for structural transitions in incommensurate alkane/urea inclusion compounds. Royal Society Open Science, 2019, 6 (8), pp.182073. <10.1098/rsos.182073>. <hal-02303232>

HAL Id: hal-02303232

<https://univ-rennes.hal.science/hal-02303232v1>

Submitted on 17 Jul 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

Comment



Cite this article: Toudic B, Guérin L, Mariette C, Frantsuzov I, Rabiller P, Ecolivet C, Hollingsworth MD. 2019 Comment on Couzi *et al.* (2018): a phenomenological model for structural transitions in incommensurate alkane/urea inclusion compounds. *R. Soc. open sci.* **6**: 182073. <http://dx.doi.org/10.1098/rsos.182073>

Received: 4 December 2018

Accepted: 1 March 2019

Subject Category:

Chemistry

Subject Areas:

chemical physics/solid-state physics

Authors for correspondence:

B. Toudic

e-mail: bertrand.toudic@univ-rennes1.fr

C. Ecolivet

e-mail: claud.ecolivet@univ-rennes1.fr

Mark D. Hollingsworth

e-mail: mdhall@ksu.edu

This article has been edited by the Royal Society of Chemistry, including the commissioning, peer review process and editorial aspects up to the point of acceptance.



Comment on Couzi *et al.* (2018): a phenomenological model for structural transitions in incommensurate alkane/urea inclusion compounds

B. Toudic¹, L. Guérin¹, C. Mariette¹, I. Frantsuzov²,
P. Rabiller¹, C. Ecolivet¹ and Mark D. Hollingsworth²

¹Univ Rennes, CNRS, IPR (Institut de Physique de Rennes) – UMR 6251, F-35000 Rennes, France

²Department of Chemistry, Kansas State University, Manhattan, KS 66506, USA

BT, 0000-0001-9918-8071; LG, 0000-0002-0509-8444; CM, 0000-0001-8067-9591; IF, 0000-0002-3589-6818; PR, 0000-0003-1566-606X; CE, 0000-0002-1929-7655; MDH, 0000-0002-1995-7182

In their recent article [1], M. Couzi *et al.* develop a standard phenomenological model of coupled order parameters, which generates one single symmetry-breaking phase transition. They apply it to the phase transitions of *n*-nonadecane/urea and *n*-hexadecane/urea, while knowing that our measurements had demonstrated, via previously published diffraction experiments [2–6], that both materials undergo at least two symmetry-breaking events revealed by systematic absences of diffraction peaks.

We have shown that in *n*-nonadecane-d₄₀/urea-d₄ and *n*-nonadecane-h₄₀/urea-h₄, there is a complex sequence of phases that follow crystallographic symmetry conditions. The first-phase transition at T_{c1} is associated with the symmetry breaking from the hexagonal high-temperature phase to a second phase (phase II); a second transition to a different space group (phase III) occurs at a lower temperature T_{c2} . These results have been discussed extensively in the literature [3,4,6–9] and were obtained using excellent spatial resolution and temperature calibration, including measurements using cold neutron scattering on triple axis spectrometers [3,4], and on a synchrotron X-ray diffractometer [7]. For *n*-nonadecane-h₄₀/urea-h₄, $T_{c1} = (158.8 \pm 0.1)$ K and $T_{c2} = (147.0 \pm 0.1)$ K, according to adiabatic measurements [10].

Instead of using these careful measurements, M. Couzi *et al.* used the results from their recently published article, which describes an X-ray diffraction study performed on *n*-nonadecane-h₄₀/urea-h₄ but with measurements reported at only two temperatures (147 K and 100 K) [11]. In a Comment concerning that article [2], which they did not contest, we demonstrated that their data have no relevance. There [2], we wrote: ‘As elaborated here, the data reported by Couzi *et al.* are not from phase II of *n*-nonadecane/urea and cannot be used to discuss the sequence of phases in this compound’.

Inexplicably, Couzi *et al.* do not address this fundamental criticism of their data in the present article [1]. They have not presented any data on pure phase II on which to base their assertion that phase II is the same space group as phase III. Furthermore, M. Couzi *et al.* are fully aware of the extraordinarily high-quality X-ray data exhibiting the unique crystallographic signature of phase II shown in fig. 2(a) of [7], (ref. 32 in [1] and ref. 11 in [11]), which was acquired at 154.5 K, in the middle of phase II of *n*-nonadecane-h₄₀/urea-h₄. This figure, in fact, is the reconstructed image that they *would* have obtained if they had actually been measuring phase II instead of phase III. However, they have chosen to ignore these data in favour of their own, which were both measurements on phase III.

In the same Comment [2], to which there was no reply, we asked Couzi *et al.* to explain how their measurements were actually made, since *only one* measurement was reported in the vicinity of phase II:

It is for Couzi *et al.* to explain why they have failed to observe the absence/presence conditions that characterize phase II, in particular the common and host superstructure Bragg peaks, which are not in phase II and whose emergence with cooling signifies phase III.

However, in their present article [1], Couzi *et al.* make the following surprising statement (where references [12,28,30] and [31] are references [5,3,4] and [11], respectively, in this Comment):

Given that the (3 + 2)-dimensional superspace groups proposed previously [12,28,30] for *n*-nonadecane/urea and *n*-hexadecane/urea have been shown [31] to be incorrect,...

We maintain that the data collected by Couzi *et al.* [11] were on the wrong phase, and as a consequence, their phenomenological description of the phase behaviour in *n*-nonadecane/urea [1] is contrary to reliable experimental measurements and does not apply to the isotopologues of *n*-nonadecane/urea. As already extensively argued in our previous Comment [2], we maintain also that two different symmetry-breaking events are indeed present in *n*-hexadecane/urea.

Data accessibility. This article contains no new data.

Authors' contributions. Each of the authors contributed substantially to the drafting and editing of the manuscript, and all authors approved of the final content.

Competing interests. We declare we have no competing interests.

Funding. No funding has been received for this article.

References

- Couzi M, Guillaume F, Harris KDM. 2018 A phenomenological model for structural phase transitions in incommensurate alkane/urea inclusion compounds. *R. Soc. open sci.* **5**, 180058. (doi:10.1098/rsos.180058)
- Toudic B, Guérin L, Mariette C, Frantsuzov I, Rabiller P, Ecolivet C, Janssen T, Hollingsworth MD. 2017 Comment on ‘The true structural periodicities and superspace group descriptions of the prototypical incommensurate composite materials: alkane/urea inclusion compounds’. *Europhys. Lett.* **119**, 66004. (doi:10.1209/0295-5075/119/66004)
- Toudic B *et al.* 2008 Hidden degrees of freedom in aperiodic materials. *Science* **319**, 69–71. (doi:10.1126/science.1146745)
- Toudic B, Rabiller P, Bourgeois L, Huard M, Ecolivet C, McIntyre GJ, Bourges P, Brezczewski T, Janssen T. 2011 Temperature–pressure phase diagram of an aperiodic host-guest compound. *Europhys. Lett.* **93**, 16003. (doi:10.1209/0295-5075/93/16003)
- Huard M, Toudic B, Rabiller P, Ecolivet C, Guérin L, Bourges P, Brezczewski T, Hollingsworth MD. 2011 Confined linear molecules inside an aperiodic supramolecular crystal: the sequence of superspace phases in *n*-hexadecane/urea. *J. Chem. Phys.* **135**, 204505. (doi:10.1063/1.3663711)
- Zerdane S, Mariette C, McIntyre GJ, Lemée-Cailleau MH, Rabiller P, Guérin L, Ameline JC, Toudic B. 2015 Neutron Laue and X-ray diffraction study of a new crystallographic superspace phase in *n*-nonadecane–urea. *Acta Crystallogr. B* **71**, 293–299. (doi:10.1107/S2052520615005442)
- Mariette C *et al.* 2013 Critical phenomena in higher dimensional spaces: the hexagonal-to-orthorhombic phase transition in aperiodic *n*-nonadecane/urea. *Phys. Rev. B* **87**, 104101. (doi:10.1103/PhysRevB.87.104101)
- Mariette C, Frantsuzov I, Wang B, Guérin L, Rabiller P, Hollingsworth MD, Toudic B. 2016 Frustrated pretransitional phenomena in aperiodic composites. *Phys. Rev. B* **94**, 184105. (doi:10.1103/PhysRevB.94.184105)
- Guérin L *et al.* 2015 Long-range modulation of a composite crystal in a five-dimensional superspace. *Phys. Rev. B* **91**, 184101. (doi:10.1103/PhysRevB.91.184101)
- López-Echarri A, Ruiz-Larrea I, Fraile-Rodríguez A, Díaz-Hernández J, Brezczewski T, Bocanegra EH. 2007 Phase transitions in the urea/*n*-nonadecane system by calorimetric techniques. *J. Phys. Condens. Matter* **19**, 186221. (doi:10.1088/0953-8984/19/18/186221)
- Couzi M, Guillaume F, Harris KDM, Palmer BA, Christensen K, Collins SP. 2016 The true structural periodicities and superspace group descriptions of the prototypical incommensurate composite materials: alkane/urea inclusion compounds. *Europhys. Lett.* **116**, 56001. (doi:10.1209/0295-5075/116/56001)