



HAL
open science

Short-term effect of pig slurry and its digestate application on biochemical properties of soils and emissions of volatile organic compounds

Cécile Monard, Laurent Jeanneau, Jean-Luc Le Garrec, Nathalie Le Bris,
Françoise Binet

► To cite this version:

Cécile Monard, Laurent Jeanneau, Jean-Luc Le Garrec, Nathalie Le Bris, Françoise Binet. Short-term effect of pig slurry and its digestate application on biochemical properties of soils and emissions of volatile organic compounds. *Applied Soil Ecology*, 2020, 147, pp.103376. 10.1016/j.apsoil.2019.103376 . hal-02355239

HAL Id: hal-02355239

<https://univ-rennes.hal.science/hal-02355239>

Submitted on 12 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Short-term effect of pig slurry and its digestate application on biochemical properties of soils and
2 emissions of volatile organic compounds

3 Cécile Monard^a, Laurent Jeanneau^b, Jean-Luc Le Garrec^c, Nathalie Le Bris^a, Françoise Binet^a

4 ^a Univ Rennes, CNRS, ECOBIO (Ecosystèmes, biodiversité, évolution) - UMR 6553, F-35000 Rennes,
5 France

6 ^b Univ Rennes, CNRS, Géosciences Rennes - UMR 6118, F-35000 Rennes, France

7 ^c Univ Rennes, CNRS, IPR (Institut de Physique de Rennes) - UMR 6251, F-35000 Rennes, France

8

9 cecile.monard@univ-rennes1.fr

10 laurent.jeanneau@univ-rennes1.fr

11 jean-luc.le-garrec@univ-rennes1.fr

12 nathalie.lebris@univ-rennes1.fr

13 francoise.binet@univ-rennes1.fr

14

15 Corresponding author: cecile.monard@univ-rennes1.fr

16 Address : Univ Rennes, CNRS, ECOBIO (Ecosystèmes, biodiversité, évolution) - UMR 6553, F-35000

17 Rennes, France

18 **Abstract**

19 Production of biogas through anaerobic digestion of organic wastes should play an important role in
20 sustainable development of energy supply, and the environmental effects of digestates have to be
21 assessed. We investigated the effect of anaerobic digestion of pig slurry (PS) on the molecular quality
22 of the digestate produced. The consequences of digested (DPS) and undigested PS use as organic soil
23 fertilizer on soil microbial and biochemical properties and C-gas emissions (CO₂ and volatile organic
24 compounds) were studied during a two-month incubation. PS and DPS differed in the amount of
25 volatile organic compounds (VOCs) emitted, in their organic C and lignin contents but not in their
26 active microbial composition. Application of both types of slurry to the soil immediately increased
27 the content of soil dissolved organic carbon (DOC) compared to the control soil. The application of
28 DPS induced few changes in the biochemical composition of soil organic matter compared to its raw
29 material (PS) that increased the amount of phenolic compounds. After 60 days, both amended and
30 control soils contained similar amounts of DOC, amended soils presenting a more diverse
31 biochemical composition of their soil organic matter. Application of both slurries to soil triggered a
32 succession of different active microbial communities, which could be attributed to the introduction
33 of new microorganisms and the input of new labile organic carbon. Changes in fungal communities
34 were stronger than those of bacteria and archaea; however, only slight differences were observed
35 between the slurries. Different fluxes and emission dynamics of five VOCs (methanol, acetone, DMS,
36 2-pentanone and phenol) were observed during the incubation time following application of PS or
37 DPS to soil while no differences in CO₂ emissions were observed. The present study calls for long-
38 term field studies with VOC analyses as a promising tool to differentiate organic fertilization
39 practices.

40

41 **Keywords**

42 Soil archaea, bacteria and fungi; Anaerobic digestion; Digestate; Biochemical organic matter
43 composition; Organic fertilizer; Soil quality.

44

45 **1. Introduction**

46 Production of biogas from organic wastes represents a major pillar of climate protection
47 measures and the European Council endorsed the new target for 2030 to increase the proportion of
48 renewable energy sources to 45% of the EU's final energy consumption (EREC, 2011). Bioenergy from
49 biogas production will thus play an important role in facilitating a sustainable development of energy
50 supply (Weiland, 2010). Biogas is produced through anaerobic digestion (AD) of wastes and leaves a
51 liquid, nutrient-rich, fermentation by-product (digestate) (Insam et al., 2014). Anaerobic digestates
52 are used as organic fertilizers applied on soil to improve its properties for crop production. Thus,
53 before we can consider the production of biogas as a sustainable practice, the environmental effects
54 of digestates have to be assessed.

55 During the AD process, the more labile organic fractions are rapidly degraded leading to an
56 increase in the biological stability of the digestate that is concentrated in recalcitrant organic
57 molecules compared with the initial raw material (Insam et al., 2015; Nkoa, 2014; Tambone et al.,
58 2009). However, there is almost no research on the molecular changes induced to the organic matter
59 (OM) during AD of slurry and their consequences once digestates are mixed with soil. Moreover, the
60 digestate is a mixture of undigested material and a microbial consortium specific to the anaerobic
61 process (Insam et al., 2015; Monlau et al., 2015), both capable of impacting the indigenous soil
62 microbiota. While the quality of organic fertilizers is defined by its OM, C and N contents, the soil
63 microbial compartment is the most commonly suggested biological indicator of soil quality. Bacteria,
64 fungi and archaea are responsible for most of the soil biochemical processes, and changes in their
65 communities can occur more quickly than changes in other soil characteristics (Doran and Parkin,
66 1994; Giacometti et al., 2013; Winding et al., 2005). Some literature studies reported differences in
67 the responses (abundance, diversity, activity) of soil microorganisms to different kinds of organic
68 fertilisers, included digestates, indicating a disturbance of soil biota (Abubaker et al., 2013; Barra
69 Caracciolo et al., 2015; Odlare et al., 2008; Sapp et al., 2015; Singla et al., 2014).

70 The composition of digestate strongly depends upon the composition of the raw material from
71 which they derived (Provenzano et al., 2011), but few studies compared the effects of digestate on
72 soil to those of its raw material. Moreover, prior to the wide-promotion of using digestates as crop
73 fertilizers, there is a need to assess their impact on C balance between soil and atmosphere. C
74 balance is mainly monitored by measuring CO₂ and CH₄ emissions but it has been demonstrated that,
75 by degrading soil OM, microorganisms also produce volatile organic compounds (VOCs) (Isidorov and
76 Jdanova, 2002; Stahl and Parkin, 1996; Stotzky et al., 1976). VOCs could thus constitute an additional
77 carbon loss from soil to the atmosphere and should be taken into account in the global C balance. It
78 has previously been shown that soil organic amendments including digestates and N-mineral
79 fertilization impact the VOC fluxes at the soil-atmosphere interface (Gray and Fierer, 2012; Seewald
80 et al., 2010). Recently, using field monitoring, Potard et al. (2017) observed that digested pig slurry
81 amendments differently impacted the soil bacterial communities and reduced the C-VOC fluxes from
82 soil to the atmosphere compared to the amendments of undigested pig slurry. Thus by modifying the
83 microbial decomposer communities and/or the soil OM status, undigested and digested organic
84 amendments impact the VOC emissions from soil.

85 Despite growing literature aiming to evaluate the impact of digestates on C balance (CH₄ and CO₂
86 emissions) (e.g. Eickenscheidt et al., 2014; Johansen et al., 2013; Møller et al., 2009; Pampillon-
87 Gonzalez et al., 2017; Pezzolla et al., 2012), there are still large uncertainties on how and to what
88 extent soil microbiota and soil biochemical quality would be influenced in the long-term, and
89 whether soil VOCs emissions would indicate such changes. The aim of the present study was thus to
90 determine i) if anaerobic microbial digestion affects the biochemical composition of the organic
91 substrate and ii) to what extent soil fertilisation with pig slurry digestate impacts soil microbial and
92 biochemical properties which in turn could change the C balance between C stabilisation in soil and C
93 loss through C-gas emissions. We compared our results using pig slurry digestate with its original raw
94 material and a control treatment without any organic fertilizer. We hypothesized that OM quality of
95 fertilizers changed due to the anaerobic digestion and that introduction of anaerobic microorganisms

96 through soil fertilization with digestate results in different patterns of VOCs emission from fertilized
97 soils. The strength of the present study rests on the combined analyses of soil OM biochemical
98 properties with the active soil microbial communities and the VOC emissions from soil. We observed
99 that the fungal communities and the biochemical composition of soil OM were less modified
100 following application of the digestate compared to that of the raw pig slurry. Overall, different
101 dynamics of VOC emissions in the two-month study were observed between the two slurries.

102

103 **2. Material and methods**

104 *2.1. Soil sampling and organic manures*

105 The soil was collected from the top 10 cm of a long-term experimental observatory for
106 environmental research (EFELE SOERE-PRO) of the French national institute of agronomic research
107 (INRA) located near Rennes (France: 48°05'35.9"N, 1°48'49.5"W) and dedicated to study the
108 environmental impact of the use of manures as soil fertilizers. The soil is a silt loam (71.3% silt, 12.6%
109 clay, 16.1% sand) with 1.9% of OM content, an acidic water pH_w of 5.8 and a total nitrogen content of
110 1.3 g kg⁻¹. The pig slurry (PS) and the liquid digestate derived from its own anaerobic digestion (DPS)
111 were collected from the Guernevez pig farming station and provided by the 'Chambre Régionale
112 d'Agriculture de Bretagne' (Rennes, France). The DPS was collected from a 20 m³ mesophilic (38°C)
113 tank reactor after a retention time of 20 days.

114

115 *2.2. Experimental design and set up*

116 The experimental design consisted in 27 microcosms corresponding to three treatments (control
117 without any amendment, PS and DPS amendments) replicated three times and performed for three
118 incubation times (initial time, 30 and 60 days). The fresh soil was sieved to < 4 mm, 2.6 kg (equivalent
119 dry weight) were placed into glass microcosms (cylinders: 25 cm high, 12 cm of diameter) and
120 compacted to a bulk density of 1.4 g cm⁻³. Soil moisture content was adjusted to 30% by adding
121 deionized water, a glass cover was placed on the top of each microcosm and they were incubated in

122 the dark at 15°C. The next day, the experiment was launched by adding 48 mL of PS, DPS or deionized
123 water (control soil; CS) at the microcosm surface (0.0113 m²), based on the recommended field
124 application rate of 4.25 liters per m². To mimic agricultural practices of organic amendment burying,
125 the top 4 cm of each microcosm was mixed with a sterile spatula and the microcosms were then
126 incubated at 15°C in the dark for up to 60 days. To maintain the soil moisture at 30% throughout the
127 incubation period, individual microcosms were weekly weighted and deionized water was added if
128 necessary. Samples from the two slurries (PS and the DPS) were stored at -20°C for organic matter
129 analyses and frozen in liquid nitrogen and kept at -80°C for further RNA extractions. Just after the
130 start of the experiment (day 0) and after 30 and 60 days of incubation, the top 10 cm of the
131 replicated soil microcosms were sampled and kept at -20°C for organic matter and microbial carbon
132 biomass analyses and frozen in liquid nitrogen and kept at -80°C for further RNA extractions.

133

134 *2.3. Gas sampling and CO₂ and VOCs measurements*

135 Gases were quantified just after the slurries or water addition and after 1, 2, 7, 14, 21, 28, 35, 42,
136 49, 56 and 60 days of incubation. For gas measurements, a 2 L glass dome including an inlet and
137 outlet airflow both composed of a gas-tight cap with silicone septum was placed at the top of each
138 microcosm and tightened using a dedicated clamp and a Teflon O-ring joint located between both of
139 them. The VOCs naturally emitted by the slurries (PS and the DPS) were also analyzed by pouring 48
140 mL of slurry in a glass beaker placed in an stainless steel base designed to be adapted to the glass
141 dome. After flushing the glass dome volume with ultra-pure air (Air zero alphagaz 2, Air Liquide,
142 France), the gases emitted by either the soil core or the slurries accumulated for one hour in the
143 glass dome. Prior to each gas sampling, the air in the headspace was homogenized using a 50 mL gas-
144 tight syringe and a charcoal cartridge air filter (Supelpure® HC Hydrocarbon trap, Sigma-Aldrich,
145 Germany) was then placed at the inlet airflow in order to purify the entering air. Three different gas
146 samplings and measurements were successively performed: CO₂ using a micro Gas Chromatograph
147 (µGC, Agilent), VOCs using both a micro Gas Chromatograph – Mass Spectrometer (microGC–MS, SRA

148 Instrument, France) after concentration on a tenax cartridge and a Proton Transfer Reaction-Mass
149 Spectrometer (PTR-MS; Ionicon GmbH, Innsbruck, Austria) (Lindinger et al., 1998) following gas
150 sampling in a Tedlar bag.

151 Precisely, CO₂ concentrations were first measured by a μ GC (Agilent) connected to the outlet
152 airflow of the glass dome. Hourly CO₂ emission rates (F_{CO_2} ; mgC-CO₂ m⁻² h⁻¹) were calculated based on
153 the following equation:

$$154 F_{CO_2} = \frac{C_{CO_2} \times V \times M_C}{VM \times A}$$

155 where C_{CO_2} is the concentration of CO₂ measured (ppm Mol), V is the volume of the headspace (0.002
156 m³), M is the molecular weight of C (12.01 g mol⁻¹), VM is the molecular gas volume (23.233 L.mol⁻¹ at
157 15°C and atmospheric pressure) and A is the soil area in the microcosms (0.0113 m²). A micro
158 Thermal Desorption unit (μ TD, AIRSENSE Analytics, Germany) was then placed at the outlet airflow of
159 the glass dome to trap the VOCs emitted in the headspace on a tenax cartridge at a flow rate of 200
160 mL.min⁻¹ at 35°C for 450 sec. Compounds were released by the trap at a helium flow rate of 300
161 mL.min⁻¹ at 180°C for 70 sec and then transferred to the micro gas chromatograph for 30 sec.
162 Compounds were separated using a capillary GC column (Stabilwax 10m) heated at 55°C and
163 detected by a 5975C mass detector Agilent (microGC-MS, SRA Instrument, France). The carrier gas
164 was helium at a constant pressure of 25 psi. The chromatograms were analyzed using the
165 ChemStation software (MSD ChemStation version).

166 Finally, 0.5 L of the headspace was sampled from the outlet airflow using a gas-tight syringe and
167 injected in a 2 L Tedlar gas collection bag (Interchim, Montluçon, France) for further VOC
168 measurements using a PTR-MS (QMS300, Ionicon GmbH, Innsbruck, Austria) (Lindinger et al., 1998).
169 Each bag was cleaned with successive flushes of ultra pure air before collection.

170 For PTR-MS analysis, Tedlar bags were connected directly to the apparatus and, operating at 135
171 Townsends (Td) (1 Td = 10⁻¹⁷ V cm² molecule⁻¹) in order to reduce VOC fragmentation. The "MID
172 mode" was selected during these experiments and consisted in scanning five masses (33, 59, 63, 87
173 and 95) in the range 21-95 amu at 0.2 s dwell time. These masses have been selected based on

174 literature, as they were previously detected as emitted by slurries (Feilberg et al., 2015; Sutton et al.,
175 1999) and taking part in both C and S cycles. Thus, mass 33 was attributed to methanol (Aaltonen et
176 al., 2013), mass 59 to acetone (Aaltonen et al., 2013), mass 63 to dimethyl sulfide (DMS) (Veres et al.,
177 2014), mass 87 to 2,3-butanedione (Feilberg et al., 2015) and mass 95 to phenol (Feilberg et al.,
178 2015). The ionization of VOCs either by O₂⁺ or NO⁺ was negligible as the O₂⁺:H₃O⁺ and NO⁺:H₃O⁺
179 ratios were ≤3.5% and 5% respectively. Five measurements of each mass concentration were
180 performed and averaged across treatments. To calculate the molecular density of each VOC, [R], the
181 following equation was used, accounting for drift tube temperature (333 K) and drift tube pressure
182 (2.20 mbar) (Zhao and Zhang 2004; de Gouw et al. 2003).

$$183 \quad [RH^+] = k_1[H_3O^+][R]t + k_2[H_3O^+(H_2O)][R]t$$

184 where [H₃O⁺], [H₃O⁺(H₂O)] and [RH⁺] are the ion signals (cps) at mass 21 (9500), mass 39 (9250), and
185 the target VOC, respectively; t is the average time (t) spent by reactants in drift tube (105 ms); k₁ and
186 k₂ are the rate coefficients between target VOC and H₃O⁺ or H₃O⁺(H₂O). Including H₃O⁺(H₂O) into the
187 reaction is a modification by Lindinger et al. (1998). Except for the slurries, each VOC flux (F_{VOC}) from
188 soil was calculated using the following equation:

$$189 \quad F_{VOCs} = \frac{C_{VOC} \times V \times M_c}{VM \times A}$$

190 where C_{VOC} is the concentration of VOC measured (ppb), V is the volume of the headspace (0.002 m³),
191 M is the molecular weight of the VOC (Mass-1), VM is the molecular gas volume (23.233 L.mol⁻¹ at
192 15°C and atmospheric pressure) and A is the soil area in the microcosms (0.0113 m²).

193

194 2.4. Soil water extractable organic matter (WEOM) composition analyses

195 The soil WEOM was extracted from 20 g of soil in 200 mL of deionized water. After orbital mixing
196 during one hour, the extracts were filtered through a 0.7 μm glass microfiber filter (whatman
197 International, Maidstone, England) and divided into two sub-samples. The first one was used for
198 dissolved organic carbon (DOC) quantification on a Shimadzu TOC-5050A total carbon analyzer and
199 the other 100 mL were lyophilized to analyze the composition of the WEOM. Pig slurry and its

200 digestate were filtered at 0.7 μm , aliquots were used for DOC quantification and 20 mL were
201 lyophilized. The lyophilisates were analysed by thermally assisted hydrolysis and methylation using
202 tetramethylammonium hydroxide coupled to a gas chromatograph and mass spectrometer (THM-
203 GC-MS) as previously described by Jeanneau et al. (2014). The detected compounds were classified
204 into five chemical classes: fatty acids (FA), carbohydrates biomarkers (CAR), lignin biomarkers (LIG),
205 phenolic compounds (PHE) including benzoic, phenylacetic, methylphenylacetic and phenylpropanoic
206 acids, and dimethyl indole (IND). Among the lignins, 8 compounds were detected: the vanillyl
207 phenols (vanillic acid, acetovanillone and vanillaldehyde), syringyl phenols (syringaldehyde, syringic
208 acid, acetosyringone) and cinnamyl phenols (*p*-coumaric acid and ferulic acid). The ratio of cinnamyl
209 to vanillyl phenols (C/V) was calculated; it reflects either the botanic source of lignin or, for a given
210 source of lignin, its biodegradation state (Hedges and Mann, 1979). FA can come from microbial and
211 plant-derived OM and these contributions can be differentiated on the basis of the length of the
212 aliphatic chain. FA with less than 19 C atoms mainly derived from microbial inputs, while FA with
213 more than 19 C and an even-over-odd predominance are characteristic of plant-derived OM
214 (Cranwell, 1974; Matsuda and Koyama, 1977). The proportion of microbial FA was calculated as the
215 percentage of low molecular weight FA (< C19) by excluding C16:0 and C18:0, which can derive from
216 both plant and microbial residues. The bacterial FA were thus composed of C12:0, C13:0, C14:0,
217 C15:0, C17:0, *anteiso* and *iso* C15:0 and C17:0, *iso* C16:0, C16:1 and C18:1, which are commonly used
218 as bacterial indicators (Frostegård et al., 1993). Among carbohydrates, the ratio of deoxyhexoses to
219 pentoses (deoxyC6/C5) was used as a balance between plant-derived (< 0.5) and microbial-derived (>
220 2) carbohydrates (Murayama, 1984; Oades, 1984).

221

222 2.5. Soil microbial biomass and community analyses

223 Soil microbial C biomass was determined by the chloroform fumigation and extraction method
224 (Vance et al., 1987) using a total organic carbon analyzer (1010 wet oxidation TOC - OI-Analytical -
225 USA). The microbial biomass C was calculated using the following equation:

226 Microbial C biomass = $(C_f - C_{nf}) \times K_c$

227 Where C_f is the DOC in fumigated soils, C_{nf} is the DOC in non-fumigated soil and K_c is a correction
228 factor of 2.64.

229 RNA was extracted from 2 g of soil using Power Soil MoBio kit (MoBio laboratories Inc., NY)
230 according to the manufacturer's instructions. The RNA extracts were electrophoresed over a 1%
231 agarose gel to assess RNA quality. DNA was removed using RQ1 DNase treatment (Promega,
232 Madison, WI) by mixing 35 μ L of RNA extract, 15 units of DNase I, 5.5 μ L of 10X DNase buffer, and the
233 mixture was incubated at 37°C for 30 min. reactions were terminated by addition of 5.55 μ L of DNase
234 I stop solution and incubation at 65°C for 10 min.

235 Terminal restriction fragment polymorphism (T-RFLP) analyses were performed for each soil
236 sample to determine the bacterial, archaeal and fungal community structures. Reverse transcription-
237 polymerase chain reaction (RT-PCR) were carried out using the Titan one-tube reverse transcription-
238 PCR kit (Roche Applied Science) to amplify the bacterial and archaeal 16S rRNA and the fungal ITS
239 with the corresponding forward primer to which was attached the 6-carboxyfluorescein (FAM) label
240 to the 5' end and reverse primer (Table 1). Reaction mixtures (25 μ L) contained soil sample RNA (3
241 μ L), 8 mM dNTPs, 1.25 μ L dithiothreitol (DTT), 2.5 U RNase inhibitor, 5 μ g bovine serum albumin
242 (BSA), 0.8 μ M of each primer and 0.5 μ L of enzyme mix in 1X reaction buffer. The following program
243 was used for the one step RT-PCR: 30 min at 51°C, 2 min at 94°C, 10 cycles of 30 s at 94°C, 30 s at the
244 specific primer's pair hybridization temperature (Table 1) and 1 min at 68°C, followed by 10 cycles for
245 bacteria, 15 for archaea and 25 for fungi of 30 s at 94°C, 30 s at the specific primer's pair
246 hybridization temperature (Table 1) and 1 min added to 5 more seconds per cycle at 68°C, and finally
247 30 s at 94°C, 30 s at the specific primer's pair hybridization temperature (Table 1) and 7 min at 68°C.
248 RT-PCR reactions were performed in duplicates for each sample, combined and the RT-PCR products
249 were purified with the GenElute PCR Clean-Up kit (Sigma-aldrich). The Labeled PCR products were
250 digested with the restriction enzyme MspI for bacteria and fungi and AluI (Promega, Madison, WI) for
251 archaea in 50 μ l separate reactions, at 37°C for 3 h. DNA was precipitated overnight and the T-RFs

252 were separated and accurately sized on the Biogenouest platform using an ABI system analyser
253 (3730; Applied Biosystems).

254 Raw data from Peak Scanner 1.0 software (Applied Biosystems) were exported to T-REX online
255 software (<http://trex.biohpc.org>) for the processing and analysis of T-RFLP data (Culman et al., 2009).
256 A clustering threshold of 0.5 bp was applied and only the length of the T-RFs between 50 and 500 bp
257 was taken into account for data analysis. Each T-RF was coded as a discrete variable (0 for its absence
258 or 1 for its presence) and the matrix generated was used in principal component analysis (PCA) to
259 compare the microbial community structures in the different treatments at the different incubation
260 times. PCA were generated using R statistical software version 3.4.1 ("<http://www.r-project.org/>")
261 and the ADE4 package (Dray and Dufour, 2007).

262 Microbial richness was estimated as the number of T-RFs for each sample and the richness
263 proportion of each microbial taxa among the overall richness (sum of the richness of bacteria,
264 archaea and fungi) was calculated for each sample and sampling time. These proportions were
265 represented in a ternary plot based on Graham and Midgley (2000).

266

267 *2.6. Statistical analyses*

268 All the statistical analyses were conducted using R statistical software version 3.4.1 ([http://www.r-](http://www.r-project.org)
269 [project.org](http://www.r-project.org)). Differences in i) the molecular composition of the DOM, ii) the amounts of microbial C
270 biomass, iii) the C-CO₂ emissions of the soil with or without slurries and iv) the VOC emissions by the
271 two slurries were analyzed using the non-parametric Kruskal-Wallis test (R package) followed by
272 Dunn's post hoc tests for pairwise comparisons using the Pairwise Multiple Comparisons of Mean
273 Rank Sums (PMCMR) package (<http://CRAN.R-project.org/> package=PMCMR).

274

275 **3. Results and discussion**

276 *3.1. Effects of anaerobic digestion on the biochemical and microbial properties of the slurries and*
277 *their VOC emissions*

278 The digested and undigested pig slurries differed from each other in some of their biochemical
279 properties (Fig. 1) but not significantly in their active microbial composition (Fig. 2). During the
280 anaerobic digestion, the slurry lost its C as CO₂ and CH₄ leading to a digestate with a half-reduced
281 organic C content (7.0 and 3.8 mg.ml⁻¹ in PS and DPS, respectively) (Fig. 1). Such a decrease is in
282 accordance with the value reported in the literature with a reduction from 40% to 82.5% of total C in
283 digested manures compared with undigested ones (Abubaker et al., 2012). The anaerobic digestion
284 resulted also in a higher proportion of lignin in the digested PS compared with the PS (4.5% and 0.7%,
285 respectively). Such concentration of recalcitrant molecules as lignin during anaerobic digestion has
286 been previously reported by Tambone et al. (2009).

287 Phenolic compounds were the main components of the dissolved fraction of both slurries (96.2%
288 and 91.2% in PS and DPS, respectively, Fig.1). They are derived from lignin degradation (Bugg et al.,
289 2011; Elder and Kelly, 1994) and among these aromatic acids, benzoic acids are used in porcine
290 breeding as dietary supplement to decrease ammonia emissions (Hansen et al., 2007) and have
291 previously been detected in pig urine (Yasuhara et al., 1984). PHE are known to be key metabolic
292 intermediates in the anaerobic digestion (Levén and Schnürer, 2005). For example, phenylpropanoic
293 acid is thought to come from the metabolic degradation of tyrosine in pig feces (Spoelstra, 1978) and
294 phenylacetic acid has been proposed as an indicator of the overall stability of the anaerobic process
295 (Iannotti et al., 1986).

296 During anaerobic digestion, the establishment of successive microbial consortia (Insam et al.,
297 2015; Manyi-Loh et al., 2013; Monlau et al., 2015) might lead to differences in the microbial
298 communities of PS and DPS. However, in the present study we only observed slight differences in the
299 active bacterial communities of PS and DPS using T-RFLP on RNA but surprisingly, no changes were
300 detected for fungi and archaea (Fig. 2). Although archaea are responsible for the production of CH₄
301 that takes place at the end of the anaerobic metabolic pathway of organic matter degradation, they
302 should have been stimulated and we could have expected changes in their community structure. It
303 has previously been shown that, based on DNA analysis, anaerobic digestion leads to an enrichment

304 of archaea from the genus *Methanosarcina* (Sun et al., 2015). By using RNA-based analysis, we did
305 not observe such changes suggesting that a common part of the community representing a core of
306 dominant microorganisms was active in both slurries. Even if the explained variations of the PCAs
307 were low (Fig. 2), we can speculate that the initial raw material rather than the AD process controlled
308 the diversity of active microorganisms in the digestate.

309 The emission profiles of five selected VOCs (methanol, acetone, DMS, 2-pentanone and phenol)
310 were determined for the slurries themselves and their concentrations varied from 1.8 up to 6.2 times
311 higher in PS than in DPS, excepting for the 2-pentanone which emissions were 1.2 times higher in
312 DPS than in PS (Table 2). The mineralization that occurred during the AD, by reducing the OM
313 available for microbial degradation, should have limited further emissions of VOCs in digestates.
314 Moreover, the emission of the studied VOCs might have been stimulated during the AD, as previously
315 observed during the composting of biowaste (Smet et al., 1999), and thus limiting their detection in
316 the digested slurry.

317

318 3.2. Impact of slurry application on soil WEOM

319 In the control soil receiving no organic fertilizer (CS), the amount of DOC increased with
320 incubation time at 15°C from 69 to 90 $\mu\text{g g}^{-1}$ dry soil, although these differences were not significant
321 due to high variability between replicates after 60 days of incubation (Kruskal Wallis, $H = 4.33$, $P =$
322 0.12; Fig. 1). The WEOM in the CS at the beginning of the incubation was mainly composed of FA
323 (87.2%) from microbial origin (84.9%) with few proportions of CAR (4.9%), LIG (4.6%) and PHE (3.0%)
324 (Fig. 1). This initial composition varied through incubation time, the proportion of microbial fatty
325 acids being the highest at 30 days (93.4 %), and finally resulted, after 60 days of incubation, in an
326 enrichment in carbohydrates (10.7%) to the detriment of the initial microbial fatty acids content
327 (73.9%) (Fig. 1).

328 As expected, both slurry additions to soil immediately increased the soil DOC content that was 2.8
329 and 2.6 times higher in soils receiving PS and DPS than in the control soil at the start of the

330 incubation, respectively (Fig. 1). Moreover, the application of PS to soil was linked to an increase in
331 the proportion of PHE (34.6%), LIG (7.8%) and IND (5.9%) associated with a decrease in FA (47%)
332 (Fig.1). This was not observed following DPS application to soil since it had a weaker impact on the
333 composition of the soil WEOM that only exhibited higher proportions of LIG (10.0%) and IND (2.2%)
334 than those observed in CS. Interestingly, the proportion of PHE in the soil that received DPS (SDPS) at
335 the start of the experiment was similar to the one in CS (3.6% vs 3.0% in SDPS and CS, respectively)
336 (Fig. 1) while it was a main component of both slurries. Such differences between the two amended
337 soils might be explained by the low initial content of DOC and thus PHE compounds observed in DPS
338 (6.7 and 3.4 mg C-PHE. L⁻¹ in PS and DPS, respectively).

339 The effect of PS application to soil was observed earlier than the one of DPS and, after 30 days,
340 the amount of DOC was still 1.5 times higher in the soil that received PS (SPS) than in CS or DPS
341 (SDPS)(Fig. 1). At that time, both slurries altered similarly the composition of soil DOC by increasing
342 its content in carbohydrates, lignin and phenolic compounds to the detriment of the microbial FA.
343 Finally, at the end of the experiment, the amended soils did not contain more DOC than the control
344 soil but they exhibited a more diverse composition of their WEOM with less microbial fatty acids and
345 higher carbohydrates and lignin contents (Fig. 1).

346 Both the lignin and carbohydrates in the WEOM of the unamended control soil mainly derived
347 from degraded material and plant inputs, as indicated by the very low C/V lignin and deoxy C6/C5
348 ratios, respectively (Fig. 3). In accordance with a mixing between the inherent soil WEOM and the
349 fine organic fraction (<0.7 µm) of PS (C/V lignin = 2.1 and deoxy C6/C5 = 2.3) and DPS (C/V lignin = 2.4
350 and deoxy C6/C5 = 1.6), these two indicators of lignin and carbohydrates sources temporarily
351 increased significantly or not after the application of slurries (Fig. 3). This transient changes in the
352 C/V lignin ratio reflected a new source of lignin brought by the slurries that was less degraded than
353 the one already present in soil. The increase in the deoxy C6/C5 ratio clearly indicated a higher
354 proportion of microbial carbohydrates (Rumpel and Dignac, 2006) linked to the ephemeral increase
355 in the C biomass of soil microorganisms observed just after the application of both slurries (Fig. 4).

356 However, both C/V and deoxy C6/C5 ratios recovered similar low values as the one observed in the
357 control soil after 60 days of incubation. Overall, the C/V lignin ratio was significantly correlated to
358 both soil DOC and microbial C biomass ($R^2 = 0.49$, $P < 0.001$; $R^2 = 0.47$, $P < 0.001$, respectively)
359 indicating the importance of the OM state of biodegradation in the dynamic of the soil C
360 compartment.

361 Altogether our results indicate that the slurries brought a new microbial source that rapidly but
362 briefly fueled the dissolved organic fraction of the soil. As the majority of soil microorganisms are
363 heterotroph, they rely on the availability of organic C, addition of exogenous OM as PS and DPS
364 should thus have supported a larger microbial biomass. In the present study no such stimulation was
365 observed during the experiment lifespan (Fig. 4) while the labile organic C was initially up to three
366 times higher in the amended soils than in the control soil (Fig. 1). The time scale of our experiment
367 might not have been adapted and a short-term impact of the slurry application might have been
368 visible by analyses performed within the first 30 days of incubation. Indeed, Potard et al. (2017),
369 based on field monitoring and next generation sequencing analysis, detected changes in soil
370 microbial communities seven days after either PS or DPS application that were not detected anymore
371 after 36 days. However, Pezzolla et al. (2015) observed fluctuating trend in the C biomass of soil
372 microorganisms up to 30 days following application of digestate from pig slurry to an agricultural soil
373 in controlled conditions. The impact of organic amendments on soil microorganisms is thus not as
374 obvious as expected and might also depend on the quantity and quality of the OM added. In the
375 present study, the fractions of dissolved organic matter ($< 0.7 \mu\text{m}$ fractions) of both slurries were
376 mainly composed of phenolic compounds and also of lignin for the DPS that are recalcitrant to
377 biodegradation and might have constrained microorganisms by stimulating only a few part of the
378 community able to degrade such compounds.

379

380 *3.3. Response of soil microbial communities*

381 The community structures of active bacteria, archaea and fungi were analyzed by T-RFLP just after
382 slurry application to soil and after 30 and 60 days of incubation (Fig. 2). Specific changes in the soil
383 microbial communities were observed for each microbial taxa and slurry following slurry application
384 to soil. At the start of the experiment, just after the application of slurries to soil (SPS0, SDPS0),
385 changes observed in fungal, archaeal and to a lesser extent in bacterial communities are likely due to
386 the introduction of microbial communities specific to PS and DPS (Fig. 2) that also increased the
387 microbial C-biomass (Fig. 4). Cropped soils are well known to sustain bacteria-dominant microbial
388 communities (Hendrix et al., 1990; Strickland and Rousk, 2010) that might have buffered the impact
389 of the introduction of active bacteria from slurries. Interestingly, the active fungal communities in the
390 soil receiving both slurries (SPS0, SDPS0) were closer to the one in the slurries themselves (PS, DPS)
391 than to one in the initial soil (CS0) (Fig. 2C). This might be due to the low abundance of fungi usually
392 observed in cropped soils (Hendrix et al., 1990; Strickland and Rousk, 2010), allowing their
393 community to be greatly shaped by the application of slurries (as substrates) and their proper fungal
394 communities. Indeed, while fungal communities have been poorly studied during anaerobic
395 digestion, a recent study highlighted their high diversity in several digesters supplied with different
396 plants and livestock manures (Langer et al., 2019).

397 Together with temporal dynamics of soil microbial communities observed in the control soil,
398 slurry application to soil led to successive active microbial communities. Several studies observed
399 differences in the response of soil microorganisms following the application of slurries and digestates
400 (Abubaker et al., 2013; Barra Caracciolo et al., 2015; Chen et al., 2012; Walsh et al., 2012). These
401 differences were mainly attributed to the composition and availability of the OM added, the digested
402 material being enriched in more stable compounds while the undigested one was concentrated in
403 labile organic C easily degradable by microorganisms (Abubaker et al., 2013; Marcato et al., 2009).
404 Such differences in the two slurries may induce shifts in the structure of the microbial community,
405 the digestate stimulating the slow-growing microorganisms compared with application of undigested
406 organic matter (Chen et al., 2012). In the present study, the application of both slurries greatly

407 impacted the composition of the active bacterial and archaeal communities after 30 days of
408 incubation (SPS30 and SDPS30), the disturbance generated by DPS being higher than by PS. At 60
409 days, the composition of active bacterial and archaeal communities in amended soils was close to the
410 one observed in control soils with the exception of active archaea in soil amended with DPS, the
411 effect of the digestate being subtle but still detected (Fig. 2A and 2B). The application of PS strongly
412 constrained the fungal community that was specific and stable with time in this treatment (SPS0,
413 SPS30 and SPS60) whilst it was highly variable in the SDPS treatment (Fig. 2C). The differences
414 observed in the fungal community in SPS and SDPS might result from their interaction with the
415 bacterial community. It is suggested that bacteria tend to dominate the ecological interactions with
416 fungi in conditions of rich resources such as in the SPS where labile organic C is available, whilst fungi
417 grow when conditions are unfavorable for bacteria (Rousk and Bååth, 2011; Walsh et al., 2012).
418 Moreover, fungi are more efficient for the degradation of complex substrates that should make them
419 better competitors in SDPS treatments (Meidute et al., 2008).

420 The species richness, estimated as the number of T-RFs, within the three microbial taxa was
421 investigated (Fig. 5). The disturbance observed in microbial community structures 30 days after
422 slurry application corresponded to a homogeneous distribution of the richness across the microbial
423 taxa whatever the treatment, with no taxa representing more than 50% of the total T-RFs (Fig. 5B).
424 At the end of incubation, the slurry application induced changes in the distribution of species
425 richness that finally led to distinct communities, CS being enriched in fungi and depleted in archaea,
426 soils amended with SPS being depleted in archaea and SDPS being slightly enriched in bacteria.

427

428 *3.4. Soil volatile organic compounds and Carbon dioxide emissions*

429 C-CO₂ fluxes were measured in the different soil microcosms during the incubation time. The
430 respiration kinetics over 60 days showed a clear time effect (Kruskal Wallis test, $H = 42.8$, $p < 0.001$)
431 but no significant impact of the slurry application despite it represented an additional C source to soil
432 compared with the control soil (Kruskal Wallis test, $H = 1.4$, $p = 0.49$). The C-CO₂ fluxes similarly

433 varied, from 11.7 ± 1.1 to 18.4 ± 0.4 in CS, from 11.0 ± 3.2 to 20.6 ± 0.1 in SPS and from 12.3 ± 0.1 to
434 19.3 ± 1.3 mg C-CO₂ m⁻² h⁻¹ in SDPS, leading to cumulative emissions of 3.7, 3.9 and 4.0 g C-CO₂ m⁻²
435 after 60 days of incubation in CS, SPS and SDPS, respectively (Fig. 6). The loss of C by aerobic
436 respiration was thus similar whatever the treatment, while different contents of DOC were observed
437 for amended soils and control soil (Fig. 1). It is likely that the amount of DOC in soil does not reflect
438 the available C to be mineralized, probably due to limitation in the availability of other nutrients for
439 soil microorganisms.

440 The present study demonstrates changes in both microbial communities and WEOM composition
441 following application of either PS or DPS to soil. We hypothesized that such modifications of soil
442 biochemical properties induced changes in the emissions of the VOCs through SOM biodegradation
443 by microorganisms. It is known that VOCs produced by soil microorganisms strongly depend on both
444 quality of the substrate degraded and on microbial diversity and activity (McNeal and Herbert, 2009;
445 Peñuelas et al., 2014; Stotzky et al., 1976). The VOC emissions were thus measured using both a μ GC-
446 MS and a PTR-MS.

447 Due to its high detection limit and to our experimental conditions providing a restricted volume
448 of soil, the μ GC-MS did not detect any VOC in CS whatever the incubation time. However, up to two
449 days after the application of both PS and DPS, emissions of acetone, 2-butanone, dichloromethane,
450 2-pentanone and toluene were identified. These molecules have already been described as emitted
451 directly by different kinds of manures or by soil following their application (Abis et al., 2018; Feilberg
452 et al., 2015; Trabue et al., 2010; Woodbury et al., 2006). Moreover, we found that these molecules
453 were emitted by the slurries (Table 2) and thus we cannot exclude their emissions by the slurries
454 themselves following their application to soil.

455 Thanks to the lower detection limit of the PTR-MS we measured the emission of five masses all
456 along the incubation time and corresponding to methanol (m33), acetone (m59), DMS (m63), 2-
457 pentanone (m87) and phenol (m95) (Fig.7). Contrarily to CO₂, distinct fluxes and dynamics of VOC

458 emissions were observed after application of slurries to soil, but we mainly observed marginal
459 responses and no specific VOC emission patterns were identified.

460 Acetone, 2-pentanone and DMS emissions were the most abundant C-VOC fluxes. Just after slurry
461 application, the soil amended with PS systematically emitted higher concentrations of any VOC
462 compared to the SDPS and C-VOCs fluxes from SDPS were even lower than those from control soil for
463 2-pentanone and methanol (Fig. 7). One day after both slurry applications, the C-VOC fluxes from the
464 microcosms with slurries were higher than those in CS, except for the phenol emission, and, this
465 increase lasted from one day for the acetone in SDPS up to 49 days for the DMS. Distinctive VOC
466 emissions following the application of PS and DPS to soil have already been observed, especially for
467 methanol fluxes that were induced after PS fertilization of soil compared with DPS fertilization
468 (Potard et al., 2017). Interestingly, for all the VOCs measured, the emissions could be lower than
469 those detected in CS (Fig. 7). Once produced, the VOCs can either be degraded by soil
470 microorganisms that use them as C source (Owen et al., 2007; Ramirez et al., 2009) or adsorbed on
471 soil that acts as a sink of VOCs (Asensio et al., 2007), two processes that reduce their emission from
472 soil to the atmosphere and make it difficult to identify specific VOC emission patterns depending on
473 amendment practices. This specificity between VOC emission patterns and amendment practices
474 have been already observed on soils from long-term experimental sites, suggesting that repeated
475 inputs are needed to allow to differentiate fertilization practices according to the VOC spectra
476 emitted by soil (Abis et al., 2018; Seewald et al., 2010).

477

478 **4. Conclusions**

479 Anaerobic digestion of pig slurry increased dissolved organic C by a factor of two, without affecting
480 microbial community structures. The application of both slurries to soil led to a similar increase in soil
481 DOC contents, microbial C-biomass and soil respiration, and disturbed soil microbial communities.
482 However, instantaneous changes in the quality of the WEOM were mainly observed following the
483 application of PS only. After 60 days, the biochemical composition of the soil WEOM was more

484 diverse in the amended soils but the DOC content was not affected anymore by the two slurry
485 applications. This disappearance of soil DOC could not be attributed to either an enhanced
486 mineralization or an increase in VOC emissions. It is likely that the amount of DOC in soil did not
487 reflect the available C to be mineralized. Further analyses of soil available nutrients (e.g. nitrogen,
488 phosphorus...) might have helped to better understand the effect of slurry application on C balance
489 and the associated biogeochemical cycles. Overall, the application of PS induced greater changes in
490 the biochemical composition of the soil organic matter and fungal communities compared to its
491 digestate (DPS), and different fluxes and emission dynamics of VOCs were observed following the
492 application of the two slurries. The present study i) calls for strengthening long-term on-field studies
493 to better predict the future impact of slurry and digestate amendments on soil quality and C-balance
494 and ii) supports the use of VOC analyses to differentiate organic fertilization practices.

495

496 **Acknowledgements**

497 We thank the editor and reviewer for their comments and suggestions. We thank A. Prinzing and M.
498 Bormans for English editing. This research was supported by the CNRS-INSU 'Ecosphère Continentale
499 et Côtière' program. The EFELE field experiment forms part of the SOERE-PRO (network of long-term
500 experiments dedicated to the study of impacts of organic waste product recycling) certified by
501 ALLENVI (Alliance Nationale de Recherche pour l'Environnement) and integrated as a service of the
502 'Investment d'Avenir' infrastructure AnaEE-France, overseen by the French National Research Agency
503 (ANR-11-INBS-0001). The authors are grateful to F. Gaillard and P. Le Roy for their contribution to
504 field experiment, to T. Morvan for the access to the EFELE site and the supply of soil and organic
505 products, to T. Ceau, S. Mahé, Q. Allard, A. Ola and J-P. Caudal for their technical supports. The
506 authors also wish to thank the European Union for financial support for the PTR-MS instrument.

507 **References**

- 508 Aaltonen, H., Aalto, J., Kolari, P., Pihlatie, M., Pumpanen, J., Kulmala, M., Nikinmaa, E., Vesala, T.,
509 Bäck, J., 2013. Continuous VOC flux measurements on boreal forest floor. *Plant Soil* 369, 241–
510 256.
- 511 Abis, L., Loubet, B., Ciuraru, R., Lafouge, F., Dequiedt, S., Houot, S., Maron, P.A., Bourgeteau-Sadet,
512 S., 2018. Profiles of volatile organic compound emissions from soils amended with organic
513 waste products. *Sci. Total Environ.* 636, 1333–1343.
- 514 Abubaker, J., Cederlund, H., Arthurson, V., Pell, M., 2013. Bacterial community structure and
515 microbial activity in different soils amended with biogas residues and cattle slurry. *Appl. Soil*
516 *Ecol.* 72, 171–180.
- 517 Abubaker, J., Risberg, K., Pell, M., 2012. Biogas residues as fertilisers – Effects on wheat growth and
518 soil microbial activities. *Appl. Energy* 99, 126–134.
- 519 Asensio, D., Peñuelas, J., Filella, I., Llusà, J., 2007. On-line screening of soil VOCs exchange responses
520 to moisture, temperature and root presence. *Plant Soil* 291, 249–261.
- 521 Barra Caracciolo, A., Bustamante, M.A., Nogues, I., Di Lenola, M., Luprano, M.L., Grenni, P., 2015.
522 Changes in microbial community structure and functioning of a semiarid soil due to the use of
523 anaerobic digestate derived composts and rosemary plants. *Geoderma* 245–246, 89–97.
524 doi:10.1016/j.geoderma.2015.01.021
- 525 Bugg, T.D.H., Ahmad, M., Hardiman, E.M., Rahmanpour, R., 2011. Pathways for degradation of lignin
526 in bacteria and fungi. *Nat. Prod. Rep.* 28, 1883. doi:10.1039/c1np00042j
- 527 Chen, R., Blagodatskaya, E., Senbayram, M., Blagodatsky, S., Myachina, O., Dittert, K., Kuzyakov, Y.,
528 2012. Decomposition of biogas residues in soil and their effects on microbial growth kinetics
529 and enzyme activities. *Biomass and Bioenergy* 45, 221–229.
- 530 Cranwell, P.A., 1974. Monocarboxylic acids in lake sediments: Indicators, derived from terrestrial and
531 aquatic biota, of paleoenvironmental trophic levels. *Chem. Geol.* 14, 1–14.
- 532 Culman, S.W., Bukowski, R., Gauch, H.G., Cadillo-Quiroz, H., Buckley, D.H., 2009. T-REX: software for

533 the processing and analysis of T-RFLP data. *BMC Bioinformatics* 10, 171.

534 Doran, J.W., Parkin, T., 1994. Defining and assessing soil quality, in: Doran, J.W., Coleman, D.C.,
535 Bezdicsek, D., Stewart, B. (Eds.), *Defining Soil Quality for Sustainable Environment*. American
536 Society of Agronomy, Madison, pp. 3–21.

537 Dray, S., Dufour, A.-B., 2007. The ade4 package: implementing the duality diagram for ecologists. *J.*
538 *Stat. Softw.* 22, 1–20.

539 Eickenscheidt, T., Freibauer, A., Heinichen, J., Augustin, J., Drösler, M., 2014. Short-term effects of
540 biogas digestate and cattle slurry application on greenhouse gas emissions affected by N
541 availability from grasslands on drained fen peatlands and associated organic soils.
542 *Biogeosciences* 11, 6187–6207.

543 Elder, D., Kelly, D., 1994. The bacterial degradation of benzoic acid and benzenoid compounds under
544 anaerobic conditions: unifying trends and new perspectives. *FEMS Microbiol. Rev.* 13, 441–468.

545 Feilberg, A., Bildsoe, P., Nyord, T., 2015. Application of PTR-MS for measuring odorant emissions
546 from soil application of manure slurry. *Sensors (Basel)*. 15, 1148–1167.

547 Frostegård, A., Tunlid, A., Bååth, E., 1993. Phospholipid Fatty Acid composition, biomass, and activity
548 of microbial communities from two soil types experimentally exposed to different heavy metals.
549 *Appl. Environ. Microbiol.* 59, 3605–17.

550 Gardes, M., Bruns, T.D., 1993. ITS primers with enhanced specificity for Basidiomycetes - application
551 to the identification of mycorrhizae and rusts. *Mol. Ecol.* 2, 113–118.

552 Giacometti, C., Demyan, M.S., Cavani, L., Marzadori, C., Ciavatta, C., Kandeler, E., 2013. Chemical and
553 microbiological soil quality indicators and their potential to differentiate fertilization regimes in
554 temperate agroecosystems. *Appl. Soil Ecol.* 64, 32–48.

555 Graham, D.J., Midgley, N.G., 2000. Graphical representation of particle shape using triangular
556 diagrams: an Excel spreadsheet method. *Earth Surf. Process. Landforms* 25, 1473–1477.

557 Gray, C.M., Fierer, N., 2012. Impacts of nitrogen fertilization on volatile organic compound emissions
558 from decomposing plant litter. *Glob. Chang. Biol.* 18, 739–748.

559 Hansen, C.F., Sørensen, G., Lyngbye, M., 2007. Reduced diet crude protein level, benzoic acid and
560 inulin reduced ammonia, but failed to influence odour emission from finishing pigs. 10th Int.
561 Symp. Dig. Physiol. Pigs, Denmark 2006, Part 2 109, 228–231.

562 Hedges, J.I., Mann, D.C., 1979. The characterization of plant tissues by their lignin oxidation products.
563 *Geochim. Cosmochim. Acta* 43, 1803–1807.

564 Hendrix, Crossley, D., Jr, Blair, J., Coleman, D., 1990. Soil biota as components of sustainable
565 agroecosystems, in: Edwards, C.A., Lal, R., Madden, P., Miller, R.H., House G. (Eds.), *Sustainable*
566 *Agricultural Systems*. Soil and Water Conservation Society, Ankeny, Iowa, pp. 637–654.

567 Iannotti, E.L., Mueller, R.E., Sievers, D.M., Georgacakis, D.G., Gerhardt, K.O., 1986. Phenylacetic acid
568 in an anaerobic swine manure digester. *J. Ind. Microbiol.* 1, 57–61.

569 Insam, H., Franke-whittle, I.H., Podmirseg, S.M., 2014. Agricultural waste management in Europe,
570 with an emphasis on anaerobic digestion. *J. Integr. F. Sci.* 11, 13–17.

571 Insam, H., Gómez-Brandón, M., Ascher, J., 2015. Manure-based biogas fermentation residues –
572 Friend or foe of soil fertility? *Soil Biol. Biochem.* 84, 1–14.

573 Isidorov, V., Jdanova, M., 2002. Volatile organic compounds from leaves litter. *Chemosphere* 48,
574 975–9.

575 Jeanneau, L., Jaffrezic, A., Pierson-Wickmann, A.-C., Gruau, G., Lambert, T., Petitjean, P., 2014.
576 Constraints on the sources and production mechanisms of dissolved organic matter in soils
577 from molecular biomarkers. *Vadose Zo. J.* 13, 0.

578 Johansen, A., Carter, M.S., Jensen, E.S., Huggard-Nielsen, H., Ambus, P., 2013. Effects of digestate
579 from anaerobically digested cattle slurry and plant materials on soil microbial community and
580 emission of CO₂ and N₂O. *Appl. Soil Ecol.* 63, 36–44.

581 Langer, S.G., Gabris, C., Einfalt, D., Wemheuer, B., Kazda, M., Bengelsdorf, F.R., 2019. Different
582 response of bacteria, archaea and fungi to process parameters in nine full-scale anaerobic
583 digesters. *Microb. Biotechnol.* 1751–7915.13409.

584 Levén, L., Schnürer, A., 2005. Effects of temperature on biological degradation of phenols, benzoates

585 and phthalates under methanogenic conditions. *Int. Biodeterior. Biodegradation* 55, 153–160.

586 Lindinger, W., Hansel, A., Jordan, A., 1998. On-line monitoring of volatile organic compounds at pptv
587 levels by means of proton-transfer-reaction mass spectrometry (PTR-MS) medical applications,
588 food control and environmental research. *Int. J. Mass Spectrom. Ion Process.* 173, 191–241.

589 Lueders, T., Friedrich, M., 2000. Archaeal population dynamics during sequential reduction processes
590 in rice field soil. *Appl. Environ. Microbiol.* 66, 2732–42.

591 Manyi-Loh, C.E., Mamphweli, S.N., Meyer, E.L., Okoh, A.I., Makaka, G., Simon, M., 2013. Microbial
592 anaerobic digestion (bio-digesters) as an approach to the decontamination of animal wastes in
593 pollution control and the generation of renewable energy. *Int. J. Environ. Res. Public Health* 10,
594 4390–4417.

595 Marcato, C.-E., Mohtar, R., Revel, J.-C., Pouech, P., Hafidi, M., Guiresse, M., 2009. Impact of
596 anaerobic digestion on organic matter quality in pig slurry. *Int. Biodeterior. Biodegradation* 63,
597 260–266.

598 Marchesi, J.R., Sato, T., Weightman, A.J., Martin, T.A., Fry, J.C., Hiom, S.J., Dymock, D., Wade, W.G.,
599 1998. Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding
600 for bacterial 16S rRNA. *Appl. Environ. Microbiol.* 64, 795–9.

601 Matsuda, H., Koyama, T., 1977. Early diagenesis of fatty acids in lacustrine sediments—II. A statistical
602 approach to changes in fatty acid composition from recent sediments and some source
603 materials. *Geochim. Cosmochim. Acta* 41, 1825–1834.

604 McNeal, K.S., Herbert, B.E., 2009. Volatile organic metabolites as indicators of soil microbial activity
605 and community composition shifts. *Soil Sci. Soc. Am. J.* 73, 579–588.

606 Meidute, S., Demoling, F., Bååth, E., 2008. Antagonistic and synergistic effects of fungal and bacterial
607 growth in soil after adding different carbon and nitrogen sources. *Soil Biol. Biochem.* 40, 2334–
608 2343.

609 Møller, J., Boldrin, A., Christensen, T.H., 2009. Anaerobic digestion and digestate use: accounting of
610 greenhouse gases and global warming contribution. *Waste Manag. Res.* 27, 813–824.

611 Monlau, F., Sambusiti, C., Ficara, E., Aboulkas, A., Barakat, A., Carrère, H., Rulkens, W., Xu, G., Dijk, L.
612 Van, Cornelissen, G., 2015. New opportunities for agricultural digestate valorization: current
613 situation and perspectives. *Energy Environ. Sci.* 8, 2600–2621.

614 Murayama, S., 1984. Changes in the monosaccharide composition during the decomposition of
615 straws under field conditions. *Soil Sci. plant Nutr.* 30, 367–381.

616 Nkoa, R., 2014. Agricultural benefits and environmental risks of soil fertilization with anaerobic
617 digestates: A review. *Agron. Sustain. Dev.* 34, 473–492.

618 Oades, J.M., 1984. Soil organic matter and structural stability: mechanisms and implications for
619 management, in: Tinsley, J., Darbyshire, J.F. (Eds.), *Biological Processes and Soil Fertility,*
620 *Developments in Plant and Soil Sciences.* Springer Netherlands, pp. 319–337.

621 Odlare, M., Pell, M., Svensson, K., 2008. Changes in soil chemical and microbiological properties
622 during 4 years of application of various organic residues. *Waste Manag.* 28, 1246–1253.

623 Osborn, A.M., Moore, E.R.B., Timmis, K.N., 2000. An evaluation of terminal-restriction fragment
624 length polymorphism (T-RFLP) analysis for the study of microbial community structure and
625 dynamics. *Environ. Microbiol.* 2, 39–50.

626 Owen, S.M., Clark, S., Pompe, M., Semple, K.T., 2007. Biogenic volatile organic compounds as
627 potential carbon sources for microbial communities in soil from the rhizosphere of *Populus*
628 *tremula*. *FEMS Microbiol. Lett.* 268, 34–39.

629 Pampillon-Gonzalez, L., Luna-Guido, M., Ruiz-Valdiviezo, V.M., Franco-Hernandez, O., Fernandez-
630 Luqueno, F., Paredes-Lopez, O., Hernandez, G., Dendooven, L., 2017. Greenhouse gas emissions
631 and growth of wheat cultivated in soil amended with digestate from biogas production.
632 *Pedosphere* 27, 318–327.

633 Peñuelas, J., Asensio, D., Tholl, D., Wenke, K., Rosenkranz, M., Piechulla, B., Schnitzler, J.P., 2014.
634 Biogenic volatile emissions from the soil. *Plant, Cell Environ.* 37, 1866–1891.

635 Pezzolla, D., Bol, R., Gigliotti, G., Sawamoto, T., López, A.L., Cardenas, L., Chadwick, D., 2012.
636 Greenhouse gas (GHG) emissions from soils amended with digestate derived from anaerobic

637 treatment of food waste. *Rapid Commun. Mass Spectrom.* 26, 2422–2430.

638 Pezzolla, D., Marconi, G., Turchetti, B., Zadra, C., Agnelli, A., Veronesi, F., Onofri, A., Benucci, G.M.N.,
639 Buzzini, P., Albertini, E., Gigliotti, G., 2015. Influence of exogenous organic matter on
640 prokaryotic and eukaryotic microbiota in an agricultural soil. A multidisciplinary approach. *Soil*
641 *Biol. Biochem.* 82, 9–20.

642 Potard, K., Monard, C., Le Garrec, J.-L., Caudal, J.-P., Le Bris, N., Binet, F., 2017. Organic amendment
643 practices as possible drivers of biogenic Volatile Organic Compounds emitted by soils in
644 agrosystems. *Agric. Ecosyst. Environ.* 250, 25–36.

645 Provenzano, M.R., Iannuzzi, G., Fabbri, C., Senesi, N., 2011. Qualitative characterization and
646 differentiation of digestates from different biowastes using FTIR and fluorescence
647 spectroscopies. *J. Environ. Prot. (Irvine, Calif.)* 2, 83–89.

648 Ramirez, K.S., Lauber, C.L., Fierer, N., 2009. Microbial consumption and production of volatile organic
649 compounds at the soil-litter interface. *Biogeochemistry* 99, 97–107.

650 Rousk, J., Bååth, E., 2011. Growth of saprotrophic fungi and bacteria in soil. *FEMS Microbiol. Ecol.* 78,
651 17–30.

652 Rumpel, C., Dignac, M.-F., 2006. Gas chromatographic analysis of monosaccharides in a forest soil
653 profile: Analysis by gas chromatography after trifluoroacetic acid hydrolysis and reduction–
654 acetylation. *Soil Biol. Biochem.* 38, 1478–1481.

655 Sapp, M., Harrison, M., Hany, U., Charlton, A., Thwaites, R., 2015. Comparing the effect of digestate
656 and chemical fertiliser on soil bacteria. *Appl. Soil Ecol.* 86, 1–9.

657 Seewald, M.S.A., Singer, W., Knapp, B.A., Franke-Whittle, I.H., Hansel, A., Insam, H., 2010. Substrate-
658 induced volatile organic compound emissions from compost-amended soils. *Biol. Fertil. Soils* 46,
659 371–382.

660 Singla, A., Dubey, S.K., Singh, A., Inubushi, K., 2014. Effect of biogas digested slurry-based biochar on
661 methane flux and methanogenic archaeal diversity in paddy soil. *Agric. Ecosyst. Environ.* 197,
662 278–287.

663 Smet, E., Van Langenhove, H., De Bo, I., 1999. The emission of volatile compounds during the aerobic
664 and the combined anaerobic/aerobic composting of biowaste. *Atmos. Environ.* 33, 1295–1303.

665 Spoelstra, S.F., 1978. Degradation of tyrosine in anaerobically stored piggery wastes and in pig feces.
666 *Appl. Environ. Microbiol.* 36, 631–8.

667 Stahl, P.D., Parkin, T.B., 1996. Microbial production of volatile organic compounds in soil microcosms.
668 *Soil Sci. Soc. Am. J.* 60, 821–828.

669 Stotzky, G., Schenck, S., Papavizas, G.C., 1976. Volatile organic compounds and microorganisms. *Crit.*
670 *Rev. Microbiol.* 4, 333–382.

671 Strickland, M.S., Rousk, J., 2010. Considering fungal:bacterial dominance in soils – Methods, controls,
672 and ecosystem implications. *Soil Biol. Biochem.* 42, 1385–1395.

673 Sun, L., Pope, P.B., Eijsink, V.G.H., Schnürer, A., 2015. Characterization of microbial community
674 structure during continuous anaerobic digestion of straw and cow manure. *Microb. Biotechnol.*
675 8, 815–27.

676 Sutton, A.L., Kephart, K.B., Verstegen, M.W.A., Canh, T.T., Hobbs, P.J., 1999. Potential for reduction
677 of odorous compounds in swine manure through diet modification. *J. Anim. Sci.* 77, 430–439.

678 Tambone, F., Genevini, P., D’Imporzano, G., Adani, F., 2009. Assessing amendment properties of
679 digestate by studying the organic matter composition and the degree of biological stability
680 during the anaerobic digestion of the organic fraction of MSW. *Bioresour. Technol.* 100, 3140–
681 3142.

682 Trabue, S., Scoggin, K., Li, H., Burns, R., Xin, H., Hatfield, J., 2010. Speciation of volatile organic
683 compounds from poultry production. *Atmos. Environ.* 44, 3538–3546.

684 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil microbial
685 biomass C. *Soil Biol. Biochem.* 19, 703–707.

686 Veres, P.R., Behrendt, T., Klaphor, A., Meixner, F.X., Williams, J., 2014. Volatile Organic Compound
687 emissions from soil: using Proton-Transfer-Reaction Time-of-Flight Mass Spectrometry (PTR-
688 TOF-MS) for the real time observation of microbial processes. *Biogeosciences Discuss.* 11,

689 12009–12038.

690 Walsh, J.J., Rousk, J., Edwards-Jones, G., Jones, D.L., Williams, A.P., 2012. Fungal and bacterial growth
691 following the application of slurry and anaerobic digestate of livestock manure to temperate
692 pasture soils. *Biol. Fertil. Soils* 48, 889–897.

693 Weiland, P., 2010. Biogas production: current state and perspectives. *Appl. Microbiol. Biotechnol.* 85,
694 849–60.

695 White, T.J., Bruns, T.D., Lee, S.B., Taylor, J.W., 1990. Amplification and direct sequencing of fungal
696 ribosomal RNA genes for phylogenetics, in: Innis, M.A., Gelfand, D.H., Sninsky, J.J., White, T.J.
697 (Eds.), *PCR Protocols - a Guide to Methods and Applications*. Academic Press, San Diego, CA, pp.
698 315–322.

699 Winding, A., Hund-Rinke, K., Rutgers, M., 2005. The use of microorganisms in ecological soil
700 classification and assessment concepts. *Ecotoxicol. Environ. Saf.* 62, 230–248.

701 Woodbury, B.L., Miller, D.N., Eigenberg, R.A., Nienaber, J.A., 2006. An inexpensive laboratory and
702 field chamber for manure volatile gas flux analysis. *Trans. ASABE* 49, 767–772.

703 Yasuhara, A., Fuwa, K., Jimbu, M., 1984. Identification of odorous compounds in fresh and rotten
704 swine manure. *Agric. Biol. Chem.* 48, 3001–3010.

705

706 Table 1

707 Description of primer sets and hybridization temperatures used for RT-PCR

708

Target group	primer	sequence	hybridization temperature	References
Bacteria	63F	5'-CAGGCCTAACACATGCAAGTC-3'	58°C	Marchesi et al., 1998; Osborn et al., 2000
	1389R	5'-ACGGGCGGTGTGTACAAG-3'		Marchesi et al., 1998; Osborn et al., 2000
Archaea	Ar109F	5'-ACKGCTCAGTAACACGT-3'	56°C	Lueders and Friedrich, 2000
	Ar912R	5'-CTCCCCGCCAATTCCTTA-3'		Lueders and Friedrich, 2000
Fungi	ITS1F	5'-CTTGGTCATTAGAGGAAGTAA-3'	55°C	White et al. 1990; Gardes and Bruns, 1993
	ITS4	5'-TCCTCCGCTTATTGATATGC-3'		White et al. 1990; Gardes and Bruns, 1993

709 Table 2


710 Concentrations (ppb) of the main VOC emitted by the pig slurry (PS) and digested pig slurry (DPS)
711 themselves (mean of three replicates with standard deviation, asterisks represent significant
712 differences between slurries, Kruskal Wallis test).

	PS	DPS
Methanol (m33)	33.6 ± 2.7	5.4 ± 1.4
Acetone (m59)	399.7 ± 176.2	88.9 ± 59.2
DMS (m63)*	19.1 ± 10.6	6.5 ± 1.9
2-Pentanone (m87)	12.8 ± 6.3	16.0 ± 9.7
Phenol (m95)	4.8 ± 3.7	2.6 ± 0.5

713

714 Fig. 1


715 Soil dissolved organic carbon concentration (DOC per mL of slurry or g of dry soil - values on top of
 716 the circles, mean of three replicates with standard deviation) and water extractable organic matter
 717 (WEOM) composition in fatty acids, carbohydrates, lignin, phenolic compounds and methylindole
 718 (proportion of analyzed compounds) in the < 0.7 μm fractions of the pig slurry and digestate pig
 719 slurry and in the top 10 cm soil of the microcosms that received either water (control soil), pig slurry
 720 (soil + PS) or digested pig slurry (soil + DPS) just after (0), 30 and 60 days following the application of
 721 slurries.



722

723 Fig. 2

724 Principal component analysis of T-RFs of bacterial (A), archaeal (B) and fungal (C) communities in the
725 pig slurry (PS) and digested pig slurry (DPS) in blue, in the control soils in black (CS0, CS30, CS60), the
726 soils with pig slurry in red (SPS0, SPS30, SPS60) and with digested pig slurry in green (SDPS0, SDPS30,
727 SDPS60) at the initial time and after 30 and 60 days of incubation, respectively.




728

729

730 Fig. 3


731 C/V lignin (A) and deoxy C6/C5 (B) ratios in the top 10 cm of soil in the microcosms that received
732 either water (control soil CS), pig slurry (SPS) or digested pig slurry (SDPS) just after (0), 30 and 60
733 days following application of the slurries. Error bars represent the standard deviation of mean values
734 (three replicates). Different letters correspond to significant differences between samples (Kruskal
735 wallis post hoc Dunn's tests).

736



737 Fig. 4


738 Microbial carbon biomass ($\mu\text{g C g}^{-1}$ dry soil) in the top 10 cm soil of the microcosms that received either
739 water (control soil), pig slurry (Soil + PS) or digested pig slurry (Soil + DPS) just after (0), 30 and 60
740 days following application of the slurries. Error bars represent the standard deviation of mean values
741 (three replicates). Different letters correspond to significant differences between samples (Kruskal
742 wallis post hoc Dunn's tests).



743


744 Fig. 5

745 Microbial richness and its proportion within the active bacterial, archaeal and fungal taxa in the
746 control soil (in blue) and in soils that received pig slurry (in orange) or digested pig slurry (in green)
747 just after (0 day) and 30 and 60 days after application of the slurries. Values in the ternary plots
748 correspond to the total species richness (mean of three replicates \pm standard deviation).



750 Fig. 6

751 Cumulative C-CO₂ emissions from the microcosms with control soil (CS), with soils that received pig
752 slurry (SPS) and digested pig slurry (SDPS) just after (0 day) and 30 and 60 days after application of
753 the slurries. Error bars indicate the standard deviation (n = 3)



754
755

756 Fig. 7

757 Net C-VOCs fluxes of the main VOCs detected in soil with pig slurry (Soil + PS) and digested pig slurry
758 (Soil + DPS) compared to the control microcosms soil just after (T0) and up to 60 days after
759 application of the slurries.

Accepted Manuscript

