



HAL
open science

Silica Exposure and Scleroderma: More Bridges and Collaboration Between Disciplines Are Needed

Alain Lescoat, Catherine Cavalin, Alice Ballerie, Valérie Lecqueur, Lucile Sesé, Claire Cazalets, Mathieu Lederlin, Guillaume Coiffier, Nicolas Belhomme, Christophe Paris, et al.

► To cite this version:

Alain Lescoat, Catherine Cavalin, Alice Ballerie, Valérie Lecqueur, Lucile Sesé, et al.. Silica Exposure and Scleroderma: More Bridges and Collaboration Between Disciplines Are Needed. American Journal of Respiratory and Critical Care Medicine, 2020, 201 (7), pp.880-882. 10.1164/rccm.201911-2218LE . hal-02440641

HAL Id: hal-02440641

<https://univ-rennes.hal.science/hal-02440641>

Submitted on 13 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Silica Exposure and Scleroderma:

More Bridges and Collaboration Between Disciplines Are Needed

Alain Lescoat MD^{1,2}, Catherine Cavalin PhD^{3,4,5}, Alice Ballerie MD^{1,2}, Valérie Lecureur PhD²,
Lucile Sesé MD^{6,7,8}, Claire Cazalets MD¹, Mathieu Lederlin MD PhD^{9,10,11}, Guillaume Coiffier¹²,
¹³ Nicolas Belhomme MD¹, Christophe Paris MD PhD^{14,2}, Ronan Garlantézec MD PhD^{15,2},
Stéphane Jouneau MD PhD¹⁶, Patrick Jégo MD PhD^{1,2}

1-Department of Internal Medicine and Clinical Immunology, CHU Rennes, University of Rennes 1, Rennes, France.

2-University of Rennes, CHU Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085, F-35000, Rennes, France.

3-Institut de Recherche Interdisciplinaire en Sciences Sociales (IRISSO), UMR CNRS-INRA 7170-1427, Université Paris-Dauphine, Paris, France

4-Laboratoire interdisciplinaire d'évaluation des politiques publiques (LIEPP) de Sciences Po, Paris, France

5-Centre d'études de l'emploi et du travail (CEET, CNAM), Noisy-le-Grand, France

6-EPAR, Université Pierre et Marie, Paris, France

7-Service de Pneumologie, AP-HP, Hôpital Avicenne, Bobigny, France

8-Université Paris, Bobigny, France

9-INSERM, U1099, F-35000, Rennes, France.

10-LTISI, Université de Rennes 1, Bât. 22, Campus de Beaulieu, F-35000, Rennes, France.

11-CHU Rennes, Department of Radiology, F-35000, Rennes, France.

12- Department of Rheumatology, CHU Rennes, Univ Rennes, France

13- Inserm U 1241, Univ Rennes, France

14-Consultations de pathologies professionnelles et environnementales, CHU Rennes, F-35000, Rennes, France.

15- Département de Santé Publique, CHRU Rennes, Rennes, France

16- Department of Respiratory Diseases, Rennes University Hospital, 35000, Rennes, France

Conflict of Interest Disclosures: Alain Lescoat, Catherine Cavalin, Alice Ballerie, Valérie Lecureur, Lucile Sesé, Claire Cazalets, Mathieu Lederlin, Guillaume Coiffier, Nicolas Belhomme, Christophe Paris, Ronan Garlantézec, Stéphane Jouneau, Patrick Jégo have nothing to disclose.

Corresponding author: Alain Lescoat

Department of Internal Medicine & Clinical Immunology, CHU Rennes, South Hospital, 16 bvd de Bulgarie - 35203 RENNES Cedex 2 - BP 90347

E-Mail : alain.lescoat@chu-rennes.fr

Phone: +33 2 99 26 71 28 fax : +33 2 99 26 71 98

Dear editor,

We have read with great interest Turner et al.'s correspondence concerning Connective Tissue Diseases (CTD) and silica exposure in artificial stone workers (1). We totally support their call for a new awareness on silica hazards and their broadening to CTDs. Indeed, silica hazards are too often, and almost systematically, narrowed to silicosis (2).

Although LD Erasmus was historically the first to link silica exposure with the occurrence of systemic sclerosis (SSc or scleroderma), B Bramwell, a Scottish physician, had described an outbreak of scleroderma among stonemasons fifty years before (3). Despite this ancient association, SSc is still largely considered today "of unknown cause". The Pasteurian paradigm (one cause, one disease) may have participated to an over-simplified vision of causality in diseases, leading to a dichotomous vision of aetiologies: either an obvious and single cause for some diseases (especially communicable diseases) or "complex diseases" with too many causal factors to be properly captured. Nonetheless, recent insights in the pathogenesis of SSc, including cancer-associated SSc, remind us that the search for a cause may not be totally in vain. Indeed, when silica exposure is properly explored - *i.e.* prospectively assessed through dedicated occupational questionnaires and/or evaluation by experts in occupational medicine and toxicology - this exposure appears to be strikingly frequent in male patients with SSc: at least half of them having occupational silica exposure in recent European studies (4). Therefore, crystalline silica exposure may be a decisive cause or trigger, and this would be especially relevant in males, more often engaged in occupations involving silica. Interestingly, recent insights in the pathogenesis of silica-induced autoimmunity in mouse models have highlighted that the production of autoantibodies after silica exposure was significantly higher in males in comparison with female littermates (5). In humans, sex/gender in SSc may be especially crucial, as SSc is frequently more severe in men.

The current outbreak of silicosis and CTDs after exposure to high silica content artificial stone dusts both demonstrates the specificity of the association of crystalline silica with

autoimmunity in comparison with other mineral dusts, and reminds us that silica exposure does not only concern mining industry but covers a wider range of sectors. In this regard, the call for systematic screening of patients for CTDs after silica exposure appears central, both in patients with and without signs of silicosis and/or pulmonary involvement. Concerning SSc, the systematic screening strategy proposed for a “very early diagnosis of SSc” (VEDOSS) (6) may be especially relevant for these silica-exposed patients. This “VEDOSS” strategy does not only include testing for antinuclear antibodies but also emphasises the central role of capillaroscopic findings and clinical detection of “puffy fingers”, a manifestation of SSc that may precede the occurrence of sclerodactyly. Such a screening would therefore imply 1) a close collaboration of clinicians who have been trained to detect puffy fingers or other SSc early signs and who are familiar with capillaroscopic examinations with 2) specialists of occupational diseases and silica-associated respiratory disorders likely to detect the situations of silica exposure.

Beyond preventability and diagnosis, the lack of a proper understanding of the pathogenesis of silica-associated autoimmunity may also result from missing bridges between disciplines. From a toxicant viewpoint, this question of silica has been almost exclusively studied through fibrotic pulmonary diseases or cancer or, when addressing autoimmunity, has been based on mouse models of systemic lupus, whereas SSc has been considered for long as the CTD most frequently associated with silica exposure. The alarming recent outbreak of silica-associated CTDs may provide an opportunity to fill this gap and offer a timely lever to collectively better understand the pathogenesis of SSc.

References

- 1/ Turner MT, Samuel SR, Silverstone EJ, Yates DH. Silica Exposure and Connective Tissue Disease: An Under-Recognised Association in Three Australian Artificial Stone Workers. [letter] *Am J Respir Crit Care Med* [online ahead of print] 09 October 2019; <https://www.atsjournals.org/doi/abs/10.1164/rccm.201905-1057LE>.
- 2/ The Lancet Respiratory Medicine. The world is failing on silicosis. *Lancet Respir Med*. 2019 Apr;7(4):283.
- 3/ Bramwell B. Diffuse sclerodermia: its frequency, its occurrence in stone-masons, its treatment by fibrinolysin, elevation of temperature do to fibrinolysin injections. *Edinb Med J* 1914; **12**: 387–401.
- 4/ De Decker E, Vanthuyne M, Blockmans D, Houssiau F, Lenaerts J, Westhovens R, Nemery B, De Langhe E. High prevalence of occupational exposure to solvents or silica in male systemic sclerosis patients: a Belgian cohort analysis. *Clin Rheumatol*. 2018 Jul;37(7):1977-1982.
- 5/ Mayeux JM, Escalante GM, Christy JM, Pawar RD, Kono DH, Pollard KM. Silicosis and Silica-Induced Autoimmunity in the Diversity Outbred Mouse. *Front Immunol*. 2018 Apr 26;9:874.
- 6/ Avouac J, Fransen J, Walker UA, Riccieri V, Smith V, Muller C, Miniati I, Turner IH, Randone SB, Cutolo M, Allanore Y, Distler O, Valentini G, Czirjak L, Müller-Ladner U, Furst DE, Tyndall A, Matucci-Cerinic M; EUSTAR Group. Preliminary criteria for the very early diagnosis of systemic sclerosis: results of a Delphi Consensus Study from EULAR Scleroderma Trials and Research Group. *Ann Rheum Dis*. 2011 Mar;70(3):476-81.