

Incidence and risk factors for acquired colonization and infection due to extended-spectrum beta-lactamase-producing Gram-negative bacilli a retrospective analysis in three ICUs with low multidrug resistance rate

Nicolas Massart, Christophe Camus, François Benezit, Mikael Moriconi, Pierre Fillatre, Yves Le Tulzo

▶ To cite this version:

Nicolas Massart, Christophe Camus, François Benezit, Mikael Moriconi, Pierre Fillatre, et al.. Incidence and risk factors for acquired colonization and infection due to extended-spectrum betalactamase-producing Gram-negative bacilli a retrospective analysis in three ICUs with low multidrug resistance rate. European Journal of Clinical Microbiology and Infectious Diseases, 2020, 39 (5), pp.889-895. 10.1007/s10096-019-03800-y . hal-02440783

HAL Id: hal-02440783 https://univ-rennes.hal.science/hal-02440783

Submitted on 31 Mar 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Incidence and risk factors for acquired colonization and infection due to extended spectrum beta-lactamase-producing Gram-negative bacilli: a retrospective analysis in three ICUs with low multidrug resistance rate.

Running title: Risk factor of ESBL-GNB colonization and infection.

Nicolas MASSART, M.D.^{1,2}, Christophe CAMUS, M.D., P.H.D.^{1,2,3}, François BENEZIT, M.D.^{1, 2,3}, Mikael MORICONI, M.D.⁴, Pierre FILLATRE, M.D.⁵, Yves LE TULZO, M.D., P.H.D.^{1, 2, 3}.

¹ Service de maladie infectieuse et de réanimation médicale CHU de rennes, 2, rue Henri le Guilloux 35000 rennes, France.

²Université Rennes1, Faculté de Médecine, Biosit, F-35043 Rennes, France

³Inserm-CIC-1414, Faculté de Médecine, Université Rennes I, IFR 140, F-35033 Rennes,

France

⁴Service de Réanimation, CH de QUIMPER, 14bis Avenue Yves Thépot, 29107 Quimper.
⁵Service de Réanimation, CH de St BRIEUC, 10, rue Henry le Guilloux 22000 Saint-Brieuc, France.

Corresponding author : Nicolas MASSART, Service de maladie infectieuse et de réanimation médicale CHU de rennes, 2, rue Henri le Guilloux 35000 rennes, France, <u>nicolasmassart@hotmail.fr</u>,

Telephone : +33 6 20 17 59 89

ABSTRACT

Purpose:

To assess risk factors for the acquisition of extended-spectrum β -lactamase-producing Gramnegative bacilli (ESBL-GNB) colonization and infection (AI) in ICUs with low ESBL-GNB prevalence rate.

Methods:

We conducted a retrospective observational study in three ICUs in Bretagne, France. All patients admitted from January 2016 to September 2017 with a length of stay of two days or more were included. Universal screening for ESBL-GNB colonization was performed in all participating ICUs.

Results:

Of the 3250 included patients, 131 (4.0%) were colonized at admission, 59 acquired colonization while hospitalized (1.9%; 95% CI [1.5%–2.5%]) and 15 (0.5%; 95% CI [0.3%–0.8%]) acquired ESBL-GNB infections. In case of infection, the specificity and the negative predictive values of preexistent colonization for the ESBL-GNB etiology were 93.2% [91.5%–95.1%] and 95.2% [93.5%–97.1%], respectively. Colonization was the main risk factor for ESBL-GNB AI (OR=9.61; 95% CI [2.86–32.29]; p< 0.001). Antimicrobial susceptibility of non-ESBL GNB isolates responsible for AI was similar for any non-carbapenem β -lactam (95%) and imipenem (94%).

Conclusion:

ESBL-GNB AIs were rare in ICUs with low ESBL-GNB prevalence rate. Prior colonization was the main risk factor for subsequent infection. Empirical carbapenem therapy could be avoided in non ESBL-GNB colonized patients with suspected AI.

2

Keyword:

Extended-spectrum beta-lactamase-producing Enterobacteriaceae, Gram-negative bacteria,

Decontamination, Healthcare-Associated Pneumonia, Bacteremia.

INTRODUCTION

Critically ill patients in the intensive care unit (ICU) have a high risk of acquired infections (AIs) (1,2), which increase morbidity and, to some extent, mortality (3). Pneumonia and bloodstream infection (BSI) are the most common infections with a reported incidence rate of 15 per 1,000 ventilator-days (2) and 9.5 per 1,000 patient-days (1), respectively. Among multidrug-resistant organisms (MDROs) (4-7), extended-spectrum beta-lactamaseproducing Gram-negative bacilli (ESBL-GNB) have an increasing prevalence reaching critical point in some ICUs (7,8), leading to the unrestricted use of carbapenems in order to avoid inappropriate antimicrobial therapy (8,9,10).

The challenge of ESBL-GNB has been overemphasized (11) as compared with real life in specific locations. In contrast, risk factors for ESBL-GNB acquired colonization and infection have been unfrequently assessed in ICUs with lower prevalence rates (12-15). The identification of risk factors could help clinicians to target the use of broad-spectrum antibiotics in this setting. The aim of our study was to assess the incidence and risk factors for ESBL-GNB acquired colonization and infection in ICUs with low prevalence of ESBL-GNB.

MATERIALS AND METHODS

Settings and patients

We conducted a retrospective observational study in three French ICUs. The largest ICU (ICU 1) was a medical unit with 22 acute-care beds in a tertiary care university hospital. Unlike the two other centers, this ICU used multiple-site decontamination in intubated patients. It consists of the administration of topical antibiotics (tobramycin, colistin and amphotericin B) four times daily in the oro-pharynx and the gastric tube, chlorhexidine body washing and nasal mupirocin, in all intubated patients for the period of intubation (16). The other two units were polyvalent ICUs with 14 beds (ICU 2) and 12 beds (ICU3). Every ICU

followed the recommendations of the French Society for Hospital Hygiene regarding the prevention and treatment of AI (available at <u>https://sf2h.net</u>). Each center had a nosocomial infection committee for the prevention and prospective census of AI.

We screened all admitted patients from January 1st 2016 to September 10th 2017. Patients with a length of stay of two days or more were included. Because rectal screening for ESBL-GNB colonization and report of AIs are routine care, the institutional ethics committee approved the study and the requirement for written informed consent was waived.

Data collection and definitions

ESBL-GNB carriage was screened on a rectal swab at admission, every week afterwards and at ICU discharge. Samples were then inoculated onto antibiotic selective agar. The synergy between third generation cephalosporin and clavulanate was confirmed using a double-disk synergy test.

Colonization at admission was defined as a positive first rectal swab for ESBL-GNB performed within 48 hours of admission. Acquisition of colonization was defined as a subsequent positive rectal swab after a negative first swab. Patients with ESBL-GNB AI were considered colonized prior to infection if their screening rectal swab was positive either before or on the day of the microbial diagnosis of infection.

Infection was considered acquired in the ICU when it was diagnosed 48 hours after admission and was not incubating on admission. The diagnosis of pneumonia included both ventilator-associated pneumonia and ICU-acquired pneumonia that developed in nonventilated patients and was made based on clinical signs (fever), radiological findings (new infiltrate on chest-X-ray or CT scan), and a positive microbiological culture of an endotracheal aspirate ($\geq 10^6$ colony-forming units/mL) or a broncho-alveolar lavage ($\geq 10^4$ colony-forming units/mL). BSI was diagnosed following one or more positive blood culture as defined by the CDC (17).

For the purpose of this study, we estimated the number of days with colonization and the number of days without colonization before the occurrence of ESBL-GNB infection according to the method reported by Frencken et al. (13) (supplementary Figure 1).

Other collected data were age, sex, reason for admission (medical/surgical), simplified acute physiology score II (SAPS II) at admission, length of stay in ICU, intubation for more than two days, admission in ICU 1 versus ICU 2 and 3 and outcome.

Primary and secondary objectives

The primary objective of the study was to assess the incidence and risk factors of ESBL-GNB acquired colonization and infection. Spectrum of AIs and antimicrobial susceptibility were secondary objectives.

Statistical analysis

Statistical analysis was performed with the statistical software R 3.4.3. Incidence rate and prevalence were expressed with the 95 percent confidence interval (95% CI), categorical variables were expressed as number (percentage) and continuous variables as median and interquartile range (IQR). When appropriate, the chi-square test and the Fisher's exact test were used to compare categorical variables. The Man-Whitney U test and the Kruskal-Wallis test were used for continuous variables when applicable. Multivariate logistic regression analysis was used to study risk factors for ESBL-GNB acquired colonization and acquired infection. Variables with a p-value <0.2 in the univariate analysis or those clinically relevant irrespective of the p-value were included for the multivariate analysis. All tests were twosided, and a P-value less than 0.05 was considered statistically significant.

RESULTS

Study population

Among 3,861 admitted patients, 3,250 with an ICU length stay of two days or more were all included in the study (Figure 1). SAPS II was 40 [28–56] and age was 62.1 years [51.0–71.0]. One hundred and thirty-one patients (4.0%) were colonized with ESBL-GNB at admission, varying from 2.0% to 4.8% depending on the ICU (Table 1, supplementary Table 1). One thousand five hundred and seven patients (46.4%) were intubated for more than 48 hours. The average length of stay in ICU was 4 days [3–9] and overall ICU mortality was 19.8%.

ESBL-GNB acquired colonization

Among the 3,119 patients not colonized at admission, 59 (1.9%; 95% CI [1.5–2.5]) acquired ESBL-GNB colonization during their ICU stay with a median length of stay of 7 days [5–14] before colonization acquisition. These patients had a higher SAPS II score than those without acquired ESBL-GNB colonization (48 [34–62] vs 39 [28–55], p=0.01). Intubation for 48 hours or more (OR = 6.02 [2.86–12.68], p< 0.001) and admission in ICU 1 (OR = 0.39 [0.22–0.70], p=0.002) were independent risk factors for ESBL-GNB acquired colonization (Table 2).

ESBL-GNB acquired infection

Fifteen infections occurred during a total of 25,857 patient-days at risk, with calculated incidence rates of 0.46 per 100 admissions [0.26–0.76] and 0.58 per 1,000 patient-days [0.49–0.68]. Six cases (40%) had prior ESBL-GNB colonization and 9 had not (Table 3), yielding incidence density rates of 4.73 per 1,000 days with colonization and 0.37 per 1,000

days without colonization respectively (incidence rate ratio 12.86 [4.08–39.31]; p < 0.001). There were 8 pneumonias (0.3% [0.1%–0.5%]) and 7 BSIs (0.2% [0.1%–0.4%]) (Table 3). The delay to the onset was similar between cases with and cases without prior colonization (p=0.82). Mortality in the ICU was similar between patients with ESBL-GNB AI and those with other AIs (33.3% versus 32.8%, p=0.92).

One hundred and ninety-two patients (5.9%) acquired 204 infections (pneumonia: 122; BSI: 82) not due to ESBL-GNB. Among the 207 patients who acquired any type of infection, 19 had previous ESBL-GNB colonization. In these patients, 6 acquired ESBL-GNB infection (31.6%), whereas 9 over 188 non- colonized patients (4.8%) acquired ESBL-GNB infection (relative risk 6.60 [2.23–17.57]). The positive and negative predictive values of ESBL-GNB colonization for ESBL-GNB etiology of AI were 31.6% [14.5%–50.5%] and 95.2% [93.5%– 97.1%], respectively, whereas sensitivity and specificity were 40.0%, [18.4%– 64.0%] and 93.2% [91.5%–95.1%]. By logistic regression analysis, simultaneous or pre-existent ESBL-GNB colonization was the main risk factor for ESBL-GNB AI (OR = 9.61 [2.86–32.29], p<0.001) (Table 4).

Prevalence of ESBL-GNB

Taking into account 131 patients who were colonized at admission, 59 patients who acquired colonization in the ICU and 7 non-colonized patients who acquired ESBL-GNB infection, the overall prevalence rate of ESBL-GNB in was 6.1% [5.2%–6.9%] and varied from 4% to 7.5% between ICUs (Supplementary Table 1).

Micro-organisms identified in acquired infections and antimicrobial susceptibility

The microorganisms and antimicrobial susceptibility are shown in Table 5. In non-ESBL microorganisms, GNB (55%) were the most frequent (*Enterobacteriaceae*, 32.4%; *Pseudomonas aeruginosa*, 17.8%). Aerobic GNB were mostly susceptible to all tested antimicrobials, without any major difference between 3rd-generation cephalosporins and imipenem.

All ESBL-GNB responsible for AIs were susceptible to amikacin and imipenem. In 13 of the 15 patients (86.7%) with ESBL-GNB AI, antimicrobial therapy was considered to be appropriate. Among 129 colonization isolates tested, all were susceptible to amikacin and only 3 (two *Pseudomonas aeruginosa* and one *Klebsiella pneumoniae*) were resistant to imipenem (supplementary Table 2).

DISCUSSION

The ESBL-GNB prevalence rate has ranged from 5.7% to 48.0% (7, 14,15,18-22). In a recent meta-analysis, Detsis et al. (18) reported the rate of ESBL-GNB acquired colonization in 13 studies and the ESBL-GNB prevalence rate could be calculated in 10. This review suggests that a low prevalence rate is below 10 % (15, 20, 21), intermediate rate is 10%-20% and high rate is above 20% (18-20).

The main finding of this study was that very few patients (0.46%) acquired ESBL-GNB GNB AIs in ICUs with low prevalence rate of resistance. Prior or simultaneous ESBL-GNB colonization was a strong risk factor for ESBL-GNB AI. This is consistent with other studies with either low (15) or high (19) ESBL-GNB prevalence rate. In patients with suspected or confirmed infection, the relative risk for the ESBL-GNB etiology after prior colonization with ESBL-GNB has been diversely assessed depending on local epidemiology. Houard et al. (19) reported a relative risk of 2.6 [1.2–5.6] in an ICU with ESBL-GNB prevalence rate of 48%. In contrast, Bruyère et al. (15) found a much higher risk (77.5 [23.7–253.4]) in ICUs with a prevalence rate of 7.3%. In our study, the relative risk was not as high (6.6 [2.2–17.6]). Differences were were due to the lower rate of ESBL-GNB AI in non-colonized patients

reported by Bruyère (0.6% versus 4.8% in our study), also explaining the higher negative predictive value of prior colonization in this study (99.5% versus 95.2% in our study).

Another important finding in our study was that the majority of ESBL-GNB AIs occurred in non-colonized patients, suggesting an exogenous cause of AI in some cases. ESBL-GNB infection in non-colonized patients has been reported previously with various rates. Jalalzaï et al. observed exogenous infection in 2 of 6 ESBL-GNB AIs (14), Bruyère et al. in 3 among 20 ESBL-GNB AIs (15) while Prevel et al. reported no infection in noncolonized patients during a 6-months study (22). The exogenous source of multi-drug resistant GNB has been examined in the study of Harris et al. (23). From 1806 previously non-colonized patients, the authors reported 27 patients with ESBL-producing Klebsiella pneumonia acquisition among whom there were 14 patients (52%) with an isolate similar by molecular typing to an isolate from another patient with an overlapping length of stay in the hospital. The authors suggest that patient-to-patient transmission is an important and underappreciated cause of ESBL-producing Klebsiella pneumonia infection and colonization (23). In contrast, Tschudin et al. reported that only 1.5% of acquired colonization was due to patient-to-patient transmission (24). An environmental contamination as a reservoir for ESBL-GNB colonization has recently been emphasized for Klebsiella pneumoniae (25). We observed that admission to ICU 1 was protective for ESBL-GNB acquisition. A major difference between ICU 1 and ICUs 2 and 3 was the use of MSD in intubated patients. The benefit of selective decontamination regimens on acquired colonization with multidrugresistant Gram-negative bacilli has been shown in France (26) as in other European countries (27). However, selective digestive decontamination does not prevent patient-to-patient crosstransmission.

Recent recommendations support a large panel of antibiotics from ceftriaxone to carbapenems in patients with ICU-acquired pneumonia (28). However, antimicrobial use

should be based on local antimicrobial susceptibility surveillance data. Taking into account the high negative predictive value of ESBL-GNB colonization for the prediction of ESBL-GNB AI and high $3^{rd}/4^{th}$ generation susceptibility rate, we suggest that non-carbapenem β lactams could be safely used in non-colonized patients with AIs in ICUs with low ESBL-GNB prevalence rate. In ESBL-GNB colonized patients, because one third of AIs were caused by ESBL-GNB and all isolates were susceptible to amikacin, we propose that amikacin to be added to a beta-lactam agent.

There were several limitations to our study. First, due to the low number of ESBL-GNB AI, other risk factors may not have been identified. Second, antimicrobial therapy of AIs was not recorded and the impact of the use of broad-spectrum antimicrobials on the incidence of ESBL-GNB AIs could not be assessed. Third, due to the intermittent character of ESBL-GNB screening, transient colonization could have been missed.

In conclusion, in ICUs with low prevalence rate of ESBL-GNB, few ESBL-GNB AIs were reported and prior or simultaneous ESBL-GNB colonization was a major risk factor for infection. In order to treat a suspected pneumonia or BSI acquired in the ICU, our data suggests that carbapenems could be avoided in non-colonized patients and amikacin combination could be used in patients colonized with ESBL-GNB.

Acknowledgment : The authors thanks Dave MAVROVIC for his help in the English writing.

Compliance with Ethical Standards:

Funding: None

Conflict of interest: The authors declare that they have no conflict of interest for this study.

Ethical approval: The local ethic committee ("comité d'éthique du CHU de rennes") approved the protocol (n°18.56).

Informed consent: This observational retrospective study respects the French reference method MR003. Patients were informed of the study and non-opposition was needed. By the French law, there is no need for patient's written approval.

References:

1 – Prowle JR, Echeverri JE, Ligabo EV, Sherry N, Taori GC, Crozier TM, et al. Acquired bloodstream infection in the intensive care unit: incidence and attributable mortality. Crit Care. 2011;15(2):R100.

2 - Rosenthal VD, Bijie H, Maki DG, Mehta Y, Apisarnthanarak A, Medeiros EA, et al. International Nosocomial Infection Control Consortium (INICC) report, data summary of 36 countries, for 2004-2009. Am J Infect Control. 2012;40(5):396-407.

3 - Melsen WG, Rovers MM, Groenwold RH, Bergmans DC, Camus C, Bauer TT, et al. Attributable mortality of ventilator-associated pneumonia: a meta-analysis of individual patient data from randomised prevention studies. Lancet Infect Dis. 2013;13(8):665-71.

4 – Barbier F, Pommier C, Essaied W, Garrouste-Orgeas M, Schwebel C, Ruckly S, et al.
Colonization and infection with extended-spectrum β-lactamase-producing
Enterobacteriaceae in ICU patients: what impact on outcomes and carbapenem exposure? J
Antimicrob Chemother. 2016;71(4):1088-97.

5 - Goulenok T, Ferroni A, Bille E, Lécuyer H, Join-Lambert O, Descamps P, et al. Risk factors for developing ESBL E. coli: can clinicians predict infection in patients with prior colonization? J Hosp Infect. 2013;84(4):294-9.

6 - Tumbarello M, Spanu T, Sanguinetti M, Citton R, Montuori E, Leone F, et al.
Bloodstream Infections Caused by Extended-Spectrum- -Lactamase-Producing Klebsiella
pneumoniae: Risk Factors, Molecular Epidemiology, and Clinical Outcome. Antimicrob
Agents Chemother. 2006;50(2):498-504.

7– Razazi K, Derde LPG, Verachten M, Legrand P, Lesprit P, Brun-Buisson C. Frequency, associated factors and outcome of multi-drug-resistant intensive care unit-acquired pneumonia among patients colonized with extended-spectrum β-lactamase-producing Enterobacteriaceae. Ann Intensive Care 2017;7:61.

8- Bradford PA. Extended-Spectrum β-Lactamases in the 21st Century: Characterization,
Epidemiology, and Detection of This Important Resistance Threat. ClinMicrobiol Rev.
2001;14(4):933

9- Paterson DL, Bonomo RA. Extended-Spectrum -Lactamases: a Clinical Update. Clin Microbiol Rev. 2005;18(4):657-86.

10 - Girometti N, Lewis RE, Giannella M, Ambretti S, Bartoletti M, Tedeschi S, et al.Klebsiella pneumoniae Bloodstream Infection: Epidemiology and Impact of InappropriateEmpirical Therapy. Medicine (Baltimore). 2014;93(17):298-309.

11 - Tabah A, Koulenti D, Laupland K, et al. Characteristics and determinants of outcome of hospital-acquired bloodstream infections in intensive care units: the EUROBACT
International Cohort Study. Intensive Care Med 2012; 38:1930–45

12- Barbier F, Bailly S, Schwebel C, Papazian L, Azoulay É, et al. Infection-related ventilator-associated complications in ICU patients colonised with extended-spectrum β -lactamase-producing Enterobacteriaceae. Intensive Care Med. 2018;44(5):616-26.

13 - Frencken JF, Wittekamp BHJ, Plantinga NL, Spitoni C, van de Groep K, Cremer OL, et
al. Associations Between Enteral Colonization With Gram-Negative Bacteria and Intensive
Care Unit–Acquired Infections and Colonization of the Respiratory Tract. Clin Infect Dis.
2018;66(4):497-503.

14 - Jalalzaï W, Boutrot M, Guinard J, Guigon A, Bret L, Poisson D-M, et al. Cessation of screening for intestinal carriage of extended-spectrum β-lactamase-producing
Enterobacteriaceae in a low-endemicity intensive care unit with universal contact precautions.
Clin Microbiol Infect. 2018; 24(4):429.e7-429.e12.

15 - Bruyère R, Vigneron C, Bador J, Aho S, Toitot A, Quenot J, et al. Significance of prior digestive colonization with extended-Spectrum β -lactamase–producing Enterobacteriaceae in patients with ventilator-associated pneumonia. Crit Care Med. 2016; 44:699–706.

16 - Camus C, Salomon S, Bouchigny C, Gacouin A, Lavoué S, Donnio P-Y, et al. Short-Term Decline in All-Cause Acquired Infections With the Routine Use of a Decontamination Regimen Combining Topical Polymyxin, Tobramycin, and Amphotericin B With Mupirocin and Chlorhexidine in the ICU: A Single-Center Experience*. Crit Care Med. 2014; 42(5):1121-30.

17 - Garner JS, Jarvis WR, Emori TG, Horan TC, Hughes JM. CDC definitions for nosocomial infections, 1988. Am J Infect Control. 1988;16(3):128-140.

18– Detsis M, Karanika S, Mylonakis E. ICU acquisition rate, risk factors, and clinical significance of digestive tract colonization with extended-spectrum beta-lactamase-producing Enterobacteriaceae: a systematic review and meta-analysis. Crit Care Med. 2017; 45:705–714.

19- Houard M, Rouzé A, Ledoux G, Six S, Jaillette E, Poissy J, et al. Relationship between digestive tract colonization and subsequent ventilator-associated pneumonia related to ESBL-producing Enterobacteriaceae. *PLoS One*. 2018;13(8):e0201688.

20 - Martins IS, Pessoa-Silva CL, Nouer SA, Pessoa de Araujo EG, Ferreira ALP, Riley LW, et al. Endemic extended-spectrum beta-lactamase-producing Klebsiella pneumoniae at an intensive care unit: risk factors for colonization and infection. Microb Drug Resist. 2006;12:50–58

21- Troche G, Joly LM, Guibert M, Zazzo JF. Detection and treatment of antibiotic-resistant bacterial carriage in a surgical intensive care unit: a 6-year prospective survey. Infect Control Hosp Epidemiol 2005; 26: 161–5.

22 - Prevel, R., Boyer, A., M'Zali, F. *et al.* Extended spectrum beta-lactamase producing *Enterobacterales* faecal carriage in a medical intensive care unit: low rates of cross-transmission and infection. *Antimicrob Resist Infect Control. 2019;* **8**, 112.

23- Harris A, Perencevich E, Johnson J, Paterson D, Morris J, Strauss S, et al. Patient-topatient transmission is important in extended-spectrum β -lactamase–producing *Klebsiella pneumoniae* acquisition. Clin. Infect. Dis. 2007;45:1347–1350.

24 - Tschudin-Sutter S,, Frei R, Dangel M, Stranden A, Widmer AF, Rate of Transmission of Extended-Spectrum Beta-Lactamase–Producing Enterobacteriaceae Without Contact Isolation, Clin. Infect. Dis. 2015 : 11:1505–1511,

25 - Guet-Revillet H, Le Monnier A, Breton N, Descamps P, Lecuyer H, Alaabouche I, et
al. Environmental contamination with extended-spectrum β-lactamases: is there any
difference between *Escherichia coli* and *Klebsiella* spp? Am J Infect Control. 2012; 40:845–
8.

26 - Camus C, Sauvadet E, Tavenard A, Piau C, Uhel F, Bouju P, et al. Decline of multidrugresistant Gram negative infections with the routine use of a multiple decontamination regimen in ICU. J Infect. sept 2016;73(3):200-9.

27 - Plantinga NL, Wittekamp BHJ, Brun-Buisson C, Bonten MJM, Cooper BS, Coll P, et al. The effects of topical antibiotics on eradication and acquisition of third-generation cephalosporin and carbapenem-resistant Gram-negative bacteria in ICU patients; a post hoc analysis from a multicentre cluster-randomized trial. Clin Microbiol Infect. 2019; Available online 14 August 2019. 28- Torres A, Niederman MS, Chastre J, Ewig S, Fernandez-Vandellos P, Hanberger H, et al. International ERS/ESICM/ESCMID/ALAT guidelines for the management of hospitalacquired pneumonia and ventilator-associated pneumonia: Guidelines for the management of hospital-acquired pneumonia (HAP)/ventilator-associated pneumonia (VAP) of the European Respiratory Society (ERS), European Society of Intensive Care Medicine (ESICM), European Society of Clinical Microbiology and Infectious Diseases (ESCMID) and Asociación Latinoamericana del Tórax (ALAT). Eur Respir J. 2017;50(3):1700582.

Caption Fig.1



Table 1. Baseline characteristics and outcome of study patients.

	All patients	Patients never colonized with Patients colonized with ESBL-GNB at		Patients with acquired ESBL-GNB	Patients with non ESBL-GNB acquired	Patients with ESBL-GNB acquired
	n = 3250	ESBL-GNB	admission	colonization	infection	infection
		n = 3060	n = 131	n = 59	n = 192	n = 15
Age, years [IQR]	62.1 [51.0 - 71.0]	62.0 [50.0 - 70.9]	65.8 [56.0 - 72.0]	66.0 [56.3 – 75.0]	64.8 [55.0 - 71.5]	70.7 [64.5 – 75.5]
Simplified acute physiology score II [IQR]	40 [28 - 56]	39 [28 - 55.0]	46 [3 3–58]	48 [34 – 62]	51 [38 - 64]	61 [46 - 67]
Reason for admission – no. (%)						
Surgical	443 (13.6)	412 (13.5)	17 (13.9)	13 (22.0)	33 (17.2)	2 (13.3)
Medical	2445 (75.2)	2346 (76.6)	105 (80.2)	41 (69.5)	132 (68.8)	13 (86.6)
Not precised	362 (11.1)	302 (9.9)	9 (6.9)	5 (8.5)	27 (14.1)	-
Male – no. (%)	2074 (63.8)	1936 (63.3)	92 (70.2)	45 (76.3)	144 (75.0)	12 (80.0)
ESBL- GNB colonization at admission - no. (%)	131 (4.0)	0	131 (100)	-	3 (1.6)	4 (26.6)
ESBL- GNB acquired colonization - no. (%)	59 (1.8)	-	-	59 (100)	10 (5.2)	4 (26.7) §
Days after admission, days [IQR]*	7 [5-14]	-	-	7 [5-14]	14 [11-25]	10 [8-14]
ESBL- GNB acquired infection - no. (%)	15 (0.5)	7 (0.2)	4 (3.1)	4 (6.8) §	-	15 (100)
Intubation duration > 48 hours, $-$ no. (%)	1507 (46.4)	1395 (45.6)	61 (46.6)	51 (86.4)	163 (84.9)	13 (86.6)
Length of stay in ICU, days [IQR]	4 [3-9]	4 [2-8]	5 [3 – 9]	16 [9 – 33]	20 [9-35]	25 [12 - 41]
Death in ICU- no. (%)	645 (19.8)	596 (19.5)	37 (28.2)	12 (20.3)	63 (32.8)	5 (33.3)

Note. ICU : Intensive Care Unit ; ESBL-GNB : extended-spectrum beta-lactamase-producing Gram-negative bacilli. * In patients with acquired colonization only.

Table 2. Risk factors for acquired colonization with extended-spectrum β-lactamase-producing Gram-negative bacilli.

Variable		Univariate	Multivariate			e
	OR	95% CI	P-value	OR	95% CI	P-value
Age > 65 years	1.33	0.80 - 2.23	0.27			
Simplified Acute Physiology Score II > 40	1.97	1.15 - 3.40	0.014	1.31	0.71 - 2.38	0.38
Male	1.77	0.97 - 3.25	0.06	1.47	0.79 - 2.73	0.22
Surgical reason for admission	1.81	0.96 - 3.40	0.07	1.80	0.91 - 3.31	0.09
Intubation duration > 48 hours	6.62	3.24 - 13.50	< 0.001	4.80	2.26 - 10.24	< 0.001
Admission in ICU 1	0.33	0.19 - 0.58	< 0.001	0.39	0.22 - 0.70	0.002

Note. OR : odds ratio. 95% CI : 95 percent confidence interval.

Table 3. Characteristics of infection due to extended-spectrum β-lactamase-producing Gram-negative bacilli acquired infection.

	Total	Previously colonized	Non-previously colonized		
		patients	patients		
Number	15	6	9		
Delay of onset, days *	9 [2-44]	9 [2-22]	8 [2-44]		
Pneumonia – no. (%)	8 (53.3)	3 (50.0)	5 (55.5)		
Bloodstream infection – no. (%)	7 (46.6)	3 (50.0)	4 (44.4)		
Death in ICU – no. (%)	5 (33.3)	2 (33.3)	3 (33.3)		

Note. ICU: Intensive Care Unit. *Results are expressed with median and [minimal – maximal]

Table 4: Risk factors for extended s	spectrum <i>B</i> -lactamase	-producing Gram	-negative bacilli acq	uired infection in 207	patients with acquired infection.
	speech will p incention	producing oran	Barre caeini aed		

Variable		Univariate		Multivariate			
	OR	95% CI	P-value	OR	95% CI	P-value	
Age > 65 years	2.99	0.92 - 9.72	0.07	3.17	0.92 - 10.87	0.07	
Simplified Acute Physiology Score $II > 40$	1.82	0.49 - 6.68	0.90				
Male	1.35	0.36 - 4.96	0.44				
Surgical reason for admission	0.82	0.17 - 3.85	0.80				
Intubation duration > 48 hours	1.16	0.25 - 5.40	0.85				
ESBL-GNB colonization	9.18	2.83 - 29.77	< 0.001	9.61	2.86 - 32.29	< 0.001	

Note. 95% CI: 95 percent confidence interval; ESBL-GNB : extended spectrum β-lactamase-producing Gram negative bacilli.

Micro-organisms		Antimicrobials: no. susceptible/no. tested (%)						
	Number of	Any non-carbapenem β-	Cefepime	Ceftazidime	Piperacillin-tazobactam	Imipenem	Amikacin	Methicillin
	isolates	lactam	_					
Non-ESBL producing microorganisms								
Enterobacteriaceae	69	52/53 (98.1)	49/50 (98.0)	47/50 (94.0)	44/53 (83.0)	53/55 (96.4)	37/41 (90.2)	-
Pseudomonas aeruginosa	38	35/38 (92.1)	34/38 (89.5)	34/38 (89.5)	35/38 (92.1)	35/38 (92.1)	37/38 (97.4)	-
Acinteobacter species	6	5/6 (83.3)	5/6 (83.3)	5/6 (83.3)	5/6 (83.3)	5/6 (83.3)	5/6 (83.3)	-
Anaerobes	5	4/4 (100)	-	-	4/4 (100)	-	-	-
Staphylococcus aureus	35	-	-	-		-	-	32/35 (91.4)
Coagulase negative Staphylococcus	32	-	-	-	_	-	-	13/28 (46.4)
Streptococcus species	14	-	-	-	-	-	-	-
Enterococcus species*	6	-	-	-	-	-	-	-
Others	8	-	-	0/1	0/1	1/1 (100)	1/2 (50.0)	-
Total	213	96/101 (95.0)	88/94 (93.6)	86/95 (90.5)	88/102 (86.3)	94/100 (94.0)	80/87 (91.9)	45/63 (71.4)
ESBL-GNB					*			
E. coli	8	5/8 (62.5)	-	1/8 (12.5)	5/8 (62.5)	8/8 (100)	8/8 (100)	-
Enterobacter cloacae	4	1/4 (25.0)	-	0/4	1/4 (25.0)	4/4 (100)	4/4 (100)	-
Klebsiella pneumoniae	3	2/3 (66.6)	-	0/3	2/3 (66.6)	3/3 (100)	3/3 (100)	-
Total	15	8/15 (53.3)	-	1/15 (6.7)	8/15 (53.3)	15/15 (100)	15/15(100)	-

Table 5: Microorganisms involved in 219 acquired infections and antimicrobial susceptibility.

Note. ESBL-GNB : extended-spectrum beta-lactamase-producing Gram-negative bacilli. * no glycopeptide resistant enterococcus was isolated.

<u>β1-</u> .se-producing Gran.