



# Epigenetic mechanisms driving tumor supportive microenvironment differentiation and function a role in cancer therapy?

Marvin Sylvestre, Karin Tarte, David Roulois

## ► To cite this version:

Marvin Sylvestre, Karin Tarte, David Roulois. Epigenetic mechanisms driving tumor supportive microenvironment differentiation and function a role in cancer therapy?. *Epigenomics*, 2020, 12 (2), pp.157-169. 10.2217/epi-2019-0165 . hal-02442385

**HAL Id: hal-02442385**

**<https://univ-rennes.hal.science/hal-02442385>**

Submitted on 13 Feb 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **Epigenetic mechanisms driving tumor supportive microenvironment differentiation and function a role in cancer therapy?**

Marvin Sylvestre<sup>1,2</sup>, Karin Tarte<sup>1,3</sup> & David Roulois<sup>1,2,4</sup>

1 - MICMAC - Microenvironment, Cell Differentiation, Immunology and Cancer

2 - Etablissement Français du Sang Bretagne

3 - CHU Pontchaillou [Rennes]

4 - Cancéropôle Grand-Ouest [Bretagne-Centre-Pays de Loire]

## Abstract

The tumor microenvironment (TME) plays a central role in tumor development and drug resistance. Within TME, the stromal cell subset, called cancer-associated fibroblasts (CAFs), is a heterogeneous population originating from poorly characterized precursors. Since CAFs do not acquire somatic mutations, other mechanisms like epigenetic regulation, could be involved in the development of these cells and in the acquisition of tumor supportive phenotypes. Moreover, such epigenetic modulations have been correlated to the emergence of an immunosuppressive microenvironment facilitating tumor evasion. These findings underline the need to deepen our knowledge on epigenetic mechanisms driving TME development and function, and to understand the impact of epigenetic drugs, that could be used in future to target both tumor cells and their TME.

**KEY WORDS:** Epigenetic, Cancer, Microenvironment, Cancer-Associated Fibroblasts

## I) Introduction

### Epigenetic mechanisms

The concept of epigenetics emerged in the middle of the 20<sup>th</sup> century, with Conrad Waddington; who proposed the word, 'epigenetic', to describe biological cell diversity that cannot be attributed to alterations in the deoxyribonucleic acid (DNA) sequence [1].

Epigenetics is now defined as modifications of DNA or factors associated with DNA, carrying an information that can be heritable but without inducing any changes into the DNA sequence.

These chromatin modifications are well described and summarized in many reviews [2-5].

They essentially include:

i) Methylation of DNA in particular; 5-methylcytosine (5mC) enriched CpG islands: This pattern, especially when found at gene promoters, is associated in most cases to inhibition of gene expression. DNA methylation is catalyzed by DNA methyltransferases (DNMT), including DNMT3A/3B that mediates the *de novo* methylation and DNMT1 that mediate maintenance of DNA methylation on the newly synthesized DNA strand during DNA replication [6].

ii) Chromatin modifications, based on the post-translational regulation of histones tails: They can ultimately modulate the compaction of the chromatin and can be associated with gene expression or repression [7]. Among histone tail modifications, acetylation and methylation of histones are probably the most studied histone marks. Histone acetylation is associated with a decompaction of the chromatin and an increased accessibility of gene promoters and enhancers, allowing transcription factor binding and regulation of gene expression [8]. Chromatin acetylation is carried out by histone acetyltransferases (HATs) and deacetylation by histone deacetyltransferases (HDACs). Histone methylations could be associated with both activation (H3K4me3, H3K36me3) or repression (H3K9me3, H3K27me3) of gene expression and is regulated by histone methyltransferases (HMTs or KMTs), adding 1 to 3 methyl groups on histone lysine or arginine. These modifications can be reversed by histones demethylases (HDMs or KDMs) that specifically recognize mono, di, or tri-methyl marks [9].

iii) Non-coding ribonucleic acid (RNA), including micro RNA (miRNA) and long non-coding RNA (lncRNA): miRNAs are small RNA molecules, acting at post-transcriptional level to block translation by degrading messenger RNA (mRNA) [10]. lncRNA regulate gene expression by regulating chromatin structure, mRNA stability, splicing, and post-translational regulation processes [11].

Altogether, these epigenetic mechanisms are involved in many biological processes, such as development, gene imprinting, X inactivation, cell fate decision and cell identity control [9,12-14]. They are also shown to be deregulated in many diseases including cancer. Nonetheless, epigenetic deregulation in cancer is not restricted to cancer cells, but can also be observed in non-malignant cells of the tumor-microenvironment (TME), suggesting that tumor niche could also be targeted by epigenetic therapy [15-18]. Through this review, we have presented an exhaustive account of the epigenetic deregulations observed in the TME, especially of the stromal cell population corresponding to the cancer-associated fibroblasts (CAFs), the suppressive immune cells namely; the regulatory T cells (Tregs) and the myeloid-derived suppressor cells (MDSCs). Finally, we have deliberated the potential impact of epigenetic therapies on the immunosuppressive properties of TME.

### **Genetic and epigenetic deregulation in cancer cells**

Cancer is a genetic and epigenetic disease, where genetic alterations and epigenetic deregulations are entangled from the beginning to the end of the oncogenic process [19]. Genetic alterations in cancer cells can occur in both oncogenes and tumor suppressors genes (TSGs). Interestingly, some additional mutations are not oncogenic *per se* but favor the crosstalk with TME thus linking tumor genetics and tumor niche features, as observed in follicular lymphoma (FL) [20]. In addition, in many cancers, genetic alterations can occur through gene coding for epigenetic regulators. As an example, mutations in genes with a role of catalyzing the post-translational modification of histones, such as the histone H3 lysine 4 (H3K4) methyltransferases KMT2D and KTM2C, the histone acetyltransferases CREBBP and EP300, and the histone H3 lysine 27 (H3K27) methyltransferase EZH2, are a hallmark of FL [21]. About 70% of FL patients harbor at least 2 mutations in chromatin-modifying genes, making the targeting of epigenetic modifiers an attractive therapeutic target in this disease. Whilst, the majority of EZH2 mutations are subclonal events, mutations of CREBBP probably arise as early driver genetic events residing within tumor cell progenitors [22]. It is also the case for *de novo* acute myeloid leukemia (AML) where genetic alteration of epigenetic regulators (DNMT3A and ten-eleven-translocation 2 (TET2)) can be observed at the beginning of the oncogenic process [23]. Besides mutations, epigenetic deregulations are observed in virtually all cancer types, with disruptions of DNA methylation, histone modifications and non-coding RNAs mechanisms and are already extensively described [24,25]. Interestingly like

genetic alterations, epigenetic deregulations in cancer cells could also impact the establishment of a supportive TME. As an example, overexpression of the H3K27 demethylase KDM6B impacts the regulation of the NF- $\kappa$ B pathway in melanoma cancer, with the overexpression of stanniocalcin 1 (STC1) and chemokine (C-C motif) ligand 2 (CCL2) by melanoma tumor cells, leading simultaneously to macrophage infiltration, angiogenesis, and lung metastases [26].

### **Tumor microenvironment leads tumor progression**

Tumor cells live in a complex ecosystem formed by infiltrating immune cells, endothelial cells, and stromal cells [27]. There is increasing evidence suggesting the involvement of TME in many tumorigenic processes including tumor cell proliferation and survival, immune escape, metastatic process, angiogenesis, and resistance to therapies [28-30]. As an example, in GC-derived B-cell lymphomas, neutrophils recruited through production of IL-8 by stromal cells, could provide supportive effect to FL B-cells *in-vitro* [31]. In addition, tumor associated macrophages (TAM), could also lead to tumor progression through the establishment of an immune-suppressive microenvironment notably via the production of chemokines like CCL17, CCL18 and CCL22, with the consequence to sequester Tregs and inhibits immune responses [32-34]. Immune suppression in the TME is also mediated by stromal cells such as CAFs and myeloid cells like MDSCs and will be further elaborated in the next sections. Interestingly, the emergence of immune-checkpoint therapy has revealed a major role of TME in the resistance to immune-checkpoint inhibitors. In particular through physical blockade of access by immune cells to tumor bed (immune-excluded tumors) and inhibition of immune cell activation/cytotoxicity (inflamed tumors), revealing a key role of immune cell number, localization, and activation in patient clinical outcome [35,36]. Recently, there have been scientific studies that describe the potential for combining epigenetic inhibitors with immune checkpoint inhibitors, as a mechanism of cancer therapy [37-39], This emphasizes the desideratum to satisfactorily discern the role of epigenetic deregulations in the emergence of immunomodulating cellular properties of the TME as well as the impact of epigenetic therapies on these cells.

## II) Epigenetic deregulation mechanisms in CAFs

### Stromal cell compartment

Cancer-associated fibroblasts (CAFs), represents one of the most abundant stromal cell populations in the tumor microenvironment. They are frequently described as myofibroblast characterized by the expression of  $\alpha$ -sma and have been shown to enhance tumor phenotypes, with critical role in tumor initiation, progression, dissemination, immune escape, and drug resistance [40]. Based on these observations, CAFs emerged as a key player in the TME with numerous reports describing the strong pro-tumorigenic properties of CAFs [41]. These pro-tumorigenic properties are mediated as an example, by the expression of stromal cell-derived factor 1 (SDF-1), also called CXCL12, which could lead to the recruitment and activation of Tregs [42,43]. CAFs also create an immunosuppressive microenvironment through the production of TGF $\beta$  and the establishment of a particular extra-cellular matrix (ECM) environment that blocks immune cell trafficking associated with immune-checkpoint therapy failure [35,44]. To date, it is quite difficult to study native CAF subpopulations due to their low numbers and lack of characterization for specific surface markers. These difficulties in describing common CAF markers can be ascribed to their multiple putative origins, including mesenchymal stromal cells, activation of resident fibroblasts, pericytes and epithelial cells that follow epithelial-to-mesenchymal transition (EMT), or conversion of endothelial cells [45]. Moreover, a recent study elegantly demonstrates that CAFs are a heterogeneous population *in situ* with different subsets of CAFs carrying specific phenotypes, gene expression profiles, and functions in a given TME [42].

Another layer of complexity could come from the fact that not only are tumors spatially heterogeneous, but CAFs themselves are a heterogeneous population, with different phenotypes and functions depending on their localization, as well as their origins [46]. In particular; solid tumors frequently include tertiary lymphoid structures, that contain CAFs with features of lymphoid stromal cells that regulate immune cell recruitment and activation within lymphoid organs. Presence of these tertiary lymphoid structures are at least in colon cancer, associated with a better outcome [47], suggesting possible anti-tumoral activities [48]. Conversely, infiltration by draining lymph nodes of tumor cells was shown to trigger reprogramming of resident lymphoid stromal cells into CAFs leading to a reduced capacity to promote leukocyte recruitment, migration, and activation; a functional phenotype associated

with a pro-tumoral activity [49]. Similar observation was made in FL where the primary tumor site is the lymph node and where infiltrating lymphoid stromal cells present a specific phenotype. These cells notably overexpress CXCL12 under the influence of T-cell-derived IL-4 and trigger FL B cell activation, migration, and adhesion [50]. However, CAFs origin in tumor-invaded lymph node is still a matter of debate and is not fully elucidated. Interestingly, bone marrow is also penetrated in about 70% of FL patients and CXCL12 is also overexpressed in FL bone marrow stromal cells. Moreover, bone marrow stromal cells also overexpress CCL2 and IL-8 that contributes to the recruitment and activation of tumor-supportive monocyte and neutrophils [31,51].

As discussed above, CAFs are heterogeneous, arising from different cells of origin and displaying both pro and anti-tumoral activities. To date, there is no clear report showing that genetic alterations could be the drivers of CAF phenotype [45]. Conversely, studies showing that CAFs in culture retain some of their properties; suggesting that epigenetic mechanisms could be involved in changes occurring in their transcriptional and phenotypic profiles [15,49].

#### **DNA methylation**

It was recently shown that CAFs could be involved in the epigenetic reprogramming of cancer cells, as observed in breast cancer [52]. Moreover, it was proposed that CAFs could present the same DNA methylation pattern as observed in tumor cells, including a global DNA hypomethylation and a local DNA hypermethylation. Such profiles have been observed in CAFs from various cancer types such as gastric cancers [53], colorectal cancer [54], lung cancer [55]. However, these previous results are not in agreement with a recent study in prostate cancer [18]. In this study, the authors by using whole-genome bisulfite sequencing, showed that CAFs from prostate cancer did not exhibit global hypomethylation but rather changed (both increased and decrease) at discrete loci, suggesting that DNA global hypomethylation in CAFs is probably, dependent of the cancer type and should be assessed more carefully with resolutive techniques. These recent observations concerning the whole genome DNA methylation profile of CAFs, both highlights the importance of analyzing DNA methylation for reprogramming of CAFs [56]. Interestingly, local hypermethylation in CAFs was shown to be involved in the conversion of normal fibroblasts into pro-invasive fibroblasts in several cancers (head and neck, lung, and breast cancer). This conversion was mediated through an increased expression of DNMT3b and a local hypermethylation of SHP-1 [57]. In this context, DNMT3b



overexpression was induced by the activation of the JAK1/JAK3 (Janus kinase) signaling pathway by P300 histone acetylation in response to the proinflammatory cytokine leukemia inhibitory factor (LIF), secreted by tumor cells [57]. In addition, inhibition of the RAS inhibitor RASAL1 by DNMT1 was observed in CAFs from renal cancer [58], whereas in prostate cancer aberrant DNA methylation targets the Ras inhibitor RASAL3 [59]. Both lead to the oncogenic activation of Ras and to the modification of CAF metabolism (glutamine synthesis) which are associated with an increase in cancer cell survival and proliferation. DNMT1 was also shown to be up-regulated in breast cancer and its up-regulation is critical for the conversion of normal fibroblasts into CAFs [60]. This conversion was mediated by HuR protein which stabilizes DNMT1 mRNA leading to enhanced pro-inflammatory properties of CAFs via an increased expression of CXCL12, TGF $\beta$ , and IL-6 [60].

### **Histone modifications and chromatin remodeling**

Histone marks were shown to be involved in the regulation of the functional properties of CAFs. These regulations can be separated in three different observations:

i) Change in histone modifying enzyme gene expression among CAFs. Indeed, it was described in breast cancer that CAFs can overexpress HDAC6, leading to the activation of prostaglandin E2/cyclooxygenase-2 (PGE2/COX2) expression in association with signal transducer and activator of transcription 3 (STAT3) and enhancing the recruitment of MDSCs and Tregs cells [61]. These observations accent that CAF epigenetic deregulations could impact not only cancer cells but also other cells of the TME [62]. HDAC1/3/8 were also shown to be involved in CAF differentiation upon TGF $\beta$  exposure, increasing tumor growth and ECM secretion [63]. Interestingly, the use of an inhibitor of colony stimulating factor-1 receptor (CSF-1R); the JNJ-40346527, to target the tumor-associated macrophage, was shown to have a pro-tumoral effect mediated by CAFs. Indeed this paradoxical pro-tumoral effect relies on the blockade of the CSF-1-dependant recruitment of HDAC2 to the promoter of granulocyte-specific chemokines in CSF1R-expressing CAF, thus increasing the release of MDSC-recruiting chemokines, that ultimately leads to an increase in polymorphonuclear MDSC (PMN-MDSC) infiltration, ultimately, explaining the limited clinical efficacy of CSF1-R inhibitor [64].

ii) Chromatin remodeling in CAFs. The transcriptional repressor ATF3 and CSL were shown to control CAF activation and are repressed in CAFs. Re-expression of these factors triggers chromatin remodeling and suppression of CAF tumor-promoting properties in mouse models

[65]. In gastric cancer, it was recently shown that CAFs have a distinct H3K27me3 profile compared to normal fibroblasts. This loss was mostly observed with genes involved in stem cell niche, cell growth and tissue development like WNT5A, GREM1, NOG and IGF2 [66]. In addition, the chromatin remodeler HMGA2 enhances the tumor supportive properties of stromal cells in mouse prostate cancer [67].

iii) CAFs can also affect epigenetic landscape in cancer cells. In ovarian cancer, CAFs induce an overexpression of EZH2 leading to an increase of cancer cell migration [68]. Interestingly, epigenetic changes induced by fibroblasts on cancer cells could also be associated with antitumor properties. In particular, it has been described that normal fibroblasts could inhibit breast cancer cell proliferation, through mechanosensitive downregulation and nuclear exit of the H3K9 demethylase JMJD1a leading to a downregulation of YAP/TAZ expression [69]. This observation paves the way to the idea that reversing and inhibiting epigenetic mechanisms involved in CAFs conversion/differentiation could promote generation of CAFs that carry anti-tumor properties.

#### **Non-coding RNA**

Many studies highlight the importance of miRNA regulation in CAF pro-tumoral properties [70]. As an example, in prostate cancer, miR-15a and miR-16 are downregulated in CAFs thus reducing the post-transcriptional repression of fibroblast growth factor 2 (Fgf-2) and Fgfr1 and enhancing cancer cell survival, proliferation, and invasiveness [71]. It was also shown that upregulation of miR-409 in CAFs from prostate cancer is sufficient for the differentiation of normal stroma into CAFs [72]. Interestingly, miR-409 can then be released by CAFs via extracellular vesicles and result in enhanced tumor progression and EMT. In ovarian cancer, miR-200 supports the up-regulation of CXCL12 beta isoform expression in a specific CAF subtype and is associated with immunosuppressive cell recruitment [42]. However, in non-mesenchymal ovarian tumors, another subset of CAFs are present which express a miRNA cluster that represses CXCL12 expression, miR141/200c and decreases Tregs recruitment in tumor niche [73], highlighting the bivalent role of CAFs and miRNA regulation.

### **III) Epigenetic deregulation of the suppressive immune cell subsets**

Besides CAFs, the TME also include several pro-tumoral and anti-tumoral immune cell subsets, and epigenetic deregulation of the immune cells in cancer is already described by others [16,62]. However, the crosstalk between CAFs and the main suppressive immune cells, namely Tregs and MDSCs, plays a key role in organizing tumor-promoting microenvironment, highlighting the need to have a holistic comprehension on the epigenetic deregulation of the suppressive cells found in TME. Moreover, understanding epigenetic mechanisms sustaining the development of this immunosuppressive network would be useful in predicting the impact of epigenetics drugs on the TME (Figure 1).

#### **Immune cells in the tumor microenvironment**

Many immune cell subsets could be identified within TME and participate in the establishment of an immunosuppressive microenvironment [62]. Among them, Tregs secrete immunosuppressive cytokines like IL-10, IL-35 and tumor growth factor beta (TGF $\beta$ ) [74] and block activation of effector T cells. This blockade is partially mediated through cytotoxic T lymphocyte associated protein 4 (CTLA-4) expression and consumption of IL-2, limiting IL-2 availability [74]. MDSCs are another heterogeneous subset of immune cells also found in TME of many tumors [75]. MDSCs belong to the monocytic (M-MDSC) or granulocytic (PMN-MDSC) lineages and display strong immunosuppressive functions mediated through diverse mechanisms including production of nitric oxide (NO), reactive oxygen species (ROS), immunosuppressive enzymes including arginase 1 (ARG1) or indoleamine 2,3-dioxygenase (IDO), and immunosuppressive cytokines like IL-10 and TGF $\beta$ , that will altogether impact T cells and naturel Killer (NK) functions [76]. As observed on Tregs, MDSCs immunosuppressive properties are also mediated by expression of the inhibitory immune checkpoint PD-L1 [77].

#### **DNA methylation**

DNA methylation was shown to be involved in the regulation of Tregs, the transcription factor forkhead box P3 (FOXP3) is crucial for the development and function of Tregs and its expression is strongly dependent on the Treg-specific demethylated region (TSDR), an epigenetic marker for natural Tregs (nTregs) [78]. DNA demethylation agents used in cancer cell therapy could thus impact Treg population. As it was shown for CAFs and cancer cells,

MDSCs also present a global DNA hypomethylation profile with a local gain of DNA hypermethylation. These are notably shown in ovarian cancer, where MDSCs presents a global DNA hypomethylation profile compared to dendritic cells, with a specific gain of DNA methylation and repression of genes associated with an immunogenic phenotype like S1PR4, RUNX1 and FAS. This loci specific methylation, could be related to an increase of DNMT3A (a *de novo* DNA methyltransferase) expression in MDSC [79]. This neoteric observation seems in contradiction with other studies shown in mice models where Cannabinoid ( $\Delta^9$ -Tetrahydrocannabinol) could induce a global hypomethylation of MDSCs and a decrease of DNMT3A and DNMT3B expression by DNA methylation of their promotor. This leads to a higher expression of Arg1 and STAT3, that ultimately promote MDSCs immunosuppressive properties [80,81], suggesting that a common epigenetic enzyme deregulation could have different ramifications on acquisition of tumor supportive properties.

#### **Histone modifications**

As observed in CAFs, histone modifications are also found altered in immune cells of tumor niche. As an example, HDAC11 was shown to be a negative regulator of MDSCs expansion in mice, in addition the same study described that MDSC isolated from HDAC11-KO tumor-bearing mice were more suppressive than MDSC, purified from the wild type mice [82]. HDAC11 in co-operation with HDAC6 was also shown to be involved in the regulation of IL-10 expression (a cytokine known to recruit immune-suppressive cells like MDSCs) by antigen presenting cells (APC). This suggested the involvement of this mechanism during tumorigenesis to promote an immunosuppressive TME. Besides HDAC11, HDAC2 was shown to be involved in the conversion of M-MDSCs to PMN-MDSCs with higher pro-tumoral properties through the inhibition of the retinoblastoma gene 1 (RB1) [83]. Moreover, the H3K4 methyltransferase SETD1B activates nitric oxide synthase 2 (nos2) expression in MDSCs, leading to an inhibition of T cell-activation, and is associated with an anti-tumor immune response [84].

Epigenetic deregulation in tumor cells can also indirectly lead to an increase of the immunosuppressive properties of TME. In ovarian cancers, EZH2 and DNMT1 are involved in gene repression of T helper 1 chemokines CXCL9 and CXCL10, associated with a decrease in CD8<sup>+</sup> effector T cell infiltration [85]. In colon cancer, T cell recruitment is also impacted by the

epigenetic regulation of CXCL9 and CXCL10 expression by H3K27me3 repression marks which is modulated by an EZH2/KDM6B balance [86].

#### **Non-coding RNA**

As observed for CAFs, it is well described that miRNA are involved in MDSCs identity and acquisition of their pro-tumoral properties [16,76]. Exosomes from glioma cells produced in hypoxic condition induce MDSCs activation by transferring miR-29a and miR-92a. These miRs enhance the proliferation and function of MDSCs by targeting HMG-box transcription factor 1 (Hbp1), a mitosis inhibitor protein and protein kinase CAMP-dependent type 1 regulatory subunit alpha (Prkar1 $\alpha$ ), an inhibitor of the STAT3 pathway activation [87]. In glioma as well, miR-10a and miR-21 that target RAR-related orphan receptor alpha (RORA) and phosphatase and tensin homolog (PTEN) respectively are also transmitted to MDSCs via exosomes from cancer cells, leading to an enhanced MDSCs differentiation and activation [88]. Moreover, in melanoma, a set of miRs (miR-146a, miR-155, miR-125b, miR-100, let-7e, miR-125a, miR-146b, miR-99b) have been associated with MDSC differentiation and poor clinical response to immune checkpoint therapy [89]. Interestingly, MDSCs in epithelial ovarian cancer can also influence TME polarization and function through the transmission of miR-21 and miR-29a to T cells, leading to the blockade of STAT3 signaling pathway and increase in immunosuppressive Tregs [90].

#### **IV) Epigenetic therapies**

Epigenetic deregulations in tumor cells relies on epigenetic enzymes that can be targeted by epigenetic drugs; many epigenetic drugs (epi-drugs) have been developed, essentially targeting the DNA-methylation machinery and HDACs [91]. The development of epi-drugs is still an intense area of research [4,5,92]. We will not discuss here the direct impacts of these epi-drugs on cancer cells but how these epi-drugs could directly and indirectly impact cells present in the TME (Table1).

## **TME mediated anti-tumoral impacts of epi-drugs**

### **-DNA methyltransferase inhibitors**

DNA demethylating agents (DNMTi), such as 5-aza-2'-deoxycytidine (5-Aza-Cdr), can be used to inhibit DNMTs and are currently approved for the treatment of myelodysplastic syndromes (MDS) and AML [93]. These DNMTi through modulations of the epigenome of the cancer cells could indirectly impact the TME of the cancer cells. As an example, DNMTi could mimic a viral infection in cancer cells called "viral mimicry", [94,95] through the re-expression of endogenous retrovirus in cancer cells and formation of double stranded RNA. This "viral mimicry" by creating an inflammatory context, favor the activation and recruitment of T lymphocytes and in consequence, lead to an increased efficacy of immunotherapy strategies [95,96]. Interestingly, DNMTi can also impact directly the TME. In particular, it was recently proposed that DNMTi could prevent CD8+ T cell exhaustion, by allowing them to retain their effectors functions [97]. Moreover, in a mice model of breast cancer, It was shown that DNMTi could impact MDSCs by reducing their expansion and potentially diminishing their immunosuppressive properties, thus favoring adoptive T cell transfer [98]. This observation confirmed previous results describing a direct impact of 5-Aza-CdR on MDSCs proliferation, in mice models of prostate cancer adenocarcinoma and lung cancer [99]. In addition, it was shown that DNMT3a genetic inhibition, is sufficient to suppress MDSCs immunosuppressive properties abrogating their capacity to suppress CD8+ T cell proliferation and the production of IFN $\gamma$  in the context of ovarian cancer [79]. Finally, to date, only one study describes the impact of DNMTi on CAF. In this study, done on human fibroblasts from various cancers (head and neck, breast, lung) the author described a constitutive activation of the JAK1/STAT3 pathway, involving both the de novo methyltransferase (DNMT3b) and DNMT1 for the maintenance of DNA methylation to stably repress the PTPN6 tyrosine phosphatase leading to the acquisition of the pro-tumoral properties of CAFs [57]. These pro-tumoral properties could be then reversed by DNMTi treatment in combination with JAK1/2 inhibitors.

### **-Histone modifying enzyme inhibitors**

Only few studies, have established the impacts of histone modifying enzyme on CAFs. One study describes the role of Scriptaid ,a selective inhibitor of HDACs, 1, 3 and 8, on CAFs, in an *in-vitro* and *in-vivo* model of melanoma [63]. They observed, that blocking of the HDACs (1,3 and 8) supresses the activation of the TGFbeta pathway in CAFs leading to a reduction of the

CAFs tumor supportive properties. In a PDX model of pancreatic ductal adenocarcinoma (PDAC), JQ1, an inhibitor of chromatin readers which contain bromo- and extra-terminal domain (BET) can reverse CAF phenotype by downregulating key pathways involved in CAF activation, in particular the activation of the TGF $\beta$  pathway [100]. These two studies highlight the importance of blocking the TGFbeta pathway on CAFs, a pathway that was recently shown to be associated with the immune-suppressive properties of CAFs [44]. In addition, a recent study, described that methyltransferase inhibitors (targeting both histones and DNA methylation), could block the capacity of CAFs to remodel the ECM and prevent metastasis formation in a model of breast cancer, through the interaction of SNAIL with the methyltransferase PRMT1 and PRMT4 [101].

To finish, indirect impact of histone modifying enzyme inhibitors on the TME, mediated by the cancer cells were described. As an example, in murine ovarian cancer, 5-Aza-CdR induces a type I IFN response, leading to the activation of cytotoxic T cells and NK cells and reducing the percentage of MDSCs. These anti-tumor effects are enhanced by a combination of HDACi and immune checkpoint inhibitors [102]. EZH2 inhibitors were also shown to inhibit Tregs, and to improve anti-CTLA-4 therapy, highlighting our incomplete knowledge of the impact of epi-drugs on anti-tumor immune response [103]. Finally, Immune checkpoint inhibitors in association with the two HDACi, entinostat or mocetinostat, decrease MDSCs recruitment and increase CD8+ T cell infiltrations in TME of breast, pancreatic cancer and non-small cell lung cancer [104,105].

### **Epi-drugs promote a pro-tumoral microenvironment**

However, epi-drugs could have a dual effect on the stromal compartments. Indeed, DNMTi were also shown to induce immunosuppressive MSC through upregulation of COX2 [106], an effect that could be beneficial in immune diseases where immune cells need to be tempered, but could have inverse effects in cancer. Moreover, DNMTi also increases the immunosuppressive properties of MDSCs through STAT3 and ARG1 activation [80]. the EZH2 inhibitor GSK126 can induce an increase of MDSCs while CD4 + and CD8+ T cells are decreased [107]. In addition, some studies suggest that HDACi could enhance MDSCs proliferation [108,109] as well as Treg differentiation [110-112]. In addition, in PDAC, HDACi were shown to induce a supportive stroma and inhibition of HDAC2 in CAFs leading to an increased secretion of tumor-supportive cytokines and chemokines [113]. These were also confirmed in

breast tumors, where HDAC1 inhibition in CAFs leads to an increased expression of osteopontin and promote tumor growth [114]. These observations highlight that epi-drugs could have a dual impact on TME, that need to be cautiously analyzed by further researches.

## **V) Future perspectives**

In conclusion, epigenetic deregulation in TME, especially in CAFs and MDSCs, are involved in the establishment of an immunosuppressive microenvironment. Moreover, epigenetic therapies targeting cancer cells, such as DNMTi and HDACi could favor or repress the tumor-supportive activities of TME. To date, there is a scarcity of studies addressing the direct impact of these epigenetic therapies on cells present in the TME. As discoveries in the role of TME towards tumor progression and resistance to therapy advances, it will become inevitable to address how these treatments could impact cells of the TME. Especially, one challenge that surfaces, would be the specific targeting of the TME to deliver epigenetic drugs. Development of strategies to deliver epigenetic drugs is an intense area of research [115,116]. Targeting of the TME is currently being developed and achievements thus far, in this field of research have been epitomized in this review [116]. However, identification of the right ligand to target specifically CAFs will be a prerequisite and is still a matter of research [116]. In addition, with the development of the single-cell epigenomics technique [117-119], it will be possible to have a better understanding of the role of epigenetic modifications for the fine-tuning of TME differentiation and functions. Such techniques, will be a prerequisite for a better design of adequate epigenetic therapeutic strategies.



## **Executive summary**

### **Epigenetic mechanisms deregulation in CAFs**

- Cancer associated fibroblasts (CAFs) are key players of the tumor microenvironment (TME) and lead to tumor progression.
- CAFs differ from normal fibroblasts, in contrary to cancers cells. CAFs do not present genomic alterations, highlighting others mechanisms involved in their modification and associated with their pro-tumorigenic properties.
- CAFs present epigenetic deregulation, like global DNA hypomethylation and local DNA hypermethylation. DNA methylation seems to be critical for CAF conversions. In addition, histone modifications and chromatin remodeling are also observed in CAFs and support their pro-tumorigenic properties.

### **Epigenetic deregulation of the suppressive immune compartment**

- TME include beside CAFs, others immunosuppressive cells and especially regulatory T cells (Tregs) and myeloid-derived suppressor cells (MDSCs).
- Immunosuppressive properties of Tregs is mediated in part by DNA demethylation of the Treg-specific demethylated region. In addition, as observed in CAFs and Cancer cells, MDSCs present a global DNA-methylation profile and a local hypermethylation.
- Epigenetic changes observed in MDSCs also include histone modification and Non-coding RNA that are both involved in the acquisition of their immunosuppressive properties.

### **Epigenetic therapies**

- Epigenetic drugs (epi-drugs) were first developed to target cancer cells. However, these epi-drugs can impact directly and indirectly on cells present in the TME
- Epi-drugs could have a dual impact on the immunosuppressive properties of cells present in TME. These Epi-drugs can both favor or repress the tumor-supportive activities of TME
- In future, a better understanding of the role of epigenetic modification in the fine-tuning of TME is a prerequisite for a better design of adequate epigenetic therapeutic strategies.

450

451 **Conflicts of interest:** The authors declare no conflicts of interest.

452

Revised manuscript

## Bibliography

1. Slack JMW. Conrad Hal Waddington: the last Renaissance biologist? *Nature Publishing Group*.
2. Goldberg AD, Allis CD, Bernstein E. Epigenetics: A Landscape Takes Shape. *Cell*. 128(4), 635–638 (2007).
3. Allis CD, Jenuwein T. The molecular hallmarks of epigenetic control. *Nat. Rev. Genet.* 17(8), 487–500 (2016).
4. Dzobo K. Epigenomics-Guided Drug Development: Recent Advances in Solving the Cancer Treatment "jigsaw puzzle". *OMICS*. 23(2), 70–85 (2019).
5. Esteller M. Epigenetic drugs: More than meets the eye. *undefined*. 12(5), 307–307 (2017).
6. Schübeler D. Function and information content of DNA methylation. *Nature*. 517(7534), 321–326 (2015).
7. Zhou VW, Goren A, Bernstein BE. Charting histone modifications and the functional organization of mammalian genomes. *Nat. Rev. Genet.* 12(1), 7–18 (2011).
8. Kouzarides T. Chromatin Modifications and Their Function. *Cell*. 128(4), 693–705 (2007).
9. Michalak EM, Burr ML, Bannister AJ, Dawson MA. The roles of DNA, RNA and histone methylation in ageing and cancer. *Nat. Rev. Mol. Cell Biol.* 15(6), 12–17 (2019).
10. Ambros V. The functions of animal microRNAs. *Nature*. 431(7006), 350–355 (2004).
11. Fernandes JCR, Acuña SM, Aoki JI, Floeter-Winter LM, Muxel SM. Long Non-Coding RNAs in the Regulation of Gene Expression: Physiology and Disease. *Noncoding RNA*. 5(1), 17 (2019).
12. Sun Y-C, Wang Y-Y, Ge W, Cheng S-F, Dyce PW, Shen W. Epigenetic regulation during the differentiation of stem cells to germ cells. *Oncotarget*. 8(34), 57836–57844 (2017).
13. V Subramaniam A, Yehya AHS, Cheng WK, Wang X, Oon CE. Epigenetics: The master control of endothelial cell fate in cancer. *Life Sciences*. 232, 116652 (2019).
14. Jambhekar A, Dhall A, Shi Y. Roles and regulation of histone methylation in animal development. *Nat. Rev. Mol. Cell Biol.* 403, 1–17 (2019).
15. Du H, Che G. Genetic alterations and epigenetic alterations of cancer-associated fibroblasts. *Oncol Lett*. 13(1), 3–12 (2017).
16. Liu M, Zhou J, Chen Z, Cheng AS-L. Understanding the epigenetic regulation of tumours and their microenvironments: opportunities and problems for epigenetic

- 487 therapy. *J. Pathol.* 241(1), 10–24 (2017).
- 488 **\*\* This review underlines the impact of epigenetic control on the tumor**  
 489 **microenvironment and in particular myeloid compartment.**
- 490 17. Marks DL, Olson RL, Fernandez-Zapico ME. Epigenetic control of the tumor  
 491 microenvironment. <http://dx.doi.org/10.2217/epi-2016-0110>. 8(12), 1671–1687  
 492 (2016).
- 493 18. Pidsley R, Lawrence MG, Zotenko E, *et al.* Enduring epigenetic landmarks define the  
 494 cancer microenvironment. *Genome Res.* 28(5), 625–638 (2018).
- 495 19. You JS, Jones PA. Cancer Genetics and Epigenetics: Two Sides of the Same Coin?  
 496 *Cancer Cell.* 22(1), 9–20 (2012).
- 497 20. Boice M, Salloum D, Mourcin F, *et al.* Loss of the HVEM Tumor Suppressor in  
 498 Lymphoma and Restoration by Modified CAR-T Cells. *Cell.* 167(2), 405–418.e13  
 499 (2016).
- 500 21. Pasqualucci L. Molecular pathogenesis of germinal center-derived B cell lymphomas.  
 501 *Immunological Reviews.* 288(1), 240–261 (2019).
- 502 22. Green MR. Chromatin modifying gene mutations in follicular lymphoma. *Blood.*  
 503 131(6), 595–604 (2018).
- 504 23. Martignoles J-A, Delhommeau F, Hirsch P. Genetic Hierarchy of Acute Myeloid  
 505 Leukemia: From Clonal Hematopoiesis to Molecular Residual Disease. *International*  
 506 *Journal of Molecular Sciences* 2018, Vol. 19, Page 3850. 19(12), 3850 (2018).
- 507 24. Jones PA, Laird PW. Cancer-epigenetics comes of age. *Nature Genetics.* 21(2), 163–  
 508 167 (1999).
- 509 25. Jones PA, Baylin SB. The Epigenomics of Cancer. *Cell.* 128(4), 683–692 (2007).
- 510 26. Park W-Y, Hong B-J, Lee J, Choi C, Kim M-Y. H3K27 Demethylase JMJD3 Employs the  
 511 NF-κB and BMP Signaling Pathways to Modulate the Tumor Microenvironment and  
 512 Promote Melanoma Progression and Metastasis. *Cancer Res.* 76(1), 161–170 (2016).
- 513 27. Hanahan D, Coussens LM. Accessories to the Crime: Functions of Cells Recruited to  
 514 the Tumor Microenvironment. *Cancer Cell.* 21(3), 309–322 (2012).
- 515 28. Hirata E, Sahai E. Tumor Microenvironment and Differential Responses to Therapy.  
 516 *Cold Spring Harb Perspect Med.* 7(7), a026781 (2017).
- 517 29. Senthebane DA, Jonker T, Rowe A, *et al.* The Role of Tumor Microenvironment in  
 518 Chemoresistance: 3D Extracellular Matrices as Accomplices. *International Journal of*  
 519 *Molecular Sciences* 2018, Vol. 19, Page 3850. 19(10), 2861 (2018).
- 520 30. Senthebane DA, Rowe A, Thomford NE, *et al.* The Role of Tumor Microenvironment  
 521 in Chemoresistance: To Survive, Keep Your Enemies Closer. *International Journal of*

522 *Molecular Sciences* 2018, Vol. 19, Page 3850. 18(7), 1586 (2017).

523 31. Guilloton F, Caron G, Ménard C, *et al.* Mesenchymal stromal cells orchestrate  
524 follicular lymphoma cell niche through the CCL2-dependent recruitment and  
525 polarization of monocytes. *Blood*. 119(11), 2556–2567 (2012).

526 32. Zhou J, Ding T, Pan W, Zhu LY, Li L, Zheng L. Increased intratumoral regulatory T cells  
527 are related to intratumoral macrophages and poor prognosis in hepatocellular  
528 carcinoma patients. *International Journal of Cancer*. 125(7), 1640–1648 (2009).

529 33. Mamrot J, Balachandran S, Steele EJ, Lindley RA. Molecular model linking Th2  
530 polarized M2 tumour-associated macrophages with deaminase-mediated cancer  
531 progression mutation signatures. *Scandinavian Journal of Immunology*. 89(5),  
532 e12760 (2019).

533 34. Wang D, Yang L, Yue D, *et al.* Macrophage-derived CCL22 promotes an  
534 immunosuppressive tumor microenvironment via IL-8 in malignant pleural effusion.  
535 *Cancer Letters*. 452, 244–253 (2019).

536 35. Mariathasan S, Turley SJ, Nickles D, *et al.* TGF $\beta$  attenuates tumour response to PD-L1  
537 blockade by contributing to exclusion of T cells. *Nature*. 554(7693), 544–548 (2018).

538 36. Kather JN, Suarez-Carmona M, Charoentong P, *et al.* Topography of cancer-  
539 associated immune cells in human solid tumors. *Elife*. 7, 16878 (2018).

540 37. Terranova-Barberio M, Thomas S, Munster PN. Epigenetic modifiers in  
541 immunotherapy: a focus on checkpoint inhibitors. *Immunotherapy*. 8(6), 705–719  
542 (2016).

543 38. Jones PA, Ohtani H, Chakravarthy A, De Carvalho DD. Epigenetic therapy in immune-  
544 oncology. *Nature Reviews Cancer*. 22, 1 (2019).

545 39. Toor SM, Sasidharan Nair V, Decock J, Elkord E. Immune checkpoints in the tumor  
546 microenvironment. *Semin. Cancer Biol.* (2019).

547 40. Costa A, Scholer-Dahirel A, Mechta-Grigoriou F. The role of reactive oxygen species  
548 and metabolism on cancer cells and their microenvironment. *Semin. Cancer Biol.* 25,  
549 23–32 (2014).

550 41. Monteran L, Erez N. The Dark Side of Fibroblasts: Cancer-Associated Fibroblasts as  
551 Mediators of Immunosuppression in the Tumor Microenvironment. *Front Immunol.*  
552 10, 1835 (2019).

553 42. Givel A-M, Kieffer Y, Scholer-Dahirel A, *et al.* miR200-regulated CXCL12 $\beta$  promotes  
554 fibroblast heterogeneity and immunosuppression in ovarian cancers. *Nature*  
555 *Communications*. 9(1), 1056 (2018).

556 \* **This study showed CAFs heterogeneity in breast cancers and the impact of a miR,**  
557 **miR-141/200a, on the immunosuppressive properties of CAFs with CXCL12b**  
558 **secretion.**

- 559 43. Costa A, Kieffer Y, Scholer-Dahirel A, *et al.* Fibroblast Heterogeneity and  
560 Immunosuppressive Environment in Human Breast Cancer. *Cancer Cell.* 33(3), 463–  
561 479.e10 (2018).
- 562 44. Chakravarthy A, Khan L, Bensler NP, Bose P, De Carvalho DD. TGF- $\beta$ -associated  
563 extracellular matrix genes link cancer-associated fibroblasts to immune evasion and  
564 immunotherapy failure. *Nature Communications.* 9(1), 4692 (2018).
- 565 45. Madar S, Goldstein I, Rotter V. “Cancer associated fibroblasts” – more than meets  
566 the eye. *Trends in Molecular Medicine.* 19(8), 447–453 (2013).
- 567 46. Chang JE, Turley SJ. Stromal infrastructure of the lymph node and coordination of  
568 immunity. *Trends Immunol.* 36(1), 30–39 (2015).
- 569 47. Colbeck EJ, Ager A, Gallimore A, Jones GW. Tertiary Lymphoid Structures in Cancer:  
570 Drivers of Antitumor Immunity, Immunosuppression, or Bystander Sentinels in  
571 Disease? *Front Immunol.* 8, 867 (2017).
- 572 48. Bindea G, Mlecnik B, Tosolini M, *et al.* Spatiotemporal dynamics of intratumoral  
573 immune cells reveal the immune landscape in human cancer. *Immunity.* 39(4), 782–  
574 795 (2013).
- 575 49. Riedel A, Shorthouse D, Haas L, Hall BA, Shields J. Tumor-induced stromal  
576 reprogramming drives lymph node transformation. *Nat. Immunol.* 17(9), 1118–1127  
577 (2016).
- 578 \*\* **This transcriptomic study during tumorigenesis show a conversion of stromal cells**  
579 **populations in CAFs-like cells in mice models.**
- 580 50. Pandey S, Mourcin F, Marchand T, *et al.* IL-4/CXCL12 loop is a key regulator of  
581 lymphoid stroma function in follicular lymphoma. *Blood.* 129(18), 2507–2518 (2017).
- 582 51. Grégoire M, Guilloton F, Pangault C, *et al.* Neutrophils trigger a NF- $\kappa$ B dependent  
583 polarization of tumor-supportive stromal cells in germinal center B-cell lymphomas.  
584 *Oncotarget.* 6(18), 16471–16487 (2015).
- 585 52. Mathot P, Grandin M, Devailly G, *et al.* DNA methylation signal has a major role in  
586 the response of human breast cancer cells to the microenvironment. *Oncogenesis.*  
587 6(10), e390 (2017).
- 588 53. Le Jiang, Gonda TA, Gamble MV, *et al.* Global Hypomethylation of Genomic DNA in  
589 Cancer-Associated Myofibroblasts. *Cancer Res.* 68(23), 9900–9908 (2008).
- 590 54. Ling E, Ringel A, Sigal-Batikoff I, *et al.* Human Colorectal Cancer Stage-dependent  
591 Global DNA Hypomethylation of Cancer-associated Fibroblasts. *Anticancer Res.*  
592 36(9), 4503–4507 (2016).
- 593 55. Vizoso M, Puig M, Carmona FJ, *et al.* Aberrant DNA methylation in non-small cell  
594 lung cancer-associated fibroblasts. *Carcinogenesis.* 36(12), 1453–1463 (2015).

- 595 56. Zeisberg EM, Zeisberg M. The role of promoter hypermethylation in fibroblast  
596 activation and fibrogenesis. *J. Pathol.* 229(2), 264–273 (2013).
- 597 57. Albrengues J, Bertero T, Grasset E, *et al.* Epigenetic switch drives the conversion of  
598 fibroblasts into proinvasive cancer-associated fibroblasts. *Nature Communications.*  
599 6(1), 10204 (2015).
- 600 \* **This study demonstrated the role of local hypermethylation in CAFs phenotype in**  
601 **breast cancers.**
- 602 58. Bechtel W, McGoohan S, Zeisberg EM, *et al.* Methylation determines fibroblast  
603 activation and fibrogenesis in the kidney. *Nat. Med.* 16(5), 544–550 (2010).
- 604 59. Mishra R, Haldar S, Placencio V, *et al.* Stromal epigenetic alterations drive metabolic  
605 and neuroendocrine prostate cancer reprogramming. *J. Clin. Invest.* 128(10), 4472–  
606 4484 (2018).
- 607 60. Al-Kharashi LA, Al-Mohanna FH, Tulbah A, Aboussekhra A. The DNA methyl-  
608 transferase protein DNMT1 enhances tumor-promoting properties of breast stromal  
609 fibroblasts. *Oncotarget.* 9(2), 2329–2343 (2018).
- 610 61. Li A, Chen P, Leng Y, Kang J. Histone deacetylase 6 regulates the immunosuppressive  
611 properties of cancer-associated fibroblasts in breast cancer through the STAT3-  
612 COX2-dependent pathway. *Oncogene.* 37(45), 5952–5966 (2018).
- 613 62. Garcia-Gomez A, Rodríguez-Ubreva J, Ballestar E. Epigenetic interplay between  
614 immune, stromal and cancer cells in the tumor microenvironment. *Clinical*  
615 *Immunology.* 196, 64–71 (2018).
- 616 63. Kim DJ, Dunleavey JM, Xiao L, *et al.* Suppression of TGFβ-mediated conversion of  
617 endothelial cells and fibroblasts into cancer associated (myo)fibroblasts via HDAC  
618 inhibition. *British Journal of Cancer.* 118(10), 1359–1368 (2018).
- 619 64. Kumar V, Donthireddy L, Marvel D, *et al.* Cancer-Associated Fibroblasts Neutralize  
620 the Anti-tumor Effect of CSF1 Receptor Blockade by Inducing PMN-MDSC Infiltration  
621 of Tumors. *Cancer Cell.* 32(5), 654–668.e5 (2017).
- 622 65. Kim DE, Procopio M-G, Ghosh S, *et al.* Convergent roles of ATF3 and CSL in  
623 chromatin control of cancer-associated fibroblast activation. *J. Exp. Med.* 214(8),  
624 2349–2368 (2017).
- 625 66. Maeda M, Takeshima H, Iida N, *et al.* Cancer cell niche factors secreted from cancer-  
626 associated fibroblast by loss of H3K27me3. *Gut.* gutjnl–2018–317645 (2019).
- 627 67. Zong Y, Huang J, Sankarasharma D, *et al.* Stromal epigenetic dysregulation is  
628 sufficient to initiate mouse prostate cancer via paracrine Wnt signaling. *Proc. Natl.*  
629 *Acad. Sci. U.S.A.* 109(50), E3395–404 (2012).
- 630 68. Xu L, Deng Q, Pan Y, *et al.* Cancer-associated fibroblasts enhance the migration  
631 ability of ovarian cancer cells by increasing EZH2 expression. *International Journal of*

- 632 *Molecular Medicine*. 33(1), 91–96 (2014).
- 633 69. Kaukonen R, Mai A, Georgiadou M, *et al*. Normal stroma suppresses cancer cell  
634 proliferation via mechanosensitive regulation of JMJD1a-mediated transcription.  
635 *Nature Communications*. 7(1), 12237 (2016).
- 636 70. Schoepp M, Ströse AJ, Haier J. Dysregulation of miRNA Expression in Cancer  
637 Associated Fibroblasts (CAFs) and Its Consequences on the Tumor  
638 Microenvironment. *Cancers (Basel)*. 9(6), 54 (2017).
- 639 71. Musumeci M, Coppola V, Addario A, *et al*. Control of tumor and microenvironment  
640 cross-talk by miR-15a and miR-16 in prostate cancer. *Oncogene*. 30(41), 4231–4242  
641 (2011).
- 642 72. Jossen S, Gururajan M, Sung SY, *et al*. Stromal fibroblast-derived miR-409 promotes  
643 epithelial-to-mesenchymal transition and prostate tumorigenesis. *Oncogene*. 34(21),  
644 2690–2699 (2015).
- 645 73. Batista L, Bourachot B, Mateescu B, Reyat F, Mechta-Grigoriou F. Regulation of miR-  
646 200c/141 expression by intergenic DNA-looping and transcriptional read-through.  
647 *Nature Communications*. 7(1), 8959 (2016).
- 648 74. Togashi Y, Shitara K, Nishikawa H. Regulatory T cells in cancer immunosuppression  
649 — implications for anticancer therapy. *Nature Reviews Clinical Oncology*. 155(Suppl),  
650 1 (2019).
- 651 75. Safari E, Ghorghanlu S, Ahmadi-Khiavi H, Mehranfar S, Rezaei R, Motallebnezhad M.  
652 Myeloid-derived suppressor cells and tumor: Current knowledge and future  
653 perspectives. *J. Cell. Physiol*. 117(1), 7021 (2018).
- 654 76. Zhang C, Wang S, Liu Y, Yang C. Epigenetics in myeloid derived suppressor cells: a  
655 sheathed sword towards cancer. *Oncotarget*. 7(35), 57452–57463 (2016).
- 656 77. Heine A, Held SAE, Schulte-Schrepping J, *et al*. Generation and functional  
657 characterization of MDSC-like cells. *Oncoimmunology*. 6(4), e1295203 (2017).
- 658 78. Lal G, Zhang N, van der Touw W, *et al*. Epigenetic Regulation of Foxp3 Expression in  
659 Regulatory T Cells by DNA Methylation. *The Journal of Immunology*. 182(1), 259–273  
660 (2009).
- 661 79. Rodríguez-Ubreva J, Català-Moll F, Obermayer N, *et al*. Prostaglandin E2 Leads to the  
662 Acquisition of DNMT3A-Dependent Tolerogenic Functions in Human Myeloid-  
663 Derived Suppressor Cells. *Cell Reports*. 21(1), 154–167 (2017).
- 664 80. Sido JM, Yang X, Nagarkatti PS, Nagarkatti M.  $\Delta^9$ -Tetrahydrocannabinol-mediated  
665 epigenetic modifications elicit myeloid-derived suppressor cell activation via  
666 STAT3/S100A8. *J. Leukoc. Biol*. 97(4), 677–688 (2015).
- 667 81. Sido JM, Nagarkatti PS, Nagarkatti M.  $\Delta^9$ -Tetrahydrocannabinol attenuates  
668 allogeneic host-versus-graft response and delays skin graft rejection through



- 669 activation of cannabinoid receptor 1 and induction of myeloid-derived suppressor  
670 cells. *J. Leukoc. Biol.* 98(3), 435–447 (2015).
- 671 82. Sahakian E, Powers JJ, Chen J, *et al.* Histone deacetylase 11: A novel epigenetic  
672 regulator of myeloid derived suppressor cell expansion and function. *Mol. Immunol.*  
673 63(2), 579–585 (2015).
- 674 83. Youn J-I, Kumar V, Collazo M, *et al.* Epigenetic silencing of retinoblastoma gene  
675 regulates pathologic differentiation of myeloid cells in cancer. *Nat. Immunol.* 14(3),  
676 211–220 (2013).
- 677 84. Redd PS, Ibrahim ML, Klement JD, *et al.* SETD1B Activates iNOS Expression in  
678 Myeloid-Derived Suppressor Cells. *Cancer Res.* 77(11), 2834–2843 (2017).
- 679 85. Peng D, Kryczek I, Nagarsheth N, *et al.* Epigenetic silencing of TH1-type chemokines  
680 shapes tumour immunity and immunotherapy. *Nature.* 527(7577), 249–253 (2015).
- 681 \* **The epigenetic silencing of EZH2 and DNA methylation in TH1 cells allow re-**  
682 **expression of CXCL9 and CXCL10 and reduce tumor immunosuppression.**
- 683 86. Nagarsheth N, Peng D, Kryczek I, *et al.* PRC2 Epigenetically Silences Th1-Type  
684 Chemokines to Suppress Effector T-Cell Trafficking in Colon Cancer. *Cancer Res.*  
685 76(2), 275–282 (2016).
- 686 87. Guo X, Qiu W, Wang J, *et al.* Glioma exosomes mediate the expansion and function  
687 of myeloid-derived suppressor cells through microRNA-29a/Hbp1 and microRNA-  
688 92a/Prkar1a pathways. *International Journal of Cancer.* ijc.32052 (2018).
- 689 88. Guo X, Qiu W, Liu Q, *et al.* Immunosuppressive effects of hypoxia-induced glioma  
690 exosomes through myeloid-derived suppressor cells via the miR-10a/ Rora and miR-  
691 21/ Pten Pathways. *Oncogene.* 37(31), 4239–4259 (2018).
- 692 89. Huber V, Vallacchi V, Fleming V, *et al.* Tumor-derived microRNAs induce myeloid  
693 suppressor cells and predict immunotherapy resistance in melanoma. *J. Clin. Invest.*  
694 128(12), 5505–5516 (2018).
- 695 90. Zhou J, Li X, Wu X, *et al.* Exosomes Released from Tumor-Associated Macrophages  
696 Transfer miRNAs That Induce a Treg/Th17 Cell Imbalance in Epithelial Ovarian  
697 Cancer. *Cancer Immunol Res.* 6(12), 1578–1592 (2018).
- 698 91. Yoo CB, Jones PA. Epigenetic therapy of cancer: past, present and future. *Nature*  
699 *Reviews Drug Discovery* 2006 5:1. 5(1), 37–50 (2006).
- 700 92. Gherardini L, Sharma A, Capobianco E, Cinti C. Targeting Cancer with Epi-Drugs: A  
701 Precision Medicine Perspective. *Curr Pharm Biotechnol.* 17(10), 856–865 (2016).
- 702 93. Issa J-PJ, Kantarjian HM. Targeting DNA Methylation. *Clin Cancer Res.* 15(12), 3938–  
703 3946 (2009).
- 704 94. Roulois D, Loo Yau H, Singhania R, *et al.* DNA-Demethylating Agents Target

705 Colorectal Cancer Cells by Inducing Viral Mimicry by Endogenous Transcripts. *Cell*.  
706 162(5), 961–973 (2015).

707 \*\* **This study demonstrated the decreasing of colorectal cancers cells proliferation by**  
708 **DNA-demethylating agents which induce a viral mimicry response.**

709 95. Chiappinelli KB, Strissel PL, Desrichard A, *et al.* Inhibiting DNA Methylation Causes an  
710 Interferon Response in Cancer via dsRNA Including Endogenous Retroviruses. *Cell*.  
711 162(5), 974–986 (2015).

712 96. Zhao H, Ning S, Nolley R, *et al.* The immunomodulatory anticancer agent, RRx-001,  
713 induces an interferon response through epigenetic induction of viral mimicry. *Clin*  
714 *Epigenetics*. 9(1), 4 (2017).

715 97. Ghoneim HE, Fan Y, Moustaki A, *et al.* De Novo Epigenetic Programs Inhibit PD-1  
716 Blockade-Mediated T Cell Rejuvenation. *Cell*. 170(1), 142–157.e19 (2017).

717 98. Terracina KP, Graham LJ, Payne KK, *et al.* DNA methyltransferase inhibition increases  
718 efficacy of adoptive cellular immunotherapy of murine breast cancer. *Cancer*  
719 *Immunol. Immunother.* 65(9), 1061–1073 (2016).

720 99. Mikyšková R, Indrová M, Vlková V, *et al.* DNA demethylating agent 5-azacytidine  
721 inhibits myeloid-derived suppressor cells induced by tumor growth and  
722 cyclophosphamide treatment. *J. Leukoc. Biol.* 95(5), 743–753 (2014).

723 100. Yamamoto K, Tateishi K, Kudo Y, *et al.* Stromal remodeling by the BET bromodomain  
724 inhibitor JQ1 suppresses the progression of human pancreatic cancer. *Oncotarget*.  
725 7(38), 61469–61484 (2016).

726 101. Sala L, Franco-Valls H, Stanisavljevic J, *et al.* Abrogation of myofibroblast activities in  
727 metastasis and fibrosis by methyltransferase inhibition. *International Journal of*  
728 *Cancer*. 19, 1423 (2019).

729 102. Stone ML, Chiappinelli KB, Li H, *et al.* Epigenetic therapy activates type I interferon  
730 signaling in murine ovarian cancer to reduce immunosuppression and tumor  
731 burden. *Proc. Natl. Acad. Sci. U.S.A.* 114(51), E10981–E10990 (2017).

732 103. Goswami S, Apostolou I, Zhang J, *et al.* Modulation of EZH2 expression in T cells  
733 improves efficacy of anti-CTLA-4 therapy. *J. Clin. Invest.* 128(9), 3813–3818 (2018).

734 104. Christmas BJ, Rafie CI, Hopkins AC, *et al.* Entinostat Converts Immune-Resistant  
735 Breast and Pancreatic Cancers into Checkpoint-Responsive Tumors by  
736 Reprogramming Tumor-Infiltrating MDSCs. *Cancer Immunol Res.* 6(12), 1561–1577  
737 (2018).

738 105. Briere D, Sudhakar N, Woods DM, *et al.* The class I/IV HDAC inhibitor mocetinostat  
739 increases tumor antigen presentation, decreases immune suppressive cell types and  
740 augments checkpoint inhibitor therapy. *Cancer Immunol. Immunother.* 67(3), 381–  
741 392 (2018).

- 742 106. Lee S, Kim H-S, Roh K-H, *et al.* DNA methyltransferase inhibition accelerates the  
743 immunomodulation and migration of human mesenchymal stem cells. *Sci Rep.* 5,  
744 8020 (2015).
- 745 107. Huang S, Wang Z, Zhou J, *et al.* EZH2 inhibitor GSK126 suppresses anti-tumor  
746 immunity by driving production of myeloid-derived suppressor cells. *Cancer Res.*  
747 *canres.2395.2018* (2019).
- 748 108. Rosborough BR, Castellaneta A, Natarajan S, Thomson AW, Turnquist HR. Histone  
749 deacetylase inhibition facilitates GM-CSF-mediated expansion of myeloid-derived  
750 suppressor cells in vitro and in vivo. *J. Leukoc. Biol.* 91(5), 701–709 (2012).
- 751 109. Reddy P. Editorial: HDAC inhibition begets more MDSCs. *J. Leukoc. Biol.* 91(5), 679–  
752 681 (2012).
- 753 110. Tao R, de Zoeten EF, Ozkaynak E, *et al.* Deacetylase inhibition promotes the  
754 generation and function of regulatory T cells. *Nat. Med.* 13(11), 1299–1307 (2007).
- 755 111. Doñas C, Fritz M, Manríquez V, *et al.* Trichostatin A promotes the generation and  
756 suppressive functions of regulatory T cells. *Clin. Dev. Immunol.* 2013(67), 679804–8  
757 (2013).
- 758 112. Akimova T, Ge G, Golovina T, *et al.* Histone/protein deacetylase inhibitors increase  
759 suppressive functions of human FOXP3+ Tregs. *Clin. Immunol.* 136(3), 348–363  
760 (2010).
- 761 113. Nguyen AH, Elliott IA, Wu N, *et al.* Histone deacetylase inhibitors provoke a tumor  
762 supportive phenotype in pancreatic cancer associated fibroblasts. *Oncotarget.* 8(12),  
763 19074–19088 (2017).
- 764 114. Pazolli E, Alspach E, Milczarek A, Prior J, Piwnica-Worms D, Stewart SA. Chromatin  
765 remodeling underlies the senescence-associated secretory phenotype of tumor  
766 stromal fibroblasts that supports cancer progression. *Cancer Res.* 72(9), 2251–2261  
767 (2012).
- 768 115. Cramer SA, Adjei IM, Labhasetwar V. Advancements in the delivery of epigenetic  
769 drugs. *Expert Opinion on Drug Delivery.* 12(9), 1501–1512 (2015).
- 770 116. Zhu Y, Yu F, Tan Y, Yuan H, Hu F. Strategies of targeting pathological stroma for  
771 enhanced antitumor therapies. *Pharmacological Research.* 148, 104401 (2019).
- 772 117. Schwartzman O, Tanay A. Single-cell epigenomics: techniques and emerging  
773 applications. *Nat. Rev. Genet.* 16(12), 716–726 (2015).
- 774 118. Lee D-S, Luo C, Zhou J, *et al.* Simultaneous profiling of 3D genome structure and DNA  
775 methylation in single human cells. *Nat. Methods.* 62(10), 1–8 (2019).
- 776 119. Kelsey G, Stegle O, Reik W. Single-cell epigenomics: Recording the past and  
777 predicting the future. *Science.* 358(6359), 69–75 (2017).

778

779

Revised manuscript

780

Treatments	Effects	References
DNA methyltransferase inhibitors	<ul style="list-style-type: none"><li>- Activation of “viral mimicry” mechanism</li><li>- Enhances tumor antigens, reverse CAFs phenotype</li><li>- Up regulation of cytokines in CD8 T cells</li><li>- Reactivation of CXCL9/10 chemokine in tumor cells</li></ul>	[57,85,94,95]
Histone methyltransferase inhibitors	<ul style="list-style-type: none"><li>- De-repress CXCL9/10 and effector T cells trafficking</li><li>- Increases immune checkpoint therapy</li></ul>	[85,86,103]
Histone deacetylase inhibitors	<ul style="list-style-type: none"><li>- Decreases of MDSCs and increase CD8+ T cells infiltration in association of immune checkpoint inhibitor</li><li>- Reverses CAFs phenotype</li><li>- Enhances tumor antigen expression</li></ul>	[63,102,104,105]
Chromatin reader inhibitors	<ul style="list-style-type: none"><li>- Reverses CAFs phenotype</li></ul>	[100]

781 **Table 1. Epigenetic drugs impact both tumor and tumor microenvironment**

782

783

**Figure 1: Impact of epigenetic regulation on crosstalk between tumor cells and CAFs, MDSCs and T cells of the microenvironment.** Immunosuppressive properties and tumor supportive effects of microenvironment cells are under the control of epigenetic mechanisms that are deregulated. These deregulations impact: cytokines, chemokines and exosomes secretion. Altogether they favor tumor progression and metastasis, escape to the immune system and resistance to immune checkpoint therapy. Green box contains the up-regulated epigenetic factors and red box the down-regulated.

**Figure 1**

