



HAL
open science

A stiffly stable fully discrete scheme for the damped wave equation using discrete transparent boundary condition

Benjamin Boutin, Thi Hoai Thuong Nguyen, Nicolas Seguin

► **To cite this version:**

Benjamin Boutin, Thi Hoai Thuong Nguyen, Nicolas Seguin. A stiffly stable fully discrete scheme for the damped wave equation using discrete transparent boundary condition. 2020. hal-02566911

HAL Id: hal-02566911

<https://univ-rennes.hal.science/hal-02566911>

Preprint submitted on 7 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A STIFFLY STABLE FULLY DISCRETE SCHEME FOR THE DAMPED WAVE EQUATION USING DISCRETE TRANSPARENT BOUNDARY CONDITION.

BENJAMIN BOUTIN, THỊ HOÀI THUƠNG NGUYỄN, AND NICOLAS SEGUIN

ABSTRACT. We study the stability analysis of the time-implicit central differencing scheme for the linear damped wave equation with boundary. In [23], Xin and Xu prove that the initial-boundary value problem (IBVP) for this model is well-posed, uniformly with respect to the stiffness of the damping, under the so-called stiff Kreiss condition (SKC) on the boundary condition. We show here that the (SKC) is also a sufficient condition to guarantee the uniform stability of the discrete IBVP for the relaxation system independently of the stiffness of the source term, of the space step and of the time step. The boundary is approximated using discrete transparent boundary conditions and the stiff stability is proved using energy estimates and the \mathcal{Z} -transform.

AMS classification: 35F46, 35L50, 65M06, 65M12.

Keywords: hyperbolic relaxation system, damped wave equation, stiff stability, discrete transparent boundary condition, central schemes, implicit scheme, energy estimates, \mathcal{Z} -transform.

1. INTRODUCTION

1.1. Context and motivation. Hyperbolic systems of partial differential equations with relaxation terms [2] are important in many physical situations, such as kinetics theories [6], gases not in thermodynamic equilibrium [21], phase transitions with small transition time [17], water waves [20, 22], reactive flows [8], river flows traffics flows and more general waves [22]. The study of the zero relaxation limit for such systems has caught much interest, both from a theoretical and numerical point of view, after the works of Liu [18], Chen, Levermore and Liu [7], Hanouzet and Natalini [13], Yong [24, 26]. The major issue in the theory of the relaxation approximations to equilibrium system of conservation laws is the appearance of stiff boundary layers in the presence of physical or numerical boundary conditions due to the additional characteristic speeds introduced in the relaxation systems. In this article, we are concerned the development of grid algorithms for solving initial boundary value problem, which involves the question of formulating boundary conditions to get a stable scheme. Due to the presence of boundary layers and to the possible interaction of the boundary and initial layers, numerical schemes have to be properly designed so as to provide accurate approximations and consistent behaviors. One of the simplest linear hyperbolic systems with relaxation is linear damped wave equation in one space dimension

$$(1.1) \quad \begin{cases} \partial_t u^\varepsilon(x, t) + \partial_x v^\varepsilon(x, t) = 0, \\ \partial_t v^\varepsilon(x, t) + a \partial_x u^\varepsilon(x, t) = -\varepsilon^{-1} v^\varepsilon(x, t), \end{cases}$$

where $a > 0$ and the relaxation time $\varepsilon > 0$ may be introduced to characterize the stiffness of the relaxation. When ε goes to zero, the model may be simplified. We expect indeed that for any position x and time t , the solution $(u^\varepsilon, v^\varepsilon)(x, t)$ tends to some $(u(x), 0)$, which is the solution of the corresponding equilibrium system [7, 23].

In order to determine a unique solution to the problem (1.1) in the quarter plane $x > 0, t > 0$, the values of the solution at initial time are prescribed

$$(1.2) \quad u^\varepsilon(x, 0) = u^0(x), \quad v^\varepsilon(x, 0) = v^0(x).$$

In some cases, the suitable boundary condition comes from physical considerations. At a solid wall that bounds the flow of a fluid, for example, one sets the normal component of the fluid velocity equal to zero (if the effects of viscosity are considered, the tangential component must also vanish). In other situations,

Date: May 7, 2020.

The authors B. Boutin and T.H.T. Nguyễn were partially supported by ANR project NABUCO, ANR-17-CE40-0025.

the choice of boundary conditions is not so obvious. This is the case when considering artificial boundaries, which do not correspond to a well-identified physical phenomenon. In general, not any boundary condition is suitable for a given hyperbolic problem. In the case of the problem (1.1), which is a particular case of the Jin-Xin relaxation model in one space dimension [14], the solution at the boundary is imposed

$$(1.3) \quad B_u u^\varepsilon(0, t) + B_v v^\varepsilon(0, t) = b(t),$$

where B_u and B_v are constants. For simplicity, we also assume the initial data $f(x) = (u^0, v^0)(x)$ and the boundary data $b(t)$ to be compatible at the space-time corner $x = 0, t = 0$, i.e.

$$f(0) = f'(0) = 0, \quad b(0) = b'(0) = 0.$$

It is easy to see that the hyperbolic structure is related to the Riemann invariants $\sqrt{a}u^\varepsilon \pm v^\varepsilon$ and to the characteristic velocities $\pm\sqrt{a}$. Therefore, the boundary condition (1.3) has to satisfy the Uniform Kreiss Condition (UKC)

$$(1.4) \quad B_u + \sqrt{a}B_v \neq 0.$$

Only under this assumption, the incoming flow $\sqrt{a}u + v$ at the boundary $x = 0$ can be deduced from the outgoing flow $\sqrt{a}u^\varepsilon - v^\varepsilon$ and the data $b(t)$. Therefore, the initial boundary value problem (IBVP) (1.1)-(1.3) is well-posed for each fixed ε (see [2, 23, 24]).

In [23], Xin and Xu study the asymptotic equivalence of a general linear system of one-dimensional conservation laws and the corresponding relaxation model proposed by Jin and Xin [14] in the limit of a small relaxation rate ε . The main issue is to extend and precise this asymptotic equivalence in the presence of physical boundaries. Within the same problematic, Yong in [24] proposed a Generalized Kreiss Condition (GKC) for general multi-dimensional linear constant coefficient relaxation systems, or one-dimensional nonlinear systems, with non-characteristic boundaries. This condition enables uniform stability estimates and a reduced boundary condition for the corresponding equilibrium system. For the special Jin-Xin system (1.1) with boundary condition (1.3) but with stiff source terms of the form $\varepsilon^{-1}(\lambda u^\varepsilon - v^\varepsilon)$ for some $\lambda \in \mathbb{R}$, Xin and Xu identify and rigorously justify a necessary and sufficient condition (which they call the Stiff Kreiss Condition, or SKC in short) on the boundary condition to guarantee the uniform well-posedness of the IBVP, independently of the relaxation parameter. In addition to the work in [26], their study also covers the characteristic case and provides optimal asymptotic expansions for the limit process, handling with boundary and/or initial layers. In the case of our system (1.1), the parameter $\lambda = 0$ so that the boundary is characteristic for limit equation, and the SKC simply reduces to

$$B_v = 0, \quad \text{or} \quad \frac{B_u}{B_v} \notin [-\sqrt{a}, 0].$$

It is classical to notice that, by linearity, the IBVP (1.1)-(1.3) can be broken up into two simpler problems, one with homogeneous initial condition

$$(1.5) \quad \begin{cases} \partial_t U(x, t) + A \partial_x U(x, t) = \varepsilon^{-1} S U(x, t), & x > 0, t > 0, \\ U(x, 0) = 0, & x > 0, \\ B U(0, t) = b(t), & t > 0, \end{cases}$$

and the other with homogeneous boundary condition

$$(1.6) \quad \begin{cases} \partial_t U(x, t) + A \partial_x U(x, t) = \varepsilon^{-1} S U(x, t), & x > 0, t > 0, \\ U(x, 0) = f(x), & x > 0, \\ B U(0, t) = 0, & t > 0, \end{cases}$$

where

$$U(x, t) = \begin{pmatrix} u(x, t) \\ v(x, t) \end{pmatrix}, \quad A = \begin{pmatrix} 0 & 1 \\ a & 0 \end{pmatrix}, \quad S = \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix}, \quad B = (B_u \ B_v).$$

The main difficult part of the proof of stability under the SKC appears with the study of (1.6). This is due to the complex interactions between the initial data, the boundary condition and the stiff relaxation term. By means of the energy method, Xin and Xu [23] show that the time evolution of the energy

$\|U(t)\|_{L^2(\mathbb{R}^+, \mathbb{R}^2)}$ decreases if the boundary condition (1.3) satisfies $B_u B_v > 0$, which is only the subclass of the SKC, so that this analysis is not satisfactory. Actually, under the SKC, they also find explicitly the solution $U(x, t)$ of the IBVP (1.6) by using the Laplace transform. The solution is decomposed into two parts, by assembling the solution for the case of the Cauchy problem without boundary and another one for the case of the IBVP but with homogeneous initial condition [23, Section 5]. It means that the issue of stiff well-posedness of the IBVP (1.1)-(1.3) is solved if we can prove the Cauchy problem and the IBVP with homogeneous initial condition, that is to say (1.5), themselves are stiffly well-posed.

The motivation of the present study is to analyze the counterpart of the above results for the fully discrete difference approximation of the IBVP (1.1)-(1.3). Because of the propagation of waves of the problem (1.1) with the characteristic velocities $\pm\sqrt{a}$, one of the waves at the boundary $x = 0$ is coming into the computation domain while the other one is going out of it. Thus, the way of formulating boundary conditions for the relaxation systems so as to guarantee the uniform stability and to minimize the artificial boundary layer is a crucial issue to the success of the schemes. In [5], the boundary is approximated using a summation-by-parts method. Using energy estimates and Laplace transforms, the semi-discrete approximation of the IBVP (1.1)-(1.3) is proved to be stiffly stable if the boundary condition (1.3) satisfies $B_u B_v > 0$, which is only the subclass of the SKC. In this article, we consider the discrete transparent boundary technique to construct a stiffly stable boundary condition. The technique and its analysis has been proposed by Arnold and Ehrhardt in [1]. Besse, Noble and co-authors apply the tools to dispersive problems [3, 4, 16]. We also refer the reader to [11] for non-reflecting methods in the context of wave problems. The recent work of Coulombel [9] proposes a systematic study of transparent boundary conditions for evolution problems. Our aim here is to prove that the SKC derived in [23] is a sufficient condition for the stiff stability of the proposed fully discrete of the IBVP (1.1)-(1.3).

1.2. Description of the numerical scheme. Let $\Delta t > 0$ being the time step. The space step $\Delta x > 0$ will always be chosen so that the parameter $\lambda_{xt} = \Delta x \Delta t^{-1}$ is kept fixed. Letting now $U_j^n = (u_j^n, v_j^n)^T$ denotes the approximation of the exact solution to (1.1)-(1.3) at the grid point $(x_j, t^n) = (j\Delta x, n\Delta t)$, for any $(j, n) \in \mathbb{N} \times \mathbb{N}$ (where we omit the explicit dependence on ε). We focus in this paper on the fully discrete approximation of the IBVP (1.1)-(1.3) obtained by the central differencing scheme in space and the implicit scheme in time.

A first step towards the fully discrete approximation of the IBVP (1.1)-(1.3) is the following system

$$(1.7) \quad \begin{cases} \frac{U_j^{n+1} - U_j^n}{\Delta t} + \frac{1}{2\Delta x} (U_{j+1}^{n+1} - U_{j-1}^{n+1}) = \frac{1}{\varepsilon} S U_j^{n+1}, & j \geq 1, n \geq 0, \\ U_j^0 = f_j, & j \geq 0, \\ B U_0^n = b^n, & n \geq 0, \end{cases}$$

where the approximations of the initial condition f_j and of the boundary data b^n are defined for example by setting $f_j = f(j\Delta x)$ for $j \geq 0$ and $b^n = b(n\Delta t)$ for $n \geq 0$.

Let us emphasize that the numerical scheme (1.7) still needs one more scalar equation at the boundary point $j = 0$ so as to be fully defined, due to the fact that the matrix B has rank one only. This is actually a discrete feature only, since in the continuous case this single equation is exactly complemented by the only incoming characteristic (at least under UKC). An additional relation to define U_0^{n+1} is thus needed. We want to use the central scheme at the boundary point $j = 0$, so that the modification of the ghost value U_{-1}^{n+1} can also be interpreted as the use of an extra boundary condition. From a mathematical point of view, the problem is set, in both cases, as follows: given an initial data compactly supported, one can construct boundary condition at $j = 0$ with the objective to approximate the exact solution of the whole space problem $\{j \in \mathbb{Z}\}$, restricted to $\{j \in \mathbb{N}\}$. If the approximate solution on $\{j \in \mathbb{N}\}$ coincides with the exact solution, one refers to these boundary conditions as transparent boundary conditions. Of course, these boundary condition should lead to a well-posed initial boundary value problem. It means that we use the discrete transparent boundary condition at $j = 0$ that determines a ghost value U_{-1}^{n+1} through the

identity

$$U_{-1}^{n+1} = \sum_{k=0}^{n+1} \mathcal{C}_{n+1-k} U_0^k,$$

where the coefficients \mathcal{C}_k will be precised explicitly in the forthcoming Definition 2.4. The extra boundary condition determines U_{-1}^{n+1} as a linear function of U_0^k for past step times only: $0 \leq k \leq n+1$. We propose the following numerical approximation at the boundary:

$$\frac{1}{\Delta t} \Gamma (U_0^{n+1} - U_0^n) + \frac{1}{2\Delta x} \Gamma A \left(U_1^{n+1} - \sum_{k=0}^{n+1} \mathcal{C}_{n+1-k} U_0^k \right) = \frac{1}{\varepsilon} \Gamma S U_0^{n+1}$$

with the matrix $\Gamma = (-aB_v \quad B_u)$. Under the SKC, this choice for the matrix Γ will be useful to construct the numerical solution $(U_j^n)_{j \in \mathbb{N}}$ in the Propositions 2.3 and 3.1.

To summarize, we study all along the paper the following fully discrete approximation of the IBVP (1.1)-(1.3):

$$(1.8) \quad \begin{cases} \frac{U_j^{n+1} - U_j^n}{\Delta t} + \frac{1}{2\Delta x} A \left(U_{j+1}^{n+1} - U_{j-1}^{n+1} \right) = \frac{1}{\varepsilon} S U_j^{n+1}, & j \geq 1, n \geq 0, \\ U_j^0 = f_j, & j \geq 0, \\ B U_0^n = b^n, & n \geq 0, \\ \frac{1}{\Delta t} \Gamma (U_0^{n+1} - U_0^n) + \frac{1}{2\Delta x} \Gamma A \left(U_1^{n+1} - \sum_{k=0}^{n+1} \mathcal{C}_{n+1-k} U_0^k \right) = \frac{1}{\varepsilon} \Gamma S U_0^{n+1}, & n \geq 0. \end{cases}$$

1.3. Main result. Dealing with the continuous IBVP (1.1)-(1.3), the UKC (1.4) is not enough and a more stringent restriction has to be imposed. Our aim is to prove that the SKC derived in [23] is then a sufficient condition for the stiff stability of the fully discrete IBVP (1.8), in other words the uniform stability with respect to the stiffness of the relaxation term.

Theorem 1.1 (Main result). *Assume that $(B_u, B_v) \in \mathbb{R}^2$ satisfies the SKC*

$$(1.9) \quad B_v = 0 \quad \text{or} \quad \frac{B_u}{B_v} \notin [-\sqrt{a}, 0].$$

Let $\lambda_{xt} \leq 3\sqrt{a}/8$ be a positive number. For any $T > 0$, there exists a constant $C_T > 0$ such that for all $\Delta t > 0$ and $\Delta x = \lambda_{xt} \Delta t$, any $(f_j)_{j \in \mathbb{N}} \in \ell^2(\mathbb{N}, \mathbb{R}^2)$ and $(b^n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N}, \mathbb{R})$, the solution $(U_j^n)_{j \in \mathbb{N}}$ to the scheme (1.8) satisfies

$$(1.10) \quad \sum_{n=0}^N \sum_{j \geq 0} \Delta x \Delta t |U_j^n|^2 + \sum_{n=0}^N \Delta t |U_0^n|^2 \leq C_T \left(\sum_{j \geq 0} \Delta x |f_j|^2 + \sum_{n=0}^N \Delta t |b^n|^2 \right),$$

where $N := T/\Delta t$ and C_T is independent of $\varepsilon \in (0, +\infty)$.

In [23], Xin and Xu considered the IBVP for the Jin-Xin relaxation model [14] and derived the SKC (1.9) to characterize its stiff well-posedness. They show in particular that the IBVP (1.1)-(1.3) is well-posed if and only if (1.9) holds. In the discrete IBVP (1.8), it seems that the SKC is also sufficient to derive uniform stability estimates. Besides, by linearity, the numerical scheme of the IBVP (1.8) can be broken up into two simpler problems, one with homogeneous initial condition $(f_j)_{j \in \mathbb{N}} \equiv 0$ and the other with homogeneous boundary $b^n \equiv 0$, for any $n \in \mathbb{N}$. The proof of Theorem 1.1 is based on two main ingredients, by assembling a result for the case of the following Cauchy problem

$$(1.11) \quad \begin{cases} \frac{(U_j^I)^{n+1} - (U_j^I)^n}{\Delta t} + \frac{1}{2\Delta x} A \left((U_{j+1}^I)^{n+1} - (U_{j-1}^I)^{n+1} \right) = \frac{1}{\varepsilon} S (U_j^I)^{n+1}, & j \in \mathbb{Z}, n \geq 0, \\ (U_j^I)^0 = f_j, & j \in \mathbb{Z}. \end{cases}$$

and another one for the problem (1.8) with homogeneous initial data. We state hereafter these two statements.

Proposition 1.2 (Cauchy problem). *For any $T > 0$, there exists $C_T > 0$ such that for all $\Delta t > 0$, any $(f_j)_{j \in \mathbb{N}} \in \ell^2(\mathbb{N}, \mathbb{R}^2)$, the solution $(U_j^I)_{j \in \mathbb{Z}}$ to (1.11) satisfies*

$$(1.12) \quad \sum_{j \in \mathbb{Z}} \Delta x |(U_j^I)^n|^2 \leq C_T \sum_{j \in \mathbb{Z}} \Delta x |f_j|^2, \quad n \in \mathbb{N},$$

where C_T is independent of $\varepsilon \in (0, +\infty)$ and $\Delta x = \lambda_{xt} \Delta t$.

Proposition 1.3 (Homogeneous initial condition). *Assume that the SKC (1.9) is satisfied. Then, there exists a constant $C > 0$ such that for any $\gamma > 0$ and any positive constant $\lambda_{xt} \leq 3\sqrt{a}/8$, the following property holds. For any $\Delta t > 0$ together with $\Delta x = \lambda_{xt} \Delta t$, and any boundary data $(b^n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N}, \mathbb{R})$, the solution $(U_j^n)_{j \in \mathbb{N}}$ to (1.8) with $(f_j)_{j \in \mathbb{N}} \equiv 0$ satisfies*

$$(1.13) \quad \frac{\gamma}{\gamma \Delta t + 1} \sum_{n \geq 0} \sum_{j \geq 0} e^{-2\gamma n \Delta t} \Delta t \Delta x |U_j^n|^2 + \sum_{n \geq 0} e^{-2\gamma n \Delta t} \Delta t |U_0^n|^2 \leq C \sum_{n \geq 0} e^{-2\gamma n \Delta t} \Delta t |b^n|^2,$$

where C is independent of $\varepsilon \in (0, +\infty)$.

To isolate the effects of a possible boundary layer and avoid the complicated interaction of boundary and initial layers, in Section 2, we consider the IBVP (1.8) with homogeneous initial data and nonzero boundary data b^n , for any $n \geq 0$. The numerical solution $(U_j^n)_{j \in \mathbb{N}}$ is constructed in Section 2.2 thanks to the \mathcal{Z} -transform [15, 19]. Furthermore, we follow the discrete transparent boundary condition at $j = 0$ as proposed in [1, 4, 16] to find the explicit formula of the sequence $(\mathcal{C}_m)_{m \geq 0}$. By using the Plancherel's theorem, under the SKC, the Proposition 1.3 is proved in Section 2.3. In order to illustrate the relevance of the SKC (1.9), we present in Section 2.4 some numerical results, for various values of the parameters (B_u, B_v) and show that the numerical solution at the boundary $x = 0$ increases quickly if the SKC (1.9) does not hold. Besides, by the decrease of the error $\|U(\cdot, t^n) - U^n\|_{\ell^2(\mathbb{N}, \mathbb{R}^2)}^2$, we can observe the convergence of the discrete solution U_j^n to the exact one $U(x_j, t^n)$. After that, we observe the behavior of the energy terms $\|U\|_{\ell^2(\mathbb{N} \times [0, T], \mathbb{R}^2)}$ and $\|U\|_{\ell^2(\{0\} \times [0, T], \mathbb{R}^2)}$ corresponding to whether or not the SKC (1.9) is valid. The nonzero initial data case is much more difficult with the sufficiency proof. This is due to the complicated interactions between the initial data, the boundary condition and the stiff relaxation term. Under the SKC, the numerical solution is again described by means of the \mathcal{Z} -transform in Section 3.1. It is decomposed into three parts, by assembling the solution for the case of Cauchy problem (1.11), a numerical error term $(U_j^{II})_{j \in \mathbb{N}}^n$ and the solution for the case IBVP (1.8) with homogeneous initial data. Since the coefficients for computing the boundary value U_{-1}^{n+1} are defined for homogeneous initial data, this numerical error $(U_j^{II})_{j \in \mathbb{N}}^n$ is due to the interaction between the Cauchy problem and the IBVP with zero initial data. For the Cauchy problem, the Proposition 1.2 is studied in Section 3.2 by means of the discrete energy method. By an application of the Plancherel's theorem for \mathcal{Z} -transform, the numerical error term $(U_j^{II})_{j \in \mathbb{N}}^n$ will be estimated in Section 3.3. In Section 3.4, we get the expected result of the Theorem 1.1 in the case IBVP with nonzero initial condition. In Section 3.5, we also look at the behavior of the numerical solution $(U_j^n)_{j \in \mathbb{Z}}$ and the energy terms $\|U\|_{\ell^2(\mathbb{N} \times [0, T], \mathbb{R}^2)}$ and $\|U\|_{\ell^2(\{0\} \times [0, T], \mathbb{R}^2)}$ corresponding to whether or not the SKC (1.9) is valid. It seems that the SKC (1.9) is also necessary condition to guarantee the uniform stability of the IBVP (1.8) independent of the effect of the relaxation source term and the boundary dissipation.

2. STIFF STABILITY OF THE IBVP WITH HOMOGENEOUS INITIAL CONDITION

In this section, we consider the discrete IBVP (1.8) with nonzero boundary condition $(b^n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N}, \mathbb{R})$ and homogeneous Cauchy data $(f_j)_{j \in \mathbb{N}} \equiv 0$. Assuming that the SKC is satisfied, the numerical solution $(U_j^n)_{j \in \mathbb{N}}$ is obtained by using the \mathcal{Z} -transform [15, 19]. Thanks to the Plancherel's theorem, we then are able to get the expected result of the Proposition 1.3.

2.1. Notations and preliminary results. Before we enter the important proofs, let us introduce some notations and preliminary results. All along the paper, the complex values z and ξ are related through the formula

$$\xi = (1 - z^{-1}) \Delta t^{-1}, \quad z = R e^{i\theta}, \quad \text{with } R > 1, \theta \in (-\pi, \pi].$$

Then, ξ obeys the inequalities

$$(2.1) \quad (1 - R^{-1}) \Delta t^{-1} \leq \text{Re } \xi \leq 2\Delta t^{-1}.$$

Besides, one also introduces the following matrix, already concerned with the continuous case [23]:

$$M(\varepsilon\xi) = A^{-1}(S - \varepsilon\xi I) = \frac{1}{a} \begin{pmatrix} 0 & -(1 + \varepsilon\xi) \\ -a\varepsilon\xi & 0 \end{pmatrix},$$

We recall that the eigenvalues and eigenvectors of $M(\varepsilon\xi)$ can be easily found to be respectively

$$\mu_{\pm}(\varepsilon\xi) = \pm \sqrt{\frac{\varepsilon\xi(1 + \varepsilon\xi)}{a}}, \quad r_{\pm}(\varepsilon\xi) = \begin{pmatrix} 1 \\ \frac{a\mu_{\mp}(\varepsilon\xi)}{1 + \varepsilon\xi} \end{pmatrix}.$$

In the above formula and all along the paper, the complex square root is defined with the branch cut along the negative real axis. Applying Lemma A.1 with the property $\text{Re } \xi > 0$ and $\varepsilon > 0$, we can prove

$$(2.2) \quad \text{Re}(\mu_{-}(\varepsilon\xi)) \leq -\frac{\varepsilon \text{Re } \xi}{\sqrt{a}} < 0,$$

while, as a consequence,

$$\text{Re}(\mu_{+}(\varepsilon\xi)) \geq \frac{\varepsilon \text{Re } \xi}{\sqrt{a}} > 0.$$

Let us introduce

$$(2.3) \quad \kappa_{\pm}(\varepsilon\xi) = \mu_{\pm}(\varepsilon\xi)\lambda_{x\varepsilon} + \sqrt{(\mu_{\pm}(\varepsilon\xi)\lambda_{x\varepsilon})^2 + 1},$$

with the notation $\lambda_{x\varepsilon} = \Delta x/\varepsilon$. According to Lemma A.2 together with the properties $\text{Re}(\mu_{-}(\varepsilon\xi)) < 0$ for $\varepsilon > 0$ and $\text{Re } \xi > 0$, we can prove $|\kappa_{-}(\varepsilon\xi)| < 1$. Besides, since $\mu_{-}(\varepsilon\xi) = -\mu_{+}(\varepsilon\xi)$, we get $\kappa_{+}(\varepsilon\xi)\kappa_{-}(\varepsilon\xi) = 1$. As a consequence, for any $\varepsilon > 0$ and $\text{Re } \xi > 0$, one has the separation property $|\kappa_{+}(\varepsilon\xi)| > 1$.

We further define the following spectral projections

$$(2.4) \quad \begin{aligned} \Phi_{+}(\varepsilon\xi) &= \frac{1}{2g(\varepsilon\xi)} \begin{pmatrix} 1 & \\ & -g(\varepsilon\xi) \end{pmatrix} \begin{pmatrix} g(\varepsilon\xi) & -1 \\ & \end{pmatrix}, \\ \Phi_{-}(\varepsilon\xi) &= \frac{1}{2g(\varepsilon\xi)} \begin{pmatrix} 1 & \\ & g(\varepsilon\xi) \end{pmatrix} \begin{pmatrix} g(\varepsilon\xi) & 1 \\ & \end{pmatrix}, \end{aligned}$$

where we set

$$(2.5) \quad g(\varepsilon\xi) = \frac{a\mu_{+}(\varepsilon\xi)}{1 + \varepsilon\xi}.$$

We also set $\Phi(\varepsilon\xi)$ the 2×2 matrix whose columns are composed by the components of the eigenvectors of the matrix $M(\varepsilon\xi)$. We recall these matrices thus satisfy the following usefull identities

$$(2.6) \quad \Phi_{+}(\varepsilon\xi) = \Phi(\varepsilon\xi) \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \Phi^{-1}(\varepsilon\xi) \quad \text{and} \quad \Phi_{-}(\varepsilon\xi) = \Phi(\varepsilon\xi) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \Phi^{-1}(\varepsilon\xi)$$

and

$$(2.7) \quad \Phi_{+}^2(\varepsilon\xi) = \Phi_{+}(\varepsilon\xi), \quad \Phi_{-}^2(\varepsilon\xi) = \Phi_{-}(\varepsilon\xi), \quad \Phi_{+}(\varepsilon\xi)\Phi_{-}(\varepsilon\xi) = \Phi_{-}(\varepsilon\xi)\Phi_{+}(\varepsilon\xi) = 0.$$

In order later on to construct and estimate the numerical solution $(U_j^n)_{j \in \mathbb{N}}$ by the \mathcal{Z} -transform, the following lemmas are usefull:

Lemma 2.1. [From [23]] Consider $\mathbb{C}_{+} = \{\zeta \in \mathbb{C}, \text{Re } \zeta \geq 0\}$ the closed complex right half-plane. Under the SKC (1.9), the quantity $g(\zeta)$ is uniformly bounded in \mathbb{C}_{+} and the quantity $B_u + g(\zeta)B_v$ is uniformly bounded away from 0 in \mathbb{C}_{+} .

We omit the proof that the reader can find in the work by Xin and Xu [23].

Lemma 2.2. *Let us consider the 4×4 matrix*

$$(2.8) \quad M_1(\varepsilon\xi) = \begin{pmatrix} 2\lambda_{x\varepsilon}M(\varepsilon\xi) & I \\ I & 0 \end{pmatrix}.$$

Then, the k -th power of $M_1(\varepsilon\xi)$ reads also

$$(2.9) \quad M_1^k(\varepsilon\xi) = -\frac{1}{\kappa_+(\varepsilon\xi) + \kappa_-(\varepsilon\xi)} \begin{pmatrix} \widehat{\kappa}_{k+1}(\varepsilon\xi)\widehat{\Psi}_k(\varepsilon\xi) & \widehat{\kappa}_k(\varepsilon\xi)\widehat{\Psi}_{k+1}(\varepsilon\xi) \\ \widehat{\kappa}_k(\varepsilon\xi)\widehat{\Psi}_{k+1}(\varepsilon\xi) & \widehat{\kappa}_{k-1}(\varepsilon\xi)\widehat{\Psi}_k(\varepsilon\xi) \end{pmatrix},$$

where

$$(2.10) \quad \begin{aligned} \widehat{\kappa}_k(\varepsilon\xi) &= (-1)^k \kappa_+^k(\varepsilon\xi) - \kappa_-^k(\varepsilon\xi), \\ \widehat{\Psi}_k(\varepsilon\xi) &= \Phi_-(\varepsilon\xi) + (-1)^k \Phi_+(\varepsilon\xi). \end{aligned}$$

Proof. In this algebraic proof, we skip the dependence on $\varepsilon\xi$. Since the columns of the matrix Φ are composed by the components of the eigenvectors of the matrix M , the considered matrix M_1^k can be reformulated simply as

$$(2.11) \quad M_1^k = \widehat{\Phi} M_2^k \widehat{\Phi}^{-1},$$

where

$$\widehat{\Phi} = \begin{pmatrix} \Phi & 0 \\ 0 & \Phi \end{pmatrix}, \quad M_2 = \begin{pmatrix} D_1 & I \\ I & 0 \end{pmatrix}, \quad D_1 = 2\lambda_{x\varepsilon} \text{diag}(\mu_-, \mu_+).$$

Let Ψ is the 4×4 matrix whose columns are composed by the components of the eigenvectors of the matrix M_2

$$\Psi = \begin{pmatrix} -\kappa_+ & \kappa_- & 0 & 0 \\ 0 & 0 & -\kappa_- & \kappa_+ \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$

so that $M_2^k = \Psi D_2^k \Psi^{-1}$ with $D_2 = \text{diag}(-\kappa_+, \kappa_-, -\kappa_-, \kappa_+)$. Therefore, the formula M_1^k in (2.11) reads

$$(2.12) \quad M_1^k = \widehat{\Phi} \Psi D_2^k \Psi^{-1} \widehat{\Phi}^{-1}.$$

By using the properties Φ_{\pm} in (2.6) and (2.12), one obtains

$$M_1^k = -\frac{1}{\kappa_+ + \kappa_-} \begin{pmatrix} \widehat{\kappa}_{k+1}\widehat{\Psi}_k & \widehat{\kappa}_k\widehat{\Psi}_{k+1} \\ \widehat{\kappa}_k\widehat{\Psi}_{k+1} & \widehat{\kappa}_{k-1}\widehat{\Psi}_k \end{pmatrix}$$

with $\widehat{\kappa}_k$ and $\widehat{\Psi}_k$ are the same as in (2.10). □

2.2. Solution by \mathcal{Z} -transform. Firstly, we apply the \mathcal{Z} -transform with respect to time index $n \in \mathbb{N}$, which is discrete analogue of the Laplace transform in time $t \in \mathbb{R}_+$. This method enables the representation and estimations of the numerical solution $(U_j^n)_{j \in \mathbb{N}}$. The definition reads as follows (see [15, 19] for more details)

$$\widehat{U}_j(z) = \mathcal{Z}\{U_j^n\}(z) = \sum_{n \geq 0} U_j^n z^{-n}, \quad |z| > 1.$$

Since we assume $(U_j^0)_{j \in \mathbb{N}} \equiv 0$, observe that the \mathcal{Z} -transform of the time-shifted numerical solution reads

$$\sum_{n \geq 0} U_j^{n+1} z^{-n} = z\widehat{U}_j(z) - zU_j^0 = z\widehat{U}_j(z).$$

Therefore, the IBVP (1.8) with zero initial data becomes

$$(2.13a) \quad \begin{cases} \widehat{U}_{j+1}(z) - \widehat{U}_{j-1}(z) = 2\lambda_{x\varepsilon}M(\varepsilon\xi)\widehat{U}_j(z), & j \geq 1, \\ B\widehat{U}_0(z) = \widehat{b}(z), \\ \Gamma A \left(\widehat{U}_1(z) - z^{-1}\Upsilon(\widehat{U}_0(z)) - 2\lambda_{x\varepsilon}M(\varepsilon\xi)\widehat{U}_0(z) \right) = 0, \end{cases}$$

where $\Upsilon(\widehat{U}_0(z))$ is the \mathcal{Z} -transform of the sequence $\left\{ \sum_{k=0}^{n+1} c_{n+1-k}U_0^k \right\}_{n \geq 0}$ and \widehat{b} stands for the \mathcal{Z} -transform of the scalar boundary data: $\widehat{b}(z) = \mathcal{Z}\{b^n\}(z) = \sum_{n \geq 0} b^n z^{-n}$.

Secondly, we look at the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ to (2.13a)-(2.13c). This is the object of the following proposition:

Proposition 2.3. *Assume that the SKC (1.9) is satisfied. Assume that Γ and Υ in the boundary condition (2.13c) are defined by*

$$(2.14) \quad \Gamma = \begin{pmatrix} -aB_v & B_u \end{pmatrix}, \quad \Upsilon(\widehat{U}_0(z)) = \kappa_+(\varepsilon\xi)z\widehat{U}_0(z).$$

Then the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z) \in \ell^2(\mathbb{N}, \mathbb{C}^2)$ to (2.13a)-(2.13c) takes the form

$$(2.15) \quad \widehat{U}_j(z) = \frac{\widehat{b}(z)}{B_u + g(\varepsilon\xi)B_v} \kappa_-^j(\varepsilon\xi)r_-(\varepsilon\xi).$$

Proof. Before we prove the above result, let us notice that we omit the explicit dependence in $\varepsilon\xi$. Firstly, we look at the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ to (2.13a) and consider the two-dimensional problem (2.13a) under the following one-step recurrence form

$$(2.16) \quad W_{j+1}(z) = M_1 W_j(z),$$

where M_1 is given by (2.8) and

$$(2.17) \quad W_j(z) = \begin{pmatrix} \widehat{U}_j(z) \\ \widehat{U}_{j-1}(z) \end{pmatrix}.$$

The solution $(W_j)_{j \in \mathbb{N}}(z)$ to (2.16) is simply $W_j(z) = M_1^j W_0(z)$. Together with the the explicit formula of M_1^j in Lemma 2.2, the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ to (2.13a) is therefore given by

$$\widehat{U}_j(z) = -\frac{1}{\kappa_+ + \kappa_-} \times \left(\widehat{\kappa}_{j+1} \widehat{\Psi}_j \widehat{U}_0(z) + \widehat{\kappa}_j \widehat{\Psi}_{j+1} \widehat{U}_{-1}(z) \right).$$

By using the definition of $\widehat{\kappa}_k$ and $\widehat{\Psi}_k$ in (2.10), the above formula is now equivalent to

$$(2.18) \quad \begin{aligned} \widehat{U}_j(z) = & -\frac{(-1)^j \kappa_+^j}{\kappa_+ + \kappa_-} \times \left[\Phi_- \left(-\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) \right) + (-1)^{j+1} \Phi_+ \left(\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) \right) \right] \\ & + \frac{\kappa_-^j}{\kappa_+ + \kappa_-} \times \left[\kappa_- \left(\Phi_- + (-1)^j \Phi_+ \right) \widehat{U}_0(z) + \left(\Phi_- + (-1)^{j+1} \Phi_+ \right) \widehat{U}_{-1}(z) \right]. \end{aligned}$$

Since we expect $(\widehat{U}_j)_{j \in \mathbb{N}}(z) \in \ell^2(\mathbb{N}, \mathbb{C}^2)$, we need a natural boundary condition at $x = +\infty$. Besides, one gets $|\kappa_+| > 1$ and $|\kappa_-| < 1$. Thus, the natural boundary condition takes the form

$$(2.19) \quad \begin{cases} \Phi_- \left(-\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) \right) = 0, \\ \Phi_+ \left(\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) \right) = 0. \end{cases}$$

By the definition of Φ_{\pm} in (2.4), the system (2.19) is equivalent to

$$\begin{cases} (g, 1) \left(-\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) \right) = 0, \\ (g, -1) \left(\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) \right) = 0. \end{cases}$$

Then, we have

$$\widehat{U}_{-1}(z) = \frac{\kappa_+}{g} \times \begin{pmatrix} 0 & 1 \\ g^2 & 0 \end{pmatrix} \widehat{U}_0(z).$$

Furthermore, we can see that

$$\Phi_- - \Phi_+ = \frac{1}{g} \begin{pmatrix} 0 & 1 \\ g^2 & 0 \end{pmatrix}.$$

Thus,

$$(2.20) \quad \widehat{U}_{-1}(z) = \kappa_+(\Phi_- - \Phi_+)\widehat{U}_0(z).$$

Plugging (2.20) into (2.18), we have

$$\widehat{U}_j(z) = \frac{\kappa_-^j}{\kappa_+ + \kappa_-} \left[\kappa_- \left(\Phi_- + (-1)^j \Phi_+ \right) + \kappa_+ \left(\Phi_- + (-1)^{j+1} \Phi_+ \right) \left(\Phi_- - \Phi_+ \right) \right] \widehat{U}_0(z).$$

Under the properties of Φ_{\pm} in (2.7), the above formula becomes

$$(2.21) \quad \widehat{U}_j(z) = \kappa_-^j \left(\Phi_- + (-1)^j \Phi_+ \right) \widehat{U}_0(z).$$

Secondly, we look at the boundary condition (2.13b) and (2.13c). Under the choice $\Upsilon(\widehat{U}_0(z))$ in (2.14), the boundary condition (2.13c) becomes

$$(2.22) \quad \Gamma A \left(\widehat{U}_1(z) - (\kappa_+ I + 2\lambda_{x\varepsilon} M) \widehat{U}_0(z) \right) = 0.$$

Indeed, we can compute separately

$$(2.23) \quad \begin{aligned} \kappa_+ I + 2\lambda_{x\varepsilon} M &= \kappa_- \Phi_- + (\kappa_+ + 2\lambda_{x\varepsilon} \mu_+) \Phi_+, \\ \widehat{U}_1(z) &= \kappa_- (\Phi_- - \Phi_+) \widehat{U}_0(z). \end{aligned}$$

Substituting (2.23) into (2.22), one obtains

$$(2.24) \quad \Gamma A \Phi_+ \widehat{U}_0(z) = 0.$$

Under the choice Γ in (2.14), we have

$$\Gamma A \Phi_+ = \frac{a}{2g} \times (B_u + gB_v) \times (g, -1).$$

Thus,

$$(2.25) \quad (B_u + gB_v) \times (g, -1) \widehat{U}_0(z) = 0.$$

From the Lemma 2.1, the equation (2.25) is equivalent under the SKC to

$$(2.26) \quad (g, -1) \widehat{U}_0(z) = 0.$$

Together with the boundary condition (2.13b), the value of $\widehat{U}_0(z)$ has to satisfy

$$\begin{pmatrix} B_u & B_v \\ g & -1 \end{pmatrix} \widehat{U}_0(z) = \widehat{b}(z) \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Then, again under the SKC, we have

$$(2.27) \quad \widehat{U}_0(z) = \frac{\widehat{b}(z)}{B_u + g(\varepsilon\xi)B_v} r_-(\varepsilon\xi).$$

Plugging the value of $\widehat{U}_0(z)$ in (2.27) into (2.21), the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ to (2.13a)-(2.13c) is given by

$$\widehat{U}_j(z) = \frac{\widehat{b}(z)}{B_u + gB_v} \kappa_-^j \left(\Phi_- + (-1)^j \Phi_+ \right) r_-(\varepsilon\xi).$$

Since $\Phi_+ r_- = 0$ and $\Phi_- r_- = r_-$, the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ finally is

$$\widehat{U}_j(z) = \frac{\widehat{b}(z)}{B_u + gB_v} \kappa_-^j r_-.$$

This ends the proof of Proposition 2.3. \square

With $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ found in (2.15), the numerical solution $(U_j^n)_{j \in \mathbb{N}}$ to the IBVP (1.8) with nonzero boundary condition $(b^n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N}, \mathbb{R})$ and homogeneous Cauchy data $(f_j)_{j \in \mathbb{N}} \equiv 0$ can be obtained by inverting the \mathcal{Z} -transform [15, 19]

$$U_j^n = \frac{1}{2\pi} \int_{-\pi}^{\pi} \widehat{U}_j(Re^{i\theta}) R^n e^{in\theta} d\theta, \quad R > 1.$$

Let us remark that an important assumption of Proposition 2.3 is $\Upsilon(\widehat{U}_0(z)) = \kappa_+(\varepsilon\xi)z\widehat{U}_0(z)$. We now follow the discrete transparent boundary as proposed in [1, 4, 16] to find the explicit formula for the sequence $(\mathcal{C}_m)_{m \geq 0}$.

Definition 2.4. Let $\varepsilon > 0$, $R > 1$, $\theta \in (-\pi, \pi]$ and then $\kappa_+(\varepsilon\xi)$ be given by (2.3). The value of $(\mathcal{C}_m)_{m \geq 0}$ is defined as follows:

$$(2.28) \quad \mathcal{C}_m = \frac{1}{\pi} \int_0^{\pi} \operatorname{Re} \left(\kappa_+(\varepsilon\xi) R^m e^{im\theta} \right) d\theta.$$

Let us mention at this step that the values \mathcal{C}_m above are designed in the case of homogeneous initial data and are kept unchanged in the case of nonzero initial data in forthcoming Section 3. Thanks to the convolution property and inverting \mathcal{Z} -transform, we now show that the definition of $(\mathcal{C}_m)_{m \geq 0}$ in (2.28) is the suitable choice to get the required identity $\Upsilon(\widehat{U}_0(z)) = \kappa_+(\varepsilon\xi)z\widehat{U}_0(z)$ in Proposition 2.3. This is the object of the next lemma:

Lemma 2.5. Let $(\mathcal{C}_m)_{m \geq 0}$ be defined from Definition 2.4, then

$$\Upsilon(\widehat{U}_0(z)) = \mathcal{Z} \left\{ \sum_{k=0}^{n+1} \mathcal{C}_{n+1-k} U_0^k \right\} (z) = \kappa_+(\varepsilon\xi)z\widehat{U}_0(z).$$

Proof. Since $\mu_+(\varepsilon\bar{\xi}) = \overline{\mu_+(\varepsilon\xi)}$, one obtains $\kappa_+(\varepsilon\bar{\xi}) = \overline{\kappa_+(\varepsilon\xi)}$. Then,

$$\begin{aligned} \operatorname{Re} \left(\kappa_+(\varepsilon\xi) R^m e^{im\theta} \right) &= \frac{1}{2} \left(\kappa_+(\varepsilon\xi) R^m e^{im\theta} + \overline{\kappa_+(\varepsilon\xi)} R^m e^{-im\theta} \right) \\ &= \frac{1}{2} \left(\kappa_+(\varepsilon\xi) R^m e^{im\theta} + \kappa_+(\varepsilon\bar{\xi}) R^m e^{-im\theta} \right). \end{aligned}$$

Thus, the value of $(\mathcal{C}_m)_{m \geq 0}$ in (2.28) can be reformulated as

$$\begin{aligned} \mathcal{C}_m &= \frac{1}{2\pi} \left(\int_0^{\pi} \kappa_+(\varepsilon\xi) R^m e^{im\theta} d\theta + \int_0^{\pi} \kappa_+(\varepsilon\bar{\xi}) R^m e^{-im\theta} d\theta \right) \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \kappa_+(\varepsilon\xi) R^m e^{im\theta} d\theta \\ &= \mathcal{Z}^{-1}(\kappa_+(\varepsilon\xi))(m). \end{aligned}$$

By the convolution property and inverting \mathcal{Z} -transform, we can conclude that

$$\Upsilon(\widehat{U}_0(z)) = \mathcal{Z} \left\{ \sum_{k=0}^{n+1} \mathcal{C}_{n+1-k} U_0^k \right\} (z) = \kappa_+(\varepsilon\xi)z\widehat{U}_0(z),$$

This ends the proof of Lemma 2.5. \square

2.3. Stiff stability analysis. Under the SKC, we now consider the Proposition 1.3 with nonzero boundary condition $(b^n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{N}, \mathbb{R})$ and homogeneous Cauchy data $(f_j)_{j \in \mathbb{N}} \equiv 0$. In order to get the uniform estimate on $(U_j^n)_{j \in \mathbb{N}}$, firstly, we prove the following lemma:

Lemma 2.6. *Assume that the parameters $a, \Delta x, \Delta t > 0$ satisfy*

$$(2.29) \quad \Delta x \leq \frac{3\sqrt{a}}{8} \Delta t.$$

Let $\varepsilon > 0$, $R > 1$, $\theta \in (-\pi, \pi]$ and then $\kappa_-(\varepsilon\xi)$ be given by (2.3). Then the following property holds

$$(2.30) \quad \sum_{j \geq 0} |\kappa_-(\varepsilon\xi)|^{2j} \leq \frac{\Delta t \sqrt{a}}{\Delta x (1 - R^{-1})}.$$

Proof. Since the property of $\operatorname{Re}(\mu_-(\varepsilon\xi))$ in (2.2), we can prove

$$(2.31) \quad \left(\operatorname{Re}(\mu_-(\varepsilon\xi)) \lambda_{x\varepsilon} + \sqrt{(\operatorname{Re}(\mu_-(\varepsilon\xi)) \lambda_{x\varepsilon})^2 + 1} \right)^2 \leq \left(\eta \Delta x + \sqrt{\eta^2 \Delta x^2 + 1} \right)^2,$$

where $\eta = -a^{-1/2} \operatorname{Re} \xi$. According to Lemma A.2 and the inequality (2.31), we have

$$|\kappa_-(\varepsilon\xi)|^2 = \left| \mu_-(\varepsilon\xi) \lambda_{x\varepsilon} + \sqrt{(\mu_-(\varepsilon\xi) \lambda_{x\varepsilon})^2 + 1} \right|^2 \leq \left(\eta \Delta x + \sqrt{\eta^2 \Delta x^2 + 1} \right)^2.$$

Then, we obtain the following estimate

$$\sum_{j \geq 0} |\kappa_-(\varepsilon\xi)|^{2j} = (1 - |\kappa_-(\varepsilon\xi)|^2)^{-1} \leq \left(1 - \left(\eta \Delta x + \sqrt{\eta^2 \Delta x^2 + 1} \right)^2 \right)^{-1}.$$

Since $\operatorname{Re} \xi$ satisfies the property (2.1), we get

$$\frac{\Delta t \sqrt{a}}{2} \leq -\frac{1}{\eta} \leq \frac{\Delta t \sqrt{a}}{1 - R^{-1}}.$$

If we assume now $\Delta x \leq \frac{3\sqrt{a}}{8} \Delta t \leq -\frac{3}{4\eta}$ then we have

$$\left(1 - \left(\eta \Delta x + \sqrt{\eta^2 \Delta x^2 + 1} \right)^2 \right)^{-1} \leq -\eta^{-1} \Delta x^{-1}.$$

Thus, we conclude that

$$\sum_{j \geq 0} |\kappa_-(\varepsilon\xi)|^{2j} \leq -\eta^{-1} \Delta x^{-1} \leq \frac{\Delta t \sqrt{a}}{\Delta x (1 - R^{-1})}.$$

This ends the proof of Lemma 2.6. □

Secondly, by an application of the following Plancherel's theorem for \mathcal{Z} -transform

$$\sum_{n \geq 0} R^{-2n} |U_j^n|^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} |\widehat{U}_j(Re^{i\theta})|^2 d\theta, \quad R > 1,$$

we have

$$\begin{aligned} \sum_{n \geq 0} R^{-2n} |U_0^n|^2 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} |\widehat{U}_0(Re^{i\theta})|^2 d\theta \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \left| \frac{\widehat{b}(z)}{B_u + g(\varepsilon\xi) B_v} \right|^2 (1 + |g(\varepsilon\xi)|^2) d\theta. \end{aligned}$$

From the Lemma 2.1, under the SKC, we then obtain

$$(2.32) \quad \sum_{n \geq 0} R^{-2n} |U_0^n|^2 \lesssim \frac{1}{2\pi} \int_{-\pi}^{\pi} |\widehat{b}(Re^{i\theta})|^2 d\theta \lesssim \sum_{n \geq 0} R^{-2n} |b^n|^2.$$

Similarly, by an application of the Plancherel's theorem for \mathcal{L} -transform, we have

$$\begin{aligned} \sum_{n \geq 0} \sum_{j \geq 0} R^{-2n} |U_j^n|^2 &= \frac{1}{2\pi} \sum_{j \geq 0} \int_{-\pi}^{\pi} |\widehat{U}_j(Re^{i\theta})|^2 d\theta \\ &= \frac{1}{2\pi} \sum_{j \geq 0} \int_{-\pi}^{\pi} \left| \frac{\widehat{b}(z)}{B_u + g(\varepsilon\xi)B_v} \right|^2 |\kappa_-(\varepsilon\xi)|^{2j} (1 + |g(\varepsilon\xi)|^2) d\theta. \end{aligned}$$

Again, under the SKC, we get from Lemma 2.1

$$(2.33) \quad \sum_{n \geq 0} \sum_{j \geq 0} R^{-2n} |U_j^n|^2 \lesssim \frac{1}{2\pi} \sum_{j \geq 0} \int_{-\pi}^{\pi} |\widehat{b}(z)|^2 |\kappa_-(\varepsilon\xi)|^{2j} d\theta.$$

Following Lemma 2.6, if we assume (2.29) holds, then the inequality (2.33) becomes

$$(2.34) \quad \frac{R-1}{R} \Delta x \sum_{n \geq 0} \sum_{j \geq 0} R^{-2n} |U_j^n|^2 \lesssim \frac{\Delta t}{2\pi} \int_{-\pi}^{\pi} |\widehat{b}(Re^{i\theta})|^2 d\theta \lesssim \Delta t \sum_{n \geq 0} R^{-2n} |b^n|^2.$$

According to (2.32) and (2.34), there exists a constant $C > 0$ such that

$$\frac{R-1}{R} \sum_{n \geq 0} \sum_{j \geq 0} R^{-2n} \Delta x |U_j^n|^2 + \sum_{n \geq 0} R^{-2n} \Delta t |U_0^n|^2 \leq C \sum_{n \geq 0} R^{-2n} \Delta t |b^n|^2.$$

By setting in the above formula $R = e^{\gamma\Delta t}$ for $\gamma > 0$ and $\Delta t > 0$, and using the classical lower bound $e^{\gamma\Delta t} \geq 1 + \gamma\Delta t$, we obtain that there exists a constant $c > 0$ such that

$$\frac{\gamma}{\gamma\Delta t + 1} \sum_{n \geq 0} \sum_{j \geq 0} e^{-2\gamma n\Delta t} \Delta t \Delta x |U_j^n|^2 + \sum_{n \geq 0} e^{-2\gamma n\Delta t} \Delta t |U_0^n|^2 \leq c \sum_{n \geq 0} e^{-2\gamma n\Delta t} \Delta t |b^n|^2.$$

This ends the proof of the Proposition 1.3.

Let us observe that the scheme (1.8) together also with its boundary condition is closed to be forward-in-time, except it is one-step implicit. By this property, changing the data b to zero after some time T and unchanged before that time T , the discrete solution U_j^n is the same for $n\Delta t < T$. Therefore, there exists a constant $C_T > 0$ such that

$$(2.35) \quad \sum_{n=0}^N \sum_{j \geq 0} \Delta x \Delta t |U_j^n|^2 + \sum_{n=0}^N \Delta t |U_0^n|^2 \leq C_T \sum_{n=0}^N \Delta t |b^n|^2,$$

with $N := T/\Delta t$. This will be useful to prove the Theorem 1.1.

2.4. Numerical experiments. In this paragraph, we first provide the behavior of the numerical solution $(U_j^n)_{j \in \mathbb{N}}$ according to whether or not the SKC (1.9) is valid. We also look at the degenerate case when the UKC (1.4) does not hold (and thus, none of the other stability conditions). Following the continuous case studied by Xin and Xu in [23], the solution of the IBVP (1.1)-(1.3) with homogeneous initial condition can be constructed by the method of Laplace transform. By inverting the Laplace transform, the solution $U(x, t)$ has form

$$U(x, t) = \mathcal{L}^{-1} \widetilde{U} = \frac{1}{2\pi} \int_{-\infty}^{+\infty} e^{\zeta t} \frac{\widetilde{b}(\zeta)}{B_u + g(\varepsilon\zeta)B_v} e^{\mu - (\varepsilon\zeta)x/\varepsilon} r_-(\varepsilon\zeta) d\beta,$$

where $\zeta = \alpha + i\beta$, $\alpha > 0$. Then, we observe the error between the exact solution $U(x_j, t^n)$ and the numerical solution U_j^n of the numerical scheme (1.8) with homogeneous initial data at the grid point $(x_j, t^n) = (j\Delta x, n\Delta t)$. After that, we present some numerical experiments and observe the effective behavior of the energy terms $\|U\|_{\ell^2(\mathbb{N} \times [0, T], \mathbb{R}^2)}$ and $\|U\|_{\ell^2(\{0\} \times [0, T], \mathbb{R}^2)}$ corresponding to whether or not the SKC (1.9) is valid.

As main parameters for the experiments, we choose $a = 1$, $B_v = 1$, $\lambda_{xt} = 1/3$ and let the relaxation rate ε and the boundary data B_u vary. The test case we consider concerns the following data. The initial data is the homogeneous one $(f_j)_{j \in \mathbb{N}} \equiv 0$. The boundary data is

$$b(t) = \frac{t}{2} \sin(t).$$

Let us observe that these data are compatible in the corner $(x, t) = (0, 0)$ in the sense that $Bf(0) = b(0)$. Moreover, the Laplace transform of $b(t)$ is

$$\tilde{b}(\zeta) = \frac{\zeta}{(\zeta^2 + 1)^2}.$$

2.4.1. The behavior of the numerical solution. Let the space step $\Delta x = 10^{-2}$ and the time step $\Delta t = \lambda_{xt}^{-1} \Delta x$. Firstly, we choose the value of B_u such that the SKC (1.9) is satisfied with $\varepsilon = 10^{-2}$ and also with $\varepsilon = 10^2$. The Figures 2.1 and 2.2 show the numerical solution $(U_j^n)_{j \in \mathbb{N}}$ over the time interval $t \in [0, 1.2)$.




FIGURE 2.1. The numerical solution $u(x, t)$ (left) and $v(x, t)$ (right) for $\varepsilon = 10^{-2}$. The SKC (1.9) holds with $B_u = -4$.

In the first case, $\varepsilon = 10^{-2}$, the incoming solution at the boundary $x = 0$ go slowly. This is due to the initial relaxation of solution to the equilibrium system. In the case $\varepsilon = 10^2$, its solution seems to be faster. It is not so much influenced by relaxation source term but more by the boundary dissipation.

Secondly, we choose the value of B_u such that the SKC (1.9) is not satisfied. Besides, we also present the numerical solution when the Uniform Kreiss Condition (1.4) is wrong. The Figures 2.3 and 2.4 show the numerical solution $(U_j^n)_{j \in \mathbb{N}}$ over the time interval $t \in [0, 0.5)$.

When the SKC (1.9) fails, we observe that the numerical solution at the boundary rise gradually. This is the case for example for $\varepsilon = 10^{-2}$ together with the parameters $(B_u, B_v) = (-1/2, 1)$. The behavior is even worse when the UKC (1.4) is not satisfied (see Figure 2.4).

2.4.2. The error between the exact solution and the numerical solution. Let us begin with the notation

$$(2.36) \quad E(t^n) := \left(\Delta x \sum_{j \geq 0} |U(x_j, t^n) - U_j^n|^2 \right)^{1/2}.$$

We choose a set of values B_u such that the SKC (1.9) is satisfied with the space step Δx and the relation rate ε vary. The error, as measured in (2.36), are reported in the Tables 1 and 2.




FIGURE 2.2. The numerical solution $u(x, t)$ (left) and $v(x, t)$ (right) for $\varepsilon = 10^2$. The SKC (1.9) holds with $B_u = -4$.




FIGURE 2.3. The numerical solution $u(x, t)$ (left) and $v(x, t)$ (right) for $\varepsilon = 10^{-2}$. The SKC (1.9) does not hold with $B_u = -1/2$.

According to the experiments in Tables 1 and 2, for some $\varepsilon \in (0, +\infty)$ and (B_u, B_v) satisfying the SKC (1.9), the observed convergence rate is 1 since going down Δx by a factor 2 decreases the error of the same factor 2. It means that the behavior of the numerical solution U_j^n is the same as the evolution of the exact solution $U(x_j, t^n)$. This is the case for example for $\varepsilon = 10^{-2}$ together with the parameters $(B_u, B_v) = (-4, 1)$.

2.4.3. The effective behavior of the energy terms. Let the space step $\Delta x = 10^{-2}$ and the time step $\Delta t = \lambda_{xt}^{-1} \Delta x$. We present hereafter the behavior of the following energy terms for $\varepsilon \in (0, +\infty)$, $T = 1.2$ and $N = T/\Delta t$. The first one corresponds to the ℓ^2 in time and space energy of the discrete solution and the




FIGURE 2.4. The numerical solution $u(x, t)$ (left) and $v(x, t)$ (right) for $\varepsilon = 10^2$. The UKC (1.4) is wrong with $B_u = -1$.

Δx	$\varepsilon = 10^{-2}$	$\varepsilon = 10^{-1}$	$\varepsilon = 1$	$\varepsilon = 10$	$\varepsilon = 10^2$
5×10^{-2}	6.8×10^{-3}	1.2×10^{-2}	2.6×10^{-2}	3.1×10^{-2}	3.2×10^{-2}
25×10^{-3}	3×10^{-3}	5.9×10^{-3}	1.3×10^{-2}	1.6×10^{-2}	1.6×10^{-2}
125×10^{-4}	1.5×10^{-3}	2.9×10^{-3}	6.8×10^{-3}	8.2×10^{-3}	8.3×10^{-3}
625×10^{-5}	7.2×10^{-4}	1.5×10^{-3}	3.4×10^{-3}	4.1×10^{-3}	4.2×10^{-3}

TABLE 1. The error $E(1.2)$ for $B_u = -4$.

Δx	$\varepsilon = 10^{-2}$	$\varepsilon = 10^{-1}$	$\varepsilon = 1$	$\varepsilon = 10$	$\varepsilon = 10^2$
5×10^{-2}	8.5×10^{-3}	1.3×10^{-2}	2.1×10^{-2}	2.3×10^{-2}	2.4×10^{-2}
25×10^{-3}	3.8×10^{-3}	6.4×10^{-3}	1.1×10^{-2}	1.2×10^{-2}	1.2×10^{-2}
125×10^{-4}	1.8×10^{-3}	3.2×10^{-3}	5.4×10^{-3}	6.1×10^{-3}	6.2×10^{-3}
625×10^{-5}	9.1×10^{-4}	1.6×10^{-3}	2.7×10^{-3}	3.1×10^{-3}	3.1×10^{-3}

TABLE 2. The error $E(1.2)$ for $B_u = 3$.

second to the ℓ^2 in time energy of the numerical trace at the boundary:



$$(2.37) \quad E_1 := \|U\|_{\ell^2(\mathbb{N} \times [0, T], \mathbb{R}^2)}^2 = \sum_{n=0}^N \sum_{j \geq 0} \Delta x \Delta t |U_j^n|^2,$$

$$E_2 := \|U\|_{\ell^2(\{0\} \times [0, T], \mathbb{R}^2)}^2 = \sum_{n=0}^N \Delta t |U_0^n|^2,$$

which are shown in the Table 3 and Figures 2.5, 2.6.

B_u	$\varepsilon = 10^{-2}$	$\varepsilon = 1$	$\varepsilon = 10^2$	B_u	$\varepsilon = 10^{-2}$	$\varepsilon = 1$	$\varepsilon = 10^2$
-4	35×10^{-5}	77×10^{-5}	47×10^{-4}	-4	66×10^{-4}	8×10^{-3}	2×10^{-2}
-2	15×10^{-4}	39×10^{-4}	43×10^{-3}	-2	29×10^{-3}	4×10^{-2}	18×10^{-2}
-1	2.6×10^{17}	1.69×10^{31}	3.17×10^{56}	-1	1.24×10^{19}	7.17×10^{32}	1.89×10^{58}
-0.5	44047.9	418525.12	2837033.2	-0.5	191420.5	9910910.98	106714237.85
1	34×10^{-4}	5×10^{-3}	10^{-2}	1	46×10^{-3}	55×10^{-3}	67×10^{-3}
3	48×10^{-5}	87×10^{-5}	2×10^{-2}	3	93×10^{-4}	94×10^{-4}	12×10^{-3}

TABLE 3. The energy terms E_1 (left) and E_2 (right).

FIGURE 2.5. Energy evolution E_1 for $B_u = -4$ (left) and $B_u = -0.5$ (right).FIGURE 2.6. Energy evolution E_2 for $B_u = -4$ (left) and $B_u = -0.5$ (right).

- For some $\epsilon \in (0, +\infty)$, the values of E_1 and E_2 rise gradually when the SKC (1.9) is not satisfied. This is the case for example for $\epsilon = 10^2$ together with the parameters $(B_u, B_v) = (-1/2, 1)$. The behavior is even worse when the UKC (1.4) is not hold.
- In the case $\epsilon = 10^{-2}$, the energy term E_1 and E_2 increase slowly. This is due to the effect of incoming solution at the boundary when the initial relaxation of solution tends to the equilibrium system. In the case $\epsilon = 10^2$, those values increase fairly rapidly. It is not so much influenced by relaxation source term but more by the boundary dissipation.

Clearly, the numerical results show that the numerical solution at the boundary $x = 0$ increase quickly as soon as the SKC (1.9) does not hold. If the UKC (1.4) is not satisfied, the behavior of numerical solution is even worse. Besides, by the decrease of the error $E(t^n)$, we can see that the values of U_j^n tend to the exact solution $U(x_j, t^n)$. Indeed, it seems that the SKC (1.9) is also necessary to ensure the non-increase rapidly of the energy terms E_1 and E_2 under the effect of the relaxation source term and the boundary dissipation.

3. STIFF STABILITY OF THE IBVP WITH HOMOGENEOUS BOUNDARY CONDITION

For convenience in the forthcoming discussions, we recall that the IBVP (1.8) with homogeneous boundary condition writes

$$(3.1) \quad \begin{cases} \frac{U_j^{n+1} - U_j^n}{\Delta t} + \frac{1}{2\Delta x} A (U_{j+1}^{n+1} - U_{j-1}^{n+1}) = \frac{1}{\varepsilon} S U_j^{n+1}, & j \geq 1, n \geq 0, \\ U_j^0 = f_j, & j \geq 0, \\ B U_0^n = 0, & n \geq 0, \\ \frac{1}{\Delta t} \Gamma (U_0^{n+1} - U_0^n) + \frac{1}{2\Delta x} \Gamma A \left(U_1^{n+1} - \sum_{k=0}^{n+1} \mathfrak{C}_{n+1-k} U_0^k \right) = \frac{1}{\varepsilon} \Gamma S U_0^{n+1}, & n \geq 0. \end{cases}$$

In [23, Section 5], under the SKC, Xin and Xu find explicitly the solution $U(x, t)$ of the IBVP (1.6) by the method of Laplace transform. The solution is decomposed into two ingredients, by assembling a solution for the case of the Cauchy problem and another one for the case of the IBVP with homogeneous initial condition. In our case, assuming the SKC to hold, the numerical solution $(U_j^n)_{j \in \mathbb{N}}$ can be constructed by the method of \mathcal{Z} -transform. Since the coefficients $(\mathfrak{C}_m)_{m \geq 0}$ are defined for homogeneous initial data, the numerical solution $(U_j^n)_{j \in \mathbb{N}}$ of the IBVP (3.1) consists of not only the solutions for case of the Cauchy problem and for the IBVP with zero initial data but also another numerical error term $(U_j^H)_{j \in \mathbb{N}}$. To complete the proof of the Theorem 1.1 with homogeneous boundary condition, we first use the means of discrete energy method in order to prove the Proposition 1.2. By an application of the Plancherel's theorem for \mathcal{Z} -transform [15, 19], the numerical error term of $(U_j^H)_{j \in \mathbb{N}}$ will be estimated in Section 3.3. After that, we get the expected result of the case IBVP with homogeneous initial condition.

3.1. Solution by \mathcal{Z} -transform. Again, we follow the explicit solving of the IBVP (3.1) by using the \mathcal{Z} -transform. With

$$\widehat{U}_j(z) = \mathcal{Z}\{U_j^n\}(z) = \sum_{n \geq 0} U_j^n z^{-n}, \quad |z| > 1.$$

Importantly, we now have $(f_j)_{j \in \mathbb{N}} \neq 0$, and thus we get

$$\sum_{n \geq 0} U_j^{n+1} z^{-n} = z \widehat{U}_j(z) - z U_j^0 = z \widehat{U}_j(z) - z f_j.$$

Therefore, (3.1) becomes

$$(3.2a) \quad \begin{cases} \widehat{U}_{j+1}(z) - \widehat{U}_{j-1}(z) = 2\lambda_{x\varepsilon} M(\varepsilon\xi) \widehat{U}_j(z) + f_{j+1} - 2\lambda_{x\varepsilon} M(\varepsilon\Delta t^{-1}) f_j - f_{j-1}, & j \geq 1, \\ B \widehat{U}_0(z) = 0, \end{cases}$$

$$(3.2b)$$

$$(3.2c) \quad \left[\Gamma A \left[\widehat{U}_1(z) - \left(\kappa_+(\varepsilon\xi) I + 2\lambda_{x\varepsilon} M(\varepsilon\xi) \right) \widehat{U}_0(z) - f_1 \right] = 0. \right.$$

Let us recall that the \mathcal{Z} -transform of $\sum_{k=0}^{n+1} \mathfrak{C}_{n+1-k} U_0^k$ is given by $\kappa_+(\varepsilon\xi) z \widehat{U}_0(z)$. Firstly, we look at the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ to (3.2a)-(3.2c). This is the object of the following proposition:

Proposition 3.1. *Assume that the SKC (1.9) is satisfied. Let $(f_j)_{j \in \mathbb{N}} \in \ell^2(\mathbb{N}, \mathbb{R}^2)$ and denote $V_k^I, w^I(\varepsilon\xi)$ and $w^H(\varepsilon\xi)$ as follows:*

$$(3.3) \quad \begin{aligned} V_k^I &= f_{k+1} - 2\lambda_{x\varepsilon} M(\varepsilon\Delta t^{-1}) f_k - f_{k-1}, \\ w^I(\varepsilon\xi) &= \sum_{k=0}^{+\infty} (-1)^{-k} \kappa_+^{-k}(\varepsilon\xi) V_k^I, \\ w^H(\varepsilon\xi) &= - \sum_{k=0}^{+\infty} \kappa_+^{-k}(\varepsilon\xi) V_k^I. \end{aligned}$$

Then, the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z) \in \ell^2(\mathbb{N}, \mathbb{C}^2)$ to (3.2a)-(3.2c) takes the form

$$\begin{aligned}
(3.4) \quad \widehat{U}_j(z) &= \frac{\kappa_-^{j+1}(\varepsilon\xi)}{4g(\varepsilon\xi)} \times \left(\frac{B_u - g(\varepsilon\xi)B_v}{B_u + g(\varepsilon\xi)B_v} \times (g(\varepsilon\xi), 1) w^I(\varepsilon\xi) - (g(\varepsilon\xi), -1) w^II(\varepsilon\xi) \right) \\
&\quad \times \left(\frac{B_u - g(\varepsilon\xi)B_v}{B_u + g(\varepsilon\xi)B_v} \times r_-(\varepsilon\xi) + (-1)^{j+1} r_+(\varepsilon\xi) \right) \\
&\quad - \frac{\kappa_-^j(\varepsilon\xi)}{\kappa_+(\varepsilon\xi) + \kappa_-(\varepsilon\xi)} \times \left(\Phi_-(\varepsilon\xi) w^I(\varepsilon\xi) + (-1)^j \Phi_+(\varepsilon\xi) w^II(\varepsilon\xi) \right) \\
&\quad + \frac{1}{\kappa_+(\varepsilon\xi) + \kappa_-(\varepsilon\xi)} \times \left(\sum_{k=0}^{j-1} \kappa_-^{j-k}(\varepsilon\xi) \left(\Phi_-(\varepsilon\xi) + (-1)^{j-k-1} \Phi_+(\varepsilon\xi) \right) V_k^I \right. \\
&\quad \left. + \sum_{k=j}^{+\infty} (-1)^{j-k} \kappa_+^{j-k}(\varepsilon\xi) \left(\Phi_-(\varepsilon\xi) + (-1)^{j-k-1} \Phi_+(\varepsilon\xi) \right) V_k^I \right).
\end{aligned}$$

Proof. Before we prove the above result, let us notice that we omit the explicit dependence in $\varepsilon\xi$. Firstly, we look at the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ to (3.2a) and consider the two-dimensional problem (3.2a) under the following nonhomogeneous one-step recurrence form

$$(3.5) \quad W_{j+1}(z) = M_1 W_j(z) + V_j,$$

where W_j is the same as in (2.17) and

$$V_j = \begin{pmatrix} V_j^I \\ 0 \end{pmatrix}.$$

The solution $(W_j)_{j \in \mathbb{N}}(z)$ to (3.5) is thus given by

$$(3.6) \quad W_j(z) = M_1^j W_0(z) + \sum_{k=0}^{j-1} M_1^{j-1-k} V_k.$$

Together with the the explicit formula of M_1^j in Lemma 2.2, the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ to (3.2a) is given by

$$\widehat{U}_j(z) = -\frac{1}{\kappa_+ + \kappa_-} \times \left[\widehat{\kappa}_{j+1} \widehat{\Psi}_j \widehat{U}_0(z) + \widehat{\kappa}_j \widehat{\Psi}_{j+1} \widehat{U}_{-1} + \sum_{k=0}^{j-1} \widehat{\kappa}_{j-k} \widehat{\Psi}_{j-k-1} V_k^I \right].$$

By the definition of $\widehat{\kappa}_k$ and $\widehat{\Psi}_k$ in (2.10), the above formula is equivalent to

$$\begin{aligned}
(3.7) \quad \widehat{U}_j(z) &= -\frac{(-1)^j \kappa_+^j}{\kappa_+ + \kappa_-} \times \left[-\kappa_+ \left(\Phi_- + (-1)^j \Phi_+ \right) \widehat{U}_0(z) + \left(\Phi_- + (-1)^{j+1} \Phi_+ \right) \widehat{U}_{-1}(z) \right. \\
&\quad \left. + \sum_{k=0}^{j-1} (-\kappa_+)^{-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I \right] \\
&\quad + \frac{\kappa_-^j}{\kappa_+ + \kappa_-} \times \left[\kappa_- \left(\Phi_- + (-1)^j \Phi_+ \right) \widehat{U}_0(z) + \left(\Phi_- + (-1)^{j+1} \Phi_+ \right) \widehat{U}_{-1}(z) \right. \\
&\quad \left. + \sum_{k=0}^{j-1} \kappa_-^{-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I \right].
\end{aligned}$$

Thanks to the definition of w^I and w^II in (3.3), one has

$$(3.8) \quad \begin{aligned} & \sum_{k=0}^{j-1} (-\kappa_+)^{-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I \\ &= \Phi_- w^I + (-1)^j \Phi_+ w^{II} - \sum_{k=j}^{+\infty} (-\kappa_+)^{-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I. \end{aligned}$$

Substituting (3.8) into (3.7), we have

$$(3.9) \quad \begin{aligned} \widehat{U}_j(z) &= -\frac{(-1)^j \kappa_+^j}{\kappa_+ + \kappa_-} \times \left[\Phi_- \left(-\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) + w^I \right) + (-1)^{j+1} \Phi_+ \left(\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) - w^{II} \right) \right] \\ &+ \frac{1}{\kappa_+ + \kappa_-} \times \sum_{k=j}^{+\infty} (-1)^{j-k} \kappa_+^{j-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I \\ &+ \frac{\kappa_-^j}{\kappa_+ + \kappa_-} \times \left[\kappa_- \left(\Phi_- + (-1)^j \Phi_+ \right) \widehat{U}_0(z) + \left(\Phi_- + (-1)^{j+1} \Phi_+ \right) \widehat{U}_{-1}(z) \right. \\ &\quad \left. + \sum_{k=0}^{j-1} \kappa_-^{-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I \right]. \end{aligned}$$

Since we expect $(\widehat{U}_j)_{j \in \mathbb{N}}(z) \in \ell^2(\mathbb{N}, \mathbb{C}^2)$, we need a natural boundary condition at $x = +\infty$. Besides, one gets $|\kappa_+| > 1$ and $|\kappa_-| < 1$. Thus, the natural boundary condition takes the form

$$(3.10) \quad \begin{cases} \Phi_- \left(-\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) + w^I \right) = 0, \\ \Phi_+ \left(\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) - w^{II} \right) = 0. \end{cases}$$

By the definition of Φ_{\pm} in (2.4), the system (3.10) is equivalent to

$$\begin{cases} (g, 1) \left(-\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) + w^I \right) = 0, \\ (g, -1) \left(\kappa_+ \widehat{U}_0(z) + \widehat{U}_{-1}(z) - w^{II} \right) = 0. \end{cases}$$

Then, we have

$$(3.11) \quad \widehat{U}_{-1}(z) = \kappa_+ (\Phi_- - \Phi_+) \widehat{U}_0(z) - \Phi_- w^I + \Phi_+ w^{II}.$$

Plugging (3.11) into (3.9), we get

$$\begin{aligned} \widehat{U}_j(z) &= \frac{1}{\kappa_+ + \kappa_-} \times \sum_{k=j}^{+\infty} (-1)^{j-k} \kappa_+^{j-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I \\ &+ \frac{\kappa_-^j}{\kappa_+ + \kappa_-} \times \left[\kappa_- \left(\Phi_- + (-1)^j \Phi_+ \right) + \kappa_+ \left(\Phi_- + (-1)^{j+1} \Phi_+ \right) \left(\Phi_- - \Phi_+ \right) \right] \widehat{U}_0(z) \\ &+ \frac{\kappa_-^j}{\kappa_+ + \kappa_-} \times \left(\Phi_- + (-1)^{j+1} \Phi_+ \right) \left(-\Phi_- w^I + \Phi_+ w^{II} \right) \\ &+ \frac{1}{\kappa_+ + \kappa_-} \times \sum_{k=0}^{j-1} \kappa_-^{j-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I. \end{aligned}$$

Under the properties of Φ_{\pm} in (2.7), the above formula becomes

(3.12)

$$\begin{aligned} \widehat{U}_j(z) &= \kappa_-^j \left(\Phi_- + (-1)^j \Phi_+ \right) \widehat{U}_0(z) - \frac{\kappa_-^j}{\kappa_+ + \kappa_-} \times \left(\Phi_- w^I + (-1)^j \Phi_+ w^II \right) \\ &+ \frac{1}{\kappa_+ + \kappa_-} \times \left[\sum_{k=0}^{j-1} \kappa_-^{j-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I + \sum_{k=j}^{+\infty} (-1)^{j-k} \kappa_+^{j-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I \right]. \end{aligned}$$

Secondly, we look at the boundary data $\widehat{U}_0(z)$ and extend the initial data $(f_j)_{j \in \mathbb{N}}$ to the whole line by setting $f_j = 0$ for $j \leq 0$. Since $f_0 = f_{-1} = 0$ and $\Phi_+ + \Phi_- = I$, we can see that

$$\begin{aligned} \widehat{U}_1(z) &= \kappa_- \left(\Phi_- - \Phi_+ \right) \widehat{U}_0(z) - \frac{\kappa_-}{\kappa_+ + \kappa_-} \times \left(\Phi_- w^I - \Phi_+ w^II \right) \\ &+ \frac{1}{\kappa_+ + \kappa_-} \times \left[\kappa_- f_1 - \kappa_+ \sum_{k=1}^{+\infty} (-1)^{-k} \kappa_+^{-k} \left(\Phi_- + (-1)^{-k} \Phi_+ \right) V_k^I \right]. \end{aligned} \quad (3.13)$$

On the other hand, by the definition of w^I and w^II in (3.3), we get the following property

$$\sum_{k=1}^{+\infty} (-1)^{-k} \kappa_+^{-k} \left(\Phi_- + (-1)^{-k} \Phi_+ \right) V_k^I = \Phi_- w^I - \Phi_+ w^II - f_1. \quad (3.14)$$

Substituting (3.14) into (3.13), the value of $\widehat{U}_1(z)$ can be reformulated as

$$\widehat{U}_1(z) = \kappa_- (\Phi_- - \Phi_+) \widehat{U}_0(z) - \Phi_- w^I + \Phi_+ w^II + f_1.$$

Thus, the equation (3.2c) becomes

$$\Gamma A \left(\kappa_- (\Phi_- - \Phi_+) \widehat{U}_0(z) - \Phi_- w^I + \Phi_+ w^II - (\kappa_+ I + 2\lambda_{x\varepsilon} M) \widehat{U}_0(z) \right) = 0. \quad (3.15)$$

Indeed, we observe that

$$\kappa_+ I + 2\lambda_{x\varepsilon} M = \kappa_- \Phi_- + (\kappa_+ + 2\lambda_{x\varepsilon} \mu_+) \Phi_+.$$

Then, the equation (3.15) can be represented as

$$\Gamma A \Phi_+ \widehat{U}_0(z) = -\frac{1}{2\kappa_+} \Gamma A \left(\Phi_- w^I - \Phi_+ w^II \right). \quad (3.16)$$

Besides, one has

$$\Gamma A \Phi_+ = \frac{a}{2g} \times (B_u + gB_v) \times (g, -1).$$

From the Lemma 2.1, the equation (3.16) is equivalent under the SKC to

$$(g, -1) \widehat{U}_0(z) = -\frac{g}{a\kappa_+(B_u + gB_v)} \Gamma A \left(\Phi_- w^I - \Phi_+ w^II \right).$$

Together with the boundary condition (3.2b), one gets

$$\begin{pmatrix} B_u & B_v \\ g & -1 \end{pmatrix} \widehat{U}_0(z) = -\frac{g}{a\kappa_+(B_u + gB_v)} \left(\Gamma A \begin{pmatrix} 0 \\ \Phi_- w^I - \Phi_+ w^II \end{pmatrix} \right).$$

Then, under the SKC, we have

$$\widehat{U}_0(z) = -\frac{g}{a\kappa_+(B_u + gB_v)^2} \times \Gamma A \left(\Phi_- w^I - \Phi_+ w^II \right) \begin{pmatrix} B_v \\ -B_u \end{pmatrix}. \quad (3.17)$$

Substituting (3.17) into (3.12), the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ to (3.2a)-(3.2c) is given by

$$\begin{aligned} \widehat{U}_j(z) = & -\frac{\kappa_-^{j+1}g}{a(B_u + gB_v)^2} \times \Gamma A \left(\Phi_- w^I - \Phi_+ w^{II} \right) \times \left(\Phi_- + (-1)^j \Phi_+ \right) \begin{pmatrix} B_v \\ -B_u \end{pmatrix} \\ & -\frac{\kappa_-^j}{\kappa_+ + \kappa_-} \times \left(\Phi_- w^I + (-1)^j \Phi_+ w^{II} \right) \\ & + \frac{1}{\kappa_+ + \kappa_-} \times \left[\sum_{k=0}^{j-1} \kappa_-^{j-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I + \sum_{k=j}^{+\infty} (-1)^{j-k} \kappa_+^{j-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I \right]. \end{aligned}$$

Together with the definitions of Φ_{\pm} and Γ in (2.6) and (2.14), respectively, we have

$$\begin{aligned} & -\frac{g}{a(B_u + gB_v)^2} \times \Gamma A \left(\Phi_- w^I - \Phi_+ w^{II} \right) \times \left(\Phi_- + (-1)^j \Phi_+ \right) \begin{pmatrix} B_v \\ -B_u \end{pmatrix} \\ & = \frac{1}{4g} \times \left(\frac{B_u - gB_v}{B_u + gB_v} \times (g, 1) w^I - (g, -1) w^{II} \right) \times \left(\frac{B_u - gB_v}{B_u + gB_v} \times r_- + (-1)^{j+1} r_+ \right). \end{aligned}$$

Therefore, the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ to (3.2a)-(3.2c) can be reformulated as

$$\begin{aligned} \widehat{U}_j(z) = & \frac{\kappa_-^{j+1}}{4g} \times \left(\frac{B_u - gB_v}{B_u + gB_v} \times (g, 1) w^I - (g, -1) w^{II} \right) \times \left(\frac{B_u - gB_v}{B_u + gB_v} \times r_- + (-1)^{j+1} r_+ \right) \\ & -\frac{\kappa_-^j}{\kappa_+ + \kappa_-} \times \left(\Phi_- w^I + (-1)^j \Phi_+ w^{II} \right) \\ & + \frac{1}{\kappa_+ + \kappa_-} \times \left[\sum_{k=0}^{j-1} \kappa_-^{j-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I + \sum_{k=j}^{+\infty} (-1)^{j-k} \kappa_+^{j-k} \left(\Phi_- + (-1)^{j-k-1} \Phi_+ \right) V_k^I \right]. \end{aligned}$$

This ends the proof of Proposition 3.1. \square

Secondly, we can see that the solution $(\widehat{U}_j)_{j \in \mathbb{N}}(z)$ to (3.2a)-(3.2c) consists of three parts:

$$(3.18) \quad \widehat{U}_j(z) = \widehat{U}_j^I(z) + \widehat{U}_j^{II}(z) + \widehat{U}_j^{III}(z),$$

where

$$\begin{aligned} \widehat{U}_j^I(z) = & \frac{(-1)^j \kappa_-^{j+1}(\varepsilon\xi)}{4g(\varepsilon\xi)} \times \left((g(\varepsilon\xi), -1) w^{II}(\varepsilon\xi) \right) \times r_+(\varepsilon\xi) \\ & -\frac{\kappa_-^j(\varepsilon\xi)}{\kappa_+(\varepsilon\xi) + \kappa_-(\varepsilon\xi)} \left(\Phi_-(\varepsilon\xi) w^I(\varepsilon\xi) + (-1)^j \Phi_+(\varepsilon\xi) w^{II}(\varepsilon\xi) \right) \\ (3.19) \quad & + \frac{1}{\kappa_+(\varepsilon\xi) + \kappa_-(\varepsilon\xi)} \times \left[\sum_{k=0}^{j-1} \kappa_-^{j-k}(\varepsilon\xi) \left(\Phi_-(\varepsilon\xi) + (-1)^{j-k-1} \Phi_+(\varepsilon\xi) \right) V_k^I \right. \\ & \left. + \sum_{k=j}^{+\infty} (-1)^{j-k} \kappa_+^{j-k}(\varepsilon\xi) \left(\Phi_-(\varepsilon\xi) + (-1)^{j-k-1} \Phi_+(\varepsilon\xi) \right) V_k^I \right], \end{aligned}$$

$$(3.20) \quad \widehat{U}_j^{II}(z) = \frac{(-1)^{j+1} \kappa_-^{j+1}(\varepsilon\xi)}{4g(\varepsilon\xi)} \times \frac{B_u - g(\varepsilon\xi)B_v}{B_u + g(\varepsilon\xi)B_v} \times \left((g(\varepsilon\xi), 1) w^I(\varepsilon\xi) \right) \times r_+(\varepsilon\xi)$$

and

$$\begin{aligned} \widehat{U}_j^{III}(z) = & \frac{\kappa_-^{j+1}(\varepsilon\xi)}{4g(\varepsilon\xi)} \times \left(\frac{B_u - g(\varepsilon\xi)B_v}{B_u + g(\varepsilon\xi)B_v} \times (g(\varepsilon\xi), 1) w^I(\varepsilon\xi) - (g(\varepsilon\xi), -1) w^{II}(\varepsilon\xi) \right) \\ (3.21) \quad & \times \frac{B_u - g(\varepsilon\xi)B_v}{B_u + g(\varepsilon\xi)B_v} \times r_-(\varepsilon\xi). \end{aligned}$$

Let us extend the initial data $(f_j)_{j \in \mathbb{Z}}$ to the whole line by setting $f_j = 0$ for $j \leq 0$. It is easy to verify that $\widehat{U}_j^I(z)$ corresponds to the \mathcal{Z} -transform of the solution $(U_j^I)^n$ of the following extended Cauchy problem (3.22).

$$(3.22) \quad \begin{cases} \frac{(U_j^I)^{n+1} - (U_j^I)^n}{\Delta t} + \frac{1}{2\Delta x} A \left((U_{j+1}^I)^{n+1} - (U_{j-1}^I)^{n+1} \right) = \frac{1}{\varepsilon} S (U_j^I)^{n+1}, & j \in \mathbb{Z}, n \geq 0, \\ (U_j^I)^0 = f_j, & j \in \mathbb{Z}. \end{cases}$$

With $(\widehat{U}_j^H)_{j \in \mathbb{N}}(z)$ found in (3.20), the value of $(U_j^H)_{j \in \mathbb{N}}^n$ can be obtained by inverting the \mathcal{Z} -transform

$$(3.23) \quad (U_j^H)^n = \frac{1}{2\pi} \int_{-\pi}^{\pi} \widehat{U}_j^H(Re^{i\theta}) R^n e^{in\theta} d\theta, \quad R > 1.$$

Indeed, the value of $\widehat{U}_j^{\text{III}}(z)$ can be reformulated as

$$\widehat{U}_j^{\text{III}}(z) = \frac{-B \left(\widehat{U}_0^I(z) + \widehat{U}_0^H(z) \right)}{B_u + g(\varepsilon\xi)B_v} \times \kappa_-(\varepsilon\xi)r_-(\varepsilon\xi).$$

Following Section 2.2, $\widehat{U}_j^{\text{III}}(z)$ corresponds to the \mathcal{Z} -transform of the solution $(U_j^{\text{III}})^n$ of the IBVP with the homogeneous initial data

$$(3.24) \quad \begin{cases} \frac{(U_j^{\text{III}})^{n+1} - (U_j^{\text{III}})^n}{\Delta t} + \frac{1}{2\Delta x} A \left((U_{j+1}^{\text{III}})^{n+1} - (U_{j-1}^{\text{III}})^{n+1} \right) = \frac{1}{\varepsilon} S (U_j^{\text{III}})^{n+1}, & j \geq 1, n \geq 0, \\ (U_j^{\text{III}})^0 = 0, & j \geq 0, \\ B (U_0^{\text{III}})^n = -B \left((U_0^I)^n + (U_0^H)^n \right), & n \geq 0, \\ \frac{1}{\Delta t} \Gamma \left((U_0^{\text{III}})^{n+1} - (U_0^{\text{III}})^n \right) + \frac{1}{2\Delta x} \Gamma A \left((U_1^{\text{III}})^{n+1} - \sum_{k=0}^{n+1} c_{n+1-k} (U_0^{\text{III}})^k \right) = \frac{1}{\varepsilon} \Gamma S (U_0^{\text{III}})^{n+1}, & n \geq 0. \end{cases}$$

3.2. The energy method for the Cauchy problem. In this paragraph, we prove the Proposition 1.2 by means of the discrete energy method. The energy estimate in the continuous case are obtained using the integration by parts rule. Therefore, we need the corresponding summation by parts rules for the discrete approximations of ∂/∂_x [12]. The idea is to find a symmetric positive definite matrix H , such that HA is symmetric and HS is negative definite. Therefore, we choose

$$H = \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}.$$

Now, let us multiply the first equation in (3.22) by $((U_j^I)^{n+1})^T H$ and sum over \mathbb{Z} , one obtains

$$(3.25) \quad \begin{aligned} \sum_{j \in \mathbb{Z}} \langle (U_j^I)^{n+1} - (U_j^I)^n, H(U_j^I)^{n+1} \rangle + \frac{\Delta t}{2\Delta x} \sum_{j \in \mathbb{Z}} \langle A \left((U_{j+1}^I)^{n+1} - (U_{j-1}^I)^{n+1} \right), H(U_j^I)^{n+1} \rangle \\ = \frac{\Delta t}{\varepsilon} \sum_{j \in \mathbb{Z}} \langle S(U_j^I)^{n+1}, H(U_j^I)^{n+1} \rangle, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ denotes the usual Euclidean inner product. Since H is a symmetric positive definite matrix, we have

$$\sum_{j \in \mathbb{Z}} \langle (U_j^I)^{n+1} - (U_j^I)^n, H(U_j^I)^{n+1} \rangle \geq \frac{1}{2} \sum_{j \in \mathbb{Z}} \left(\langle (U_j^I)^{n+1}, H(U_j^I)^{n+1} \rangle - \langle (U_j^I)^n, H(U_j^I)^n \rangle \right).$$

Together with the symmetric matrix HA , the second flux term in (3.25) becomes

$$\sum_{j \in \mathbb{Z}} \langle A((U_{j+1}^I)^{n+1} - (U_{j-1}^I)^{n+1}), H(U_j^I)^{n+1} \rangle = 0.$$

Thus, we directly get the inequality

$$(3.26) \quad \sum_{j \in \mathbb{Z}} (\langle (U_j^I)^{n+1}, H(U_j^I)^{n+1} \rangle - \langle (U_j^I)^n, H(U_j^I)^n \rangle) \leq \frac{2\Delta t}{\varepsilon} \sum_{j \in \mathbb{Z}} \langle S(U_j^I)^{n+1}, H(U_j^I)^{n+1} \rangle.$$

Let us remind that HS is negative definite. Then, from the inequality (3.26), for any $n \in \mathbb{N}$, the following inequality holds

$$(3.27) \quad \sum_{j \in \mathbb{Z}} \langle (U_j^I)^n, H(U_j^I)^n \rangle \leq \sum_{j \in \mathbb{Z}} \langle f_j, Hf_j \rangle.$$

Furthermore, since H is a symmetric positive definite matrix, the following inequality holds for some constants $m, k > 0$

$$(3.28) \quad m \langle (U_j^I)^n, H(U_j^I)^n \rangle \leq \langle (U_j^I)^n, (U_j^I)^n \rangle \leq k \langle (U_j^I)^n, H(U_j^I)^n \rangle.$$

According to (3.27) and (3.28), there exists a constant $C > 0$ such that

$$(3.29) \quad \sum_{j \in \mathbb{Z}} \Delta x \left| (U_j^I)^n \right|^2 \leq C \sum_{j \in \mathbb{Z}} \Delta x |f_j|^2, \quad \text{for any } n \in \mathbb{N},$$

with the constant C independent of ε and Δx .

This ends the proof of the Proposition 1.2.

To complete the proof of the Theorem 1.1 for the numerical scheme of the IBVP (3.1), observe that from (3.29) and setting $f_j = 0$ for $j < 0$, for any $T > 0$, there exists $C_T > 0$ such that

$$(3.30) \quad \sum_{n=0}^N \sum_{j \geq 0} \Delta x \Delta t \left| (U_j^I)^n \right|^2 \leq C_T \sum_{j \geq 0} \Delta x |f_j|^2,$$

with $N = T/\Delta t$. Furthermore, from the inequality (3.29) and $\Delta x = \Delta t \lambda_{xt}$, one obtains

$$(3.31) \quad \sum_{n=0}^N \Delta t \left| (U_0^I)^n \right|^2 \leq C_T \sum_{j \geq 0} \Delta x |f_j|^2.$$

3.3. The uniform estimate on $(U_j^H)^n$. The following lemma concerns the estimate on $(U_j^H)^n$:

Lemma 3.2. *Assume that the SKC (1.9) is satisfied and let $\lambda_{xt} \leq 3\sqrt{a}/8$ be a positive number. Then, for any $T > 0$, there exists a constant $C_T > 0$ such that for any $\Delta t > 0$ together with $\Delta x = \lambda_{xt}\Delta t$, for any $(f_j)_{j \in \mathbb{N}} \in \ell^2(\mathbb{N}, \mathbb{R}^2)$, the values of $(U_j^H)_{j \in \mathbb{N}}$ defined in (3.23) satisfy*

$$(3.32) \quad \sum_{n=0}^N \sum_{j \geq 0} \Delta t \Delta x \left| (U_j^H)^n \right|^2 + \sum_{n=0}^N \Delta t \left| (U_0^H)^n \right|^2 \leq C_T \sum_{j \geq 0} \Delta x |f_j|^2$$

where $N := T/\Delta t$ and C_T is independent of $\varepsilon \in (0, +\infty)$.

Proof. By an application of the following Plancherel's theorem for \mathcal{Z} -transform, we have

$$\begin{aligned} \sum_{n \geq 0} R^{-2n} \left| (U_0^H)^n \right|^2 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} |\widehat{U}_0^H(Re^{i\theta})|^2 d\theta, \quad R > 1 \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{|\kappa_-(\varepsilon\xi)|^2}{16|g(\varepsilon\xi)|^2} \times \left| \frac{B_u - g(\varepsilon\xi)B_v}{B_u + g(\varepsilon\xi)B_v} \right|^2 \times |(g(\varepsilon\xi), 1)w^I(\varepsilon\xi)|^2 \times (1 + |g(\varepsilon\xi)|^2) d\theta. \end{aligned}$$

From the Lemma 2.1, still under the SKC, $B_u + g(\varepsilon\xi)B_v$ is uniformly bounded away from 0 in $\varepsilon\xi \in \mathbb{C}_+$, and $g(\varepsilon\xi)$ is uniformly bounded in $\varepsilon\xi \in \mathbb{C}_+$. Moreover one has $|\kappa_-(\varepsilon\xi)| < 1$, we therefore obtain

$$(3.33) \quad \sum_{n \geq 0} R^{-2n} |(U_0^H)^n|^2 \lesssim \sum_{k \geq 0} |f_k|^2.$$

Similarly, by an application of the Plancherel's theorem for \mathcal{Z} -transform, under the SKC, we have

$$(3.34) \quad \sum_{n \geq 0} \sum_{j \geq 0} R^{-2n} |(U_j^H)^n|^2 \lesssim \sum_{j \geq 0} |\kappa_-(\varepsilon\xi)|^{2j} \sum_{k \geq 0} |f_k|^2.$$

Following Lemma 2.6, since we assume that the condition (2.29) holds, one gets the following property

$$\sum_{j \geq 0} |\kappa_-(\varepsilon\xi)|^{2j} \leq \frac{\Delta t \sqrt{a}}{\Delta x (1 - R^{-1})}.$$

Together with $\lambda_{xt} = \Delta x / \Delta t$, the inequality (3.34) becomes

$$(3.35) \quad \frac{R-1}{R} \sum_{n \geq 0} \sum_{j \geq 0} R^{-2n} \Delta x |(U_j^H)^n|^2 \lesssim \sum_{k \geq 0} \Delta x |f_k|^2.$$

Assembling the estimates (3.33) and (3.35), there exists $C > 0$ such that

$$\frac{R-1}{R} \sum_{n \geq 0} \sum_{j \geq 0} R^{-2n} \Delta x |(U_j^H)^n|^2 + \sum_{n \geq 0} \sum_{j \geq 0} R^{-2n} \Delta t |(U_0^H)^n|^2 \leq C \sum_{k \geq 0} \Delta x |f_k|^2.$$

By setting in the above formula $R = e^{\gamma \Delta t}$ for $\gamma > 0$ and $\Delta t > 0$, and using the classical lower bound $e^{\gamma \Delta t} \geq 1 + \gamma \Delta t$, we obtain that there exists a constant $c > 0$ such that

$$\frac{\gamma}{\gamma \Delta t + 1} \sum_{n \geq 0} \sum_{j \geq 0} e^{-2\gamma n \Delta t} \Delta t \Delta x |(U_j^H)^n|^2 + \sum_{n \geq 0} e^{-2\gamma n \Delta t} \Delta t |(U_0^H)^n|^2 \leq C \sum_{k \geq 0} \Delta x |f_k|^2.$$

Then, for all $T > 0$, there exists a constant $C_T > 0$ such that

$$\sum_{n=0}^N \sum_{j \geq 0} \Delta t \Delta x |(U_j^H)^n|^2 + \sum_{n=0}^N \Delta t |(U_0^H)^n|^2 \leq C_T \sum_{k \geq 0} \Delta x |f_k|^2,$$

with $N = T / \Delta t$. □

3.4. Stiff stability analysis. Following Section 2.3, for any $T > 0$, there exists $C_T > 0$ such that the solution $(U_j^{III})_{j \in \mathbb{N}}^n$ to (3.24) satisfies

$$\sum_{n=0}^N \sum_{j \geq 0} \Delta x \Delta t |(U_j^{III})^n|^2 + \sum_{n=0}^N \Delta t |(U_0^{III})^n|^2 \leq C_T \sum_{n=0}^N \Delta t \left| B \left((U_0^I)^n + (U_0^H)^n \right) \right|^2.$$

Furthermore, from the inequalities (3.31) and (3.32), one obtains

$$\sum_{n=0}^N \Delta t \left| B \left((U_0^I)^n + (U_0^H)^n \right) \right|^2 \leq C_T \sum_{k \geq 0} \Delta x |f_k|^2.$$

Therefore, we show the uniform estimate on $(U_j^{III})_{j \in \mathbb{N}}^n$

$$(3.36) \quad \sum_{n=0}^N \sum_{j \geq 0} \Delta x \Delta t |(U_j^{III})^n|^2 + \sum_{n=0}^N \Delta t |(U_0^{III})^n|^2 \leq C_T \sum_{j \geq 0} \Delta x |f_j|^2,$$

with the positive constant C_T independent of ε , Δx and Δt .

To complete the proof of the Theorem 1.1 for the numerical scheme of the IBVP (3.1), observe that from the inequalities (3.30) and (3.36), for any $T > 0$, there exists $C_T > 0$ such that for any $(f_j)_{j \in \mathbb{N}} \in \ell^2(\mathbb{N}, \mathbb{R}^2)$ and $N := T/\Delta t$, the solution $(U_j^n)_{j \in \mathbb{N}}$ to (3.1) satisfies

$$(3.37) \quad \sum_{n=0}^N \sum_{j \geq 0} \Delta x \Delta t |U_j^n|^2 + \sum_{n=0}^N \Delta t |U_0^n|^2 \leq C_T \sum_{j \geq 0} \Delta x |f_j|^2,$$

where the constant C_T independent of $\varepsilon, \Delta x$ and Δt . This is the last step to prove the Theorem 1.1.

3.5. Numerical experiments. In this paragraph, we present some numerical experiments for the behavior of the numerical solution $(U_j^n)_{j \in \mathbb{N}}$ corresponding to whether or not the SKC (1.9) holds. We also look at the numerical solution when the UKC (1.4) is wrong. After that, we observe the effective behavior of the energy terms E_1 inside the domain and E_2 along the boundary, which are defined in (2.37).

In our numerical experiments, we choose $a = 1$, $B_v = 1$, $\lambda_{xt} = 1/3$, fix the space step $\Delta x = 5 \times 10^{-3}$, the time step $\Delta t = \lambda_{xt}^{-1} \Delta x$, and let the relaxation rate ε and the boundary data B_u vary. The boundary data is the homogeneous one $b^n \equiv 0$, for any $n \in \mathbb{N}$. The initial data is

$$f_j = \begin{cases} 100 \times \begin{pmatrix} \frac{13}{30} - x_j \\ x_j - \frac{1}{4} \end{pmatrix} \times \begin{pmatrix} 1 & -1 \end{pmatrix}^T, & \text{if } x_j \in \left[\frac{1}{4}, \frac{13}{30} \right], \\ \begin{pmatrix} 0 & 0 \end{pmatrix}^T, & \text{otherwise.} \end{cases}$$

Let us first observe that these data are compatible in the corner $(x, t) = (0, 0)$ in the sense that $Bf_0 = 0$. Moreover, the choice of an initial data with support in $[1/4, 13/30]$ is motivated by the property of finite speed of propagation available at the continuous side (1.1). More precisely, the exact solution we approximate has characteristic velocities ± 1 and therefore vanishes outside some space interval $[0, 0.63]$ for small times in $[0, 0.2]$. Thus, we choose for our experiments the space interval $[0, 1]$ and the time interval $[0, T]$ with $T = 0.2$. Let us mention that the numerical experiments are performed with another discrete right boundary condition at $x = 1$. This is chosen to be the classical homogeneous first order Neumann extrapolation boundary condition $U_{j+1}^n = U_j^n$, for any $n \in \mathbb{N}$, at the rightmost cell J . That boundary condition indeed exhibits convenient stability features for both the inflowing and the outflowing transport equation [10].

3.5.1. The behavior of the numerical solution. Firstly, we choose a set of values B_u such that the SKC (1.9) is satisfied with $\varepsilon = 10^{-2}$ and also with $\varepsilon = 10^2$. The Figures 3.1 and 3.2 show numerical solution $(U_j^n)_{j \in \mathbb{N}}$ over the time interval $t \in [0, 0.2]$.

In the first case, $\varepsilon = 10^{-2}$, due to the initial relaxation of solution to the equilibrium system, the numerical solution descends over time (see Figure 3.1). In the case $\varepsilon = 10^2$, at time $t < 0.2$, its solution seems to translate to the left and the ghost solutions do not go backward in space for the implicit scheme. After that, the initial condition re-enters the domain from the left boundary (see Figure 3.2). It is not so much influenced by relaxation source term but more by the boundary dissipation.

Secondly, we choose the value of B_u such that the SKC (1.9) is not satisfied. Besides, we also present the numerical solution when the Uniform Kreiss Condition (1.4) is wrong. The Figures 3.3 and 3.4 show numerical solution $(U_j^n)_{j \in \mathbb{N}}$ over the time interval $t \in [0, 0.2]$.

We can observe that the numerical solution at the boundary rise gradually when the SKC (1.9) fails. This is the case for example for $\varepsilon = 10^{-2}$ together with the parameters $(B_u, B_v) = (-1/2, 1)$. When the UKC (1.4) does not hold, the behavior is even worse (see Figure 3.4).

3.5.2. The effective behavior of the energy terms. We present hereafter the effective behavior of the energy terms E_1 and E_2 for $\varepsilon \in (0, +\infty)$, $T = 0.2$ and $N = T/\Delta t$.

According to Table 4 and Figures 3.5, 3.6, we can see that

- For any $\varepsilon \in (0, +\infty)$, the values of E_1 and E_2 rise gradually when the SKC (1.9) is not satisfied. This is the case for example for $\varepsilon = 10^2$ together with the parameters $(B_u, B_v) = (-1/2, 1)$. The behavior is even worse when the UKC (1.4) is not hold.




FIGURE 3.1. The numerical solution $u(x, t)$ (left) and $v(x, t)$ (right) for $\varepsilon = 10^{-2}$. The SKC (1.9) is valid with $B_u = -2$.

B_u	$\varepsilon = 10^{-2}$	$\varepsilon = 1$	$\varepsilon = 10^2$
-4	0.038009	0.083375	0.160487
-2	0.038014	0.191338	0.440354
-1	7.54×10^{19}	8.94×10^{30}	3.149×10^{41}
-0.5	696.71	16628.4	101893.6
1	0.030684	0.03537	0.038031
3	0.035051	0.038022	0.046627

B_u	$\varepsilon = 10^{-2}$	$\varepsilon = 1$	$\varepsilon = 10^2$
-4	2.13×10^{-5}	1.05×10^{-3}	1.46×10^{-3}
-2	2.69×10^{-5}	0.43×10^{-2}	0.71×10^{-2}
-1	5.3×10^{21}	8.68×10^{32}	3.58×10^{43}
-0.5	19235.1	125624.5	437872.2
1	2.87×10^{-5}	0.0515	0.0715
3	1.9×10^{-5}	0.0620	0.0894

TABLE 4. The energy terms E_1 (left) and E_2 (right).

- On the boundary $x = 0$, the value of E_2 for $\varepsilon = 10^{-2}$ increase slowly. This is due to the effect of incoming solution at the boundary when the initial relaxation of solution tends to the equilibrium system. In the case $\varepsilon = 10^2$, its value increase fairly rapidly. It is not so much influenced by relaxation source term but more by the boundary dissipation.

Clearly, in our numerical experiment, the numerical solution at the boundary $x = 0$ increase quickly as soon as the SKC (1.9) is not valid. The behavior of numerical solution is even worse if the UKC (1.4) is not satisfied. Indeed, it seems that the SKC (1.9) is also necessary condition to ensure the non-increase rapidly of the energy terms E_1 and E_2 under the effect of the relaxation source term and the boundary dissipation.

APPENDIX A. TECHNICAL LEMMAS

Lemma A.1. *Let $\zeta \in \mathbb{C}$ with $\operatorname{Re} \zeta > 0$ and $h(\zeta) = \sqrt{\zeta(1 + \zeta)}$, then $\operatorname{Re} \zeta \leq \operatorname{Re} h(\zeta)$.*

Proof. In the half plane $\{\zeta \in \mathbb{C} : \operatorname{Re} \zeta \geq 0\}$, the complex function $h(\zeta)$ is analytic. As usual, we take $\sqrt{\zeta}$ to be the principal branch with the branch cut along the negative real axis.

Let $\zeta = x + yi$ with $x \geq 0$, $y \in \mathbb{R}$ and

$$p = x(1 + x) - y^2, \quad q = (1 + 2x)y.$$

Then,

$$\operatorname{Re} h(\zeta) = \operatorname{Re} \sqrt{p + qi} = \sqrt{\frac{p + \sqrt{p^2 + q^2}}{2}}.$$




FIGURE 3.2. The numerical solution $u(x,t)$ (left) and $v(x,t)$ (right) for $\varepsilon = 10^2$. The SKC (1.9) is valid with $B_u = -2$.

Now, we observe that

$$\begin{aligned} \sqrt{p^2 + q^2} &= \sqrt{(x(1+x) - y^2)^2 + (1+2x)^2 y^2} \\ &= \sqrt{(x(1+x) + y^2)^2 + y^2} \\ &\geq x(1+x) + y^2. \end{aligned}$$

Therefore,

$$\operatorname{Re} h(\zeta) \geq \sqrt{x(1+x)} \geq x.$$

This ends the proof of Lemma A.1. □

Lemma A.2. Let $\zeta \in \mathbb{C}$ with $\operatorname{Re} \zeta < 0$, then $|\zeta + \sqrt{\zeta^2 + 1}| \leq \operatorname{Re} \zeta + \sqrt{(\operatorname{Re} \zeta)^2 + 1} < 1$.

Proof. Assume that $\zeta = x + yi$ with $x < 0$ and $y \in \mathbb{R}$.

Case 1: Consider first the easy case $y = 0$. Then

$$\left| \zeta + \sqrt{\zeta^2 + 1} \right| = x + \sqrt{x^2 + 1},$$




FIGURE 3.3. The numerical solution $u(x,t)$ (left) and $v(x,t)$ (right) for $\varepsilon = 10^{-2}$. The SKC (1.9) is not valid with $B_u = -0.5$.




FIGURE 3.4. The numerical solution $u(x,t)$ (left) and $v(x,t)$ (right) for $\varepsilon = 10^2$. The UKC (1.4) is wrong with $B_u = -1$.

but since $x < 0$, one obtains by simple considerations the inequality $x + \sqrt{x^2 + 1} < 1$.

Case 2: In the general case $y \neq 0$, let us begin with some notations:

$$\zeta^2 + 1 = p_1 + q_1 i, \quad \text{with } p_1 = x^2 - y^2 + 1 \text{ and } q_1 = 2xy,$$



$$\sqrt{\zeta^2 + 1} = a_1 + b_1 i, \quad \text{with } a_1 = \sqrt{\frac{p_1 + \sqrt{p_1^2 + q_1^2}}{2}} \text{ and } b_1 = \text{sgn}(q_1) \sqrt{\frac{-p_1 + \sqrt{p_1^2 + q_1^2}}{2}}.$$

Together with these notations, some algebraic identities are available:

$$(A.1) \quad x^2 + b_1^2 + 1 = a_1^2 + y^2 \quad \text{and} \quad y = \frac{a_1 b_1}{x},$$

Firstly, we prove the next inequality

$$(A.2) \quad a_1 x^2 + b_1^2 x + a_1 b_1^2 \geq 0.$$

FIGURE 3.5. Energy evolution E_1 for $B_u = -4$ (left) and $B_u = -0.5$ (right).FIGURE 3.6. Energy evolution E_2 for $B_u = -4$ (left) and $B_u = -0.5$ (right).

We can see that the inequality (A.2) is equivalent to $a_1(x^2 + b_1^2) \geq -xb_1^2$ and since $x < 0$, the latter is now equivalent to its squared version, that reads

$$a_1^2 x^2 (x^2 + 2b_1^2) \geq b_1^4 (x^2 - a_1^2).$$

By the definition of a_1, b_1 above, the previous inequality is

$$\begin{aligned} & 4x^2 \left(p_1 + \sqrt{p_1^2 + q_1^2} \right) \left(x^2 - p_1 + \sqrt{p_1^2 + q_1^2} \right) \geq \left(-p_1 + \sqrt{p_1^2 + q_1^2} \right)^2 \left(2x^2 - p_1 - \sqrt{p_1^2 + q_1^2} \right) \\ \Leftrightarrow & 4x^4 \left(p_1 + \sqrt{p_1^2 + q_1^2} \right) + 2x^2 q_1^2 \geq \left(p_1 - \sqrt{p_1^2 + q_1^2} \right) (4x^2 p_1 + q_1^2) \\ \Leftrightarrow & 4x^4 \left(p_1 + \sqrt{p_1^2 + q_1^2} \right) + 2x^2 \times 4x^2 y^2 \geq \left(p_1 - \sqrt{p_1^2 + q_1^2} \right) (4x^2 (x^2 - y^2 + 1) + 4x^2 y^2) \\ \Leftrightarrow & 2x^2 \left(\sqrt{p_1^2 + q_1^2} + y^2 \right) \geq p_1 - \sqrt{p_1^2 + q_1^2}. \end{aligned}$$

But, for any $p_1, q_1 \in \mathbb{R}$, this is easy to see that $p_1 - \sqrt{p_1^2 + q_1^2} \leq 0$ and thus any of the previous inequalities and so the expected one (A.2) follow.

Now let us observe that the required inequality $|\zeta + \sqrt{\zeta^2 + 1}| \leq x + \sqrt{x^2 + 1}$ is fully equivalent to

$$(A.3) \quad (x + a_1)^2 + (y + b_1)^2 \leq (x + \sqrt{x^2 + 1})^2,$$

that we prove now. According to the algebraic identities in (A.1), by eliminating the occurrences of y , the previous formula is equivalent to

$$a_1x + b_1^2 + \frac{a_1b_1^2}{x} \leq x\sqrt{x^2 + 1}.$$

In addition, we observe that $x^2 + 1 = x^{-2}(a_1^2x^2 + a_1^2b_1^2 - b_1^2x^2)$, and thus the previous inequality is equivalent to

$$(A.4) \quad a_1x + b_1^2 + \frac{a_1b_1^2}{x} \leq -\sqrt{a_1^2x^2 + a_1^2b_1^2 - b_1^2x^2}.$$

Since $x < 0$ and from the inequality (A.2), the formula (A.4) reads also

$$\begin{aligned} (a_1x^2 + b_1^2x + a_1b_1^2)^2 &\geq x^2(a_1^2x^2 + a_1^2b_1^2 - b_1^2x^2) \\ \Leftrightarrow (x + a_1)^2(x^2 + b_1^2) &\geq 0. \end{aligned}$$

This ends the proof of the inequality (A.3). Now since $\operatorname{Re} \zeta < 0$, the analysis of the first easy case again applies to get $\operatorname{Re} \zeta + \sqrt{(\operatorname{Re} \zeta)^2 + 1} < 1$.

This ends the proof of Lemma A.2. □

REFERENCES

- [1] A. Arnold, M. Ehrhardt, and I. Sofronov. Discrete transparent boundary conditions for the Schrödinger equation: fast calculation, approximation, and stability. *Communications in Mathematical Sciences*, 1(3):501–556, 2003.
- [2] S. Benzoni-Gavage and D. Serre. *Multi-dimensional hyperbolic partial differential equations, First-order systems and applications*. Oxford Mathematical Monographs, 2006.
- [3] C. Besse, M. Ehrhardt, and I. Lacroix-Violet. Discrete artificial boundary conditions for the linearized Korteweg–de Vries equation. *Numerical Methods for Partial Differential Equations. An International Journal*, 32(5):1455–1484, 2016.
- [4] C. Besse, P. Noble, and D. Sanchez. Discrete transparent boundary conditions for the mixed KDV–BBM equation. *Journal of Computational Physics*, 345:484–509, 2017.
- [5] B. Boutin, T. H. T. Nguyen, and N. Seguin. A stiffly stable semi-discrete scheme for the characteristic linear hyperbolic relaxation with boundary. *ESAIM: Mathematical Modelling and Numerical Analysis*, 2020.
- [6] C. Cercignani. *The Boltzmann Equation*. Applied Mathematical Sciences. Springer, New York, NY, 1988.
- [7] G. Q. Chen, C. D. Levermore, and T.-P. Liu. Hyperbolic conservation laws with stiff relaxation terms and entropy. *Communications on Pure and Applied Mathematics*, 47(6):787–830, 1994.
- [8] P. Colella, A. Majda, and V. Roytburd. Theoretical and Numerical Structure for Reacting Shock Waves. *SIAM Journal on Scientific and Statistical Computing*, 7(4):1059–1080, 1986.
- [9] J.-F. Coulombel. Transparent numerical boundary conditions for evolution equations: derivation and stability analysis. *Annales de la Faculté des Sciences de Toulouse. Mathématiques. Série 6*, 28(2):259–327, 2019.
- [10] M. Goldberg. On a boundary extrapolation theorem by Kreiss. *Mathematics of Computation*, 31(138):469–477, 1977.
- [11] M. J. Grote and J. B. Keller. Exact Nonreflecting Boundary Conditions for the Time Dependent Wave Equation. *SIAM Journal on Applied Mathematics*, 55(2):280–297, 1995.
- [12] B. Gustafsson, H.-O. Kreiss, and J. Oliger. *Time dependent problems and difference methods*. John Wiley & Sons, 2013.
- [13] B. Hanouzet and R. Natalini. Global Existence of Smooth Solutions for Partially Dissipative Hyperbolic Systems with a Convex Entropy. *Archive for Rational Mechanics and Analysis*, 169(2):89–117, 2003.
- [14] S. Jin and Z. Xin. The relaxation schemes for systems of conservation laws in arbitrary space dimensions. *Communications on Pure and Applied Mathematics*, 48(3):235–276, 1995.
- [15] E. I. Jury. *Theory and Application of the Z-Transform Method*. Wiley and Sons, New York, 1964.
- [16] M. Kazakova and P. Noble. Discrete Transparent Boundary Conditions for the Linearized Green–Naghdi System of Equations. *SIAM Journal on Numerical Analysis*, 58(1):657–683, 2020.
- [17] E. M. L. L. D. Landau. *Statistical Physics*. Elsevier (Singapore) Pte Ltd, 1980.
- [18] T.-P. Liu. Hyperbolic conservation laws with relaxation. *Communications in Mathematical Physics*, 108(1):153–175, 1987.
- [19] A. V. Oppenheim, A. S. Willsky, and S. H. Nawab. *Signals and Systems (2Nd Ed.)*. Prentice-Hall, Inc., Upper Saddle River, NJ, USA, 1996.

- [20] J. J. Stoker. *Water waves. The mathematical theory with applications*. New York, NY: Wiley, reprint of the 1957 original edition, 1992.
- [21] W. G. Vincenti and C. H. Kruger. *Introduction to Physical Gas Dynamics*, volume 70. John Wiley and Sons, 1966.
- [22] G. B. Whitham. *Linear and nonlinear waves*. John Wiley & Sons, Hoboken, NJ, 1974.
- [23] Z. Xin and W.-Q. Xu. Stiff Well-Posedness and Asymptotic Convergence for a Class of Linear Relaxation Systems in a Quarter Plane. *Journal of Differential Equations*, 167(2):388–437, 2000.
- [24] W.-A. Yong. Boundary conditions for hyperbolic systems with stiff source terms. *Indiana University Mathematics Journal*, 48(1):115–137, 1999.
- [25] W.-A. Yong. Singular Perturbations of First-Order Hyperbolic Systems with Stiff Source Terms. *Journal of Differential Equations*, 155(1):89–132, 1999.
- [26] W.-A. Yong. Entropy and Global Existence for Hyperbolic Balance Laws. *Archive for Rational Mechanics and Analysis*, 172(2):247–266, 2004.

IRMAR (UMR CNRS 6625), UNIVERSITÉ DE RENNES, CAMPUS DE BEAULIEU, 35042 RENNES CEDEX, FRANCE.

Email address: `benjamin.boutin@univ-rennes1.fr`

Email address: `thi-hoai-thuong.nguyen@univ-rennes1.fr`

Email address: `nicolas.seguin@univ-rennes1.fr`