



**HAL**  
open science

# Nitro End Groups Remarkable Vibrational Reporters for Charge Transfer in the Excited States of Oligo(p-phenyleneethynylene)-Bridged Donor-Acceptor Dyads

J. Kubicki, Maciej Lorenc, P. Cochelin, O. Mongin, A. Amar, A. Boucekkine,  
A. Gaje, M.G. Humphrey, M. Morshedi, S. Lorenzen, et al.

► **To cite this version:**

J. Kubicki, Maciej Lorenc, P. Cochelin, O. Mongin, A. Amar, et al.. Nitro End Groups Remarkable Vibrational Reporters for Charge Transfer in the Excited States of Oligo(p-phenyleneethynylene)-Bridged Donor-Acceptor Dyads. *Journal of Physical Chemistry C*, 2020, 124 (18), pp.9755-9764. 10.1021/acs.jpcc.0c01532 . hal-02862821

**HAL Id: hal-02862821**

**<https://univ-rennes.hal.science/hal-02862821>**

Submitted on 15 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Nitro End-Groups: Remarkable Vibrational Reporters for Charge Transfer in the Excited States of Oligo(*p*-Phenyleneethynylene)-Bridged Donor-Acceptor Dyads

*Jacek Kubicki,<sup>\*,[a]</sup> Maciej Lorenc,<sup>[b]</sup> Pierre Cochelin,<sup>[a,b]</sup> Olivier Mongin,<sup>[c]</sup> Anissa Amar,<sup>[d]</sup> Abdou Boucekkine,<sup>[c]</sup> Arnold Gaje,<sup>[a,b]</sup> Mark G. Humphrey,<sup>[e]</sup> Mahbod Morshedi,<sup>[e]</sup> Sabine Lorenzen,<sup>[f]</sup> Florian Rauch,<sup>[f]</sup> Charlotte Scheufler,<sup>[f]</sup> Todd B. Marder<sup>[f]</sup> and Frédéric Paul<sup>\*,[c]</sup>*

[a] Adam Mickiewicz University, Faculty of Physics, Uniwersytetu Poznańskiego 2, 61-614 Poznań, Poland, [b] Univ Rennes, CNRS, IPR (Institut de Physique de Rennes) - UMR 6251, F-35000 Rennes, France, [c] Univ Rennes, CNRS, ISCR (Institut des Sciences Chimiques de Rennes) - UMR 6226, F-35000, Rennes, France, [d] Département de Chimie, Faculté des Sciences, Université Mouloud Mammeri, 15000 Tizi-Ouzou, and Faculté de Chimie, Université des Sciences et de la Technologie Houari-Boumediene, 16111 Bab-Ezzouar (Algeria). [e] Research School of Chemistry, Australian National University, Canberra, ACT 2601, Australia. [f] Institut für Anorganische Chemie and Institute for Sustainable Chemistry & Catalysis with Boron (ICB), Julius-Maximilians-Universität Würzburg, Am Hubland, 97074 Würzburg (Germany).

## AUTHOR INFORMATION

**Corresponding Author**


\* jacek.kubicki@amu.edu.pl, frederic.paul@univ-rennes1.fr

**Abstract.** The D- $\pi$ -A dyads [(*n*-Hex)<sub>2</sub>N(1,4-C<sub>6</sub>H<sub>4</sub>)]C $\equiv$ C[(1,4-C<sub>6</sub>H<sub>4</sub>)NO<sub>2</sub>] (**1**) and [(*n*-Hex)<sub>2</sub>N(1,4-C<sub>6</sub>H<sub>4</sub>)]C $\equiv$ C(1,4-C<sub>6</sub>H<sub>4</sub>)C $\equiv$ C[(1,4-C<sub>6</sub>H<sub>4</sub>)NO<sub>2</sub>] (**2**) have been studied by ultrafast time-resolved infrared absorption spectroscopy. After excitation into their singlet charge-transfer (<sup>1</sup>CT) state, a fast decay (ca. 6 ps for **1** and 1 ps for **2**) of the initially populated singlet state into a ground state (major pathway) and a longer-lived excited state (possibly the triplet state) is observed. The nitro and alkyne groups were used as vibrational reporters to probe the changes induced by the charge-transfer process. For the first time, we confirm experimentally that these changes are consistent with expectations based on the traditional valence-bond representations of the CT states of these push-pull chromophores. An almost identical charge transfer takes place in the two dyads, despite different length  $\pi$ -bridges between the donor and acceptor groups. Complementary DFT calculations support the experimental assignments and have helped clarifying the photophysical behavior of **1** and **2**.

## 1 Introduction.

Donor-acceptor (D-A) dyads are very important chromophores that play a pivotal role in an increasing number of key societal applications<sup>1</sup> such as solar light harvesting,<sup>2-4</sup> molecular sensing,<sup>5</sup> bioimaging,<sup>6</sup> or ultrafast information treatment.<sup>7,8</sup> Evaluation of their performance in each of these fields is crucial, and such a task often relies on a detailed knowledge of the donor-acceptor charge-transfer state which usually constitutes the lowest allowed excited state of these compounds.<sup>1</sup> While theoretical modelling of such excited states can nowadays be undertaken routinely, the exact degree of charge transfer (CT) between the donor and acceptor end-groups has remained elusive.<sup>9</sup> Ultrafast transient vibrational spectroscopies can greatly facilitate the optimization of compounds<sup>10,11</sup> because charge-transfer leads to changes in various bond strengths in the excited molecule. Monitoring vibrational modes after photo-excitation can therefore provide reliable experimental insight into the structural changes underlying the charge-transfer process. A ubiquitous class of dyads are those in which the donor and acceptor groups are separated by oligo(1,4-phenyleneethynylene)-containing bridges. This type of bridge is rigid, highly versatile from the synthetic perspective, and chemically very resistant.<sup>12-20</sup> Furthermore, alkyne vibrations are found in a specific range in the infrared (IR) region (1900-2200 cm<sup>-1</sup>), so these can provide a convenient probe to monitor changes in the bridge, as recently shown by Vauthey and coworkers<sup>10</sup> and others.<sup>21-23</sup> Herein, we report our results on related nitro-terminated dyads (Scheme 1), and show that the nitro group is a remarkable vibrational reporter on the acceptor side of the charge-transfer. The ground states (GSs) of such molecules typically exhibit two N-O stretching modes that are easily recognizable in their IR vibrational spectra, namely an antisymmetric stretch in the 1500-1600 cm<sup>-1</sup> region and a symmetric stretch in the 1400-1300 cm<sup>-1</sup> region.<sup>24</sup> These vibrational modes are directly affected by any change in electron

1  
2  
3 density at the nitro group and, as such, they should provide an indication of the amount of charge  
4 transferred to this group. Indeed, upon population of the CT state, the increase in electron density  
5 at the nitro group induces a weakening of the N-O bonds, which results in red-shifts of both the  
6 antisymmetric and symmetric stretching modes. Importantly, because the nitro group is redox-  
7 active, the magnitude of the vibrational changes induced by transfer of one electron to this group  
8 may be experimentally estimated by spectroelectrochemistry, by monitoring the specific  
9 vibrational  $\nu(\text{N}=\text{O})$  modes upon one-electron reduction. Based on existing literature,<sup>25-27</sup> selected  
10 experiments reveal shifts to lower wavenumbers of ca.  $280\text{ cm}^{-1}$  for the antisymmetric and  
11 symmetric stretching modes of nitrobenzene, upon one-electron reduction of the nitro group  
12 attached to a phenyl ring when these modes are modelled by DFT computations and tracked by  
13 projection analysis.<sup>28,29</sup>



34 Scheme 1. Molecular structures of **1** and **2**.

35  
36  
37 As we shall now demonstrate with  $[(n\text{-Hex})_2\text{N}(1,4\text{-C}_6\text{H}_4)]\text{C}\equiv\text{C}[(1,4\text{-C}_6\text{H}_4)\text{NO}_2]$  (**1**) and  $[(n\text{-Hex})_2\text{N}(1,4\text{-C}_6\text{H}_4)]\text{C}\equiv\text{C}(1,4\text{-C}_6\text{H}_4)\text{C}\equiv\text{C}[(1,4\text{-C}_6\text{H}_4)\text{NO}_2]$  (**2**) (Scheme 1), two examples of  
38 organic oligo(1,4-phenyleneethynylene)-bridged D-A dyads featuring a terminal nitro group,  
39 time-resolved infrared spectroscopy (TRIR) studies of the specific infrared signatures of the  
40 alkyne and nitro groups afford a comprehensive picture of the electronic changes taking place  
41 subsequent to excitation into the CT band. These dyads were chosen for this study because the  
42 dialkylamino moiety is a commonly encountered donor group, the hexyl chains being employed  
43 here to ensure sufficient solubility in the (organic) solvent in which the studies were undertaken.

1  
2  
3 Furthermore, comparison between **1** and **2** might facilitate a quantitative measure of the impact  
4  
5 of bridge extension upon charge separation in the CT state.  
6  
7

## 8 9 **2. Methods.**

10  
11  
12 *2.1. Ultrafast mid-IR spectrometer.* The ultrafast laser system consists of a femtosecond  
13  
14 titanium sapphire oscillator (Mai Tai, Spectra Physics) and a high energy titanium sapphire  
15  
16 regenerative amplifier (Spitfire Ace, Spectra Physics, 100 fs, 1 kHz, 5 mJ) pumped by Empower  
17  
18 (1 kHz, 20 W). The 800 nm beam coming out of Spitfire Ace was divided by beam splitters in  
19  
20 order to simultaneously pump Topas Prime with a NirUVis frequency mixer (420 nm excitation  
21  
22 wavelength was chosen to selectively excite the CT state in both compounds) and Topas Prime  
23  
24 with a DFG mixer (mid-IR probe wavelength with the central wavelength tuned from about 1000  
25  
26  $\text{cm}^{-1}$  until about 3300  $\text{cm}^{-1}$ ). 2DQuick Transient spectrometer (PhaseTech Spectroscopy) was  
27  
28 used to record transient spectra. As mid-IR detector MCT array was applied (FPAS-01444, 2 x  
29  
30 64 pixels, Infrared System Development). The dispersion of the spectra were obtained by  
31  
32 iHR320 (Horiba) monochromator equipped with set of three grating with different blaze angles  
33  
34 and different number of grooves per mm (8  $\mu\text{m}$ , 75 grooves/mm; 6  $\mu\text{m}$ , 100 grooves/mm; 4  $\mu\text{m}$ ,  
35  
36 75 grooves/mm) in order to cover the whole spectral range of interest (1000 – 3300  $\text{cm}^{-1}$ ) and to  
37  
38 achieve the proper spatial width of the spectrum illuminating MCT array. The angle between  
39  
40 polarizations of the pump beam and the probe beam was set at the magic angle (54.7°) in order to  
41  
42 avoid rotational effects on measured kinetics. The transient mid-IR experiments were performed  
43  
44 with a flow cell (Harrick Scientific) with 0.5 mm optical path equipped by 2 mm thick  $\text{BaF}_2$  front  
45  
46 window and the  $\text{CaF}_2$  2 mm thick back window and the sample volume was about 20  $\mu\text{L}$ .  
47  
48 Absorbance of the solution (0.5 mm solution layer) was around 0.3 at the pump wavelength. The  
49  
50 instrument response function (IRF) was about 200 fs (measured as the FWHM). The solution  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 was purged by dry nitrogen to avoid presence of the oxygen in the solution. Typical pump  
4 energy was about 1.5  $\mu$ J. The entire set of pump probe delay positions was repeated at least four  
5  
6 times in order to improve signal-to-noise ratio and also to check data reproducibility. Ultrafast  
7  
8 experiments were repeated on other day to additionally check data reproducibility.  
9

10  
11  
12  
13 *2.2. Synthesis and solvents.* Unless otherwise noted in SI, the following conditions apply. All  
14 syntheses were carried out using standard Schlenk and glovebox techniques under an argon  
15 atmosphere. The solvents used were dried using a solvent purification system (SPS) from  
16 Innovative Technology and were degassed and stored over molecular sieves under argon.  
17  
18 Deuterated solvents ( $\text{CDCl}_3$ ,  $\text{C}_6\text{D}_6$ ) used for NMR spectroscopy were purchased from Cambridge  
19  
20 Isotope Laboratories.  $\text{C}_6\text{D}_6$  was dried over molecular sieve and stored under argon atmosphere  
21  
22 before use. *N,N*-Dihexyl-4-iodoaniline<sup>30</sup> and 1-ethynyl-4-nitrobenzene<sup>31</sup> were prepared  
23  
24 according to the literature. All other starting materials were purchased from commercial sources  
25  
26 and were used without further purification. Characterization and photophysical and  
27  
28 photochemical studies of **1** and **2**, together with important remarks regarding experiments in  
29  
30  $\text{CH}_2\text{Cl}_2$ , are described in the Supporting Information (SI;  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR: Figure S1-S2;  
31  
32 photophysical studies: Table S1, Figure S3-S6). The solvents for ultrafast studies were  
33  
34 spectrophotometric grade from Sigma-Aldrich.  
35  
36  
37  
38  
39  
40  
41  
42  
43

44 *2.3. Calculations.* All calculations reported in this work have been carried out using the  
45 Gaussian09-D01 suite of programs.<sup>32</sup> In order to simplify calculations the hexyl chains from  
46  
47 compound **1** and **2** were replaced by methyl groups and the compounds were named as **1'** and **2'**,  
48  
49 respectively. Their geometry optimizations have been performed employing the MPW1PW91<sup>33</sup>  
50  
51 functional in combination with the 6-31G\* basis set. Solvent effects, namely  $\text{CH}_2\text{Cl}_2$ , have been  
52  
53 simulated by means of the polarizable continuum model (PCM).<sup>34</sup> Next, TD-DFT calculations  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 have been carried out not only to compute the electronic spectra. The geometries of the singlet  $S_1$   
4 and  $S_2$  states of the molecules under consideration were then optimized. The calculations of the  
5 normal modes of vibration of the  $S_0$ ,  $S_1$  and  $S_2$  states have subsequently been performed. The  
6 UV-visible spectra have been simulated not only at the MPW1PW91 level but also with the  
7 CAM-B3LYP<sup>35</sup> hybrid functional, using the previously optimized  $S_0$  geometries. Drawings of  
8 molecular structures and molecular orbitals were done using GaussView<sup>36</sup> program.  
9

### 10 11 12 13 14 15 16 17 18 **3. Results and discussion.** 19

20  
21 As confirmed by TD-DFT calculations (see SI) performed on computationally-simpler model  
22 compounds (**1'** and **2'**), in which the hexyl chains have been replaced by methyl groups,  
23 compounds **1** and **2** should possess an (easily discernible) low-energy CT band, while the higher-  
24 energy absorption bands should correspond to  $\pi^* \leftarrow \pi$  transitions (Figure 1a and Table S2). These  
25 assignments are consistent with literature reports on similar push-pull compounds.<sup>37,38</sup> Thus, in  
26  $\text{CH}_2\text{Cl}_2$ , the CT bands are experimentally observed at 423 nm and 395 nm while the  $\pi^* \leftarrow \pi$   
27 transitions are located at 295 nm and 327 nm for **1** and **2**, respectively (Figure 1a, Table S1). The  
28 TD-DFT study is satisfactory for **1** but underestimates the energy of the CT state in **2** (Table S2);  
29 specifically, it fails to reproduce the experimentally-observed overlap of the two transitions seen  
30 with **2** (Figure 1a).<sup>39</sup> In addition, DFT optimizations (PCM formalism) in the ground state (GS)  
31 reveal that the most stable conformation of both compounds in  $\text{CH}_2\text{Cl}_2$  is the fully planar one,  
32 i.e. the one optimizing interaction of donor and acceptor end-groups through the  $\pi$ -electron  
33 system. However, given the low activation barriers for rotation around the molecular axis  
34 defined by the triple bonds,<sup>40,41</sup> various conformers might nevertheless be present in solution.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60







Figure 1. (a) Steady state UV-vis absorption spectra of **1** and **2** dissolved in CH<sub>2</sub>Cl<sub>2</sub>. (b) Steady state IR absorption spectra of **1** and **2** dispersed in a KBr pellet.

The reporter vibrational modes, the  $\nu(\text{C}\equiv\text{C})$  band and the two  $\nu(\text{N}=\text{O})$  bands (antisymmetric and symmetric modes) experimentally observed at ca. 2205, 1590 and 1345  $\text{cm}^{-1}$  for **1** and **2** (Figure 1b and Figure S12e) are fairly accurately modelled in the GS of the DFT studies of **1'** and **2'** (Table S3). Thus, of the two distinct  $\nu(\text{C}\equiv\text{C})$  modes calculated for **2'** and separated by 20  $\text{cm}^{-1}$  (2187  $\text{cm}^{-1}$ , 2265  $\text{cm}^{-1}$ ; 2207  $\text{cm}^{-1}$ , 84  $\text{cm}^{-1}$ ), only one has a significant intensity, *viz.* the mode at lowest energy corresponding to the alkyne located closer to the amino group. The second stretching mode expected for **2** might thus correspond to the shoulder on the high energy side of the  $\nu(\text{C}\equiv\text{C})$  band. For both compounds **1** and **2**, additional bands are also present on the low energy side of the main  $\nu(\text{C}\equiv\text{C})$  band (Figure 1b, Figure S12e). Due to their significantly weaker intensity, these bands (which are not modelled by DFT) might correspond either to overtones of vibrational modes at lower energy, Fermi resonance bands<sup>42</sup> or  $\nu(\text{C}\equiv\text{C})$  mode(s) of (minor) rotamers of **1** or **2** present in solution at 20 °C.<sup>41</sup> They were therefore not considered in

our kinetic modelling. The symmetric and asymmetric  $\nu(\text{N}=\text{O})$  modes of the nitro group of **1** and **2** are also fairly well modelled by DFT in the GS using the model compounds **1'** and **2'**. Based on our calculations for these compounds (Table S3), it appears that the asymmetric  $\nu(\text{N}=\text{O})$  mode is far less useful than the symmetric mode as an IR-marker. Indeed, besides the fact that it might be experimentally difficult to differentiate the antisymmetric mode from the aromatic  $\nu(\text{C}=\text{C})$  modes (Figure S12e), which will also be shifted following photoexcitation into the first excited states, its intensity should be significantly weaker than that of the symmetric stretching mode.<sup>43</sup> In the following, we have therefore decided to focus our attention on the  $\nu(\text{C}\equiv\text{C})$  and symmetric  $\nu(\text{N}=\text{O})$  modes which should be easier to track than the asymmetric  $\nu(\text{N}=\text{O})$  mode in TRIR experiments.






Figure 2. (a,b) TRIR spectra in the spectral range of the C≡C vibrational marker following electronic excitation into the CT state of **1**; time delays specified in the legend. (c) Raw experimental kinetic data recorded for **1** at the three selected wavenumbers specified in the legend.

Starting with **1**, TRIR absorption spectra were recorded in CH<sub>2</sub>Cl<sub>2</sub> following electronic excitation at 420 nm, i.e. at a wavelength corresponding to excitation to the CT state of this dyad. The resulting transient spectra recorded in the spectral range of the C≡C stretching vibrations are presented in Figure 2 and Figure S13. Following excitation, a strong positive band develops at 2128 cm<sup>-1</sup> within the duration of the laser pulse, and a relatively weak negative band forms at 2205 cm<sup>-1</sup> on a comparable timescale (Figure 2a), the latter corresponding to partial depopulation of the GS caused by the electronic excitation. The 2205 cm<sup>-1</sup> band is well-resolved because it does not overlap spectrally with the positive band of much higher intensity. This negative band, formed just after excitation, has a similar amplitude (of the order of  $1.3 \times 10^{-3}$ : Figure S14a) to that of many other organic compounds,<sup>44</sup> including that of a recently investigated triazine-based dyad.<sup>10</sup> As the intensity of an IR band is correlated with the square of the change

1  
2  
3 in the local dipole moment, the much stronger positive band (ca. 14 times more intense than the  
4 negative band: Figure S14a) must be related to a stronger local change of the dipole moment for  
5 this C≡C group after electronic excitation. The computational study afforded a similar outcome;  
6 the calculated intensity for the C≡C vibration in the ground state (1995 km/mol: Table S3) being  
7 far less intense than that of the first singlet excited state (11041 km/mol: Table S3). Note,  
8 though, that the calculated intensity in the singlet excited state is for the relaxed structure, while  
9 experimentally the initial positive signal must correspond to the Franck–Condon (FC) state. The  
10 77 cm<sup>-1</sup> shift to low-energy of the 2205 cm<sup>-1</sup> band to afford the 2128 cm<sup>-1</sup> band provides clear  
11 evidence of a weakening of this C≡C bond in the singlet CT state, suggesting a partial population  
12 (depopulation) of the antibonding  $\pi^*$  (bonding  $\pi$ ) alkyne-based molecular orbitals (MOs) in the  
13 excited CT state. Subsequent to its rise, the 2128 cm<sup>-1</sup> band undergoes a simultaneous partial  
14 decay and blue shift to 2138 cm<sup>-1</sup> (Figure 2a). Finally, the 2138 cm<sup>-1</sup> band decays in an  
15 exponential way (Figure 2c, Figure S15, Table S7) with a time constant of about 5.6 ps. The  
16 initial spectral shift and narrowing of the nascent 2128 cm<sup>-1</sup> band is consistent with vibrational  
17 cooling of the newly formed excited state. As the 2128/2138 cm<sup>-1</sup> band experiences a fast initial  
18 decay (time constant of the order of 1 ps), a new (positive) band forms with a maximum at 2173  
19 cm<sup>-1</sup> (Figure 2a, Figure S15b, Table S7). After a further ca. 1 ps, during which the 2173 cm<sup>-1</sup>  
20 band reaches its maximum, both bands (2138/2173 cm<sup>-1</sup>) decay with similar time constants of  
21 ca. 6 ps (Figure 2b,c, Figure S15b,d, Table S7). These observations suggest that the underlying  
22 excited species are in rapid equilibrium. Because the second band is at a slightly higher energy,  
23 which is consistent with a comparatively greater multiple C-C bond order, we tentatively  
24 attribute this second band to the formation of a non-planar rotamer that possibly corresponds to a  
25 twisted intramolecular charge-transfer (TICT) state; we note that a TICT state will facilitate  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 charge separation in the excited state.<sup>45</sup> This second band is still significantly more intense than  
4 the negative band at 2205 cm<sup>-1</sup>. According to our calculations (Table S3), the latter corresponds  
5 to the  $\nu(\text{C}\equiv\text{C})$  mode expected for the fully relaxed state of the lowest singlet state (calculated to  
6 be at 2106 cm<sup>-1</sup> for **1'**), derived from population of the singlet state after relaxation. In contrast  
7 with this result, a planar conformation was calculated for the first excited state of tolane,<sup>46,47</sup> but  
8 in the present case the twisted conformation adopted for the S<sub>1</sub> state following relaxation may  
9 result from the presence of the push-pull substituents at the para-positions of the terminal phenyl  
10 rings. While exact correspondence of computational and experimental data cannot be expected  
11 given the use of a slightly different model compound (**1'**), both the increase in intensity of the  
12  $\nu(\text{C}\equiv\text{C})$  mode in the relaxed singlet state and its shift to lower energy are qualitatively supported  
13 by the DFT calculations.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29


30 After about 18 ps, when both of these positive bands have almost decayed (Figure S13c), the  
31 spectral features of the remaining transients are consistent with (i) the formation of a  
32 vibrationally-excited (hot) GS (band at ca. 2190 cm<sup>-1</sup>) and with (ii) the formation of a longer-  
33 lived excited state exhibiting a C≡C stretching mode at even lower energy that that  
34 corresponding to its initially populated CT (S<sub>1</sub>) state (ca. 2100 cm<sup>-1</sup>: Figure S13c). The negative  
35 signal at ca. 2205 cm<sup>-1</sup> corresponds to depopulation of the GS, while the positive signal (ca.  
36 2190 cm<sup>-1</sup>) is related to the formation of the aforementioned hot ground state as the recovery of  
37 the ground state takes place.<sup>48,49</sup> These spectral changes are direct evidence of the presence of a  
38 fast and efficient internal conversion (IC) following electronic excitation. All spectral changes  
39 resulting from recovery of the GS are completed after ca. 50 ps, a period typical for vibrational  
40 cooling (VC) of organic compounds in solution,<sup>48,49</sup> although some depopulation of the ground  
41 state is still visible (Figure S13c, Figure S14b), indicating that not all excited molecules have  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 relaxed to the GS. The negative band (amplitude ca.  $1.5 \times 10^{-4}$ ; Figure S14b) persists beyond 106  
4 ps, suggesting partial decay of the initially populated singlet states into longer-lived transients  
5 which probably correspond to triplet states.<sup>10</sup> The ratio of the final amplitude of the negative  
6 band at  $2205 \text{ cm}^{-1}$  ( $1.5 \times 10^{-4}$ ; Figure S14b) to the initial amplitude of this band ( $1.3 \times 10^{-3}$ ;  
7 Figure S14a) is 0.12, so the estimated quantum yield of the IC is 0.88.

8  
9  
10  
11  
12  
13  
14  
15  
16 Apart from these spectral changes corresponding to IC and to VC to the GS, the relatively weak  
17 positive band which develops at ca.  $2100 \text{ cm}^{-1}$  (Figure S13c, Figure S14b) can be assigned to a  
18 longer-lived state. This relatively weak, positive band, initially masked by the strong band  
19 formed by the singlet excited state, becomes apparent after 18 ps, by which time the singlet  
20 excited state has largely decayed. This new band persists up to 100 ps and longer (Figure S13c,  
21 Figure S14b). Based on the extended lifetime of the excited state giving rise to this new C≡C  
22 stretching mode and on our calculations, this signal may correspond to the formation of the first  
23 triplet state(s) of **1**.<sup>10</sup> The C≡C stretch of the first triplet state is calculated to be at  $2048 \text{ cm}^{-1}$  by  
24 DFT i.e. at a lower energy than the C≡C stretch of the relaxed singlet state ( $2106 \text{ cm}^{-1}$ ; see Table  
25 S3).

26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40 Our next goal was to detect the fate of the NO<sub>2</sub> symmetric vibration following the electronic  
41 excitation of **1**. After the photoinduced charge transfer to the nitro group, this vibration should be  
42 shifted to lower energy. The strong NO<sub>2</sub> symmetric vibration is observed at  $1342 \text{ cm}^{-1}$  in the GS,  
43 a spectroscopic region devoid of other intense vibrations (Figure S12e). Calculations for **1**<sup>1</sup>  
44 predict that this IR marker is located at  $1042 \text{ cm}^{-1}$  in the first singlet excited state after  
45 vibrational relaxation (Table S3). The corresponding experimental signal was eventually  
46 detected at  $1048 \text{ cm}^{-1}$  (Figure 3a), following a kinetic evolution (increase and decrease) with  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 similar time constants to those of the  $\nu(\text{C}\equiv\text{C})$  mode attributed to the singlet excited state (Figure  
4 3b, Figure S16 and Table S8). The shorter time constant corresponding to its fast decay ( $\sim 0.85$   
5 ps) is attributed to relaxation of the Franck-Condon state, and the longer time constant ( $\sim 5.9$  ps)  
6 is suggested to correspond to the lifetime of this excited singlet state, once relaxed. Thus, we can  
7  
8 assign the  $1048\text{ cm}^{-1}$  band as the  $\text{NO}_2$  symmetric vibration of the vertical singlet state. The  
9  
10 increase in intensity of this signal appears to be delayed relative to the corresponding increase in  
11  
12 the signal of the singlet state  $\nu(\text{C}\equiv\text{C})$  vibration (Figure S17). We will return to this particular  
13  
14 observation later. Similar to the  $\nu(\text{C}\equiv\text{C})$  vibration, a new band assignable to the longer-lived  
15  
16 transients was also detected in the spectral range of the  $\text{NO}_2$  symmetric vibration (Figure S18)  
17  
18 after all spectral changes related to the decay of the excited singlet state and VC were complete.  
19  
20 Based on its persistence, this very weak band is suggested to correspond to the triplet state, for  
21  
22 which the symmetric N-O mode is expected in this spectral range based on our computations  
23  
24 ( $1049\text{ cm}^{-1}$ , Table S3).  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36






Figure 3. (a) TRIR spectra in the spectral range of the symmetrical  $\text{NO}_2$  vibrational marker, following electronic excitation into the CT state of **1**; time delays specified in the legend. The sharp features on the time-resolved spectra are due to an experimental artefact caused by the detection system employed. (b) Kinetic data recorded at  $1048\text{ cm}^{-1}$ .

The presence of an efficient IC process, depopulating the excited singlet state(s) for **1**, is also suggested by the spectroscopic data gathered in the spectral range of the  $\text{NO}_2$  symmetric vibration in the GS (Figure S19). A negative band at  $1342\text{ cm}^{-1}$ , formed within the duration of the laser pulse, is related to the depopulation of the GS of **1** due to the electronic excitation. The temporal spectral evolution recorded in this spectral range supports the presence of an efficient IC process, which first forms the hot ground state, and which subsequently, via VC, recovers the population of the GS. The time constants of these spectral changes are consistent with those reported for other organic compounds (Figure S19c,d).<sup>48,49</sup> After ca. 80 ps, when all processes related to VC are over, there still exists a weak and negative signal (that persists beyond 3 ns) at  $1342\text{ cm}^{-1}$  evincing some GS depopulation (Figure S19b). A weak positive band with a maximum at  $1347\text{ cm}^{-1}$  (overlapping partially with the negative band) is also detected, but its origin has not yet been clearly established.






Figure 4. (a-b) TRIR spectra in the spectral range of the C≡C vibrational marker following electronic excitation into the CT state of **2**; time delays specified in the legend. (c) Raw experimental kinetic data recorded for **2** at the three selected wavenumbers specified in the legend. (d) Normalized experimental kinetic data recorded for **2** at the three selected wavenumbers specified in the legend

The longer homologue **2** was then investigated by TRIR spectroscopy. In the  $\nu(\text{C}\equiv\text{C})$  spectral range, the situation with this compound was expected to be slightly more complex, because two stretching modes instead of one were expected to be infrared-active in the excited states. Indeed,

1  
2  
3 DFT calculations predict two distinct  $\nu(\text{C}\equiv\text{C})$  modes separated by ca.  $10\text{ cm}^{-1}$  for  $\mathbf{2}'$  in the first  
4  
5 singlet state (following relaxation) (Table S3:  $2134\text{ cm}^{-1}$ ,  $13006\text{ km/mol}$ ;  $2145\text{ cm}^{-1}$ ,  $62780$   
6  
7  $\text{km/mol}$ ). These modes are calculated to be at slightly lower energy than the corresponding  
8  
9 modes for the GS (Table S3,  $2187\text{ cm}^{-1}$  and  $2207\text{ cm}^{-1}$ ). There is a significant increase in  
10  
11 intensity for each  $\nu(\text{C}\equiv\text{C})$  mode in the first singlet state of  $\mathbf{2}'$  (following relaxation). In contrast  
12  
13 to the ground state, the most intense of these two modes in the singlet excited state is that at  
14  
15 lowest energy, a mode which is dominated by the stretching motion of the triple bond closer to  
16  
17 the nitro group. This observation is consistent with this triple bond being the more polarized  
18  
19 when the charge-transfer state is populated. The first singlet state of  $\mathbf{2}'$  is computed to adopt a  
20  
21 fully planar conformation after vibrational relaxation, in contrast to the  $S_1$  state of  $\mathbf{1}'$  which was  
22  
23 shown to adopt a twisted (perpendicular) conformation around the central  $\text{C}\equiv\text{C}$  bond. This  
24  
25 computational result for  $\mathbf{2}'$  resembles that reported for the “parent” 1,4-  
26  
27 bis(phenylethynyl)benzene.<sup>21</sup>  
28  
29  
30  
31  
32  
33


34  
35 Close inspection of the spectral changes in the  $\nu(\text{C}\equiv\text{C})$  region reveals the growth of one rather  
36  
37 than two  $\nu(\text{C}\equiv\text{C})$  bands (maximum at  $2117\text{ cm}^{-1}$ : Figure 4a). Note, though, that two modes  
38  
39 separated by  $10\text{ cm}^{-1}$  are predicted by the calculations, and these might not have been resolved  
40  
41 under our experimental conditions, especially if one of them is far less intense than the other. As  
42  
43 observed for  $\mathbf{1}$ , this new positive band at  $2117\text{ cm}^{-1}$  shifts to  $2129\text{ cm}^{-1}$  and then decays, giving  
44  
45 rise to another band at ca.  $2159\text{ cm}^{-1}$  (Figure 4a) which increases in intensity and then decays at a  
46  
47 similar rate to the band at  $2129\text{ cm}^{-1}$  (Figure 4b, Figure S21, Table S9). Analogous to  $\mathbf{1}$ , the  
48  
49 second band at slightly higher energy ( $2159\text{ cm}^{-1}$ ) can be attributed to the formation of the TICT  
50  
51 state. The observed spectral changes suggest that this transient species also coexists in rapid  
52  
53 equilibrium with the planar CT state (which in this case is the relaxed  $S_1$  state, as indicated by  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 DFT calculations). The decay of these initially-populated singlet states is significantly faster for  
4  
5 **2** than for **1** (ca. 1 ps: Figure 4c-d, Figure S21, Table S9).  
6  
7

8  
9 A larger spectral shift of the (positive) band in the  $S_1$  state from the position of this vibration in  
10 the GS is observed for the  $\nu(\text{C}\equiv\text{C})$  mode(s) of **2** ( $88\text{ cm}^{-1}$ ) compared to that of **1** ( $77\text{ cm}^{-1}$ ).  
11  
12 Similar to observations with **1**, a transient at ca.  $1049\text{ cm}^{-1}$  is observed for **2** which exhibits  
13  
14 kinetic behavior closely related to that of the  $\nu(\text{C}\equiv\text{C})$  band attributed to the  $S_1$  state (Figure S22-  
15  
16 S23b). Although somewhat wider than the band previously observed for **1** in the same spectral  
17  
18 range (Figure S25), this signal can be similarly assigned to the symmetric stretching mode of the  
19  
20 nitro group in the  $S_1$  state of **2**. Analogous to **1**, its growth occurs slightly later than that of the  
21  
22 signals corresponding to the  $\nu(\text{C}\equiv\text{C})$  stretching mode (Figure S23b). In qualitative agreement  
23  
24 with these experimental observations, DFT calculations on the relaxed  $S_1$  state of **2'** indicate that  
25  
26 the  $\nu(\text{C}\equiv\text{C})$  mode(s) should be more intense and should occur at lower energy than in the GS.  
27  
28 Calculations of the  $\nu(\text{N}=\text{O})$  symmetric mode ( $1040\text{ cm}^{-1}$ : Table S3) correspond closely with the  
29  
30 energy observed for this transient of **2** ( $1049\text{ cm}^{-1}$ , Figure S22a).  
31  
32  
33  
34  
35  
36  
37

38  
39 Finally, and also reminiscent of observations with **1**, the overall decay process of the CT state  
40 of **2** affords strong evidence for a very efficient IC process followed by hot GS formation and  
41 then recovery of the GS (Figure S20d). As with **1**, some depopulation of the GS persists (Figure  
42 S20f) beyond 3 ns (the time range of our spectrometer); the ratio of the final amplitude of the  
43 negative band at  $2205\text{ cm}^{-1}$  ( $6 \times 10^{-5}$ : Figure S20f) to the initial amplitude of this band ( $5 \times 10^{-4}$ :  
44  
45 Figure S20e) is also ca. 0.12, so the estimated quantum yield of the IC is 0.88. As previously  
46  
47 observed with **1**, transient spectra recorded in the spectral range of the  $\nu(\text{C}\equiv\text{C})$  vibrations do not  
48  
49 completely decay (at least within the 3 ns available in our spectrometer), but instead give rise to  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


another and longer-lived excited state, evidenced by the appearance of a new C≡C absorption at 2109 cm<sup>-1</sup> (Figure S20d). This weak positive band, already visible after 5 ps, is even more apparent after 9 ps (by which time the singlet excited state has completely decayed), and it persists for more than 50 ps, suggesting a partial decay of the initially populated singlet states into this longer-lived transient. This persistent signal also exhibits a spectral evolution typical of the VC of a nascent state (Figure S20d). By analogy with **1**, this new C≡C stretching mode might therefore be related to the formation of a triplet state(s).<sup>10</sup> Similar to observations with **1**, a very weak and persistent band in the N-O symmetric vibration region was observed over longer time delays (Figure S24) which, based on our DFT calculations (Table S3), might be attributed to the N-O symmetric mode of the triplet state. Once again, and similar to observations with **1**, the presence of an IC process and depopulation of the excited singlet state(s) of **2** is well evidenced in the spectral range corresponding to the symmetric vibration of the nitro group in the GS (Figure S26).



Scheme 2. Proposed processes underlying the spectroscopic IR changes observed for **1** and **2** subsequent to excitation into the singlet CT manifold.

From the above observations, we propose Scheme 2 to rationalize the spectroscopic behavior of **1** and **2** subsequent to excitation at 420 nm. Some of the structural changes undergone in the first excited states of **1** and **2** can now be addressed based on the available IR data.

The position of the C≡C vibrational mode is the same for **1** and **2** in the GS (2205 cm<sup>-1</sup>). In contrast, the relaxed singlet excited-state bands are located at 2138 cm<sup>-1</sup> and 2129 cm<sup>-1</sup> while the second positive bands are located at 2173 cm<sup>-1</sup> and 2159 cm<sup>-1</sup> for **1** and **2**, respectively (Figure S27-S28).<sup>50</sup> Overall, despite small differences, these shifts reflect a fairly comparable decrease in the triple bond order of the alkyne bridge, suggesting that some of the electron density delocalized from the amino donor group during the CT process remains localized in the π\* molecular orbitals (MO) of the C≡C bonds, while some of the electronic density in the π MO of the C≡C bond is delocalized toward the nitro acceptor group. These data are consistent with the valence-bond (VB) representation of the CT state usually proposed for these dyads (Scheme 3),<sup>1,51</sup> although the decrease in the C-C bond order is less pronounced than the simplistic VB representation would suggest. This phenomenon is more qualitatively illustrated by the differential density plots between S<sub>1</sub> and S<sub>0</sub> for **1'** and **2'** (Figure S7), although these plots correspond to the Franck-Condon state, not the relaxed CT state.



Scheme 3. Valence-bond representation of the CT state for **1** ( $n = 1$ ) and **2** ( $n = 2$ ).

1  
2  
3 In contrast to observations made with the  $\nu(\text{C}\equiv\text{C})$  modes, a much larger decrease in energy is  
4  
5 observed for the symmetric  $\nu(\text{N}=\text{O})$  mode of the nitro group for both compounds (ca.  $295\text{ cm}^{-1}$ )  
6  
7 upon proceeding to the excited state, reflecting a more significant decrease in the  $\text{N}=\text{O}$  bond  
8  
9 order than the  $\text{C}\equiv\text{C}$  bond order (Table S4-S5, Figure S9-S10). This experimental shift is  
10  
11 supported by DFT calculations on **1'** and **2'** (Table S3). The electron density changes suggested  
12  
13 by the VB representation of the CT state are also reflected in the differential density plots  
14  
15 between the  $S_1$  and  $S_0$  states that were obtained for **1'** and **2'** (Figure S7). Both plots clearly  
16  
17 reveal that most of the charge is transferred from the nitrogen lone pair of the amino group and  
18  
19 the  $\pi$  system of the adjacent aryl group toward the (most) remote aryl ring and the oxygen atoms  
20  
21 of the nitro group. The similar shifts of the symmetric  $\nu(\text{N}=\text{O})$  modes found for **1** and **2**  
22  
23 following excitation to the CT state suggest that similar electron density resides on the acceptor  
24  
25 group after Franck-Condon relaxation, in spite of the fact that the electronic coupling between  
26  
27 the donor and acceptor groups in these dyads should significantly decrease upon progressing  
28  
29 from **1** to **2**.<sup>52</sup> This conclusion is also consistent with the close similarity between these shifts (ca.  
30  
31  $295\text{ cm}^{-1}$ ) and that found for the symmetric  $\nu(\text{N}=\text{O})$  mode of the reduced nitro group following  
32  
33 one-electron reduction of closely-related nitroaromatics.<sup>27,28</sup> The latter observation confirms that  
34  
35 the  $\text{N}-\text{O}$  stretching modes are perturbed in a very similar way in the CT state (after vibrational  
36  
37 cooling) to that when one additional electron is transferred to the nitroaryl unit, although less  
38  
39 than one electron is actually localized on the “reduced” nitro group in all of these cases.<sup>51</sup> The  
40  
41 electronic couplings between donor and acceptor will determine the amount of charge instantly  
42  
43 transferred between the two sides of the molecule in the Franck-Condon state, whereas it is the  
44  
45 change in local charge (Tables S4-S5, Figure S9) residing on the nitrogen and oxygen atoms of  
46  
47 the nitro group after vibrational relaxation that determines the shift of the  $\nu(\text{N}=\text{O})$  modes (i.e. the  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 change in bond order between the nitrogen and oxygen atoms). Consistent with this  
4 interpretation, the density differential plots between  $S_0$  and  $S_1$  for **1'** and **2'** (Figure S7) reveal  
5 that the amount of charge transferred during the vertical CT process depends on the bridge  
6 length. This amount decreases upon bridge extension (0.826 e for **1'** vs. 0.661 e for **2'**). In  
7 contrast, the N-O computed bond-order changes between the GS and the vibrationally relaxed  
8 CT state remain fairly constant for **1'** and **2'** (Figure S10), in line with the constancy of the IR  
9 shifts observed for the N-O stretching modes of **1** and **2**. These findings confirm that the  
10 symmetric  $\nu(\text{N}=\text{O})$  mode of the nitro group, likewise to the  $\nu(\text{C}\equiv\text{C})$  mode, constitutes a reliable  
11 IR marker for monitoring the bonding changes between specific atoms of the nitro group induced  
12 by the CT process in these dyads.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

27 Compared to the initial transients detected for **1** and **2** (believed to correspond to the lowest  
28 singlet states), much less spectroscopic data is available on the longer-lived transients (believed  
29 to correspond to the triplet states). The slower decay and lower concentration of the species at  
30 the origin of these signals make them more difficult to detect by TRIR spectroscopy. The weak  
31 red-shifted  $\nu(\text{C}\equiv\text{C})$  mode indicates a further weakening of the bond order within their alkyne  
32 linkers, in qualitative agreement with calculations made for the lower triplet states of model  
33 complexes **1'** and **2'**. The latter are consistent with diradical states featuring unpaired spins partly  
34 delocalized on the bridge.<sup>53</sup> Our calculations predict a weak signal located at ca.  $1049\text{ cm}^{-1}$  for **1'**  
35 and ca.  $1079\text{ cm}^{-1}$  for **2'** corresponding to the symmetric  $\nu(\text{N}=\text{O})$  mode of these triplet states. In  
36 accordance with these theoretical predictions, weak signals were experimentally detected in this  
37 spectral range for both **1** and **2**.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 The difference in the relaxation rates for the  $S_1$  states of **1** and **2** is still unclear, as is the origin  
4 of the slight delay observed between the increase in the intensity of the  $\nu(\text{C}\equiv\text{C})$  modes and the  
5  $\nu(\text{N}=\text{O})$  modes in the singlet states. The discrepancy in relaxation rates may stem from the  
6 different conformations adopted by these dyads in their respective  $S_1$  states after vibrational  
7 relaxation. Both adopt a fully planar conformation in their ground and  $T_1$  states, and the slower  
8 kinetic behavior for **1** may result from the need for rotation from the TICT state to the required  
9 planar geometry for internal conversion and intersystem crossing, a motion not needed with **2** for  
10 which the relaxed CT state is also planar. The delay observed in the singlet states between the  
11 increase in intensity of the  $\nu(\text{C}\equiv\text{C})$  and  $\nu(\text{N}=\text{O})$  modes (Figure S23) could manifest vibrational  
12 energy transfer. This could occur via vibrational cascade (percolation down the energy levels) or  
13 “through-bond” transfer (anharmonic coupling with solvent), but in either case the peak  
14 population of the  $\nu(\text{N}=\text{O})$  mode would be delayed with respect to that of the  $\nu(\text{C}\equiv\text{C})$  mode. The  
15 observed delay of 0.5 ps is in fact surprisingly close to the value reported in the literature for  
16 modes separated by a few hundred wavenumbers.<sup>54</sup> These important observations will be the  
17 subject of future investigations.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

#### 39 **4. Conclusions**

40  
41  
42 In summary, two nitro-substituted donor-acceptor oligo(phenyleneethynylene)-bridged dyads  
43 **1** and **2** were studied by time-resolved infrared (TRIR) spectroscopy. The nitro and alkyne  
44 functional groups were used as vibrational reporters to study the photo-induced charge-transfer  
45 process in these archetypal push-pull dyads. Overall, very similar spectral changes were  
46 experimentally found for these two compounds. Following excitation into their charge-transfer  
47 (CT) states, a fast decay ( $\sim 6$  ps for **1** and  $\sim 1$  ps for **2**) due to the presence of the very efficient  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 and fast IC process (major pathway) is observed, while simultaneously the formation of longer-  
4 lived transient species (minor pathway) is observed, probably corresponding to triplet states.  
5  
6 Although the nitro group is often avoided in model compounds for ultrafast transient studies  
7  
8 because it quenches luminescence in polar solvents, we have shown here that, from the  
9  
10 viewpoint of transient vibrational investigations based on absorption, the nitro group constitutes  
11  
12 an outstanding reporter for studying photo-induced charge-transfer processes. From a structural  
13  
14 perspective, the shifts to lower wavenumbers observed for both the  $\nu(\text{C}\equiv\text{C})$  and  $\nu(\text{N-O})_{\text{sym}}$   
15  
16 modes in the present study reveal a weakening of these modes in the singlet excited states, in line  
17  
18 with expectations based on classic VB representations traditionally advanced for CT states.  
19  
20 Consistent with the computed bond orders in the singlet excited states (Figure S9), the observed  
21  
22 shifts of the symmetric N-O stretching mode suggest that a similar degree of charge transfer to  
23  
24 the nitro group in the CT state for both dyads, despite the different length bridges connecting the  
25  
26 donor to the acceptor groups. The spectroscopic changes of the nitro group during this process  
27  
28 resemble those observed upon one-electron reduction, consistent with the changes having  
29  
30 resulted from the addition of (nearly) one electron to the acceptor, and despite the decrease in  
31  
32 charge transferred from donor to acceptor group across a lengthened molecular bridge. We  
33  
34 conclude that the nitro group is a reliable local marker of charge, and anticipate that this  
35  
36 underutilized resource will be more commonly used in TRIR investigations aimed at modeling  
37  
38 dyads featuring strong acceptor groups in the future. The study of further derivatives is currently  
39  
40 underway in our group to define the scope of the utility of the nitro reporter and to further  
41  
42 rationalize the various experimental decay rates for the CT states of **1** and **2**.  
43  
44  
45  
46  
47  
48  
49  
50

## 51 52 ASSOCIATED CONTENT

### 53 54 55 **Supporting Information.**

1  
2  
3 The Supporting Information is available free of charge on the ACS Publications website.  
4  
5 Femtosecond transient absorption mid-IR spectrometer used, synthesis of compounds and  
6  
7 characterization, NMR spectra of the compounds, TD-DFT calculations (PDF), transient  
8  
9 absorption spectra and kinetic studies.  
10  
11  
12

### 13 AUTHOR INFORMATION

#### 16 Notes

17  
18 The authors declare no competing financial interests.  
19  
20  
21

22 Jacek Kubicki, ORCID: 0000-0003-0790-5876  
23  
24

25 Frédéric Paul: ORCID: 0000-0002-8256-0129  
26  
27  
28  
29

### 30 ACKNOWLEDGMENT

31  
32 J.K. acknowledges financial support from the Ministry of Science and Higher Education. The  
33  
34 CNRS (LIA Redochrom - N° 1194) is acknowledged for financial support. M.G.H. thanks the  
35  
36 Australian Research Council for financial support (grant no. DP170100408). T.B.M. thanks the  
37  
38 Julius-Maximilians-Universität Würzburg for support. A.G. thanks the University of the  
39  
40 Philippines for financial support.  
41  
42  
43

### 44 REFERENCES

- 45  
46  
47 (1) Bures, F. Fundamental Aspects of Property Tuning in Push–Pull molecules. *RSC Adv.*  
48  
49 **2014**, *4*, 58826-58851.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 (2) Lyu, S.; Farré, Y.; Ducasse, L.; Pellegrin, Y.; Toupance, T.; Olivier, C.; Odobel, F. Push–  
4 Pull Ruthenium Diacetylide Complexes: New Dyes for p-Type Dye-Sensitized Solar Cells. *RSC*  
5 *Adv.* **2016**, *6*, 19928-19936.  
6  
7  
8  
9  
10 (3) Massin, J.; Lyu, S.; Pavone, M.; Muñoz-García, A. B.; Kauffmann, B.; Toupance, T.;  
11 Chavarot-Kerlidou, M.; Artero, V.; Olivier, C. Design and Synthesis of Novel Organometallic  
12 Dyes for NiO Sensitization and Photo-Electrochemical Applications. *Dalton Trans.* **2016**, *45*,  
13 12539-12547.  
14  
15  
16  
17  
18  
19 (4) Durand, R. J.; Gauthier, S.; Achelle, S.; Groizard, T.; Kahlal, S.; Saillard, J.-Y.; Barsella,  
20 A.; Le Poul, N.; Robin-Le Guen, F. Push–Pull D– $\pi$ -Ru– $\pi$ -A Chromophores: Synthesis and  
21 Electrochemical, Photophysical and Second-Order Nonlinear Optical Properties. *Dalton Trans.*  
22 **2018**, *47*, 3965-3975.  
23  
24  
25  
26  
27  
28 (5) Reichardt, C. Solvatochromic Dyes as Solvent Polarity Indicators. *Chem. Rev.* **1994**, *94*,  
29 2319-2358.  
30  
31  
32  
33 (6) Campagnola, P. J.; Loew, L. M. Second-Harmonic Imaging Microscopy for Visualizing  
34 Biomolecular Arrays in Cells, Tissues and Organisms. *Nature Biotechnology* **2003**, *21*, 1356-  
35 1360.  
36  
37  
38  
39  
40 (7) Brédas, J. L.; Adant, C.; Tackx, P.; Persoons, A. Third-Order Nonlinear Optical Response  
41 in Organic Materials: Theoretical and Experimental Aspects. *Chem. Rev.* **1994**, *94*, 243-278.  
42  
43  
44 (8) Marder, S. R. Organic Nonlinear Optical Materials: where have we been and where are we  
45 going? *Chem. Commun.* **2006**, 131-134.  
46  
47  
48  
49 (9) Adamo, C.; Jacquemin, D. The Calculations of Excited-State Properties with Time-  
50 Dependent Density Functional Theory. *Chem. Soc. Rev.* **2013**, *42*, 845-856.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 (10) Dereka, B.; Svechkarev, D.; Rosspeintner, A.; Tromayer, M.; Liska, R.; Mohs, A. M.;  
4 Vauthey, E. Direct Observation of a Photochemical Alkyne–Allene Reaction and of a Twisted  
5 and Rehybridized Intramolecular Charge-Transfer State in a Donor–Acceptor Dyad. *J. Am.*  
6 *Chem. Soc.* **2017**, *139*, 16885-16893.  
7  
8 (11) Orr-Ewing, A. J. Perspective: How can Ultrafast Laser Spectroscopy Inform the Design of  
9 New Organic Photoredox Catalysts for Chemical and Materials Synthesis? *Struct. Dyn.* **2019**, *6*,  
10 010901-010906.  
11  
12 (12) Biswas, M.; Nguyen, P.; Marder, T. B.; Khundkar, L. R. Unusual Size Dependence of  
13 Nonradiative Charge Recombination Rates in Acetylene-Bridged Compounds. *J. Phys. Chem. A*  
14 **1997**, *101*, 1689-1695.  
15  
16 (13) Nguyen, P.; Lesley, G.; Marder, T. B.; Ledoux, I.; Zyss, J. Second-Order Nonlinear  
17 Optical Properties of Push-Pull Bis(phenylethynyl)benzenes and Unsymmetric Platinum  
18 Bis(phenylacetylide) Complexes. *Chem. Mater.* **1997**, *9*, 406-408.  
19  
20 (14) Lesley, M. J. G.; Woodward, A.; Taylor, N. J.; Marder, T. B.; Cazenobe, I.; Ledoux, I.;  
21 Zyss, J.; Thornton, A.; Bruce, D. W.; Kakkar, A. K. Lewis Acidic Borane Adducts of Pyridines  
22 and Stilbazoles for Nonlinear Optics. *Chem. Mater.* **1998**, *10*, 1355-1365.  
23  
24 (15) Tour, J. M. Molecular Electronics. Synthesis and Testing of Components. *Acc. Chem. Res.*  
25 **2000**, *33*, 791-803.  
26  
27 (16) Collings, J. C.; Parsons, A. C.; Porres, L.; Beeby, A.; Batsanov, A. S.; Howard, J. A. K.;  
28 Lydon, D. P.; Low, P. J.; Fairlamb, I. J. S.; Marder, T. B. Optical properties of Donor–Acceptor  
29 Phenylene-Ethynylene Systems Containing the 6-Methylpyran-2-one Group as an Acceptor.  
30 *Chem. Commun.* **2005**, 2666-2668.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 (17) Nguyen, P.; Lesley, G.; Dai, C.; Taylor, N. J.; Marder, T. B.; Chu, V.; Viney, C.; Ledoux,  
4 I.; Zyss, J. Well-Defined Conjugated Rigid-Rods as Multifunctional Materials: Linear and  
5 Nonlinear Optical Properties and Liquid Crystalline Behavior. In *Applications of Organometallic*  
6 *Chemistry in the Preparation and Processing of Advanced Materials*; Harrod, J. F., Laine, R. M.,  
7 Eds., 1995; Vol. 297; pp 333-347.  
8  
9  
10  
11  
12  
13  
14 (18) McGrier, P. L.; Solntsev, K. M.; Zuccherro, A. J.; Miranda, O. R.; Rotello, V. M.; Tolbert,  
15 L. M.; Bunz, U. H. F. Hydroxydialkylamino Cruciforms: Amphoteric Materials with Unique  
16 Photophysical Properties. *Chem. Eur. J.* **2011**, *17*, 3112-3119.  
17  
18  
19  
20  
21 (19) Bunz, U. H. F.; K. Seehafer; Bender, M.; Porz, M. Poly(aryleneethynylene)s (PAE) as  
22 Paradigmatic Sensor Cores. *Chem. Soc. Rev.* **2015**, *44*, 4322-4336.  
23  
24  
25  
26 (20) Nojo, W.; Reingold, I. D.; Bard, J. P.; Chase, D. T.; Deng, C.-L.; Haley, M. M. Donor-  
27 /Acceptor-Substituted Tetrakis(arylethynyl) Benzenes: The Influence of Donor Group on  
28 Optoelectronic Properties. *ChemPlusChem* **2019**, *84*, 1-6.  
29  
30  
31  
32  
33 (21) Beeby, A.; Findlay, K. S.; Low, P. J.; Marder, T. B.; Matousek, P.; Parker, A. W.; Rutter,  
34 S. R.; Towrie, M. Studies of the S1 State in a Prototypical Molecular Wire using Picosecond  
35 Time-Resolved Spectroscopies. *Chem. Commun.* **2003**, 2406–2407.  
36  
37  
38  
39  
40 (22) Delor, M.; Keane, T.; Scattergood, P. A.; Sazanovich, I. V.; Greetham, G. M.; Towrie, M.;  
41 Meijer, A. J. H. M.; Weinstein, J. A. On the Mechanism of Vibrational Control of Light-Induced  
42 Charge Transfer in Donor–Bridge–Acceptor assemblies. *Nature Chem.* **2015**, *7*, 689-695.  
43  
44  
45  
46 (23) Wu, Y.; Yu, P.; Chen, Y.; Zhao, J.; Liu, H.; Li, Y.; Wang, J. Intensified CC Stretching  
47 Vibrator and Its Potential Role in Monitoring Ultrafast Energy Transfer in 2D Carbon Material  
48 by Nonlinear Vibrational Spectroscopy. *J. Phys. Chem. Lett.* **2019**, *10*, 1402-1410.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 (24) Bellamy, L. J. *The infrared spectra of complex molecules*; Methuen & Co. Ltd: London,  
4  
5 1955.  
6  
7  
8 (25) Tian, D.; Jin, B. FT-IR Spectroelectrochemical Study of the Reduction of 1,4-  
9  
10 Dinitrobenzene on Au Electrode: Hydrogen Bonding and Protonation in Proton Donor mixed  
11  
12 Media. *Electrochim. Acta* **2011**, *56*, 9144-9151.  
13  
14 (26) Ezumi, K.; Miyazaki, H.; Kubota, T. Stretching Vibration of Nitro and N-Oxide Groups of  
15  
16 the Anion Radicals of 4-Nitropyridine N-Oxide and Related Nitro Compounds. *J. Phys. Chem.*  
17  
18 **1970**, *74*, 2397-2402.  
19  
20 (27) Ma, R.; Yuan, D.; Chen, M.; Zhou, M. Infrared Spectrum of Nitrobenzene Anion in Solid  
21  
22 Argon. *J. Phys. Chem. A* **2009**, *113*, 1250-1254.  
23  
24 (28) Steill, J. D.; Oomens, J. Spectroscopically resolved Competition between Dissociation and  
25  
26 Detachment from Nitrobenzene Radical Anion. *Int. J. Mass Spectro.* **2011**, *308*, 239-252.  
27  
28 (29) Some confusion exist about the redox-induced shifts of the  $\nu(\text{N=O})_{\text{sym}}$  mode reported in  
29  
30 the extant literature for nitroaromatics<sup>26</sup> because the  $\nu(\text{N=O})$  modes and the  $\nu(\text{C-N})$  mode move  
31  
32 in opposite directions on the wavelength scale and change intensities, the  $\nu(\text{N=O})_{\text{sym}}$  mode  
33  
34 crossing the  $\nu(\text{C-N})$  mode.  
35  
36 (30) Mitzel, F.; Boudon, C.; Gisselbrecht, J.-P.; P. Seiler; Gross, M.; Diederich, F. Donor-  
37  
38 Substituted Perethynylated Dehydroannulenes and Radiaannulenes: Acetylenic Carbon Sheets  
39  
40 featuring Intense Intramolecular Charge Transfer. *Helv. Chim. Acta* **2004**, *87*, 1130-1157.  
41  
42 (31) Walters, K. A.; Kim, Y.-J.; Hupp, J. T. Experimental Studies of Light-Induced Charge  
43  
44 Transfer and Charge Redistribution in  $(\text{X}_2\text{-bipyridine})\text{ReI}(\text{CO})_3\text{Cl}$  Complexes. *Inorg. Chem.*  
45  
46 **2002**, *41*, 2909-2919.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 (32) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J.  
4 R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A. et al. Gaussian 09, Revision D.01;  
5 Gaussian, Inc.: Pittsburgh, PA, 2015.  
6  
7  
8  
9  
10 (33) Adamo, C.; Barone, V. Exchange Functionals with Improved Long-Range Behavior and  
11 Adiabatic Connection Methods without Adjustable Parameters: the mPW and mPW1PW  
12 Models. *J. Chem. Phys.* **1998**, *108*, 664-675.  
13  
14  
15  
16 (34) Tomasi, J.; Mennucci, B.; Cammi, R. Quantum Mechanical Continuum Solvation Models  
17 *Chem. Rev.* **2005**, *105*, 2999-3093.  
18  
19  
20  
21 (35) Yanai, T.; Tew, D. P.; Handy, N. C. A New Hybrid Exchange-Correlation Functional using  
22 the Coulomb-Attenuating Method (CAM-B3LYP). *Chem. Phys. Lett.* **2004**, *393*, 51-57.  
23  
24  
25  
26 (36) Dennington, R.; Keith, T.; Millam, J. GaussView, Version5; Semichem Inc.: Shawnee  
27 Mission, KS, 2009.  
28  
29  
30  
31 (37) Stiegman, A. E.; Graham, E. M.; Perry, K. J.; Kundkar, L. R.; Cheng, L.-T.; Perry, J. W.  
32 The Electronic Structure and Second-Order Nonlinear Optical Properties of Donor-Acceptor  
33 Acetylenes: A Detailed Investigation of Structure-Property Relationships. *J. Am. Chem. Soc.*  
34 **1991**, *113*, 7658-7666.  
35  
36  
37  
38 (38) Ramakrishna, G.; Bhaskar, A.; T. Goodson, I. Ultrafast Excited State Relaxation Dynamics  
39 of Branched Donor- $\delta$ -Acceptor Chromophore: Evidence of a Charge-Delocalized State. *J. Phys.*  
40 *Chem. B* **2006**, *110*, 20872-20878.  
41  
42  
43  
44  
45  
46 (39) The use of the CAM-B3LYP functional, in principle better suited to model long range  
47 charge transfers,<sup>9</sup> proved unsatisfactory here (Table S2).  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 (40) Beeby, A.; Finlay, K.; Low, P. J.; Marder, T. B. A Re-Evaluation of the Photophysical  
4 Properties of 1,4-Bis(phenylethynyl)benzene: a Model for Poly(phenyleneethynylene). *J. Am.*  
5  
6 *Chem. Soc.* **2002**, *124*, 8280-8284.  
7  
8  
9  
10 (41) A barrier of rotation of 1.7 kcal mol<sup>-1</sup> can be derived by DFT for the rotation of the  
11 aminophenyl part of the molecule in **2'** (Figure S11).  
12  
13  
14 (42) Paul, F.; Mevellec, J.-Y.; Lapinte, C. Vibrational Spectroscopic Investigations of Fe(II)  
15 and Fe(III) Organoiron sigma-Alkynyl complexes. Solid State Raman and Solution Infrared  
16 Intensity Measurements. *J. Chem. Soc., Dalton Trans.* **2002**, 1783-1790.  
17  
18  
19 (43) Calculations on **1'** and **2'** in the S<sub>1</sub> state predict this mode to appear in the range 1390-1430  
20 cm<sup>-1</sup> (Table S3).  
21  
22  
23 (44) Kubicki, J.; Zhang, Y. L.; Wang, J.; Luk, H. L.; Peng, H. L.; Vyas, S.; Platz, M. S. Direct  
24 Observation of Acyl Azide Excited States and Their Decay Processes by Ultrafast Time  
25 Resolved Infrared Spectroscopy. *J. Am. Chem. Soc.* **2009**, *131*, 4212-4213.  
26  
27  
28 (45) Rettig, W. Charge Separation in Excited States of Decoupled Systems—TICT Compounds  
29 and Implications Regarding the Development of New Laser Dyes and the Primary Process of  
30 Vision and Photosynthesis. *Angew. Chem. Int. Ed.* **1986**, *25*, 971-988.  
31  
32  
33 (46) Menning, S.; Krämer, M.; Duckworth, A.; Rominger, F.; Beeby, A.; Dreuw, A.; Bunz, U.  
34 H. F. Bridged Tolanes: A Twisted Tale. *J. Org. Chem.* **2014**, *79*, 6571–6578.  
35  
36  
37 (47) Krämer, M.; Bunz, U. H. F.; Dreuw, A. Comprehensive Look at the Photochemistry of  
38 Tolane. *J. Phys. Chem. A* **2017**, *121*, 946-953.  
39  
40  
41 (48) Hamm, P.; Ohline, S. M.; Zinth, W. Vibrational Cooling after Ultrafast Photoisomerization  
42 of Azobenzene Measured by Femtosecond Infrared Spectroscopy. *J. Chem. Phys.* **1997**, *106*,  
43 519-529.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 (49) Kubicki, J.; Zhang, Y. L.; Xue, J. D.; Luk, H. L.; Platz, M. Ultrafast Time-Resolved  
4 Studies of the Photochemistry of Acyl and Sulfonyl Azides. *Phys. Chem. Chem. Phys.* **2012**, *14*,  
5 10377-10390.  
6  
7

8  
9  
10 (50) The shift of the  $\nu(\text{C}\equiv\text{C})$  band upon population of the singlet state in **2** should be treated  
11 cautiously, because the DFT calculations on **2'** suggest an inversion of intensity between the two  
12  $\nu(\text{C}\equiv\text{C})$  modes in the first singlet state (corresponding to the CT state). Based on these  
13 calculations, both  $\nu(\text{C}\equiv\text{C})$  modes in **2'** should experience different, but smaller (40 and 77  $\text{cm}^{-1}$ )  
14 shifts than the unique  $\nu(\text{C}\equiv\text{C})$  mode in **1'** (81  $\text{cm}^{-1}$ ), contrary to the experimental observations on  
15 **1** and **2** (77  $\text{cm}^{-1}$  vs. 88  $\text{cm}^{-1}$ ). This discrepancy might result from the change in intensity  
16 between the two  $\nu(\text{C}\equiv\text{C})$  modes in the singlet state, since a shift of 103  $\text{cm}^{-1}$  is derived from  
17 computations when only the most intense  $\nu(\text{C}\equiv\text{C})$  mode is considered in the ground and  $S_1$  state.  
18  
19

20 (51) Dehu, C.; Meyers, F.; Brédas, J. L. Donor-acceptor diphenylacetylenes: geometric  
21 structure, electronic structure, and second-order nonlinear optical properties. *J. Am. Chem. Soc.*  
22 **1993**, *115*, 6198-6206.  
23  
24

25 (52) Paddon-Row, M. N. Orbital Interactions and Long-Range Electron Transfer. *Adv. Phys.*  
26 *Org. Chem.* **2003**, *38*, 1-85.  
27  
28

29 (53) Zweig, A.; Hoffmann, A. K. Chemistry of dianions. Formation and Reactivity of Anionic  
30 Species Derived from Tetraphenylbutadiene. *J. Am. Chem. Soc.* **1962**, *84*, 3278-3284.  
31  
32

33 (54) Wang, Z. H.; Pakoulev, A.; Dlott, D. D. Watching Vibrational Energy Transfer in Liquids  
34 with Atomic Spatial Resolution. *Science* **2002**, *296*, 2201-2203.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## TOC GRAPHICS

