



HAL
open science

A new primer for metabarcoding of spider gut contents

Denis Lafage, Vasco Elbrecht, Jordan P. Cuff, Dirk Steinke, Peter A.
Hambäck, Ann Erlandsson

► To cite this version:

Denis Lafage, Vasco Elbrecht, Jordan P. Cuff, Dirk Steinke, Peter A. Hambäck, et al.. A new primer for metabarcoding of spider gut contents. *Environmental DNA*, 2020, 2 (2), pp.234-243. 10.1002/edn3.62 . hal-03097420

HAL Id: hal-03097420

<https://univ-rennes.hal.science/hal-03097420>

Submitted on 22 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

A new primer for metabarcoding of spider gut contents

Denis Lafage^{1,2}  | Vasco Elbrecht^{3,4}  | Jordan P. Cuff⁵  | Dirk Steinke³ | Peter A. Hambäck⁶  | Ann Erlandsson¹ 

¹Department of Environmental and Life Sciences/Biology, Karlstad University, Karlstad, Sweden

²UMR ECOBIO, Université de Rennes 1, Rennes, France

³Centre for Biodiversity Genomics, University of Guelph, Guelph, ON, Canada

⁴Centre for Molecular Biodiversity Research, Zoological Research Museum Alexander Koenig, Bonn, Germany

⁵School of Biosciences, Cardiff University, Cardiff, UK

⁶Department of Ecology, Environment and Plant Sciences, Stockholm University, Stockholm, Sweden

Correspondence

Denis Lafage, Department of Environmental and Life Sciences/Biology, Karlstad University, Universitetsgatan 2, Karlstad 65188, Sweden.
Email: lafage.denis@gmail.com

Funding information

RIVEM Group (Karlstad University); University of Guelph's Canada First Research Excellence Fund project Food from Thought

Abstract

As a key predator group, spiders have received a lot of attention by food web ecologists. The difficulty involved in studying their diet has led to the use of new technologies such as metabarcoding of gut contents. The amplification of a broad range of spider prey without amplifying spiders themselves is challenging. Until now, an efficient universal primer for this purpose was not available. We developed a novel forward primer (NoSpi2) targeting the COI gene. The primer was designed not to amplify spiders of *Pardosa* genus while amplifying most other invertebrates. NoSpi2 was tested together with the reverse primer BR2 in silico, in vitro on single specimens of prey and spiders, on mock and malaise trap communities, and in an ecological application. In silico evaluation predicted high primer bias for *Pardosa* species and more generally for spiders of the oval calamistrum clade (Lycosidae and closely related species) and low bias for other invertebrates. These results were confirmed by in vitro tests. Additionally, some spider families were not amplified contrary to our expectations. We demonstrated a high efficiency for the primer pair NoSpi2/BR2 which recovered 94% of taxa in the mock community and 85% of the taxa detected by the best invertebrate primer pair known for the malaise trap community. The field experiment showed that Lycosidae (*Hygrolycosa*, *Pardosa*, *Piratula*, *Trochosa*) DNA is not amplified by NoSpi2/BR2. It demonstrated a broad range of detectable prey species (12 orders, 67 families, 117 species). The ability of NoSpi2/BR2 primer to reliably amplify prey species, without amplifying any predator DNA, makes it an ideal choice for gut content analysis for lycosid species and related species, even enabling the homogenization of entire specimens without dissection. Given that the detected prey species included other spiders and carabid beetles, this primer could be also used to study intraguild predation.

KEYWORDS

diet, gut content metabarcoding, lycosidae, molecular diet analysis, prey detection

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Environmental DNA* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

The ecological interactions of spiders, the most abundant terrestrial arthropod predator group found on every life-supporting landmass, are pivotal in many ecological networks (Riechert & Lockley, 1984; Turnbull, 1973). As such, they have received a lot of attention by food web ecologists of diverse fields including pest control (Holland et al., 2016), pollutant transfers (Kraus et al., 2016; Walters, Otter, Kraus, & Mills, 2018), and cross-ecosystem fluxes (Lafage et al., 2019). Spiders feed on predigested fluids of their prey, restricting field studies of spider diet to direct observation, stable isotope analysis, and molecular analysis (Birkhofer et al., 2017; Pompanon et al., 2012; Symondson, 2002). The inherent bias and the laborious nature of direct observation have led to the increased use of molecular techniques to investigate spider diet, of which DNA metabarcoding is currently among the most accurate and efficient for analysis of polyphagous generalist diets (Piñol, Senar, & Symondson, 2018).

General metabarcoding-based analyses of diet require PCR primers that amplify a broad range of potential prey species, but ideally without amplifying predator DNA. Given the degraded nature of gut content or fecal DNA in comparison with the relatively intact DNA of the predator, the latter can outcompete prey DNA in both PCR and sequencing (Vestheim & Jarman, 2008). The selection of primers is thus the most critical step for dietary metabarcoding studies (Piñol et al., 2018). Many studies use primer combinations that target specific species (Boreau de Roincé, Lavigne, Mandrin, Rollard, & Symondson, 2013; Greenstone & Shufan, 2003; Kuusk, Cassel-Lundhagen, Kvarnheden, & Ekbon, 2008), while others focus on a limited number of closely related groups (Chapman, Schmidt, Welch, & Harwood, 2013; Hambäck, Weingartner, Dalén, Wirta, & Roslin, 2016; Hosseini, Keller, Schmidt, & Li, 2011). While this approach does eliminate amplification of predator DNA, it also requires preexisting knowledge of the diet and reduces the possibility of finding unexpected prey species.

Many studies employ blocking probes, which are predator-specific oligonucleotides that bind to the DNA without extension thereby inhibiting predator amplification (Piñol, San Andrés, Clare, Mir, & Symondson, 2014; Vestheim & Jarman, 2008). This can, however, also block closely related prey species and increase amplification bias (Piñol et al., 2014). Other ways to reduce predator DNA include extraction from feces (Sint, Thurner, Kaufmann, & Traugott, 2015), spider webs (Xu, Yen, Bowman, & Turner, 2015), or regurgitates. However, these methods either reduce the concentration and quality of DNA relative to gut content extraction

(Agustí et al., 2003) or are not possible due to the fluid-feeding behavior of spiders (Kamenova et al., 2018; Waldner, Sint, Juen, & Traugott, 2013). The amount of predator DNA in spider gut content extractions can also be limited by extraction of just the abdomen, which has a higher proportion of prey DNA, but nevertheless a majority of the extracted DNA will be from the predator (Krehenwinkel, Rödder, & Tautz, 2015). Similarly, size selection with magnetic beads can be used to limit the amount of intact longer predator DNA present in an extract, leaving only shorter degraded prey DNA (Krehenwinkel et al., 2015), but this removal is based wholly on size and may limit the prevalence of more recently ingested prey and does not remove degraded predator DNA. A better approach would be to use primers which do not amplify the DNA of the predator while still amplifying prey DNA. Lineage-specific primers have previously been designed for multiplex amplification of different prey lineages (Krehenwinkel et al., 2019), but no single primer pair has yet been developed for universal amplification of spider prey while avoiding predator amplification.

In the present study, a novel forward PCR primer was designed to amplify spider prey DNA from spider gut content extracts without amplifying spider predator DNA. We specifically targeted *Pardosa* spiders (but we also investigated phylogenetically closely related species) which belongs to one of the most diverse and common spider groups (Piacentini et Ramírez, 2019). This primer together with a previously designed reverse primer is relevant for future metabarcoding-based analyses of spider diet.

2 | METHODS

2.1 | Primer development and in silico evaluation

Mitochondrial cytochrome c oxidase subunit 1 (COI) sequences for 15 freshwater invertebrate groups (Vamos, Elbrecht, & Leese, 2017) and one spider genus (*Pardosa*) were downloaded and clustered using the "PrimerMiner" package v0.18 (Elbrecht & Leese, 2017a). All following analyses were run in R 3.6 (R Core Team, 2019). Sequences were aligned in Geneious 8.1.7 (Kearse et al., 2012) using MAFFT v7.017 (Kato, 2002). PrimerMiner's "selectivetrimer" function was used to trim 26 bp of the HCO and 25 bp of the LCO binding sites, and the alignment for each group was visualized with PrimerMiner to visually identify suitable primer binding sites. Sites conserved among target spider prey taxa but differing in *Pardosa* sequences were selected, and a primer was designed based on the most optimal combination of sites (Table 1 and Figure 1). The resulting forward primer (NoSpi2) (Table 1) contains a high base degeneracy and binds

TABLE 1 Primer characteristics as determined with Primer3 2.3.7 implemented in Geneious v2019.0.4 with default settings

Primer	Sequence (5'-3')	Source	Average Tm (°C)	Average GC (%)	Length (bp)
NoSpi2	TTYCCHCGWATAAAAYAYATAAG	This study	51.9	29.7	23
BR2	TCDGGRTGNCCRAARAAYCA	Elbrecht and Leese (2017b)	59.0	49.2	20




FIGURE 1 Bar plot showing NoSpi2 binding site variability of *Pardosa* sp., spiders of the oval calamistrum clade, other spider families, and other insects, generated with PrimerMiner. Amplification of spiders of the genus *Pardosa* and the oval calamistrum clade is expected to be unlikely, due to mismatches at the 3' end of the NoSpi2 primer. Insect data from Vamos et al. (2017), base composition, green: thymine, blue: cytosine, red: adenine, yellow: guanine, blocked positions highlighted in orange

at position 233,255 of the COI Folmer region (Folmer, Black, Hoeh, Lutz, & Vrijenhoek, 1994), taking advantage of sequence polymorphisms to enable amplification of target prey species and avoiding amplification of spiders of the genus *Pardosa*. Amplicon length in combination with the reverse primer BR2 (Table 1) (Elbrecht & Leese, 2017b) was 403 bp.

PrimerMiner was also used to evaluate the primer pair (NoSpi2 and BR2) against alignments of 30 aquatic and terrestrial invertebrate groups and 103 spider families, using default tables for mismatch scoring. As a reference, penalty scores were also computed for the standard barcoding primers LCO1490 and HCO2198 (Folmer et al., 1994).

2.2 | In vitro evaluation

In vitro evaluation of the primer pair was performed using two approaches. First, we tested the primer pair on single individuals of potential prey and spiders and on mix of prey and spider DNA. The second approach consisted in testing the primer pair by metabarcoding on a known MOC community (to assess the percentage recovery of species) and on a large and very diverse malaise trap sample (to compare its performance to the best know primer pair for arthropods).

The primer pair was tested in vitro on 6 freshwater and 13 terrestrial invertebrate families (total of 28 species), and 20 spider families (32 species) separately (Appendix S1 and S2). Freshwater invertebrates were selected from samples collected in 2017 from four Swedish rivers using drift nets. Spiders were selected from samples collected in 2017 in the riparian zone of the same four Swedish rivers using a vacuum sampler. Terrestrial invertebrates were selected from

samples collected in 2016 in the United Kingdom from arable crops via vacuum sampler, and from decaying beech wood via hand sorting. For herbivorous invertebrates, whole bodies were used, and for predators only legs to restrict possible amplification of gut contents. To investigate possible competitive effects (interference of predator DNA with amplification of prey DNA), sample mixes with equal DNA concentrations (39 ng/ μ l) of starved Lycosidae (*Pardosa amentata*) and Collembola (Poduridae) or Trichoptera (Limnephiliidae) or Plecoptera (Leutridae) were also tested. In addition, DNA extracts from a starved Lycosidae (*Pardosa amentata*) at a concentration of 90.8 ng/ μ l supplemented with Trichoptera (Limnephiliidae) DNA in a decreasing concentration (lowest tested addition 0.007 ng/ μ l) were analyzed.

DNA extraction of tissue samples was performed using the QIAamp DNA Micro Kit (Qiagen), following manufacturer instructions. For amplification, the combination of NoSpi2 (Table 1) and BR2 was used (Elbrecht & Leese, 2017b). PCRs were carried out in 25 μ l reaction volumes containing 2 μ l of DNA extract, 12.5 μ l of PCR Multiplex kit (Qiagen), and 2.5 μ l of each primer (at 10 μ M). Thermocycler conditions were as follows: initial denaturation at 95°C for 15 min; 30 cycles of 30 s at 94°C, 90 s at 48°C, and 90 s at 72°C; and a final extension for 10 min at 72°C. Positive amplifications were confirmed by visual inspection of PCR products in 2% agarose gels. For positive amplifications, PCR products were purified using ExoSAP-IT™ PCR Product Cleanup Reagent (Thermo Fisher Scientific). In case of negative amplification, a second PCR with LCO1490/HCO2198 primers was run using the same conditions (different annealing temperature: 40°C) to ensure that the sample contained amplifiable DNA. Purified PCR products (positive samples with NoSpi2/BR2 and positive samples with LCO1490/HCO2198) were then Sanger-sequenced

and the resulting sequences processed using the sangeranalyse R package (v. 0.1) <https://github.com/roblanf/sangeranalyseR>. Low-quality ends of sequences were automatically trimmed based on their quality (default cutoff of 0.0001). Subsequently, forward and reverse sequences were merged into a consensus sequence. If the quality of one of the sequences was low, only the other sequence was used. Consensus sequences were queried against NCBI GenBank using the blastn algorithm (Camacho et al., 2009) to retrieve species identity.

In order to assess the arthropod detection efficiency of the primer pair, it was also used to metabarcode an insect mock sample (Braukmann et al., 2018) and a malaise trap sample from Ontario, Canada, both previously tested with 21 primer sets (Elbrecht et al., 2019). A two-step PCR was used to amplify and tag the DNA fragments. The first PCR was performed in 25 μ l reaction volumes containing 1 μ l of DNA extract (12.5 ng), 12.5 μ l of PCR Multiplex plus kit (Qiagen), 0.5 μ M of each primer (NoSpi2 + BR2), and DNA-free water. The conditions for these PCRs were as follows: initial denaturation at 95°C for 5 min; 30 cycles of 30 s at 95°C, 30 s at 48°C, and 50 s at 72°C; and a final extension of 5 min at 72°C. The second PCR was performed using fusion primers in 25 μ l reaction volumes: 1 μ l product from PCR1, 12.5 μ l of PCR Multiplex kit (Qiagen), 0.2 μ M of each primer (P7_NoSpi2 CAAGCAGAAGACGGCATAACGAGATGTGACTGGAGT TCAGACGTGTGCTCTTCCGATCTCTGATTYCCCHCGWAT AAAYAYATAAG combined with mock sample: P5_BR2_F, malaise sample: P5_BR2_X, Elbrecht & Steinke, 2018). The second PCR conditions were as follows: initial denaturation at 95°C for 5 min; 20 cycles of 30 s at 95°C, 30 s at 48°C, and 2 min at 72°C; and a final extension of 5 min at 72°C. PCR products were normalized and pooled using Sequal prep (Thermo Fisher Scientific). Primer dimer was subsequently removed using SPRIselect cleanup according to the manufacturer protocol (0.76 \times SPRIselect used) (Beckman Coulter). The final library contained 96 additional samples of mostly malaise samples and macrozoobenthos kick samples. Sequencing was carried out by the AAC Genomics Facility at the University of Guelph, Canada, on an Illumina MiSeq using the 600 v3 cycle kit with 5% PhiX spike in. As fusion primer inline tags were used, indexing was skipped and an additional 16 bp was added to the first read, leading to paired-end sequencing of 316 + 300 bp length. Bioinformatic processing was done following Elbrecht et al. (2019) using the JAMP pipeline.

2.3 | Field experiment

To ascertain its performance in an ecological application, the primer pair was used to amplify spider gut DNA from a field experiment conducted in 2018. Lycosid spiders (*Hygrolycosa*, *Pardosa*, *Piratula*, and *Trochosa*) were sampled at three riparian sites along the Klarälven River (Sweden). Samples were collected weekly between May 15 and June 19. In each location, spiders were collected from recently

flooded areas ($n = 20$) and nonflooded areas ($n = 10$). Individuals were collected using clean forceps and stored individually in 99.6% ethanol before freezing at -20°C .

Each adult was identified to species level before sterile removal of its abdomen for DNA extraction using the QIAamp DNA Micro Kit, as described above. Extra care was taken to avoid contamination: each spider was manipulated in a sterile petri dish and dissected with sterilized forceps and blades. Each step of the follow-up procedures such as DNA extraction, PCR mixes, and PCR product detection and isolation was also performed at separate locations to avoid contamination. To allow multiplex sequencing of the samples, we followed a two steps protocol: A first PCR was run to allow detection and amplification of prey DNA and to prepare sequencing by adding Illumina adaptors. A second PCR was run to allow dual tagging of the samples. The first PCR was performed with the primer pair NoSpi2/BR2 extended with universal Turesq Illumina adaptors (33 and 21 bp, respectively) using the PCR conditions described (see in vitro evaluation part). Positive amplifications were confirmed by visual inspection of PCR products in 2% agarose gels. Samples exhibiting bands of the expected size were purified using Agencourt AMPure XP beads following manufacturer specifications (Beckman Coulter). DNA concentration of the cleaned PCR products was determined using a Qubit fluorometer (Thermo Fisher). A second PCR was performed to barcode the samples with dual indexes (13 forward \times 10 reverse) extended with Illumina adapters (Hugerth et al., 2014). PCRs were carried out in 25 μ l reaction volumes containing 2 μ l of the cleaned PCR product, 12.5 μ l of KAPA Taq ReadyMix (2 \times) (Sigma-Aldrich), 1 μ l of each primer (at 10 μ M), and 8.5 μ l H₂O. Thermocycler conditions were as follows: an initial denaturation at 98°C for 2 min; 10 cycles of 20 s at 98°C, 30 s at 62°C, and 30 s at 72°C; and a final extension for 2 min at 72°C. PCR product concentrations were measured using a Qubit fluorometer before equimolar pooling (20.1 ng/ μ l). The pooled product was then purified using Agencourt AMPure XP beads following manufacturer specifications (Beckman Coulter). The DNA library was sequenced in a flow cell on an Illumina MiSeq v3, PE 2 \times 300, at the Science for Life Laboratory, Sweden (www.scilifelab.se).

Sequences were processed using the "dada2" package (Callahan et al., 2016) in R. The pipeline includes demultiplexing, filtering, trimming, dereplication, correction of errors, merging of forward and reverse sequences, and clustering into amplicon sequence variant (ASV). All ASVs produced were screened against BOLD (Ratnasingham & Hebert, 2007). Sequences with no match were subsequently queried against NCBI GenBank. Sequences were attributed at species level with a similarity match $\geq 98\%$, to the genus level with a similarity match $\geq 95\%$, to the family level with a similarity match $\geq 90\%$, and to the order level with a similarity match $\geq 85\%$ (Elbrecht, Vamos, Meissner, Aroviita, & Leese, 2017). If several species obtained identical similarity, the species with more occurrences in Sweden, based on GBIF data (<http://www.gbif.org/>), was selected. When several ASVs attributed to the same sample referred to the

same species, the number of differences between sequences was computed. If the number of differences was higher than 50, the two ASVs were considered two different individuals. Finally, only species which are likely preyed upon by spiders were kept (i.e., bacteria and fungi sequences were discarded).

3 | RESULTS

3.1 | Primer development and in silico evaluation

We designed a primer NoSpi2, which in combination with the primer BR2 (Elbrecht & Leese, 2017b) targets the Folmer region of the COI gene to generate a sequence of 403 bp. In silico evaluation of the primer pair showed high penalty scores for *Pardosa* species and more generally for all spider families of the oval calamistrum clade (Figure 2) with the strongest values obtained for Lycosidae.

In silico evaluation of the potential prey (8 aquatic and 22 terrestrial) showed mostly lower penalty values for NoSpi2 compared with LCO1490 and for BR2 compared with HCO2198 (Figure 3). Among aquatic prey, NoSpi2 and BR2 always had lower penalty scores. Among terrestrial prey, LCO showed lower penalty scores for Embioptera, Strepsiptera, and Archeognatha. BR2 showed lower penalty scores for all other terrestrial prey tested.

3.2 | In vitro evaluation

The primer pair NoSpi2/BR2 was evaluated in vitro against 19 invertebrate families (28 species) and 20 spider families (32 species). It was able to amplify the correct sequence from 25 invertebrate species and 16 spider species (Appendix S3–S9). Only one Hymenoptera species (*Tricopria* sp.) was not amplified. A sequence obtained from the PCR with LCO1490, identified as *Pardosa palustris*, was not amplified with NoSpi2/BR2. Two species (one Collembola and one Diptera) could not be identified by sequencing due to the low quality of the reads. The primer pair NoSpi2/BR2 produced contrasting results for spiders (Appendix S3). Several families were not amplified: Lycosidae, Amaurobidae, Clubionidae, Ctenidae, Dysderidae, Oonopidae, Oxyopidae, Philodromidae, Pholcidae, Pisauridae, and Thomisidae. It appears that the Dysderidae sample was also negative with LCO/HCO2198 suggesting that no amplifiable DNA was present (i.e., no DNA present or PCR inhibition). The Oonopidae and Pisauridae samples were amplified by LCO1490/HCO2198, but the species could not be identified. Among the Linyphiidae species tested, only one sample was not amplified, but the genus mismatch between the morphological and DNA identification suggests a sample error. All mixes for the assessment of a potential inhibitory effect of predator DNA and for the detection threshold of the primer were successfully amplified by NoSpi2/BR2 and provided accurate sequencing results.

The NoSpi2 + BR2 primer set did recover up to 94.17% of taxa of the mock community, with an average of 310.7 ($SD = 4.22$) and 352.2 ($SD = 2.51$) taxa recovered at 10,000 and 100,000 read sequencing

depth, respectively (subsampling with 1,000 iterations). The same primer set recovered an average of 445.7 ($SD = 9.60$) and 678.7 ($SD = 6.17$) taxa of the malaise sample (again at 10,000 and 100,000 read sequencing depth). This recovery represents approximately 10%–15% lower taxon recovery than the currently most efficient universal invertebrate primer pair (e.g., BF3 + BR2) in Elbrecht et al. (2019).

3.3 | Field experiment

Among the 388 spiders sampled, 126 led to positive amplification results. The twelve species tested gave positive amplifications. We obtained 540 ASVs corresponding to 12 orders, 67 families, and 117 species (Table 2 and Figure 4, detailed table in Appendix S10 and raw ASVs in Appendix S11). Fifteen spider samples resulted in ASVs that had no match in BOLD nor NCBI GenBank. One ASV corresponded to a spider but not to predator DNA (*Clubiona lutescens*). No reads corresponded to predator DNA.


4 | DISCUSSION

In this study, we developed a forward primer (NoSpi2) specifically designed for metabarcoding of gut contents of *Pardosa* spiders and phylogenetically closely related families of the oval calamistrum clade. Together with the previously designed reverse primer BR2 (Elbrecht & Leese, 2017b), it detected DNA from a broad range of aquatic and terrestrial arthropods. By using a mock sample and a malaise trap sample from a prior study evaluating primer performance (Elbrecht et al., 2019), we were able to confirm that NoSpi2 recovered about 94% of taxa. Taking design constraints into account, the primer should detect most prey taxa.

The performance of NoSpi2/BR2 was compared in silico to the barcoding primer pair LCO1490/HCO2198 (Folmer et al., 1994) which it outperformed for metabarcoding all of the groups tested except Embioptera, which are typically tropical insects.

In vitro tests for single specimen were also very successful for prey, with most species being amplified. In some cases, results were less good, but this is likely not only linked to primer performance itself. For example, low matching scores for some chironomids are very likely the result of the underrepresentation of inland chironomids in GenBank and BOLD (Ekrem, Willassen, & Stur, 2007). Nevertheless, a poor performance of the primer pair on the species tested cannot be excluded. As expected, NoSpi2/BR2 in vitro tests for spiders showed no amplification for *Pardosa* and some members of the oval calamistrum clade. Only the amplification of *Piratula hygrophila* was contrary to our expectation, but Sanger sequencing revealed that the DNA originated from a very common Linyphiidae (*Neriene montana*), demonstrating the ability of our primer pair to detect intraguild predation among spiders. Similarly, the amplification of *Diaea dorsata* was unexpected, but Sanger sequencing again demonstrated that the DNA originated

FIGURE 2 Summary of the phylogenetic tree of the world's spider clades/families from Wheeler et al. (2016). Colors correspond to penalty scores calculated with PrimerMiner for NoSpi2. The higher the penalty score, the lower the likelihood of amplification



from a prey species: *Sialis fuliginosa* (Megaloptera). This result further demonstrates the ability of NoSpi2/BR2 to amplify prey DNA without strong amplification of predator DNA. The detection of prey is surprising as only DNA from legs was used, but spider digestive caeca extend into femurs making the detection of prey possible, even when using only legs. We expected members of the families Amaurobidae, Clubionidae, Oonopidae, Philodromidae, and Pholcidae to be amplified by NoSpi2/BR2 given their low primer bias value (<320). This was not the case; however, only one sample was tested per family. More tests are required to verify the capacity of NoSpi2/BR2 to amplify these families. Dysderidae were neither amplified with NoSpi2/BR2 nor amplified with LCO1490/HCO2198. The low bias values (306 for NoSpi2 and 6 for HCO) suggest the presence of PCR inhibitors or DNA nucleases in the sample. In vitro tests with different mixes of spider and prey (Trichoptera) suggested no inhibitory effect or lower detection threshold through the inclusion of predator DNA. Indeed, prey was successfully amplified in all mixes and was identifiable to species level with matches greater than 99%, even at very low concentrations in the mix (2.8^{-6} ng/ μ l).

Our metabarcoding test of the primer pair against a mock community demonstrated a very satisfying recovery of 94.14% of

species. The mock community only included species from families supposed to be amplified by the primer pair. Future user must keep in mind that within each family around 6% of the species might not be detected. The opposite is true for spiders: within each family not supposed to be amplified, some species might be. The metabarcoding test on a malaise trap showed that while not amplifying *Pardosa* spiders, the primer pair performed well compared with the best primer pair known for malaise trap metabarcoding. Indeed, it detected only 15% fewer species.

The field test confirmed that NoSpi2/BR2 can detect a large number of arthropod species (up to 12 orders and 117 species). Interestingly, no predator DNA was amplified, confirming that there is no amplification for the tested Lycosidae species (genera tested: *Pardosa*, *Trochosa*, *Hygrolycosa*, *Xerolycosa*, and *Pirata*) using NoSpi2.

Studies focusing on spider diet in natural habitats using metabarcoding are rare. Wirta, Weingartner, Hambäck, and Roslin (2015) found *Pardosa glacialis* to consume mainly Diptera and Lepidoptera in the High Arctic. Hambäck et al. (2016) also found Diptera as the main prey of *Pardosa prativaga* followed by Lepidoptera, Coleoptera, and Heteroptera in the Baltic shoreline. In both cases, the authors used primers designed to amplify specifically Diptera and Lepidoptera,




FIGURE 3 Radar charts of penalty scores for each forward and reverse primer. Scores are computed using PrimerMiner (Elbrecht & Leese, 2017a). Low scores correspond to high amplification efficiency

TABLE 2 Summary table showing the number of species and reads for each order found in 126 spider gut contents

Class or order	Number of species	Number of reads
Acari	3	7,706
Araneae	1	1,027
Coleoptera	5	33,063
Collembola	4	6,568
Diptera	42	242,192
Hemiptera	32	116,518
Hymenoptera	14	137,023
Lepidoptera	6	16,838
Opiliones	2	521
Stylommatophora	1	48
Thysanoptera	2	43
Trichoptera	2	4,256

potentially resulting in an underestimation of other orders. The main orders constituting Lycosid diets in our study were Diptera, Hemiptera, Hymenoptera, and Lepidoptera. Interestingly, spiders, harvestmen, and carabid beetles were also found in spider gut contents which confirm the potential of NoSpi2 for the investigation of intraguild predation, which has often been documented among spiders and for carabid beetles feeding on spiders (Davey et al., 2012; Lang, 2003; Sitvarin & Rypstra, 2014). Although coleopterans have been reported as spider prey (Hambäck et al., 2016), predation of carabid beetles by spiders is less commonly reported than the opposite (but see: Krehenwinkel, Kennedy, Pekár, & Gillespie, 2017; Roubinet et al., 2018; Sint, Kaufmann, Mayer, & Traugott, 2019; Staudacher et al., 2018).

Snail (Stylommatophora) DNA was also detected in spider gut content, showing that despite gastropods generally being considered carabid and harvestmen prey, they are indeed a constituent of the diet of spiders (Nyffeler & Symondson, 2001).




FIGURE 4 Proportion of prey orders detected per spider species. Numbers within bars represent the total number of ASVs per species. Numbers above the bars show the number of individuals tested per species

Despite these very promising results, the fragments generated by NoSpi2/BR2 are relatively long (403 bp). This length ensures a high taxonomic resolution, but it could also limit DNA detectability (Symondson, 2002) due to the higher likelihood of missing shorter degraded sequences (Deagle, Eveson, & Jarman, 2006; Symondson, 2002). Further studies, particularly feeding experiments, could help to determine the dynamics of DNA degradation enhancing our understanding of spider diet.

5 | CONCLUSION

Molecular techniques that enable the simple and accurate analysis of spider diet are of great importance for studies investigating the impact of environmental change on ecological processes affecting the most abundant arthropod predators. The primer NoSpi2 developed and extensively validated for this study is ideal for gut content analysis of predator spider species of the genus *Pardosa* and should perform well for other species of the oval calamistrum

clade. It is enabling dietary and biological control studies as well as intraguild predation studies. NoSpi2 amplifies a large number of prey but not the predator species and thereby renders dissection unnecessary because an entire homogenized spider can be used as sample.

ACKNOWLEDGMENTS

We also thank the RIVEM Group (Karlstad University) for funding the in vitro and field tests. V.E. and D.S., as well as the metabarcoding experiments, were supported by funding through the University of Guelph's Canada First Research Excellence Fund project Food from Thought.

CONFLICT OF INTEREST

None declared.

AUTHOR CONTRIBUTIONS

D.L. conceived the idea and led the project. V.E. designed the primer. D.L., V.E., J.C., P.A.H., and A.E. contributed to the laboratory work,

data processing, and analyses. D.L., J.C., and V.E. wrote the draft manuscript. All authors contributed critically to the drafts and gave final approval for publication.

DATA AVAILABILITY STATEMENT

Trimmed sequences from the Sanger sequencing (in vitro evaluation) are available under the NOSPI project on BOLD. Raw sequence data from the mock community metabarcoding and malaise sample are available under the SRA accession numbers SRX5975749 and SRX5975748. Data from the complete run with the 96 additional samples not analyzed in this study are available under accession PRJNA546583. Raw sequence data from the field test metabarcoding are available under SRA accession number PRJNA550981. The code used for data analysis is available on GitLab under the project ID: 9419382 (<https://gitlab.com/DenisLafage/nospi>).

ORCID

Denis Lafage  <https://orcid.org/0000-0002-6205-611X>
 Vasco Elbrecht  <https://orcid.org/0000-0003-4672-7099>
 Jordan P. Cuff  <https://orcid.org/0000-0002-0198-4940>
 Peter A. Hambäck  <https://orcid.org/0000-0001-6362-6199>
 Ann Erlandsson  <https://orcid.org/0000-0002-3813-9548>

REFERENCES

- Agustí, N., Shayler, S. P., Harwood, J. D., Vaughan, I. P., Sunderland, K. D., & Symondson, W. O. C. (2003). Collembola as alternative prey sustaining spiders in arable ecosystems: Prey detection within predators using molecular markers. *Molecular Ecology*, 12(12), 3467–3475. <https://doi.org/10.1046/j.1365-294X.2003.02014.x>
- Birkhofer, K., Bylund, H., Dalin, P., Ferlian, O., Gagic, V., Hambäck, P. A., ... Jonsson, M. (2017). Methods to identify the prey of invertebrate predators in terrestrial field studies. *Ecology and Evolution*, 7(6), 1942–1953. <https://doi.org/10.1002/ece3.2791>
- Boreau de Roincé, C., Lavigne, C., Mandrin, J.-F., Rollard, C., & Symondson, W. O. C. (2013). Early-season predation on aphids by winter-active spiders in apple orchards revealed by diagnostic PCR. *Bulletin of Entomological Research*, 103, 148–154. <https://doi.org/10.1017/S0007485312000636>
- Braukmann, T. W. A., Ivanova, N. V., Prosser, S. W. J., Elbrecht, V., Steinke, D., Ratnasingham, S., ... Hebert, P. D. N. (2018). Revealing the complexities of metabarcoding with a diverse arthropod mock community. *BioRxiv*, 433607. <https://doi.org/10.1101/433607>
- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, 13(7), 581–583. <https://doi.org/10.1038/nmeth.3869>
- Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, 10(1), 421. <https://doi.org/10.1186/1471-2105-10-421>
- Chapman, E. G., Schmidt, J. M., Welch, K. D., & Harwood, J. D. (2013). Molecular evidence for dietary selectivity and pest suppression potential in an epigeal spider community in winter wheat. *Biological Control*, 65(1), 72–86. <https://doi.org/10.1016/j.biocntrol.2012.08.005>
- Davey, J. S., Vaughan, I. P., Andrew King, R., Bell, J. R., Bohan, D. A., Bruford, M. W., ... Symondson, W. O. C. (2012). Intraguild predation in winter wheat: Prey choice by a common epigeal carabid consuming spiders. *Journal of Applied Ecology*, 50(1), 271–279. <https://doi.org/10.1111/1365-2664.12008>
- Deagle, B. E., Eveson, J. P., & Jarman, S. N. (2006). Quantification of damage in DNA recovered from highly degraded samples – A case study on DNA in faeces. *Frontiers in Zoology*, 3(1), 11. <https://doi.org/10.1186/1742-9994-3-11>
- Ekrem, T., Willassen, E., & Stur, E. (2007). A comprehensive DNA sequence library is essential for identification with DNA barcodes. *Molecular Phylogenetics and Evolution*, 43(2), 530–542. <https://doi.org/10.1016/j.ympev.2006.11.021>
- Elbrecht, V., Braukmann, T. W. A., Ivanova, N. V., Prosser, S. W. J., Hajibabaei, M., Wright, M., ... Steinke, D. (2019). Validation of COI metabarcoding primers for terrestrial arthropods. *PeerJ*, 7, e7745.
- Elbrecht, V., & Leese, F. (2017a). PrimerMiner: An R package for development and *in silico* validation of DNA metabarcoding primers. *Methods in Ecology and Evolution*, 8(5), 622–626. <https://doi.org/10.1111/2041-210X.12687>
- Elbrecht, V., & Leese, F. (2017b). Validation and development of coi metabarcoding primers for freshwater macroinvertebrate bioassessment. *Frontiers in Environmental Science*, 5, 1–11. <https://doi.org/10.3389/fenvs.2017.00011>
- Elbrecht, V., & Steinke, D. (2018). Scaling up DNA metabarcoding for freshwater macrozoobenthos monitoring. *Freshwater Biology*, 64, 380–387. <https://doi.org/10.7287/peerj.preprints.3456v4>
- Elbrecht, V., Vamos, E. E., Meissner, K., Aroviita, J., & Leese, F. (2017). Assessing strengths and weaknesses of DNA metabarcoding-based macroinvertebrate identification for routine stream monitoring. *Methods in Ecology and Evolution*, 8(10), 1265–1275. <https://doi.org/10.1111/2041-210X.12789>
- Folmer, O., Black, M., Hoeh, W., Lutz, R., & Vrijenhoek, R. (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3(5), 294–299. <https://doi.org/10.1371/journal.pone.0013102>
- Greenstone, M. H., & Shufan, K. A. (2003). Spider predation: Species specific identification of gut contents by polymerase chain reaction. *Journal of Arachnology*, 31, 131–134. [https://doi.org/10.1636/0161-8202\(2003\)031\[0131:SPSI0G\]2.0.CO;2](https://doi.org/10.1636/0161-8202(2003)031[0131:SPSI0G]2.0.CO;2)
- Hambäck, P. A., Weingartner, E., Dalén, L., Wirta, H., & Roslin, T. (2016). Spatial subsidies in spider diets vary with shoreline structure: Complementary evidence from molecular diet analysis and stable isotopes. *Ecology and Evolution*, 6(23), 8431–8439. <https://doi.org/10.1002/ece3.2536>
- Holland, J. M., Bianchi, F. J. J. A., Entling, M. H., Moonen, A.-C., Smith, B. M., & Jeanneret, P. (2016). Structure, function and management of semi-natural habitats for conservation biological control: A review of European studies. *Pest Management Science*, 72(9), 1638–1651. <https://doi.org/10.1002/ps.4318>
- Hosseini, R., Keller, M. A., Schmidt, O., & Li, D.-M. (2011). Design and assessment of primers to detect insect pests of Brassica crops in the gut contents of arthropod predators. *Munis Entomology and Zoology*, 6(1), 477–492.
- Hugerth, L. W., Wefer, H. A., Lundin, S., Jakobsson, H. E., Lindberg, M., Rodin, S., ... Andersson, A. F. (2014). DegePrime, a program for degenerate primer design for broad-taxonomic-range PCR in microbial ecology studies. *Applied and Environmental Microbiology*, 80(16), 5116–5123. <https://doi.org/10.1128/AEM.01403-14>
- Kamenova, S., Mayer, R., Rubbmark, O. R., Coissac, E., Plantegenest, M., & Traugott, M. (2018). Comparing three types of dietary samples for prey DNA decay in an insect generalist predator. *Molecular Ecology Resources*, 18(5), 966–973. <https://doi.org/10.1111/1755-0998.12775>
- Katoh, K. (2002). MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research*, 30(14), 3059–3066. <https://doi.org/10.1093/nar/gkf436>

- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., ... Drummond, A. (2012). Geneious basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*, 28(12), 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Kraus, J. M., Pomeranz, J. F., Todd, A. S., Walters, D. M., Schmidt, T. S., & Wanty, R. B. (2016). Aquatic pollution increases use of terrestrial prey subsidies by stream fish. *Journal of Applied Ecology*, 53(1), 44–53. <https://doi.org/10.1111/1365-2664.12543>
- Krehenwinkel, H., Kennedy, S. R., Adams, S. A., Stephenson, G. T., Roy, K., & Gillespie, R. G. (2019). Multiplex PCR targeting lineage-specific SNP: A highly efficient and simple approach to block out predator sequences in molecular gut content analysis. *Methods in Ecology and Evolution*, 10, 982–993. <https://doi.org/10.1111/2041-210X.13183>
- Krehenwinkel, H., Kennedy, S., Pekár, S., & Gillespie, R. G. (2017). A cost-efficient and simple protocol to enrich prey DNA from extractions of predatory arthropods for large-scale gut content analysis by Illumina sequencing. *Methods in Ecology and Evolution*, 8(1), 126–134.
- Krehenwinkel, H., Rödder, N., & Tautz, D. (2015). Eco-genomic analysis of the poleward range expansion of the wasp spider *Argiope bruennichi* shows rapid adaptation and genomic admixture. *Global Change Biology*, 21, 4320–4332.
- Kuusik, A.-K., Cassel-Lundhagen, A., Kvarnheden, A., & Ekblom, B. (2008). Tracking aphid predation by lycosid spiders in spring-sown cereals using PCR-based gut-content analysis. *Basic and Applied Ecology*, 9(6), 718–725. <https://doi.org/10.1016/j.baae.2007.08.012>
- Lafage, D., Bergman, E., Eckstein, R. L., Österling, E. M., Sadler, J. P., & Piccolo, J. J. (2019). Local and landscape drivers of aquatic-to-terrestrial subsidies in riparian ecosystems: A worldwide meta-analysis. *Ecosphere*, 10(4), e02697. <https://doi.org/10.1002/ecs2.2697>
- Lang, A. (2003). Intraguild interference and biocontrol effects of generalist predators in a winter wheat field. *Oecologia*, 134(1), 144–153. <https://doi.org/10.1007/s00442-002-1091-5>
- Nyffeler, M., & Symondson, W. O. C. (2001). Spiders and harvestmen as gastropod predators. *Ecological Entomology*, 26(6), 617–628. <https://doi.org/10.1046/j.1365-2311.2001.00365.x>
- Piacentini, L. N., & Ramirez, M. J. (2019). Hunting the wolf: A molecular phylogeny of the wolf spiders (Araneae, Lycosidae). *Molecular Phylogenetics and Evolution*, 136, 227–240. <https://doi.org/10.1016/j.ympev.2019.04.004>
- Piñol, J., San Andrés, V., Clare, E. L., Mir, G., & Symondson, W. O. C. (2014). A pragmatic approach to the analysis of diets of generalist predators: The use of next-generation sequencing with no blocking probes. *Molecular Ecology Resources*, 14(1), 18–26. <https://doi.org/10.1111/1755-0998.12156>
- Piñol, J., Senar, M. A., & Symondson, W. O. C. (2018). The choice of universal primers and the characteristics of the species mixture determine when DNA metabarcoding can be quantitative. *Molecular Ecology*, 28(2), 407–419. <https://doi.org/10.1111/mec.14776>
- Pompanon, F., Deagle, B. E., Symondson, W. O. C., Brown, D. S., Jarman, S. N., & Taberlet, P. (2012). Who is eating what: Diet assessment using next generation sequencing. *Molecular Ecology*, 21(8), 1931–1950. <https://doi.org/10.1111/j.1365-294X.2011.05403.x>
- R Core Team (2019). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <http://www.r-project.org>
- Ratnasingham, S., & Hebert, P. D. (2007). BOLD: The barcode of life data system (<http://www.barcodinglife.org>). *Molecular Ecology Notes*, 7(3), 355–364.
- Riechert, S. E., & Lockley, T. (1984). Spiders as biological control agents. *Annual Review of Entomology*, 29, 299–320. <https://doi.org/10.1146/annurev.en.29.010184.001503>
- Roubinet, E., Jonsson, T., Malsher, G., Staudacher, K., Traugott, M., Ekblom, B., & Jonsson, M. (2018). High redundancy as well as complementary prey choice characterize generalist predator food webs in agroecosystems. *Scientific Reports*, 8(1), 8054. <https://doi.org/10.1038/s41598-018-26191-0>
- Sint, D., Kaufmann, R., Mayer, R., & Traugott, M. (2019). Resolving the predator first paradox: Arthropod predator food webs in pioneer sites of glacier forelands. *Molecular Ecology*, 28(2), 336–347. <https://doi.org/10.1111/mec.14839>
- Sint, D., Thurner, I., Kaufmann, R., & Traugott, M. (2015). Sparing spiders: Faeces as a non-invasive source of DNA. *Frontiers in Zoology*, 12(1), 3. <https://doi.org/10.1186/s12983-015-0096-y>
- Sitvarin, M. I., & Rypstra, A. L. (2014). The importance of intraguild predation in predicting emergent multiple predator effects. *Ecology*, 95(10), 2936–2945. <https://doi.org/10.1890/13-2347.1>
- Staudacher, K., Rennstam Rubbmark, O., Birkhofer, K., Malsher, G., Sint, D., Jonsson, M., & Traugott, M. (2018). Habitat heterogeneity induces rapid changes in the feeding behaviour of generalist arthropod predators. *Functional Ecology*, 32(3), 809–819. <https://doi.org/10.1111/1365-2435.13028>
- Symondson, W. O. C. (2002). Molecular identification of prey in predator diets. *Molecular Ecology*, 11(4), 627–641. <https://doi.org/10.1046/j.1365-294X.2002.01471.x>
- Turnbull, A. L. (1973). Ecology of the true spiders (Araneomorphae). *Annual Review of Entomology*, 18(1), 305–348. <https://doi.org/10.1146/annurev.en.18.010173.001513>
- Vamos, E., Elbrecht, V., & Leese, F. (2017). Short COI markers for freshwater macroinvertebrate metabarcoding. *Metabarcoding and Metagenomics*, 1, e14625. <https://doi.org/10.3897/mbmg.1.14625>
- Vestheim, H., & Jarman, S. N. (2008). Blocking primers to enhance PCR amplification of rare sequences in mixed samples – A case study on prey DNA in Antarctic krill stomachs. *Frontiers in Zoology*, 5(1), 12. <https://doi.org/10.1186/1742-9994-5-12>
- Waldner, T., Sint, D., Juen, A., & Traugott, M. (2013). The effect of predator identity on post-feeding prey DNA detection success in soil-dwelling macro-invertebrates. *Soil Biology and Biochemistry*, 63, 116–123. <https://doi.org/10.1016/j.soilbio.2013.03.030>
- Walters, D. M., Otter, R. R., Kraus, J. M., & Mills, M. A. (2018). Riparian spiders indicate the magnitude and sources of polychlorinated biphenyl contamination at a large contaminated sediment site. *Environmental Toxicology and Chemistry*, 37, 2467–2474. <https://doi.org/10.1002/etc.4216>
- Wheeler, W. C., Coddington, J. A., Crowley, L. M., Dimitrov, D., Goloboff, P. A., Griswold, C. E., ... Zhang, J. (2016). The spider tree of life: Phylogeny of Araneae based on target-gene analyses from an extensive taxon sampling. *Cladistics*, 33, 574–616. <https://doi.org/10.1111/cla.12182>
- Wirta, H. K., Weingartner, E., Hambäck, P. A., & Roslin, T. (2015). Extensive niche overlap among the dominant arthropod predators of the High Arctic. *Basic and Applied Ecology*, 16(1), 86–92. <https://doi.org/10.1016/j.baae.2014.11.003>
- Xu, C. C. Y., Yen, I. J., Bowman, D., & Turner, C. R. (2015). Spider web DNA: A new spin on noninvasive genetics of predator and prey. *PLoS ONE*, 10(11), e0142503. <https://doi.org/10.1371/journal.pone.0142503>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Lafage D, Elbrecht V, Cuff JP, Steinke D, Hambäck PA, Erlandsson A. A new primer for metabarcoding of spider gut contents. *Environmental DNA*. 2020;2:234–243. <https://doi.org/10.1002/edn3.62>