



HAL
open science

Shaping of soil microbial communities by plants does not translate into specific legacy effects on organic carbon mineralization

Eve Hellequin, Françoise Binet, Olivier Klarzynski, Sara Hallin, Jaanis Juhanson, Virginie Daburon, Cécile Monard

► To cite this version:

Eve Hellequin, Françoise Binet, Olivier Klarzynski, Sara Hallin, Jaanis Juhanson, et al.. Shaping of soil microbial communities by plants does not translate into specific legacy effects on organic carbon mineralization. *Soil Biology and Biochemistry*, 2021, 163, pp.108449. 10.1016/j.soilbio.2021.108449 . hal-03430174

HAL Id: hal-03430174

<https://univ-rennes.hal.science/hal-03430174>

Submitted on 16 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 Shaping of soil microbial communities by plants does not translate into 2 specific legacy effects on organic carbon mineralization

3

4 Eve Hellequin ^{1*}, Françoise Binet ¹, Olivier Klarzynski ², Sara Hallin³, Jaanis Juhanson³,
5 Virginie Daburon¹, Cécile Monard^{1*}

6 1. University of Rennes, CNRS, ECOBIO [(Ecosystèmes, biodiversité, évolution)] - UMR 6553, F-35000 Rennes, France.

7 2. BIO3G Company, 7 rue du Bourg-Neuf 22230 Merdrignac, France.

8 3. Swedish University of Agricultural Sciences, Department of Forest Mycology and Plant Pathology, Box 7026, 75007,
9 Uppsala, Sweden

10 * Corresponding authors:

11 -Eve Hellequin (current address: University of Sorbonne, CNRS, EPHE, PSL, UMR METIS, F-75005 Paris, France.
12 (eve.hellequin@upmc.fr)

13 -Cécile Monard (cecile.monard@univ-rennes1.fr)

14 Abstract

15 Plants shape soil microbial communities through their root architecture, their rhizodeposits
16 and return of dead plant material to the soil. These interactions can have a strong influence
17 on the soil organic carbon dynamics. However, it is unclear whether the plant species effects
18 on the soil microbial community could influence the organic carbon mineralization through
19 plant legacy effects. Therefore, we examined how and to what extent a short-term plant
20 growing phase affected the total and active soil microorganisms and through a possible plant
21 legacy, also the mineralization of soil organic carbon, a central ecosystem function. Using a
22 controlled pot experiment, we first showed that the two phylogenetically distinct plants,
23 *Arabidopsis thaliana* and *Triticum aestivum*, differently shaped the soil microbial
24 communities when recruiting from the same soil community. Although both plants recruited
25 plant-growth promoting bacteria in the vicinity of their roots, *A. thaliana* had a stronger
26 effect than *T. aestivum* and also recruited saprophytic fungi, while inhibiting fungal
27 pathogens. Due to plant legacy effects on the soil microbial communities, different microbial
28 successions occurred in the two previously planted soils when subjected to plant litter. By
29 contrast, plant legacy effects on soil basal respiration were not plant-specific, with basal
30 respiration increasing similarly in both cases and moreover did not translate to changes in
31 litter carbon mineralization in the short-term of 49 days. Our results suggest that the soil
32 nutrient dynamics rather than changes in soil microbial community composition drive the
33 organic carbon mineralization of added litter. The present study brings new insights in how

34 the relationships between plants, microorganisms and soil nutrient dynamics affect litter
35 carbon cycling.

36 **Key-words:** plant legacy, litter mineralization, soil bacteria, soil fungi, *Arabidopsis thaliana*,
37 *Triticum aestivum*, microbial succession

38 Introduction

39 Soil harbors a large amount of phylogenetically and functionally diverse
40 microorganisms (Torsvik and Øvreås, 2002) that interact with plants and contribute to
41 above-below ground exchanges. The “plant-soil feedbacks” concept is defined as plant-
42 mediated changes in soil microbiota and soil nutrient availability that influence the fitness of
43 the same or other plants in return (Perkins et al., 2011a; Van der Putten et al., 2013). Plants
44 shape the composition and diversity of soil microbiota through rhizodeposits and roots
45 architecture, litter C:N ratio and litter carbon quality (Heal et al., 1997; Grayston et al., 1998;
46 Zak et al., 2003; Gu et al., 2012). By providing new habitats and resources in the vicinity of
47 their roots, plants induce the colonization of microorganisms, therefore the rhizosphere is
48 known to be a microbial hotspot (Lynch and Whipps, 1990 ; Lundberg et al., 2012; Reviewed
49 in Muller et al., 2016). The microbial diversity declines sequentially from bulk soil to
50 rhizosphere, suggesting increasingly stronger competition among microorganisms (Muller et
51 al., 2016) or a selection of specific microorganisms induced by the plant. Thus, plant roots
52 shape the microbial community directly or indirectly by enriching and/or activating specific
53 microorganisms, but also by depleting and/or inhibiting other ones (Berg and Smalla, 2009).

54 The modifications induced by plants on soil microbiota may persist after their
55 disappearance (i.e. legacy effects) (Van der Putten et al., 2016), for example, via their
56 rhizodeposits or litter. Such legacy effects of plants have been shown to influence soil carbon
57 storage (Lange et al., 2015) and soil respiration (de Oliveira et al., 2020), and could therefore
58 affect the mineralization of organic carbon and nutrients from litter. The capacity of the soil
59 to mineralize organic matter is attributed to the production of specific enzymes that allow
60 specialized microorganisms to use carbon and nutrients of organic matter with low
61 nutritional value, known as “metabolic ability” (Keiser et al., 2011). Depending on the
62 functional capacity of the decomposer communities, soils have different metabolic abilities
63 (Keiser et al., 2014, Veen et al., 2018). Thus, for the same litter, differences in its

64 decomposition rate can occur depending on the microbial community present and its active
65 fraction. While active soil microorganisms have been shown to only contribute 2% of the
66 total microbial biomass in soils, they are linked to soil metabolic ability (Blagodatskaya and
67 Kuzyakov, 2013) and may be more relevant than the total microbial community to assess
68 plant legacies on soil function. Hence, by shaping the microbial communities, plants may
69 indirectly also influence the soil organic matter decomposition and, in the long-term,
70 potentially accelerate its own litter decomposition by selecting an adapted decomposing
71 microbial community (Ayres et al., 2009a; Veen et al., 2015a).

72 In the present study, we asked whether plant species shape the soil microbial
73 community and if such plant-specific effects could influence, the organic carbon
74 mineralization through plant legacy effects. For this purpose, we conducted an experiment
75 with two phases, with the first phase denoted the “plant-soil experiment”, in which we
76 aimed at examining how a short-term (two months) planted agricultural soil affects the total
77 and the active microbial communities and to what extent it depends on plant species. In the
78 second phase referred to the “litter-carbon mineralization experiment”, we determined if
79 such changes induce plant legacy effects that affect the litter-carbon mineralization. Two
80 phylogenetically distinct plants, a common plant model (*Arabidopsis thaliana*) and a crop
81 (*Triticum aestivum*) were grown in the plant-soil experiment and followed by the litter-
82 carbon mineralization experiment using the soils from phase one to which litter of *T.*
83 *aestivum* was added. In this experiment, differences in the daily and cumulative
84 mineralization rates of organic carbon from litter and subsequent nutrient release were
85 determined. To identify early plant legacy effects on soil microbiota, both DNA (total
86 microbial community) and RNA (active microbial community) were used to quantify
87 microbial abundances (quantitative PCR) and identify the changes in microbial diversity using
88 a 16S rRNA gene and ITS amplicon sequencing approach.

89 We hypothesized that i) specific microbial communities would develop according to
90 plant species, through specific recruitment from the soil and ii) specific plant-induced
91 changes in microbial communities, soil nutrient and carbon contents would then translate
92 into plant-specific legacy effects on the mineralization of added litter to soil, despite the
93 generalist feature of this function.

94 2. Materials and methods

95 2.1. Plant-soil experiment (phase 1)

96 The soil was an agricultural topsoil (0-20 cm depth) from the agronomic experimental site of
97 the Brittany regional chamber of agriculture at Kerbernez (Plomelin, France; 47°56'39.3"N /
98 4°07'47.9"O), collected in April 2018 in the field border where grass was growing. It was air-
99 dried at ambient room temperature for 48 h and sieved at 4 mm prior to storage in the dark
100 at 4 °C. The soil was used for the plant-soil experiment, performed in May 2018. Soil
101 physical, chemical and microbial properties were determined following sieving at 2 mm and
102 according to Carter and Gregorich (2008) and are presented in Table 1. The soil was planted
103 with either *A. thaliana* or *T. aestivum* or not planted (bare soil). The *A. thaliana* seeds from
104 the Ecotype Columbia-O and the *T. aestivum* L. cv. "Baldus" seeds were kindly provided by
105 IsoLife BV (Wageninigen, the Netherlands). The seeds were surface-sterilized, directly sown
106 on 10 kg equivalent dry soil in sterile planters (length: 50 cm, width: 17 cm, height 15 cm) at
107 a density of 2.6×10^{-6} and 1.22×10^{-5} seeds per m^2 for *T. aestivum* and *A. thaliana*,
108 respectively. The planters were placed in an environmental growth chamber (Percival
109 scientific, Perry, Iowa, USA) with the following conditions: 16 hours light (day), 8 hours dark
110 (night) and light intensity of $160 \mu\text{mol m}^{-2} \text{s}^{-1}$, air temperature of 20 °C and relative humidity
111 of 80%. Soils were watered with sterile water periodically to maintain WHC around 60% by
112 weighting the planters. After 63 days, which corresponded to the flowering stage of *A.*
113 *thaliana*, the bare soil from the not planted planter and the soil in the vicinity of the roots of
114 *A. thaliana* and *T. aestivum* were harvested. Four replicates of the original soil (blank) and
115 four soil replicates (30 g of dry weight equivalent for each replicate) from the same bare and
116 planted planters (Bare soil, *A. thaliana* and *T. aestivum*) were sampled and stored at -20 °C
117 and -80 °C for further chemical and microbial DNA- and RNA-based analyses, respectively
118 (n=16). The remaining soil was used for setting-up the second experimental phase.

119 2.2 Litter-carbon mineralization experiment (phase 2)

120 In the second experimental phase, we used soil microcosms to study the plant legacy
121 effects on C mineralization based on a 3 x 2 factorial design (three soils from phase 1 without
122 and with *T. aestivum* litter addition). For each of the six treatments, four replicates were
123 independently designed for each of the five incubation times (0, 3, 7, 21 and 49 days) after

124 litter addition to soil, allowing destructive soil sampling. The design corresponded to a total
125 of 120 soil microcosms. Each soil (e.g. bare soil (control), and previously planted soils with *A.*
126 *thaliana* or *T. aestivum*) was sieved at 2 mm according to the AFNOR standard (AFNOR XP U
127 44–163) and soil aliquots of 30 g weight equivalent dry soil were mixed with or without 184
128 mg of *T. aestivum* sterile chopped straw, corresponding to an input of 40 mg of organic
129 carbon and 0.3 mg of organic nitrogen. The straw was previously chopped into fine pieces of
130 1 mm. To avoid soil contamination by external microorganisms, the straw was sterilized by
131 Ionisos (Dagneux, France) with gamma irradiation at 25 KGy. The sterilization efficiency was
132 checked by none amplification of the 16S rRNA gene with a polymerase chain reaction (PCR).
133 Each soil microcosm was placed in a hermitically closed 1 L glass jar that was incubated in
134 the dark at 28 °C. At each incubation time, except the initial time point, the CO₂ produced
135 and accumulated in the headspace was measured by micro gas chromatography (Agilent,
136 Santa Clara, California, USA) and expressed as mg of C-CO₂ g⁻¹ dry soil. Before each
137 measurement, the air of the headspace was homogenized with a 50 ml syringe. After each
138 measurement, the air of the headspace was entirely renewed, to ensure that the headspace
139 did not become anoxic, and the soil moisture was maintained at 60% WHC by weighting and
140 adding sterilized water if needed. At day 0 and after each CO₂ measurement (days 3, 7, 21
141 and 49) a time series of microcosms was destroyed and the soil was stored at -20 °C and -80
142 °C for further chemical and microbial RNA-based analyses, respectively.

143 **2.3 Chemical analyses**

144 The soil chemical analyses are detailed in Hellequin et al. (2020). Briefly, the total
145 organic carbon and nitrogen were determined with an elemental analyzer (Elementar, Vario
146 PYRO Cube, Lyon, France). The dissolved organic carbon and the dissolved inorganic nitrogen
147 (NO₃⁻ and NH₄⁺) were measured by an automated photometric analyzer following K₂SO₄
148 extraction. The Olsen phosphorus concentrations were determined according to Olsen et al.
149 (1954). The microbial carbon and nitrogen biomass (MBC and MBN, respectively) were
150 determined according to Vance et al. (1987).

151 **2.4 Microbial quantification and metabarcoding**

152 The DNA and RNA extractions were performed following the protocol described by
153 Griffiths et al. (2000) with some modifications according to Nicolaisen et al., (2008) and
154 Monard et al. (2013). Further details on DNA extraction and subsequent quantitative PCR
155 (qPCR) are available in supplementary materials (section 1). Half of the extract was stored at-
156 20 °C before further DNA-based analyses (original soil and 0 day samples). The other half
157 was subjected to a DNase treatment for further RNA-based analyses (original soil and
158 samples from day 0, 3, 7, 21 and 49). RNA (200 ng) were used for reverse transcription (RT).
159 Half of the cDNA products was stored at -20 °C to prepare 16S rRNA and ITS1 amplicons for
160 the quantification of the total bacteria and fungi by qPCR and the other half was used for 2 X
161 300 Illumina sequencing. The libraries were constructed using a two-step PCR approach "16S
162 metagenomics sequencing library preparation" protocol provided by Illumina.

163 **2.5 Data analyses**

164 The microbial sequences are available on the sequence read archive (SRA) database
165 (Bioproject PRJNA540147). The microbial sequence analyses were performed according to
166 Hellequin et al. (2020) with the following changes: i) only the bacterial and fungal OTUs
167 present in at least 4 out of 128 samples and representing at least 0.0005% of all sequences
168 were retained, which corresponded to OTUs with a minimum of 12 and 29 sequences for
169 bacteria and fungi, respectively. In the final data set, a total of 1,253,087 bacterial sequences
170 corresponding to 8,911 OTUs and 3,113,807 fungal sequences corresponding to 1,351 OTUs
171 were obtained.

172 All the statistical analyses were performed using R (v3.4.3, Core Team, 2017). The
173 number of sequences per sample was randomly subsampled to the lowest number, with
174 bacterial and fungal DNA and cDNA amplicons being normalized separately using the *rarefy*
175 function from the *vegan* package version 2.5-4 (Oksanen et al., 2019); 6 399 and 15 589 for
176 bacterial and fungal DNA, and 7 000 and 6 000 for bacterial and fungal cDNA. Twenty-one
177 out of the 256 samples were below these thresholds and were removed before further
178 analyses. The rarefaction curves at each sampling date and before and after normalization
179 were performed using the *rarecurve* function from the *vegan* package (Figures S1 and S2).
180 For each sample, the abundance of each OTU was calculated based on its relative abundance
181 from the sequencing analysis and to the bacterial 16S rRNA and fungal ITS quantifications

182 performed simultaneously. The richness (number of OTUs) (Figure S11) and Shannon
183 diversity index were calculated to estimate the alpha diversity using the *phyloseq* package
184 (McMurdie and Holmes, 2013).

185 To test for statistical differences between plant-soil treatments, we used analysis of variance
186 (ANOVA) and Tukey's post-hoc test on the following endpoints: emissions, bacterial and
187 fungal Shannon diversities, microbial carbon and nitrogen biomasses, soil nutrient (N and P)
188 contents, and total and active bacterial and fungal abundances. To test for significant
189 difference in the composition of the total and active bacterial and fungal communities at the
190 end of the plant growth phase, powered partial least squares discriminant analysis (PPLS-DA)
191 were generated using DNA-based OTU or RNA-based OTU tables using the *RVAideMemoire*
192 package version 0.9-72 (Hervé, 2019) and the *p/s* package version 2.7-0 (Mevik et al., 2018).
193 ANOVAs were carried out on each OTU to identify significant differences in their
194 abundances. The significant DNA-based or RNA-based OTUs were determined as enriched or
195 activated *versus* depleted or inhibited by the plants, respectively. Over the incubation period
196 in the second experiment, PPLS-DA were generated with the RNA-based OTUs and at each
197 sampling date to compare the active bacterial and fungal community compositions between
198 soils derived from bare soil, *A. thaliana* or *T. aestivum* and amended or not with litter.
199 ANOVAs were carried out on each OTU to identify significant differences in their abundances
200 according to the PPLS-DA results. In parallel, we used co-inertia analyses to identify the
201 relationships between the OTUs that contributed significantly to the differences in
202 community composition between soils and the soil dissolved organic carbon and nutrient
203 contents. Only the sampling times (i.e. day 7 for bacteria and days 0 and 49 for fungi) at
204 which both the PPLS-DA and the co-inertia resulted in significant differences were kept to
205 discuss the results. To identify microbial successions during the incubation, additional PPLS-
206 DA on the RNA-based OTUs were generated to compare the composition of active bacterial
207 and fungal communities over time.

208

209 **3. Results**

210 **3.1 Planting effects on soil carbon and nutrients (N, P) contents**

211 Without plants, the total content of dissolved mineral nitrogen increased significantly
212 after 63 days due to a higher amount of NO_3^- , whereas the content of NH_4^+ remained the
213 same (Figure 1). In soils with plants, dissolved soil organic carbon content increased
214 significantly, whereas the dissolved NO_3^- significantly decreased compared with the bare soil.
215 Both plants tended to increase the microbial carbon (MBC) (significant for *A. thaliana*) and
216 nitrogen biomass (MBN). The soil P content was significantly depleted by *A. thaliana*, but not
217 by *T. aestivum* (Figure 1). Furthermore, *A. thaliana* increased the ratio of carbon from soil
218 microbial biomass to total soil organic carbon compared to the original soil.

219 3.2. Planting effects on total and active soil microorganisms

220 Both bacteria and fungi were affected after 63 days of incubation without plants
221 present. A decrease in diversity of the total and active community of both bacteria and fungi
222 was observed as well as a decrease in the abundances of the active bacterial and fungal
223 communities and the RNA:DNA ratio (Figure 2). The active bacteria and fungi were even less
224 abundant in soils with plants. With *T. aestivum*, the original levels of total and active
225 bacterial diversity were maintained, whereas they increased in soil with *A. thaliana*. For the
226 total fungal diversity, the original level was maintained with both plant species, but the
227 diversity of the active fungal community decreased even though *A. thaliana* showed higher
228 diversity levels than bare soil. Interestingly, the copy numbers of the 16S rRNA was 3-fold
229 higher than its gene (Figure 2).

230 All soils were dominated by the same main bacterial phyla, i.e. *Alphaproteobacteria*,
231 *Acidobacteria*, *Actinobacteria* and *Planctomycetes*, but *A. thaliana* enriched for *Zygomycota*,
232 *Ascomycota* and some unknown fungi (Figure 3). The growth of *A. thaliana* and *T. aestivum*
233 induced significant changes in the structure of the total bacterial and fungal communities
234 and the active fungal community in the vicinity of their roots (Figure 4 A, E & G, $P = 0.004$,
235 89.81 % of variance explained; $P = 0.028$, 76.36 %; $P = 0.003$, 98.17 %, respectively), while
236 the structure of the active bacterial community was similar between planted and bare soils
237 (Figure 4 C, $P = 0.215$, 88.09 % of variance explained).

238 We further identified the OTUs driving the observed differences, either because they
239 were enriched or depleted (from DNA) and activated or inhibited (from cDNA) by the plants
240 (Figure 4 B, E, D, H). While the plants induced depletion or inhibition of a high number of
241 bacterial OTUs, they mainly induced enrichment or activation of specific fungal OTUs (Figure

242 4). *Bacillus* (OTUs 1 and 7) and *Methyloligellaceae* (OTU 17) were the most depleted bacteria
243 by the two plants while the *Xanthobacteraceae* (OTUs 5, 4 and 9) was specifically depleted
244 by *A. thaliana* (Figure 4B). Among the most abundant bacterial OTUs inhibited by the two
245 plants, some were depleted by both plants (OTUs 1, 4 and 7), and the bacterial OTUs
246 specifically inhibited by *A. thaliana* were all affiliated to *Bradyrhizobium* (OTUs 5, 3 and 2)
247 (4D). For fungi, the only OTU inhibited by the two plants was affiliated to *Sordariomycetes*
248 (OTU 1) (4H).

249 On top of these general trends, we found a stronger enrichment by *A. thaliana* than *T.*
250 *aestivum* that was more pronounced for fungi than bacteria (Figure 4 B & E). A total of 46
251 and 70 DNA-based bacterial and fungal OTUs, respectively, were enriched with *A. thaliana*,
252 against 32 and 14 with *T. aestivum*. *Proteobacteria* (delta, beta and alpha) were mainly
253 enriched by *A. thaliana* while *T. aestivum* mainly enriched *Actinobacteria*, *Acidobacteria*,
254 *Bacteroidetes* and *Planctomycetes*, (data not shown). For fungi, the enriched OTUs by *A.*
255 *thaliana* were highly abundant and mainly affiliated to *Mortierella* (OTUs 5, 11 and 13) (4F).
256 We detected a stronger activation of bacteria with *T. aestivum* than with *A. thaliana*, which
257 instead activated more fungi when compared with the total microbial communities (4 D &
258 G). The bacterial and fungal OTUs that were activated were also different from those that
259 were enriched (data not shown).

260 3.3. Plant legacies on basal respiration and carbon-litter mineralization

261 The modifications observed in the bacterial and fungal communities after 63 days of plant-
262 growth were plant specific. In the following litter-carbon mineralization experiment these
263 changes were associated with functional changes in the soil basal respiration rate. In soil
264 without litter addition, the basal respiration rate in soils that previously had plants,
265 regardless of plant species, was always higher than in the bare soil (Figure 5). The daily
266 emissions were peaking at day 7, with *T. aestivum* having the strongest effect, before
267 decreasing until day 49 independent of the previous treatment in phase 1.

268 Following soil C-enrichment (litter addition), the soil respiration rate increased in all soils as
269 expected (Figure 5). By contrast to soil without litter addition, the respiration rate and
270 cumulative C-respired were higher in the bare soil than in soils that previously had plants, at
271 days 7 and 21. At day 49, these observations reverted, the respiration rate being higher in
272 the soils that had plants than in the bare soil, with no differences between plants (Figure 5).

273

274 **3.4. Plant legacies on active microbial successions**

275 The temporal dynamic of abundances and composition of active bacterial and fungal
276 communities varied according to the soils being either C-enriched or not by litter (Figure 3).
277 In the previously bare soil without litter, the abundances of active bacteria and fungi
278 gradually decreased over time. In previously planted soils without litter addition we
279 observed an increase in the abundance of active fungi and bacteria at day 3. For fungi, this
280 was due to an activation of *Ascomycota* (Figure S8). At day 49, the bacterial and fungal
281 abundances were similar to those at the beginning of the second phase experiment.

282 In soils with litter addition, bacterial and fungal abundances gradually increased until day 21,
283 but then reverted to their initial abundance at day 49. The DNA-based OTUs significantly
284 enriched by the growth of *A. thaliana* or *T. aestivum* (Figure 4 B & F) were compared to their
285 corresponding RNA-based OTUs at all time points of the litter-carbon mineralization
286 experiment. Among the bacterial OTUs enriched by *A. thaliana* or *T. aestivum*, some were
287 activated at the beginning of the incubation (days 3 and 7), but did not remain activated
288 until day 49 (Figures S4 & S5). The fungal OTUs enriched by *A. thaliana* were not more
289 activated along the litter-carbon mineralization experiment (Figure S6), but for those
290 enriched by *T. aestivum*, we observed higher abundances at day 21 (Figure S7).

291

292 The soil C-enrichment with litter did not conceal the plant legacy effects detected without
293 litter addition. In *A. thaliana* soil, similar microbial dynamics as without litter were observed,
294 however, with litter, the changes in abundances were less pronounced. The increase in
295 microbial abundances observed at day 3 was due to an activation of *Alphaproteobacteria*
296 (*Rhizobiales*) and *Ascomycota* (*Sordariomycetes*) (Figure S9). Further, we observed an
297 activation of *Deltaproteobacteria* (*Myxococcales*) and *Zygomycota* at day 49. In the *T.*
298 *aestivum* soil, the community dynamics were different with litter than without, with high
299 bacterial and fungal abundances at day 21 and lower abundances at day 49 compared to day
300 0. The decreased fungal abundance at day 49 was mainly due to a decrease in active
301 *Ascomycota* (*Sordariomycetes*) (Figure S9).

302 When the structure of the active bacterial or fungal communities was different
303 between soils, we used co-inertia analyses to identify if there was any relationships between
304 the OTUs that contributed significantly to the differences between soils and the measured
305 soil variables. The NO_3^- soil content was the sole variable explaining the differences between
306 soils. Such significant relationships between planted soil history and soil parameters were
307 identified for bacteria at day 7 with litter and for fungi at day 0 and 49 with and without
308 litter, respectively (Figure S10). For bacteria, OTUs mainly affiliated to *Alphaproteobacteria*
309 (*Caulobacteraceae* and *Sphingomonadaceae*), *Bacteroidetes* (*Cytophagaceae* and
310 *Microscillaceae*) and *Gammaproteobacteria* (*Burkholderiaceae* and *Cellvibrio*) were more
311 abundant in the bare soil and positively correlated with the NO_3^- soil content (Figure S10).
312 For fungi, the OTUs that were more abundant in the bare soil and positively correlated with
313 the NO_3^- content were affiliated to *Ascomycota* (*Sordariomycetes* and *Dothideomycetes*).
314 Lastly, at day 49, the OTUs that were more abundant in the *A. thaliana* soil compared to the
315 bare soil, were negatively correlated with NO_3^- soil content and affiliated to *Zygomycota*
316 (*Mortierella*) and *Basidiomycota* (*Solicocozyma* and *Tremellales*).

317

318 **4. Discussion**

319 **4.1. Plant-specific effects on the total and active soil microorganisms and soil nutrients**

320 As hypothesized, the growth of both *A. thaliana* and *T. aestivum* induced the
321 selection of specific soil bacteria and fungi in the vicinity of their roots. We showed that the
322 effect of *A. thaliana* was stronger, especially for fungal communities. Root architecture as
323 well as rhizodeposits are highly plant species-specific and could thus differentially act on
324 microbial communities depending on plant species, leading to specific recruitment of root-
325 and rhizosphere-associated microorganisms (Berg and Smalla, 2009). In the present study
326 the presence of plants, irrespective of plant species, resulted in an increase in the dissolved
327 soil organic carbon content and carbon in microbial biomass, likely reflecting the inflow of
328 carbon to soil microbes through plant carbon deposition. At the same time, a depletion in
329 mineral nitrogen was observed, which could be due to plant N uptake that occurred
330 concomitantly with plant carbon deposition during plant growth. Tkacz et al. (2015) showed
331 that changes in the soil microbiota following plant growth appeared quickly after only one
332 plant generation and were plant-species dependant. However, in our study we showed a

333 stronger effect of *A. thaliana* than *T. aestivum* on abundances and community composition
334 of total bacteria and fungi, while Tkacz et al. (2015) showed the opposite, probably due to its
335 extensive root system (Tkacz et al., 2015). In our experiment the plants grew longer than in
336 the study of Tkacz et al. (2015), allowing *A. thaliana* to reach its flowering state. This could
337 explain the differences as it has been shown that the plant effects on soil microbiota depend
338 on the growth stage of the plant, with the strongest effect at the flowering stage, whereas
339 the effect is weaker during earlier growth stages (Cheng and Kuzyakov, 2005). The growth
340 stage might thus have overtaken the effect of root density in shaping rhizospheric
341 microbiota leading to different results than those of Tkacz et al. (2015). Another explanation
342 could be the differences in the properties of the soils used. While we used a sandy-loam
343 agricultural soil, Tkacz et al. (2015) used artificial soils consisting of compost or sand. They
344 observed that plant selection on the rhizosphere microbiota depended strongly on soil
345 composition probably due to contrasted organic nutrient contents. Nevertheless, both
346 studies results point toward a specific plant effect on soil microbial communities that
347 depends on the properties of the original soil.

348 Differences between total and active microbial community have already been demonstrated,
349 with active microorganisms being a subset of the total community (Romanowicz et al.,
350 2016). However, for bacteria, it is well known that the copy number of the 16S rRNA gene
351 varies between taxa (Větrovský and Baldrian, 2013) and Blazewicz et al., (2013) pointed out
352 that the relationship between rRNA concentration and bacterial growth rate can differ
353 greatly among taxa. These differences could explain why we observed higher 16S rRNA copy
354 numbers than its gene. Globally, we showed that the enriched or activated OTUs differed
355 between plant species, indicating a plant-specific effect on both total or active bacteria and
356 fungi. However, most of the depleted and inhibited OTUs were common to both plants and
357 the negative interaction therefore seemed less plant-specific. The depletion or inhibition of
358 certain OTUs was more pronounced for bacteria than for fungi. Surprisingly, both plants
359 depleted or inhibited *Bacillus sp.* and *Bradyrhizobium sp.*, of which some are known to be
360 plant growth-promoting bacteria (reviewed in Govindasamy et al., 2010). However, both
361 plants also recruited plant growth-promoting bacteria, which can provide direct benefits to
362 plant growth and health by influencing the nutrient status or affecting plant-pathogen
363 interaction. *T. aestivum* selected more total and active bacteria, while *A. thaliana* recruited

364 preferentially total and active fungi and inhibited fungal pathogens like *Alternaria* sp. Some
365 of these species, e.g. *Mycrosphaerella tassiana* and *Pyrenophora teres* (similar to OTUs 9, 7
366 and 2, respectively) are known to cause leaf diseases, such as net blotch (Thomma, 2003;
367 El-said et al., 2014; Moya et al., 2018). Further, *A. thaliana* enriched *Mortierella*, *Tremellales*
368 and *Sordariomycetes*, in agreement with both Tkacz et al., (2015) and Fabianska et al.,
369 (2019). *Mortierella* and *Tremellales* are known to be root associated saprotrophic fungi
370 involved in litter decomposition (Keim et al., 2014; Kuramae et al., 2013). Regarding the
371 main bacterial phyla enriched by *A. thaliana*, the *Proteobacteria* have already been detected
372 as recruited by this plant (Micallef et al., 2009; Bulgarelli et al., 2012; Lundberg et al., 2012)
373 and members of this phylum possess several plant growth-promoting mechanisms (see
374 references in Naz et al., 2014) and/or are associated with diseases suppression (Mendes et
375 al., 2011). *T. aestivum* enriched other bacteria, such as *Planctomycetes*, *Acidobacteria*,
376 *Bacteroidetes* and *Actinobacteria*, which is in agreement with those identified by Mahoney
377 et al. (2017). *Actinobacteria* and *Bacteroidetes* are known to be plant-growth-promoting
378 bacteria through the synthesis of different phytohormones, siderophores and enzymes and
379 also their ability for solubilization of mineral nutrients such as phosphorus, potassium, and
380 zinc (reviewed in Yadav et al., 2018). The soil phosphorus contents we measured in soil
381 planted with *T. aestivum* are in accordance with higher microbial ability for providing
382 phosphorus from soil matrix due to enrichment of plant growth-promoting bacteria by *T.*
383 *aestivum*. A renewal of the soil phosphorus pool through solubilization could explain the
384 differences in the soil content of phosphorus between the two plants, with the phosphorus
385 pool decreasing in *A. thaliana* but not in *T. aestivum*. Also root architecture and mycorrhizal
386 colonization can influence the efficiency of plants phosphorus uptake (Richardson et al.,
387 2009; Zuo et al., 2014).

388 **4.2. No plant legacy effects on the mineralization of soil organic carbon despite higher** 389 **plant-induced basal respiration**

390 Based on the observed plant species-specific effects on the microbial community
391 composition in the plant-soil experiment and the difference in the quantity or quality of the
392 carbon (from rhizodeposits or litter) between plant species (Johnson et al., 2007; Gul et al.,
393 2012), we expected different basal respiration rates during the mineralization experiment.
394 However, there were no plant-species effects, but rather a plant effect. The quantity of

395 organic carbon and the respiration rates were enhanced in both soils that previously had
396 plants compared to the bare soil when no litter was added. The previously planted soils also
397 had higher diversity of active and total bacteria as well as total fungi compared to the
398 previously bare soil. These results suggest functional redundancy within the communities,
399 which could be expected for carbon respiration, which is a generalist function of central
400 importance for soil ecosystem resilience (Bender et al., 2016). Moreover, we cannot exclude
401 that the plant legacy might have been reduced by the soil disturbance and the
402 homogenization that occurred during the soil sieving for conducting the carbon litter
403 mineralization experiment.

404 We did not observe any plant-specific effect following the addition of *T. aestivum*
405 litter. The seeds of *T. aestivum* used for the plant-soil experiment and the litter used for the
406 mineralization experiment came from the same variety and batch. Hence, in the context of
407 plant-soil feedbacks and legacy effect, we could have expected some litter specificity (Veen
408 et al., 2015a). Due to previous plant uptake, the mineral nutrient availability in the soil was
409 lower and could thus have limited the degradation of high C:N litter, as previously shown
410 (Partey et al., 2014). This nutrient limitation would explain the lower mineralization rates in
411 previously planted soil compared to the bare soil when litter was added. In agreement, we
412 observed a pulse of emission at day 7 in the bare soil with litter and simultaneously, we
413 identified that diazotrophic bacteria, such as *Caulobacteraceae* (*Alphaproteobacteria*),
414 *Burkholderia* and *Cellvibrio* (*Gammaproteobacteria*) (Suarez et al., 2014; Pagé et al., 2019),
415 were activated and positively correlated to the soil NO_3^- content. Their activity would
416 thereby support the nitrogen demand needed for degradation of the high C:N ratio litter
417 from *T. aestivum*. Moreover, the limitation of the litter carbon mineralization that we
418 observed and which was linked to the inorganic nitrogen status, is in agreement with the
419 regulatory gate theory proposed by Brookes et al. (2008). This theory suggests that the rate
420 of the soil organic matter mineralization is directly governed by abiotic processes, such as
421 nutrients dynamics in the soil matrix, rather than by soil microbial communities themselves
422 (composition, abundance or activity).

423 A microbial legacy due to specific plant species was observed until the end of the
424 mineralization experiment for both plants, although microbial successions with different
425 responses to the addition of litter according to soils were demonstrated. Among the OTUs

426 enriched by *A. thaliana*, some bacterial OTUs were also activated in the beginning of the
427 incubation period of the litter experiment but, globally, these OTUs did not remain activated
428 until the end of the experiment. We previously highlighted that both plant species recruited
429 total plant growth-promoting bacteria during their growth. Due to these intimate
430 interactions, the disappearance of the plants could lead to the inactivation of these specific
431 bacteria during the mineralization experiment. The peak in abundance of active bacteria
432 observed at day 3 was potentially due to the activation of copiotrophic microorganisms
433 (*Alphaproteobacteria* and *Ascomycota*), while at the end of the incubation, the oligotrophic
434 *Deltaproteobacteria* were stimulated (Bastian et al., 2009; Verastegui et al. 2014; Bastida et
435 al., 2016). Microbial abundances shared similar temporal patterns in the *T. aestivum* and *A.*
436 *thaliana* soils without litter, but not when the soil was C-enriched with litter. The peak of
437 active fungal abundances at day 21 was mainly due to unknown fungi and their decrease at
438 day 49 was mainly due to the copiotrophic *Ascomycota*. At this timepoint, the higher
439 abundances of active microorganisms in the soil that previously had plants were
440 concomitant with higher daily emissions.

441 To conclude, the effect of plants on the soil microbiome was plant-species specific.
442 While both plants recruited plant growth-promoting bacteria in the vicinity of their roots, *A.*
443 *thaliana* had a stronger effect on fungi than *T. aestivum*. The species specific responses were
444 also different between active and total bacterial and fungal communities. However, this was
445 not reflected in the observed plant legacy effect on the carbon mineralization from litter as
446 higher basal respiration rates were detected in planted soils irrespective of plant species.
447 Our results suggest that the soil nutrient dynamics induced by the presence of plants rather
448 than changes in soil microbial communities drive the organic carbon mineralization of added
449 litter. Nevertheless, soils with a planting history either maintained or displayed a higher
450 abundance of active soil microorganisms, which was concomitant with higher daily
451 respiration rates. In addition, the present study further highlights the importance to consider
452 both total (DNA-based) and active (RNA-based) microbial communities to be able to
453 differentiate how plants control enrichment/depletion and activation/inhibition of specific
454 microorganisms.

455

456 **Acknowledgments**

457 This research was supported by the "Association Nationale de la Recherche et de la
458 Technologies » (ANRT) through the convention N° 2016/0091 to F. Binet and O. Klarzynski
459 and by the University of Rennes 1 through its international mobility programm N° 12/
460 2018 to E. Hellequin. This research was also partly funded by the 2015-2016 BiodivERsA call
461 by The Swedish Research Council Formas (contract 2016-0194). The authors thank Mikael
462 Fauchaux for sampling the soil in the field before the experiment and Nathalie Le Bris and
463 Marion Chorin from the EcoChim platform for the soil nutrient measurements
464 (<https://ecobio.univ-rennes1.fr/plateforme-ecochimie-ecochim-presentation.php>).

465 **Competing Interests**

466 The authors declare no competing interests.

467 **References**

- 468 Ayres, E., Steltzer, H., Simmons, B.L., Simpson, R.T., Steinweg, J.M., Wallenstein, M.D.,
469 Mellor, N., Parton, W.J., Moore, J.C., Wall, D.H., 2009. Homefield advantage
470 accelerates leaf litter decomposition in forests. *Soil Biology & Biochemistry* 41, 606–
471 610.
- 472 Barberán, A., McGuire, K.L., Wolf, J.A., Jones, F.A., Wright, S.J., Turner, B.L., Essene, A.,
473 Hubbell, S.P., Faircloth, B.C., Fierer, N., 2015. Relating belowground microbial
474 composition to the taxonomic, phylogenetic, and functional trait distributions of
475 trees in a tropical forest. *Ecology Letters* 18, 1397–1405.
- 476 Bastian, F., Bouziri, L., Nicolardot, B., Ranjard, L., 2009. Impact of wheat straw decomposition
477 on successional patterns of soil microbial community structure. *Soil Biology &*
478 *Biochemistry* 41, 363– 275.
- 479 Bastida, F., Torres, I.F., Moreno, J.L., Baldrian, P., Ondoño, S., Ruiz-Navarro, A., Hernández,
480 T., Richnow, H.H., Starke, R., García, C., Jehmlich, N., 2016. The active microbial
481 diversity drives ecosystem multifunctionality and is physiologically related to carbon
482 availability in Mediterranean semi-arid soils. *Molecular Ecology* 25, 4660-4673.
- 483 Bender, S.F., Wagg, C., van Der Heijden, M.G.A., 2016. An underground revolution:
484 Biodiversity and soil ecological engineering for agricultural sustainability. *Trends in*
485 *Ecology and Evolution* 31, 440–452.
- 486 Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z., Godhe, A., De Wit, P.,
487 Sánchez-García, M., Ebersberger, I., de Sousa, F., Amend, A., Jumpponen, A.,
488 Unterseher, M., Kristiansson, E., Abarenkov, K., Bertrand, Y.J.K., Sanli, K., Eriksson, M.,
489 Vik, U., Veldre, V., Nilsson, H., 2013. Improved software detection and extraction of
490 ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for

- 491 analysis of environmental sequencing data. *Methods in Ecology and Evolution* 4, 914-
492 919.
- 493 Berg, G., Smalla, K., 2009. Plant species and soil type cooperatively shape the structure and
494 function of microbial communities in the rhizosphere. *FEMS Microbiology Ecology* 68,
495 1-13.
- 496 Blagodatskaya, E., Kuzyakov, Y., 2013. Active microorganisms in soil: critical review of
497 estimation criteria and approaches. *Soil Biology & Biochemistry* 67, 192–211.
- 498 Blazewicz, S.J., Barnard, R.L., Daly, R.A., Firestone, M.K., 2013. Evaluating rRNA as an
499 indicator of microbial activity in environmental communities: limitations and uses.
500 *ISME Journal* 7, 2061-2068.
- 501 Bodenhausen, N., Horton, M.W., Bergelson, J., 2013. Bacterial communities associated with
502 the leaves and the roots of *Arabidopsis thaliana*. *PLoS ONE* 8, e56329.
- 503 Brookes, P.C., Cayuela, M.L., Contin, M., De Nobili, M., Kemmitt, S.J., Mondini, C., 2008. The
504 mineralisation of fresh and humified soil organic matter by the soil microbial
505 biomass. *Waste Management* 28, 716–722.
- 506 Bulgarelli D., Rott, M., Schlaeppi, K., Ver Loren van Themaat, E., Ahmadinejad, N., Assenza,
507 F., Rauf, P., Huettel, B., Reinhardt, R., Schmelzer, E., Peplies, J., Gloeckner, F.O.,
508 Amann, R., Eickhorst, T., Schulze-Lefert, P., 2012. Revealing structure and assembly
509 cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature* 488, 91–95.
- 510 Cayuela, M.L., Sinicco, T., Mondini, C., 2009. Mineralization dynamics and biochemical
511 properties during initial decomposition of plant and animal residues in soil. *Applied*
512 *Soil Ecology* 41, 118–27.
- 513 Cheng, W., Kuzyakov, Y., 2005. Root effects on soil organic matter decomposition. In: Wright
514 S., Zobel R. (Eds.), *Roots and soil management: interactions between roots and the*
515 *soil*. Agronomy monograph, ASA, Madison, 48, 119–143.
- 516 Csardi, G., Nepusz, T., 2006. The *igraph* software package for complex network research.
517 *International Journal of Complex Systems* 1695. <http://igraph.org>
- 518 De Oliveira, A.B., Cantarel, A.A.M., Seiller, M., Florio, A., Bérard, A., Hinsinger, P., Le Cadre,
519 E., 2020. Short-term plant legacy alters the resistance and resilience of soil microbial
520 communities exposed to heat disturbance in a Mediterranean calcareous soil.
521 *Ecological Indicators* 108, 105740.
- 522 Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., Knight, R., 2011. UCHIME improves
523 sensitivity and speed of chimera detection. *Bioinformatics* 27, 2194–2200.
- 524 El-Said, A.H.M., Saleem, A., Maghraby, T.A., Hussein, M.A., 2014. Cellulase activity of some
525 phytopathogenic fungi isolated from diseases leaves of broad bean. *Archives of*
526 *Phytopathology and Plant Protection* 47(17), 2078–2094.

- 527 Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., Maman, S.,
528 Hernandez-Raquet, G., Combes, S., Pascal, G., 2018. FROGS: Find Rapidly OTUs With
529 Galaxy Solution. *Bioinformatics* 34, 1287–1294.
530 <https://doi.org/10.1093/bioinformatics/btx791> PMID:29228191
- 531 Eilers, K.G., Lauber, C.L., Knight, R., Fierer, N., 2010. Shifts in bacterial community structure
532 associated with inputs of low molecular weight carbon compounds to soil. *Soil*
533 *Biology & Biochemistry* 42, 896–903.
- 534 Fabianska, I., Gerlach, N., Almario, J., Bucher, M., 2019. Plant-mediated effect of soil
535 phosphorus on the root-associated fungal microbiota in *Arabidopsis thaliana*. *New*
536 *Phytologist* 221, 2123–2137.
- 537 Gardes, M., Bruns, T.D., 1993. ITS primers with enhanced specificity for basidiomycetes
538 application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2, 113-
539 118. <https://doi.org/10.1111/j.1365-294X.1993.tb00005.x> PMID:8180733
- 540 Govindasamy, V., Senthilkumar, M., Magheshwaran, V., Kumar, U., Bose, P., Sharma, V.,
541 Annapurna, K., 2010. *Bacillus* and *Paenibacillus* spp.: Potential PGPR for sustainable
542 agriculture. In Maheshwari D.K., (Ed.), *Plant growth and health promoting bacteria*.
543 Berlin, Springer-Verlag, 333–364.
- 544 Grayston, S.J., Wang, S., Campbell, CD., Edwards, A.C., 1998. Selective influence of plant
545 species on microbial diversity in the rhizosphere. *Soil Biology & Biochemistry* 30, 369-
546 378.
- 547 Griffiths, R.I., Whiteley, A.S., O'Donnell, A.G., Bailey, M.J., 2000. Rapid method for
548 coextraction of DNA and RNA from natural environments for analysis of ribosomal
549 DNA- and rRNA based microbial community composition. *Applied Environmental*
550 *Microbiology* 66, 5488–5491.
- 551 Gul, S., Whalen, J.K., Ellis, B.E., Mustafa, A.F., 2012. Influence of plant residue chemistry on
552 soil CO₂-C production: A study with *Arabidopsis thaliana* cell wall mutants of KNAT7,
553 MYB75 and CCR1. *Pedobiologia* 55 (6), 349-356,
- 554 Hanson, C.A., Allison, S.D., Bradford, M.A., Wallenstein, M.D., Treseder, K.K., 2008. Fungal
555 taxa target different carbon sources in forest soil. *Ecosystems* 11, 1157–
556 1167. <https://doi.org/10.1007/s10021-008-9186-4>
- 557 Heal, O.W., Anderson, J.M., Swift, M.J., 1997. Plant litter quality and decomposition: an
558 historical overview. In: Giller, K.E. and Cadisch, G. (eds), *Driven by nature-plant litter*
559 *quality and decomposition*. CAB Int, Wallingford, 3-32.
- 560 Hellequin, E., Monard, C., Quaiser, A., Henriot, M., Klarzynski, O., Binet, F., 2018. Specific
561 recruitment of soil bacteria and fungi decomposers following a biostimulant
562 application increased crop residues mineralization. *PLoS ONE* 13, e0209089.

- 563 Hellequin, E., Monard, C., Chorin, M., Le bris, N., Daburon, V., Klarzynski, O., Binet, F., 2020.
564 Responses of active soil microorganisms facing to a soil biostimulant input compared
565 to plant legacy effects. *Scientific Reports* 10, 13727.
- 566 Henrik Nilsson, R., Larsson, K-H., Taylor, A.F.S., Bengtsson-Palme, J., Jeppesen, T.S., Schigel,
567 D., Kennedy, P., Picard, K., Glöckner, F.O., Tedersoo, L., Saar, I., Kõljalg, U.,
568 Abarenkov, K., 2018. The UNITE database for molecular identification of fungi:
569 handling dark taxa and parallel taxonomic classifications. *Nucleic acids research*. doi:
570 10.1093/nar/gky1022
- 571 Hervé, M., 2019. RVAideMemoire: Testing and Plotting Procedures for Biostatistics. R
572 package version 0, 9-72. <https://CRAN.R-project.org/package=RVAideMemoire>
573
- 574 Jackson, S., Cimentada, J., Ruiz, E., 2019. corrr: Correlations in R. *R package version*.
575 0.3.1. <https://CRAN.R-project.org/package=corrr>
576
- 577 Johnson, J.M.F., Barbour, N.W., Weyers, S.L., 2007. Chemical composition of crop biomass
578 impacts its decomposition. *Soil Science Society of America Journal* 71, 155-162.
579
580
- 581 Keim, J., Mishra, B., Sharma, R., Ploch, S., Thines, M., 2014. Root-associated fungi of
582 *Arabidopsis thaliana* and *Microthlaspi perfoliatum*. *Fungal Diversity* 66, 99–111.
- 583 Keiser, A.D., Strickland, M.S., Fierer, N., Bradford, M.A., 2011. The effect of resource history
584 on the functioning of soil microbial communities is maintained across time.
585 *Biogeosciences* 8, 1477–1486. <https://doi.org/10.5194/bg-8-1477-2011>
- 586 Keiser, A.D., Keiser, D.A., Strickland, M.S., Bradford, M.A., 2014. Disentangling the
587 mechanisms underlying functional differences among decomposer
588 communities. *Journal of Ecology* 102, 603–609. [https://doi.org/10.1111/1365-
589 2745.12220](https://doi.org/10.1111/1365-2745.12220)
- 590 Kjøller, A.H., Struwe, S., 2002. Fungal communities, succession, enzymes and decomposition.
591 In: Burns R.G., Dick R., (Eds.) *Enzymes in the environment: activity, ecology and
592 applications*. New York, Marcel Dekker, 267–284.
- 593 Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glöckner F.O., 2012.
594 Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-
595 generation sequencing-based diversity studies. *Nucleic Acids Research* 41, e1.
596 <https://doi.org/10.1093/nar/gks808PMID:22933715>
- 597 Kuramae, E.E., Hillekens, R.H.E., de Hollander, M., van der Heijden, M.G.A., van den Berg, M.,
598 van Straalen, N.M., Kowalchuk, G.A., 2013. Structural and functional variation in soil
599 fungal communities associated with litter bags containing maize leaf. *FEMS
600 Microbiology Ecology* 84, 519–531.
- 601 Lange, M., Eisenhauer, N., Sierra, C.A., Bessler, H., Engels, C., Griffiths, R.I., Mellado-Vázquez,
602 P.G., Malik, A.A., Roy, J., Scheu, S., Steinbeiss, S., Thomson, B.C., Trumbore, S.E.,

- 603 Gleixner, G., 2015. Plant diversity increases soil microbial activity and soil carbón
604 storage. *Nature Communication* 6, 6707.
- 605 Ihrmark, K., Bödeker, I.T.M., Cruz-Martinez, K., Friberg, H., Kubartova, A., Schenck, J., Strid,
606 Y., Stenlid, J., Brandström-Durling, M., Clemmensen, K.E., Lindahl, B.D., 2012. New
607 primers to amplify the fungal ITS2 region – evaluation by 454-sequencing of artificial
608 and natural communities. *FEMS Microbiology Ecology* 82, 666–677.
- 609
- 610 Lin Pedersen, T., 2018. *ggraph: An Implementation of Grammar of Graphics for Graphs and*
611 *Networks. R package version. 1.0.2.* <https://CRAN.R-project.org/package=ggraph>
612
- 613 Liu, J., Sui, Y., Yu, Z., Shi, Y., Chu, H., Jin, J., Liu, X., Wang, G., 2015. Soil carbon content drives
614 the biogeographical distribution of fungal communities in the black soil zone of
615 northeast China. *Soil Biology Biochemistry* 83, 29–39.
- 616 Lundberg, D.S., Lebeis, S.L., Paredes, S.H., Yourstone, S., Gehring, J., Malfatti, S., Tremblay, J.,
617 Engelbrekton, A., Kunin, V., Glavina del Rio, T., Edgar, R.C., Eickhorst, T., Ley, R.E.,
618 Hugenholtz, P., Green Tringe, S., Dangl, J.L., 2012. Defining the core *Arabidopsis*
619 *thaliana* root microbiome. *Nature* 488, 86–93.
- 620 Lynch, J.M., Whipps, J.M., 1990. Substrate flow in the rhizosphere. *Plant and Soil* 129, 1-10.
- 621 Magoc, T., Salzberg, S.L., 2011. FLASH: Fast Length Adjustment of Short reads to improve
622 genome assemblies. *Bioinformatics* 1–8.
- 623 Mahé, F., Rognes, T., Quince, C., de Vargas, C., Dunthorn, M., 2014. Swarm: robust and fast
624 clustering method for amplicon-based studies. *Peer Journal* 2, 593.
625 <https://doi.org/10.7717/peerj.593> PMID:25276506
- 626 Mahoney, A.K., Yin C., Hulbert, S.H., 2017. Community structure, species variation, and
627 potential functions of rhizosphere-associated bacteria of different winter wheat
628 (*Triticum aestivum*) cultivars. *Frontiers in Plant Science* 8, 132.
- 629 Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing
630 reads. *EMBnet.Journal* 2226-6089.
631 <http://journal.embnet.org/index.php/embnetjournal/article/view/200>. Date
632 accessed: 27 apr. 2019. doi:<https://doi.org/10.14806/ej.17.1.200>.
- 633 McMurdie, P.J., Holmes, S., 2013. Phyloseq: An R package for reproducible interactive
634 analysis and graphics of microbiome census data. *PLoS ONE* 8, e61217.
- 635
- 636 Mendes, R., Kruijt, M., de Bruijn, I., Dekkers, E., van der Voort, M., Schneider, J.H.M., Piceno,
637 Y.M., DeSantis, T.Z., Andersen, G.L., Bakker, P.A.H.M., Raaijmakers, J.M., 2011.
638 Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science*
639 332, 1097–1100.
- 640
- 641 Mevik, B.H., Wehrens, R., Liland, K.H., 2018. *pls: Partial Least Squares and Principal*
642 *Component Regression. 2.7-0.* <https://CRAN.R-project.org/package=pls>
643

- 644 Micallef, S.A., Shiaris, M.P., Colon-Carmona, A., 2009. Influence of *Arabidopsis thaliana*
645 accessions on rhizobacterial communities and natural variation in root exudates.
646 Journal of Experimental Botany 60, 1729–1742.
- 647
648 Monard, C., Gantner, S., Stenlid, J., 2013. Utilizing ITS1 and ITS2 to study environmental
649 fungal diversity using pyrosequencing. FEMS Microbiology Ecology 84, 165-175.
650
- 651 Moya, P., Girotti, J.R., Toledo, A.V., Sisterna, M.N., 2018. Antifungal activity of *Trichoderma*
652 VOCs against *Pyrenophora teres*, the causal agent of barley net blotch. Journal of
653 Plant Protection Research 58, 45–53.
654
- 655 Müller, D.B., Vogel, C., Bai, Y., Vorholt, J.A., 2016. The plant microbiota: systems-level insights
656 and perspectives. Annual Review of Genetics 50, 211–234.
- 657 Muzer, G., de Waal, E.C., Uitterlinden, A.G., 1993. Profiling of complex microbial
658 populations by denaturing gradient gel electrophoresis analysis of polymerase chain
659 reaction-amplified genes coding for 16S rRNA. Applied Environmental Microbiology
660 59, 695–700.
661
- 662 Naz, I., Mirza, M.S., Bano, A., 2014. Molecular characterization of rhizosphere bacterial
663 communities associated with wheat (*Triticum aestivum* L.) cultivars at flowering
664 stage. Journal of Animal and Plant Science 24, 1123–1134.
- 665 Nicolaisen, M.H., Bælum, J., Jacobsen, C.S., Sørensen, j., 2008. Transcription dynamics of the
666 functional *tfdA* gene during MCPA herbicide degradation by *Cupriavidus_necator*
667 AEO106 (pRO101) in agricultural soil. Environmental Microbiology 10, 571-579.
- 668 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P.R.,
669 O’Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H.,
670 2019. Vegan: Community Ecology Package. *R package version. 2*, 5-4.
671 <https://CRAN.R-project.org/package=vegan>
672
- 673 Olsen, S., Cole, C.V., Watanabe, F.S., Dean, L.A., 1954. Estimation of available phosphorus in
674 soils by extraction with sodium bicarbonate. USDA Circular Nr 939, US Gov.Print.
675 Office, Washington, D.C
- 676 Pagé, A.P., Tremblay, J., Masson, L., Greer, C.W., 2019. Nitrogen- and phosphorus-starved
677 *Triticum aestivum* show distinct belowground microbiome profiles. PLoS One 14,
678 e0210538.
- 679 Partey, S.T., Preziosi, R.F., Robson, G.D., 2014. Improving maize residue use in soil fertility
680 restoration by mixing with residues of low C-to-N ratio: effects on C and N
681 mineralization and soil microbial biomass. Journal of Soil Science and Plant Nutrition
682 14, 518-531.
- 683 Pascault N., Ranjard, L., Kaisermann, A., Bachar, D., Christen, R., Terrat, S., Mathieu, O.,
684 Lévêque, J., Mougél, C., Henault, C., Lemanceau, P., Péan, M., Boiry, S., Fontaine, S.,

- 685 Maron, P-A., 2013. Stimulation of different functional group of bacteria by various
686 plant residues as a driver of soil priming effect. *Ecosystems* 16, 810–822.
- 687 Perkins, L.B., Johnson, D.W., Nowak, R.S., 2011. Plant-induced changes in soil nutrient
688 dynamics by native and invasive grass species. *Plant and Soil* 345, 365–374.
- 689 Richardson, A.E., Barea, J-M, McNeill, A.M., Prigent-Combaret, C., 2009. Acquisition of
690 phosphorus and nitrogen in the rhizosphere and plant growth promotion by
691 microorganisms. *Plant and Soil* 321, 305–339.
- 692 Romanowicz, K.J., Freedman, Z.B., Upchurch, R.A., Argiroff, W.A., Zak, D.R., 2016. Active
693 microorganisms in forest soils differ from the total community yet are shaped by the
694 same environmental factors: the influence of pH and soil moisture. *FEMS
695 Microbiology Ecology* 92, fiw149.
- 696 Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open
697 source tool for metagenomics. *Peer Journal* 4, 2584. doi: 10.7717/peerj.2584
- 698 Rudrappa, T., Czymmek, K.J., Paré, P.W., Bais, H.P., 2008. Root-secreted malic acid recruits
699 beneficial soil bacteria. *Plant Physiology* 148, 1547–1556. doi:10.1104/pp.108.127613
- 700 Strehmel, N., Böttcher, C., Schmidt, S., Scheel, D., 2014. Profiling of secondary metabolites in
701 root exudates of *Arabidopsis thaliana*. *Phytochemistry* 108, 35–46
- 702 Suarez, C., Ratering, S., Kramer, I., Schnell, S., 2014. *Cellvibrio diazotrophicus* sp. Nov., a
703 nitrogen-fixing bacteria isolated from the rhizosphere of salt meadow plants and
704 emended description of the genus *Cellvibrio*. *International Journal of Systematic and
705 Evolutionary Microbiology* 64, 481–486
- 706 Sun, Q., Hu, Y., Wang, R., Guo, S., Yao, L., Duan, P., 2018. Spatial distribution of microbial
707 community composition along a steep slope plot of the Loess Plateau. *Applied Soil
708 Ecology* 130, 226-236.
- 709 Thomma, B.P.H.J., 2003. *Alternaria* spp.: from a general saprophyte to specific parasite.
710 *Molecular Plant Pathology* 4, 225–236.
- 711 Tkacz, A., Cheema, J., Chandra, G., Grant, A., Poole, P.S., 2015. Stability and succession of the
712 rhizosphere microbiota depends upon plant type and soil composition. *ISME Journal*
713 9, 2349–2359. doi: 10.1038/ismej.2015.41
- 714 Torsvik, V., Øvreås, L., 2002. Microbial diversity and function in soil: from genes to
715 ecosystems. *Current Opinion in Microbiology* 5, 240-245.
- 716 Vance, E.D., Brookes, P.C., Jenkinson, D.S., 1987. An extraction method for measuring soil
717 microbial biomass C. *Soil Biology Biochemistry* 19, 703–707.
718 [https://doi.org/10.1016/0038-0717\(87\)90052-6](https://doi.org/10.1016/0038-0717(87)90052-6)
- 719 Van der Putten, W.H., Bardgett, R.D., Bever, J.D., Bezemer, T.M., Casper, B.B., Fukami, T.,
720 Kardol, P., Klironomonos, J.N., Kulmatiski, A., Schweitzer, J.A., Suding, K.N., Van de

- 721 Voorde, T.F.J., Wardle, D.A., 2013. Plant–soil feedbacks: the past, the present and
722 future challenges. *Journal of Ecology* 101, 265–276.
- 723 Van der Putten, W.H., Bradford, M.A., Brinkman, E.P., van de Voorde, T.F.J., Veen, G.F., 2016.
724 Where, when and how plant-soil feedback matters in a changing world. *Functional*
725 *Ecology* 30, 1109–1121.
- 726 Veen, G.F., Freschet, G.T., Ordonez, A., Wardle, D.A., 2015. Litter quality and environmental
727 controls of home-field advantage effects on litter decomposition. *Oikos* 124, 187-195.
728 <http://dx.doi.org/10.1111/oik.01374>
- 729 Veen, G.F., Keiser, A.D., van der Putten, W.H., Wardle, D.A., 2018. Variation in home-field
730 advantage and ability in leaf litter decomposition across successional gradients.
731 *Functional Ecology* 32, 1563–1574.
- 732 Verastegui, Y., Cheng, J., Engel, K., Kolczynski, D., Mortimer, S., Lavigne, J., Montalibet, J.,
733 Romantsov, T., Hall, M., McConkey, B.J., Rose, D.R., Tomashek, J.J., Scott, B.R.,
734 Charles, T.C., Neufeld, J.D., 2014. Multisubstrate isotope labeling and metagenomic
735 analysis of active soil bacterial communities. *mBio* 5, e01157– 14.
- 736 Větrovský, T., Baldrian, P., 2013. The Variability of the 16S rRNA Gene in Bacterial Genomes
737 and Its Consequences for Bacterial Community Analyses. *PLoS ONE* 8(2), e5792.
- 738 White, T.J., Bruns, T., Lee, S., Taylor, J.W., 1990. Amplification and direct sequencing of
739 fungal ribosomal RNA Genes for phylogenetics. *PCR Protocols: a guide to methods*
740 *and applications*. Academic Press Incorporation, 315–322.
- 741
- 742 Wickham, H., 2017. tidyverse: Easily Install and Load the 'Tidyverse'. R package version 1.2.1.
743 <https://CRAN.R-project.org/package=tidyverse>
- 744
- 745 Yadav, A.N., Verma, P., Kumar, S., Kumar, V., Kumar, M., Sugitha, T.C.K., Singh, B.P., Saxena,
746 A.K., Dhaliwal, H.S., 2018. Actinobacteria from rhizosphere: molecular diversity,
747 distributions and potential biotechnological applications. *New and Future*
748 *Developments in Microbial Biotechnology Bioengineering*, 13-41.
- 749 Zak, D.R., Holmes, W.E., White, D.C., Peacock, A.D., Tilman, D., 2003. Plant diversity, soil
750 microbial communities, and ecosystem function: Are there any links? *Ecology* 84,
751 2042–2050
- 752 Zuo, S., Li, X., Ma, Y., Yang, S., 2014. Soil microbes are linked to the allelopathic potential of
753 different wheat genotypes. *Plant and Soil* 378, 49–58. doi: 10.1007/s11104-013-
754 2020-6

755

756 Figure legends

757 **Figure 1: Carbon, nitrogen and Olsen phosphorus content in the original soil and at the end**
758 **of the plant-growing experiment (*A. thaliana*, *T. aestivum* and bare soil).** The different
759 letters indicate significant differences between treatments according to ANOVA and Tukey's

760 test ($n=4$, $p<0.05$). The red dot in each box indicates the mean value. MBC: microbial
761 biomass carbon, MBN: microbial biomass nitrogen.

762

763 **Figure 2: Microbial community abundances and diversity in the original soil and after**
764 **terminating the plant-soil experiment (*A. thaliana*, *T. aestivum* and bare soil).** The
765 abundance and diversity (Shannon diversity index) of the total bacteria and fungi were
766 obtained from DNA extracts and the abundance and diversity of active bacteria and fungi
767 from RNA extracts. The different letters indicate significant differences between treatments
768 according to ANOVA and Tukey's tests ($n=4$, $p<0.05$).

769 **Figure 3: Legacy effects of previously cultivated *A. thaliana* and *T. aestivum* in soil without**
770 **or with litter addition on composition of the active bacterial and fungal communities.** The
771 composition of the total bacterial and fungal communities present just after the plant-
772 growth phase (63 days) is represented on the left part of the figure.

773 **Figure 4: Structure and composition of the total and active bacterial and fungal**
774 **communities in unplanted (bare soil) or planted soil with *A. thaliana* or *T. aestivum* after**
775 **63 days of plant-growth.** A, C, E, G – Powered partial least squares discriminant analysis
776 (PPLS-DA) describing the total and active bacterial and fungal community structures at the
777 OTU level after plant-growth. B, D, F, H - Ternary plots describing the distribution of the
778 individual bacterial and fungal DNA-based or RNA-based OTUs between the soil treatments
779 after plant-growth (grey circles). Color code: Enriched (DNA based) or activated (RNA based)
780 OTUs by *A. thaliana* (dark green circles) or by *T. aestivum* (blue circles), depleted (DNA
781 based) or inhibited (RNA based) OTUs by *A. thaliana* (light green) or by *T. aestivum* (light
782 blue), enriched or activated OTUs by both plants (orange) and depleted or inhibited by both
783 plants (yellow). The numbers below *A. thaliana* or *T. aestivum* corresponded to the OTUs
784 that were only present in these soils. The inserted boxes zoom on the most abundant OTUs.


785 **Figure 5: Daily and cumulative C-CO₂ emissions at different sampling times in different**
786 **soils without or with litter addition.** Different letters indicate significant differences at each
787 day between treatments according to ANOVA and Tukey's test ($n=4$, $p<0.05$). Bare soil: not
788 planted soil; *A. thaliana*: soil previously planted with *A. thaliana*; *T. aestivum*: soil previously
789 planted with *T. aestivum*; Bare soil-litter: not planted soil with plant litter addition;
790 *A.thaliana*-litter: soil previously planted with *A. thaliana* and with plant litter addition; *T.*
791 *aestivum*: soil previously planted with *T. aestivum* and with plant litter addition

792


Table 1: Physical, chemical and microbiological properties of the soil used for plant-soil experiment (phase 1).
(Means and standard errors, n=4).

Physical and chemical	Data (mean, n=4)	Microbiological properties	Data (mean, n=4)
% Clay	12.9	MBC ($\mu\text{g g dry soil}^{-1}$)	221 \pm 13
% silt	40.2	MBN ($\mu\text{g g dry soil}^{-1}$)	17 \pm 2
% sand	46.8	Total fungi TS (copy g dry soil ⁻¹)	2.29E+07 \pm 1.15E+07
% organic matter	4.9	Total bacteria 16S (copy g dry soil ⁻¹)	1.12E+09 \pm 5E+08
Tot orgC ($\text{mg g dry soil}^{-1}$)	33 \pm 1	DNA ($\text{ng g dry soil}^{-1}$)	21323 \pm 8299
tot N ($\text{g kg dry soil}^{-1}$)	3 \pm 0.12	Active fungi (ITS (copy g dry soil ⁻¹))	9.80E+07 \pm 7.81E+06
Diss tot N ($\mu\text{g g dry soil}^{-1}$)	20 \pm 23	Active bacteria 16S (copy g dry soil ⁻¹)	1.66E+12 \pm 3.84E+10
Diss NO ₃ ⁻ ($\mu\text{g g dry soil}^{-1}$)	5 \pm 2	RNA ($\text{ng g dry soil}^{-1}$)	13371 \pm 1153
Diss NH ₄ ⁺ ($\mu\text{g g dry soil}^{-1}$)	4 \pm 0.2		
Diss orgC ($\mu\text{g g dry soil}^{-1}$)	85 \pm 4		
Olsen PO ₄ ($\text{mg g dry soil}^{-1}$)	69 \pm 6		
P ($\text{mg kg dry soil}^{-1}$)	212		
Ca ($\text{mg kg dry soil}^{-1}$)	1159		
Mg ($\text{mg kg dry soil}^{-1}$)	182		
K ($\text{mg kg dry soil}^{-1}$)	392		
pH _{water}	5.5		
CEC	10.8		





Journal Pre-proof




Abundances







Diversity



Abundances




Diversity




Bacterial phyla

- Chloroflexi
- Acidobacteria
- Actinobacteria
- Firmicutes
- Alphaproteobacteria
- Gammaproteobacteria
- Gemmatimonadetes
- Deltaproteobacteria
- Planctomycetes




Fungal phyla


- Ascomycota
- Zygomycota
- Basidiomycota
- Unknown




Bacteria




Fungi



Without litter



With litter



- *A. thaliana* and *T. aestivum* differently shaped the soil microbial communities
- *A. thaliana* had a stronger effect, especially for fungal communities
- PGPB were recruited in the vicinity of the roots of both *A. thaliana* and *T. aestivum*
- Plant legacy effects traduced in changes in soil organic carbon and nutrient dynamics.
- Plant legacy effects linked to nutrient dynamics rather than to microbial community

Journal Pre-proof

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof