



## Chemical cues of identity and reproductive status in Japanese macaques

Lucie Rigai, Stefano Vaglio, Joanna Setchell, Naoko Suda-Hashimoto, Takeshi Furuichi, Cécile Garcia

### ► To cite this version:

Lucie Rigai, Stefano Vaglio, Joanna Setchell, Naoko Suda-Hashimoto, Takeshi Furuichi, et al.. Chemical cues of identity and reproductive status in Japanese macaques. American Journal of Primatology, 2022, 84 (8), 10.1002/ajp.23411 . hal-03716316

**HAL Id: hal-03716316**

**<https://univ-rennes.hal.science/hal-03716316>**

Submitted on 7 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Chemical cues of identity and reproductive status in Japanese macaques**

Lucie Rigaill<sup>1,2</sup>, Stefano Vaglio<sup>3</sup>, Joanna M. Setchell<sup>4</sup>, Naoko Suda-Hashimoto<sup>2</sup>, Takeshi Furuichi<sup>2</sup>, Cécile Garcia<sup>5</sup>

<sup>1</sup> EthoS (Éthologie animale et humaine), CNRS, Université de Rennes 1, Normandie Université, Rennes, France

<sup>2</sup> Center for the Evolutionary Origins of Human Behavior, Kyoto University, Inuyama, Japan

<sup>3</sup> School of Sciences, University of Wolverhampton, UK

<sup>4</sup> Department of Anthropology & Behaviour, Ecology and Evolution Research Centre, Durham University, Durham, UK

<sup>5</sup> Eco-anthropologie (EA), Muséum national d'Histoire naturelle, CNRS, Université Paris Cité, Paris, France

Short title: Chemical signaling in Japanese macaques

Corresponding author: Lucie Rigaill, EthoS - Ethologie animale et humaine, Campus de Beaulieu - Bâtiment 25, 263, 35042 Rennes Cedex, France. [lucie.rigaill@gmail.com](mailto:lucie.rigaill@gmail.com)

**Abstract**

Olfactory communication plays an important role in the regulation of socio-sexual interactions in mammals. There is growing evidence that both human and non-human primates rely on odors to inform their mating decisions. Nevertheless, studies of primate chemical ecology remain scarce due to the difficulty of obtaining and analyzing samples. We analyzed 67 urine samples from 5 captive female Japanese macaques (*Macaca fuscata*) and 30 vaginal swabs from 3 of these females using gas chromatography–mass spectrometry and examined the relationship between odor (compounds identified, richness, intensity, and diversity) and female identity as well as cycle phase. We found a total of 36 urine compounds of which we identified 31, and 68 vaginal compounds of which we identified 37. Our results suggest that urine and vaginal odor varied more between individuals than within cycle phases. However, we found that within a female cycle, urine samples from similar phases may cluster more than samples from different phases. Our results suggest that female odor may encode information about identity (vaginal and urine odor) and reproductive status (urine odor). The question of how conspecifics use female urine and vaginal odor remains open and could be tested using bioassays. Our results and their interpretation are constrained by our limited sample size and our study design. Nonetheless, our study provides insight into the potential signaling role of female odor in sexual communication in Japanese macaques and contributes to our understanding of how odors may influence mating strategies in primates.

Keywords: olfactory communication, gas chromatography–mass spectrometry, sexual signaling, urine and vaginal odor, *Macaca fuscata*

## Introduction

Communication plays a crucial role in mating decisions. Individuals are expected to choose or avoid potential mates based on the costs (e.g., disease transmission, energy investment, sexual competition, parental investment) and benefits (e.g., good genes, social support, increased fitness) associated with mating (Beltran-Bech & Richard, 2014; Møller et al., 1999; Thrall et al., 2000; Trivers, 1972). As a result, individuals exchange information about their attributes, including social rank, genotype, health, and fertility (e.g., sexual maturity, reproductive status) (Candolin, 2003; Snowdon, 2004), directly and indirectly benefitting both emitters and receivers. Emitters may attract potential mates and enhance intra-sexual competition, thus simultaneously increasing their mating opportunities and chances of producing attractive and strong offspring (*sexy-sons/daughters* and *good genes* hypotheses R. A. Fisher, 1915; Hamilton & Zuk, 1982). Meanwhile, receivers may balance the cost of mating by focusing their mating effort on mates that are fertile and capable of dealing with the costs of reproduction and by using information about their rivals' state to adjust their mating strategies to the level of competition.

Like many other animals, primates communicate through multiple sensory channels. Females display multiple traits that are attractive to males and correlate with their fixed (e.g., social rank, age) or variable (e.g., reproductive status) attributes (Tables 1-2). Most studies of these traits have focused on female visual traits such as sexual swellings (enlargement of the anogenital area) and skin coloration in relation to fertility, probably because the sense of vision is particularly well developed in primates (Jacobs, 2008; Osorio & Vorobyev, 2005). Females can also inform conspecifics about their fertility through auditory (vocalizations or voice), behavioral (proceptive behaviors), and olfactory (odors) traits (Table 1).

**Table 1.** Examples of the potential ovulatory signaling function of female behavioral, visual, auditory, and olfactory traits in primates.

Species	Female trait	Reproductive status	Evidence
common marmosets, <i>Callithrix jacchus</i>	Anogenital odor	Correlates with the fertile phase.	Encoded information and receivers' response (Kücklich et al., 2019)
pygmy marmosets, <i>Cebuella pygmaea</i>	Anogenital gland or secretion odor	Correlates with the fertile phase.	Receivers' response (Converse et al., 1995)
tufted capuchins, <i>Cebus apella</i>	Proceptive behaviors	Correlate with the fertile phase.	Encoded information (Carosi & Visalberghi, 2002)
crested macaques, <i>Macaca nigra</i>	Sexual swelling (size)	Correlates with the fertile phase and conception <sup>1</sup> and swellings are not expressed during pregnancy.	Encoded information (Higham et al., 2012)
	Proceptive behaviors	Correlate with the fertile phase and conception.	
	Copulation calls	Correlate with the fertile phase.	
Barbary macaques, <i>Macaca sylvanus</i>	Sexual swelling (size)	Correlates with the fertile phase and swellings are expressed during early pregnancy.	Encoded information (Brauch et al., 2007; Möhle et al., 2005; Young et al., 2013)
	Proceptive behaviors	Not clear.	Encoded information (Brauch et al., 2007; Young et al., 2013)
	Copulation calls	Do not correlate with the fertile phase.	Encoded information (Pfefferle et al., 2008, 2011; Semple & McComb, 2000) Receivers' response (Semple & McComb, 2000)
stumptailed macaques, <i>Macaca arctoides</i>	Vaginal secretions	May indicate the periovulatory period and stimulate male sexual arousal	Receivers' response (Cerdeña-Molina et al., 2006)

long-tailed macaques, <i>Macaca fascicularis</i>	Sexual swelling (size)	Does not correlate with the fertile phase and swellings are expressed during early pregnancy.	Encoded information (Engelhardt et al., 2005, 2007)
	Proceptive behaviors	Correlate with the fertile phase and are expressed during the early pregnancy.	
	Copulation calls	Do not correlate with the fertile phase.	Encoded information (Engelhardt et al., 2012)
Japanese macaques, <i>Macaca fuscata</i>	Red skin coloration (face and hindquarters)	Correlates with the pregnancy period (face coloration) but not with the fertile phase or conception.	Encoded information (Rigaill et al., 2015; Rigaill, MacIntosh, et al., 2017; Rigaill et al., 2019) Receivers' response (Rigaill & Garcia, 2021)
	Proceptive behaviors	Correlate with the pregnancy period but not with the fertile phase.	Encoded information (Garcia et al., 2009; O'Neill et al., 2004; Rigaill et al., 2015)
	Copulation and 'estrus' calls	Do not correlate with the fertile phase, and may correlate with pregnancy.	
	Urine odor	May not correlate with the fertile phase.	Receivers' response (Rigaill, Suda-Hashimoto, et al., 2017)
rhesus macaques, <i>Macaca mulatta</i>	Red skin coloration (face)	Correlates with the fertile phase.	Encoded information (Dubuc et al., 2009; Higham et al., 2010) Receivers' response (Higham et al., 2011)
olive baboons, <i>Papio anubis</i>	Sexual swelling (size or shape)	Correlates with the fertile phase and pregnancy.	Encoded information (Altmann, 1970; Higham, Heistermann, et al., 2008; Higham, MacLarnon, et al., 2008; Rigaill et al., 2013)
	Red skin coloration (sexual swelling)	Correlates with pregnancy status but not with the fertile phase.	Encoded information (Higham, MacLarnon, et al., 2008; Rigaill et al., 2013)
	Proceptive behaviors	May correlate with the fertile phase.	Encoded information (Higham et al., 2009; Rigaill et al., 2013)
	Vaginal secretions	Correlate with the fertile phase	Encoded information (Vaglio et al., 2021) Receivers' response (Rigaill et al., 2013)

yellow baboons, <i>Papio cynocephalus</i>	Copulation calls	Do not correlate with the fertile phase.	Encoded information and receivers' response (Semple, 2001; Semple et al., 2002)
chacma baboons, <i>Papio ursinus</i>	Vaginal odor	Correlates with the fertile phase	Receivers' response (P. M. R. Clarke et al., 2009)
mandrills, <i>Mandrillus sphinx</i>	Sexual swelling (size or shape)	May not correlate with conception.	Encoded information (Huchard et al., 2009; Setchell & Wickings, 2004a, 2004b)
	Red skin coloration (sexual swelling)	May not correlate with conception.	Encoded information (Setchell, Charpentier, et al., 2006; Setchell & Wickings, 2004a)
	Red skin coloration (face)	Correlates with the pregnancy status, but not with the fertile phase or conception.	Encoded information (Setchell et al., 2009; Setchell, Wickings, et al., 2006)
sooty mangbeys, <i>Cercocebus torquatus atys</i>	Sexual swelling (size)	Correlates with the fertile phase but not with pregnancy.	Encoded information (Gordon et al., 1991)
howler monkeys, <i>Alouatta pigra</i>	Proceptive behaviors	Correlate with the fertile phase.	Encoded information (Van Belle et al., 2009)
bonobos, <i>Pan paniscus</i>	Sexual swelling (size)	Does not correlate with the fertile phase or pregnancy.	Encoded information (Douglas et al., 2016; Furuichi, 1987)
chimpanzees, <i>Pan troglodytes</i>	Sexual swelling (size)	Correlates with the fertile phase but not with pregnancy.	Encoded information (Deschner et al., 2003, 2004; Emery & Whitten, 2003; Wallis & Lemmon, 1986)
	Copulation calls	Do not correlate of the fertile phase.	Encoded information (Townsend et al., 2011)
	Vaginal secretions	May correlate with the fertile phase	Encoded information (Matsumoto-Oda et al., 2003)
humans, <i>Homo sapiens</i>	Red skin coloration (cheeks and lips)	Does not correlate with the fertile phase.	Encoded information (Burriss et al., 2015; Rigaill, 2020)
	Proceptive behaviors	Not clear.	Encoded information (Gangestad et al., 2002; Haselton et al., 2007)

	Voice pitch	Correlates with the fertile phase.	Encoded information and receivers' response (Pipitone & Gallup, 2008, 2012; Pisanski et al., 2018; Shoup-Knox et al., 2019)
	Body odor	Pleasantness may correlate with the period of highest fertility.	Receivers' response (Gildersleeve et al., 2012; Haselton et al., 2007; Kuukasjärvi et al., 2004; Singh & Bronstad, 2001) but see (Mei et al., 2022)

66 <sup>1</sup> difference between conceptive and non-conceptive cycles.



In comparison to visual and auditory traits, little is known about the potential role of olfactory traits in primate sexual communication. This is probably because primates were long regarded as microsmatic (having a reduced sense of olfaction, Heymann, 2006; T. D. Smith & Bhatnagar, 2004). However, there is now good evidence that odors encode information about an individual's state and may regulate primate social interactions (Table 2). Therefore, there is ample reason to suspect that odors also influence mating decisions in both human and non-human primates. Indeed, males attend more to female odors during the period of highest fertility than at other times (Table 1).

76 **Table 2.** Examples of individual characteristics encoded by odors in primates.

Species	Olfactory traits	Individual characteristics	Evidence
crowned lemurs, <i>Eulemur coronatus</i>	Anogenital odor (males: scent marks, females: secretion)	Correlates with sex.	Encoded information (Elwell et al., 2021)
ring-tailed lemurs, <i>Lemur catta</i>	Labial and scrotal secretions	Correlate with kinship, genetic compatibility and genetic quality. Are affected by health status.	Encoded information (Boulet et al., 2009, 2010; Charpentier et al., 2008; Grogan et al., 2019) Receivers' response (Charpentier et al., 2010; Grogan et al., 2019; Harris et al., 2018)
emperor tamarins, <i>Saguinus imperator</i>	Anogenital scent marks, scent gland secretions, and skin odor	Correlate with sex and differ between reproductive and non-reproductive individuals.	Encoded information (Poirier, Waterhouse, Dunn, et al., 2021; Poirier, Waterhouse, Watsa, et al., 2021)
Weddell's saddleback tamarins, <i>Leontocebus weddelli</i>	Anogenital scent marks, scent gland secretions, and skin odor	Correlate with sex and differ between reproductive and non-reproductive individuals.	Encoded information (Poirier, Waterhouse, Dunn, et al., 2021; Poirier, Waterhouse, Watsa, et al., 2021)
common marmosets, <i>Callithrix jacchus</i>	Circumgenital scent marks	Correlate with female familiarity, identity, age, and parity.	Encoded information (T. E. Smith et al., 1997; T. E. Smith, 2006; Kücklich et al., 2019) Receivers' response (Kücklich et al., 2019)
owl monkeys, <i>Aotus nancymaae</i>	Perianal gland secretions	Correlate with sex, age, and kinship.	Encoded information (Macdonald et al., 2008; Spence-Aizenberg et al., 2018)
rhesus macaques, <i>Macaca mulatta</i>	Body odor (genital area)	Correlates with familiarity or group membership.	Receivers' response (Henkel et al., 2015)
mandrills, <i>Mandrillus sphinx</i>	Sternal gland secretions and hairs	Correlate with sex, age, social rank in males, group identity, individual genetic quality, genetic compatibility, and health	Encoded information (Poirotte et al., 2017; Setchell et al., 2010, 2011; Vaglio et al., 2016) Receivers' response (Poirotte et al., 2017)

chimpanzees, <i>Pan troglodytes</i>	Urine odor	Correlates with group membership and kinship.	Receivers' response (Henkel & Setchell, 2018) <sup>77</sup>
humans, <i>Homo sapiens</i>	Axillary or body odor	Is affected by health but does not correlate with human leucocyte antigen genotype.	Receivers' response (Olsson et al., 2014; Probst et al., 2017)

Variation in the chemical composition of female odor, such as the presence or absence of compounds and their relative abundance, across and within individuals may inform conspecifics about a female's fertility. For example, sex hormones can influence the composition of female odor (humans: Michael et al., 1974; rhesus macaques: Michael & Keverne, 1970). Across primate species, the chemical composition of female odor varies between mating and non-mating seasons (ring-tailed lemurs: Scordato & Drea, 2007; Greene & Drea, 2014; and potentially Milne-Edwards' sifakas, *Propithecus edwardsi*: Morelli et al., 2013), across cycle phases (common marmosets: Kücklich et al., 2019; olive baboons: Vaglio et al., 2021; but see for chimpanzees: Fox, 1982), and between reproductive and non-reproductive individuals (tamarins: Poirier, Waterhouse, Dunn, et al., 2021; Poirier, Waterhouse, Watsa, et al., 2021; owl monkeys: Spence-Aizenberg et al., 2018). However, only one study of naturally cycling female catarrhines (Afroeurasian monkeys and apes, including humans) has yet found evidence that fine-scale intra-cycle variation in vaginal odor intensity correlates with ovulation (Vaglio et al., 2021). This limits our understanding of whether and how female odors modulate male and female mating strategies across primates.

Here, we aimed to determine the potential signaling function (i.e., encoded information) of female odor in relation to sexual communication in Japanese macaques (*Macaca fuscata*). In this seasonal species, female behaviors, vocalizations, and skin coloration vary with female reproductive status, both between cycle phases (Garcia et al., 2009; O'Neill et al., 2004; Rigaill et al., 2019), and from pre- to post-conception periods (Rigaill et al., 2015), but do not accurately signal ovulation (Rigaill & Garcia, 2021). It is not clear yet if males have access to other information about ovulation. Males biased their mating effort to the fertile phase in one study (Garcia et al., 2009) but not in others (O'Neill et al., 2004; Rigaill & Garcia, 2021). However, females exhibit a strong vaginal or urinary odor during the mating season and males frequently inspect females' genital areas by sniffing or touching them (Garcia & Rigaill,

unpublished observations of Wakasa captive group housed at Kyoto University Primate Research Institute during the mating seasons in 2011-12 and 2015-16, and of Main Group of Kojima Island, Wildlife Research Center, Kyoto University during the mating season in 2013-14). Female odor may thus vary between individuals and cycle phases, as shown in other primates (Table 1), and guide male mating behaviors.

We aimed to determine if female odor contains information about a female's timing of ovulation. Here, we describe female urine and vaginal odor in Japanese macaques, focusing on their chemical composition, complexity, intensity, and diversity, and examine how odor varies between individuals and cycles (i.e., with a female's cycle) and within cycles (i.e., between pre-fertile, fertile, and post-fertile phases). Our sample size is small (67 samples from 5 females for urine, 30 samples from 3 females for vaginal odor), due to the difficulty inherent in obtaining samples, but our study informs two hypotheses:

1. Female vaginal or urine odor conveys information about individual characteristics. If so, we predict that the chemical composition, complexity, intensity, or diversity of female urine or vaginal odor will vary across individuals. We also predict greater variation across than within individuals.
2. Female odor conveys information about the timing of ovulation. If so, we predict that the chemical composition, complexity, intensity, or diversity of vaginal and urine odor will vary between a given cycle's cycle phases, particularly between the fertile phase and the pre- and post-fertile phases.

## Methods

### Study subjects

We sampled 5 adult and naturally cycling female Japanese macaques ( $12.2 \pm \text{SD } 3.0$  years old, range = 8–16 years) housed at the Kyoto University Center for the Evolutionary Origins of

Human Behavior (EHUB, Inuyama, Japan; previously the Primate Research Institute). During the training period, females were housed in individual cages (W90 cm, D76 cm, H85 cm) in the same room (size: W4.0 m, D6.0 m, temperature: 27°C). Females were moved to a different room (individual cage size: W78 cm, D65 cm, H80 cm, temperature: 20°C, window to the external environment) for the data collection period. In both periods, females were housed with 3 adult males with whom they had visual, auditory, and olfactory communication but no physical contact. Animals were fed twice daily (monkey pellets and sweet potatoes). Water was supplied *ad libitum*.

Our research protocol was reviewed and approved by the EHUB Center for Human Evolution Modeling Research (research protocol 2014-082-10). Our methods comply with the Guidelines for the Care and Use of Nonhuman Primates of Kyoto University EHUB and with the American Society of Primatologists Principles for the Ethical Treatment of Non-Human Primates.

## **Animal training**

We used positive reinforcement (Fernström et al., 2009) to train female macaques to present their anogenital area and allow us to collect vaginal swabs for odor sampling (methods developed by NSH for EHUB). We trained females for 26 weeks from May to October 2014. Each female was trained during a 2-min session between 10.30 and 11.30 AM, 3 to 5 times a week. One trainer (NSH or LR) conducted the training session with one observer (LR, NSH, or LD). We used 1 cm<sup>3</sup> diced apples and peanuts as the primary reinforcer, and a clicker as a secondary reinforcer. We carried out training step by step following a predetermined order of defined behaviors (see supplementary material, Table S1 adapted from Fernström et al., 2009). One female failed to reach cooperation level 2 after 60 training sessions. We thus excluded her from our training protocol and replaced her with another cycling adult female. We conducted

429 training sessions in total (mean per female = 86.8 sessions, range = 28–103 sessions). Success rates varied across individuals. All females achieved anogenital presentation training level 1 but only 2 allowed vaginal sampling. These 2 females already had experience with positive reinforcement training for similar or other purposes at EHUB. For vaginal sampling, we sampled these 2 females along with a third female, who did not complete training, but spontaneously presented her hindquarters to the experimenter and accepted odor sampling during the data collection period.

### **Collection of vaginal and urine samples**

We collected data during the 2014–2015 mating season, from early November to late January (85 days in total). One experimenter (LR) collected vaginal and urine samples between 08.30 AM and 11.00 AM. We aimed to collect samples every 2 days starting from the end of the first observed menstruation, which we determined as the day when no fresh blood was observed on the female anogenital area. Mean menstrual cycle length is  $27.3 \pm \text{SD } 5.1$  days in both wild and captive populations (Enomoto et al., 1979; Fooden & Aimi, 2005; Garcia et al., 2009; Nigi, 1975; O'Neill et al., 2004). If the experimenter failed to collect a vaginal or urine sample on a designated day, they tried again the following day. The experimenter collected vaginal samples by gently rubbing a sterile cotton swab (Rikaken, Japan) 5 times on the vaginal walls (N = 3 females). The experimenter started vaginal sample collection at least 10 min after entering the experimental room and after any observed micturition. The experimenter collected environmental controls by exposing sterile swabs to the air in the room to later identify volatiles that did not derive from the sampled females. To sample urine, the experimenter placed a plastic container, covered with mesh to prevent contamination from feces, under each female's individual cage to collect urine samples (approximately 2 ml) directly after micturition (N = 5 females). Our use of plastic containers to collect urine and the lack of matching controls such

as water collected from the same plastic containers to test for any effects of using plastic containers or vials on our urine results may affect subsequent chemical analyses (Drea et al., 2013).

We stored all samples in sterile sealed glass vials at -80 °C at EHUB to prevent chemical degradation of the volatile compounds (Drea et al., 2013). We used screw-capped clear glass vials of different diameters (D) and heights (H) (19x 10 ml vials of D 18 mm X H 50 mm, 94x 20 ml vials D 25 mm X H 50 mm, 8x 50ml vials of D 30 mm X H 80 mm). We used different sizes of vial to store different types of sample. This may have affected our results because evaporation of samples depends on the available air space, and may be confounded with our predictor variables. In parallel, we kept a fraction of each urine sample at -20 °C for hormonal analyses to determine the female's reproductive status. In total, we collected 35 vaginal samples (mean per female =  $11.7 \pm \text{SD } 4.5$ , N = 3 females), 169 urine samples (mean per female =  $33.8 \pm \text{SD } 6.7$ , N = 5 females), and 45 environmental control samples over two consecutive menstrual cycles for each female.

### **Determination of ovulation periods**

We determined sex hormone concentrations in a subset of 157 urine samples selected to determine ovulation dates (2 consecutive menstrual cycles per female, 10 menstrual cycles in total, mean per female =  $31.4 \pm \text{SD } 5.9$  samples). This sample is slightly smaller than the total because we excluded samples collected on consecutive days, and those collected just before or after menstruation. We analyzed urine samples for estrone conjugates (E1C) and pregnanediol-3-glucuronide (PdG) using previously described and validated enzyme immunoassays (EIA) (Rigaill, Suda-Hashimoto, et al., 2017). We estimated that ovulation (day 0) occurred the day following a urinary E1C peak associated with a continuous raise in PdG concentrations (Fujita et al., 2001). We labelled the day directly preceding the estimated ovulation day as day -1, the



day directly following it as day +1, and so on. We defined the fertile phase as the 5-day period covering day -2 to day +2 to account for the life span of sperm in the female tract (macaques: Behboodi et al., 1991). The pre-fertile and post-fertile phases covered the 5 days preceding (day -7 to day -3) and following (day +3 to day +7) the fertile phase. Hormonal profiles showed that one of the 5 females (female 1) had abnormal hormonal patterns after her first cycle, so we excluded her second cycle from the analyses. The cycle length of the 8 cycles for which we observed menstrual bleeding was  $29.4 \pm \text{SD } 6.9$  days.

### **Chemical analyses**

We used 67 urine samples (mean per female =  $7.4 \pm \text{SD } 0.7$  samples, 2 consecutive cycles per female except for one female, Table 3) and 30 vaginal samples (mean per female =  $7.5 \pm \text{SD } 0.6$  samples, one menstrual cycle for two females and two consecutive menstrual cycles for the remaining female, Table 4) collected during the pre-fertile, fertile, and post-fertile phases for odor analyses, along with 24 corresponding environmental control samples collected on the same day as the test samples. We did not analyze the remaining 102 urine samples (mean per female =  $21.4 \pm \text{SD } 8.0$  samples) and 5 vaginal samples (5 samples from female 1, second abnormal cycle) collected outside the 15-day period of interest.

We carried out laboratory analyses of odor at the Laboratory for Analysis and Research in Environmental Chemistry - Italian National Research Council, Florence, Italy. We shipped samples via airmail, using dry ice to keep them cold ( $-30^{\circ}\text{C}$ ). We investigated the volatile compounds found in the Japanese macaque odor using solid-phase microextraction and gas chromatography-mass spectrometry, applying the same methods used in other work on primate odor signals (reviewed in Walker & Vaglio, 2021).

We introduced a  $65 \mu\text{m}$  polydimethylsiloxane/divinylbenzene solid-phase microextraction syringe needle through the vial septum and exposed the fiber to the headspace

above the sample in the vial for 15 minutes at 40°C. We analyzed the adsorbed volatile analytes using a 5975C mass spectrometer (Agilent Technologies, Santa Clara, CA) EI, 70 eV, coupled directly to a 7890B gas chromatograph (Agilent Technologies) equipped with a fused silica HP5-MS UI capillary column (Agilent Technologies) 30 m x 0.25 mm crossbonded 5%-phenyl-95%-dimethylpolysiloxane, film thickness 0.25µm. We maintained the injector temperature at 270°C and transfer line temperature at 280°C. We made injections in splitless mode (purge valve opened after 1 minute) with a constant flow of helium carrier gas of 1 ml per minute. We started the oven temperature program at 45°C for 2 minutes, then raised it by 4°C per minute to 170°C and then by 20°C per minute to 300°C.

We assessed potential contamination from the laboratory environment using blank analyses of an empty 10 ml vial (Supelco) following the same procedure used for the samples (laboratory controls). We conditioned the fiber at 260°C for 5 minutes pre-injection and 20 minutes post-injection to avoid any possible carry-over effects. We conducted these conditioning steps every day before analyzing samples.

### **Identification of compounds**

From visual inspection of chromatograms (see examples of typical chromatograms in the supplementary material, Figure S1), we detected 36 peaks from urine samples and 68 peaks from vaginal samples that were absent in both environmental and laboratory controls. We found 25 compounds that did not derive from the animals and removed these from the swab results. We standardized peak retention times using an internal standard (alpha pinene). We integrated chromatograms to obtain retention time and peak area data using ChemStation software (Agilent Technologies). We tentatively identified the eluted compounds by comparing the experimental spectra with the spectra provided by the National Institute of Standards and Technology (NIST) mass spectral database, version MSDF.01.01.2317 (Agilent Technologies).

We considered the identification valid when the minimum matching factor exceeded 80%. If more than one compound was a good match for the same chromatographic peak, we compared Kovats' retention index with values reported in the literature for the same chromatographic column type to minimize the chances of misidentification. We also checked whether more than one compound co-eluted below the same detected peak by checking the mass fragments using ChemStation software.

We determined the relative amounts of compounds by integrating the areas of the corresponding peaks in the total ion current profile and calculated percentages with respect to the total area. We retained peaks that comprised at least 0.05% of the total area of the chromatogram to avoid problems associated with unreliable quantification at very low relative amounts (i.e., to exclude the background noise), although this may mean that we missed trace chemicals (T. E. Smith et al., 2001). We analyzed all samples over a few days to minimize interassay variability.

We describe the chemical composition of the 67 urine and 30 vaginal samples according to female identity as well as cycle phases, including the percentage of samples in each category that included each compound detected for urine (N = 36 compounds) and vaginal (N = 68 compounds) samples (Tables 3-4). For the following analyses, we excluded all compounds that co-eluted as we cannot be sure about the measure of their contribution to the total ion current (Drea et al., 2013). Two urine samples only contained co-eluted compounds, this may be due to a mistake occurring at any step between data collection and chemical evaluation. We thus removed these samples from the data set. Overall, we excluded 1 of 36 (3%) urine compounds (Table 3) and 10 of 68 (15%) vaginal compounds (Table 4).

## **Data analyses**

We carried out analyses using R software (version 4.0.3). We conducted our analyses on 35 compounds detected across 65 urine and 58 compounds detected across 30 vaginal samples.

We investigated the relationship between female odor and identity as well as cycle phase. Several quantitative or qualitative analytic tools can be used to investigate chemical data. The choice of one approach over another largely depends on the nature of the data (and the methods used for chemical analyses), the species (e.g., whether chemical profiles are composed of single or multiple compounds), the number of samples, and the research question (Drea et al., 2013). Some studies assess the presence/absence of one or few compounds or categories of compounds of interest (e.g., fatty acids or alcohols), while others reduce the dimensionality of the data set (Drea et al., 2013). However, these methods may underestimate or overestimate variance in chemical composition between samples by analyzing only a subset of the data (Drea et al., 2013).

We opted for a conservative approach, keeping all compounds of the animal swab results in the analyses and avoiding subjective thresholds such as retaining compounds or variables based on *a priori* assumptions about these compounds' roles or on mathematical rather than biological reasons. We used Nonmetric Multidimensional Scaling (NMDS), a rank-based approach relying on distance or dissimilarity matrix, to represent pairwise dissimilarity between samples in a dimensional space.

We tested for differences in both urine and vaginal samples according to female identity and cycle phases. We computed a distance matrix for the data with the Bray-Curtis dissimilarity index using the `vegdist` function in the *vegan* package (Oksanen et al., 2020). We then calculated the 2-dimensional NMDS coordinates from these Bray-Curtis indices using the `metaMDS` function in the *vegan* package and plotted the outcomes (Wickham, 2016). The stress factors, which roughly represent the goodness of fit, associated with the coordinates were 0.19 for urine and 0.18 for vaginal data, and are thus considered fair (K. R. Clarke, 1993). We then

performed a permutational multivariate analysis of variance (PERMANOVA) using distance matrices by using the *Adonis* function in (*vegan* package). This analysis tests whether the centroids and dispersion of the groups tested as defined by measure space are equivalent for all groups (Anderson, 2017). Because females contributed the data for one or two consecutive menstrual cycles, we used “a female’s cycle” in our analyses as a proxy for both “menstrual cycle” and “female identity”. Here we thus tested the effect of the interaction between a female’s cycle and cycle phase (i.e., *Are samples from the same cycle phase more or less similar than samples from another cycle phase within a female given cycle?*).

To facilitate comparison with other studies of similar research questions and substrates (e.g., vaginal secretions, Vaglio et al., 2021), we also calculated three diversity indexes to graphically investigate variation in sample richness (total number of detected peaks per sample), Shannon’s H (which accounts for the relative abundance of each compound within a sample) and odor intensity (total area of detected peaks per sample). Richness and Shannon H of both urine and vaginal samples followed a normal distribution while urine and vaginal intensity followed a lognormal distribution. We explored variation across females and cycle phases graphically by plotting urine and vaginal richness, Shannon H, and intensity against either female identity (categorical variable) or cycle phase (categorical variable).

## **Data availability statement**

The data that support the findings of this study are openly available in figshare, doi:10.6084/m9.figshare.19649766.

## **Results**

### **Identity of volatile compounds**

We detected 36 volatile compounds from the analysis of 67 urine samples. We proposed a tentative identification for 31 of these compounds, including one that may result from chromatic co-elution (Table 3). We detected 68 volatile compounds from the analysis of 30 vaginal samples. We proposed a tentative identification for 37 of these vaginal compounds, with nine compounds that may result from chromatic co-elution (Table 4). Only five volatile compounds were found in both urine and vaginal samples (2-pentanone, 2-hexanone, dimethyl disulfide, toluene, 2-heptanone, Tables 3-4). This represents 5% (5/104) of all the compounds detected in urine and vaginal samples.

#### **Inter-individual variation in female odor**

Nineteen of 36 urine compounds (53%) were unique to one of the five females (Table 3). Although the two dimensions extracted from the NMDS did not clearly cluster samples per female identity (Fig. 1), the results of the PERMANOVA suggest that a female's cycle explains 25% of the variance in the data (Table 5). Urine richness, and odor intensity varied across females (Fig. 2), with lower richness values in female 2 (Fig. 2A) and higher odor intensity in female 1 (Fig. 2C). Shannon H index showed lower inter-individual variation (Fig. 2).

Twenty-eight of 68 vaginal compounds (41%) appeared to be unique to one of the three females (Table 4). The two dimensions extracted from NMDS and the results of the PERMANOVA suggest that a female's cycle explains 29% of the variance in the data and the grouping patterns of samples (Fig. 3, Table 5). Vaginal richness, Shannon's H index, and odor intensity varied across females (Fig. 2). One female (female 1) showed lower vaginal richness, Shannon H, and intensity values than the two other females sampled (Fig. 2D, E, F).

#### **Intra-cycle variation in female odor**

Ten urine compounds (28%) were only found during the fertile phase (Table 3). The cycle phase explained only 3% of the variance in urine data (Table 5). However, the interaction between a female's cycle and cycle phase explained most of the variance in the data (27%, Table 5). Thus, within a given female cycle, samples from same phases may cluster more than samples from different phases. There was little intra-cycle variation in urine richness, Shannon H, or intensity (Fig. 4). Urine richness, but not Shannon H (Fig. 4B), values tended to decrease from pre- to post-fertile phase (Fig. 4A), while urine intensity increased slightly during the fertile phase compared to pre- and post-fertile phases (Fig. 4C).

Six vaginal compounds (9%) were only found during the fertile phase (Table 4). The cycle phase or the interaction between a female's cycle and cycle phase explained respectively 6% and 17% of the variance in vaginal data (Table 5). Vaginal richness, Shannon H, or intensity did not appear to vary across the menstrual cycle (Fig. 4).

Individual trajectories are presented in the supplementary materiel (Fig. S2-3).

**Table 3.** The 36 chemical compounds retrieved from 65 urine samples, their tentative identification, their mean peak area, and their presence in all samples and in samples from the fertile phase. Data are presented for all females and for each female's menstrual cycle. Compounds marked \* co-eluted and are not included in graphical exploration of the data. <sup>1</sup> Compounds often used as lab solvents (Drea et al., 2013) but kept in the analyses for comparison with previously published data set (Delbarco-Trillo et al., 2013; Vaglio et al., 2021).

*(Table 4 is provided as a separate file as it didn't fit in the manuscript)*

**Table 4.** The 68 chemical compounds retrieved from 30 vaginal samples, their tentative identification, their mean peak area, and their presence in all samples and in samples from the

fertile phase. Data are presented for all females and for each female's menstrual cycle. Compounds marked \* co-eluted and are not included in graphical exploration of the data.<sup>1</sup> Compounds often used as lab solvents (Drea et al., 2013) but kept in the analyses for comparison with previously published data set (Vaglio et al., 2021).

(Table 4 is provided as a separate file as it didn't fit in the manuscript)

**Table 5.** PERMANOVA results based on Bray-Curtis dissimilarities using abundance data for female odors from urine and vaginal samples

	df	Sum of Squares	Mean of Squares	F	R <sup>2</sup>
<b><i>Urine samples</i></b>					
Female's cycle	8	4.88	0.61	2.62	0.25
Cycle phase	2	0.63	0.31	1.35	0.03
Interaction	16	5.39	0.34	1.45	0.27
Residuals	38	8.84	0.23		0.45
Total	64	19.74			1.00
<b><i>Vaginal samples</i></b>					
Female's cycle	3	3.15	1.05	3.59	0.29
Cycle phase	2	0.62	0.31	1.06	0.06
Interaction	6	1.90	0.32	1.08	0.17
Residuals	18	5.27	0.29		0.48
Total	29	10.94			1.00

## Discussion

We analyzed inter-individual and intra-cycle variation in urine and vaginal samples in female Japanese macaques based on identified compounds, odor richness, diversity and intensity. We identified 31 urine and 37 vaginal compounds of potential semiochemical importance in this species that may be of interest for future work. Although our limited sample size restricts the interpretation of our results, our analyses suggest that urine and vaginal odors varied more between females than between cycle phases. However, we also found that, within a female's cycle, urine samples from the same cycle phase may cluster more than samples from different phases.



We tentatively identified 31 (86%) of 36 volatile compounds found in urine samples and 37 (54%) of 68 volatile compounds detected in female vaginal secretions. Some of the volatile compounds we identified are known to play a role in plant or insect communication or metabolism (source: <https://pubchem.ncbi.nlm.nih.gov>). Some of the compounds we detected may be from the diced apples we used as a food reward during data collection (e.g., aldehydes: hexanal, nonanal, decanal, benzaldehyde; alcohols: ethanol, 1-butanol, 3-methyl-; esters: ethyl acetate, (Espino-Díaz et al., 2016)). However, it is also possible that these compounds are widespread across plant and animal species due to their role in both intra- and inter-species communication (e.g., pollination, alarm signals, mate attraction: (Das et al., 2013; Kelliher, 2007; Leonhardt et al., 2016). Indeed, 16 (51%) and 13 (35%) of the volatile compounds we identified in urine and vaginal samples are found in female odor profiles (urine, vaginal secretions, glands, hairs) of other primate species, from strepsirrhines to hominins (Tables 3-4). Despite these general similarities across species, vaginal secretions have very dissimilar odor profiles in Japanese macaques (Table 4) and olive baboons (Vaglio et al., 2021). This is consistent with the hypothesis that (dis)similarities in odor profiles across primates relate to socio-ecological and phylogenetic factors (Delbarco-Trillo et al., 2011; delBarco-Trillo & Drea, 2014; Heymann, 2006; Jänig et al., 2019; Ueno, 1994).

We found that the relative abundance of compounds across urine and vaginal samples varied across individuals in female Japanese macaques. Urine is known to play a role in the individual recognition mechanisms of several species of mammals, including some platyrrhines (mice, *Mus domesticus*: J. L. Hurst et al., 2001; common marmosets: T. E. Smith, 2006). To our knowledge, our study is the first to show that urine odor, along with vaginal odor, may convey information about identity in a catarrhine species.

Exchanging information about identity through odors may benefit Japanese macaques, a species that lives in large cohesive groups (Fooden & Aimi, 2005). As proposed by Henkel

and colleagues for rhesus macaques (Henkel et al., 2015), the ability to identify individuals – and thus potentially kin, familiar, or higher-ranking conspecifics – may be crucial to coordinate movement and regulate social interactions. Males may pay particular attention to female odor during the mating season. Dominant or central males, which have a closer access to females, could use female anogenital (vaginal and urine) odor to follow and mate-guard fertile females during group movement and to frustrate females’ attempts to escape male monopolization. Subordinate or peripheral males may also use female odor to find mating opportunities. Urine odor may be of particular interest in this latter case: since the information can be decoupled from the emitters, receivers may follow such fingerprints in the environment to find receptive and fertile females.

Our study provides some evidence that urine, but not vaginal, odor can be informative about a female’s reproductive status in Japanese macaques. This may explain why in a previous study, male Japanese macaques did not investigate (i.e., smell or taste) urine samples from the fertile phase more than samples from other phases, as they were exposed to unknown rather than familiar females’ odors (Rigaill, Suda-Hashimoto, et al., 2017). Across animals, urine is known to play a role in sexual signaling (Rothschild’s giraffes, *Giraffa camelopardalis rothschildi*: Bercovitch et al., 2006; common squirrel monkeys, *Saimiri sciureus*: Candland et al., 1980; woolly spider monkeys, *Brachyteles arachnoides*: Milton, 1985; capuchin monkeys, *Cebus apella*: Phillips et al., 2011; Asian elephants, *Elephas maximus*: Rasmussen et al., 1982; giant pandas, *Ailuropoda melanoleuca*: Swaisgood et al., 2002). Our results suggest that this communicative process through urine may also be present in catarrhines. To investigate this hypothesis further, studies should assess male response to familiar female urine odor across different cycle phases using bioassays.

The fact that vaginal odor does not appear to encode information about the fertile period is surprising considering previous results in other primate species (Table 1 and Introduction).

It is possible that our small sample size prevented us from detecting any intra-cycle variation. Moreover, variation may occur within a few days of ovulation rather than across the broader cycle phases we studied. More and finer-scaled data are needed to better understand the relationship between vaginal odor and ovulatory signaling in Japanese macaques and other primate species.

Exciting research questions remain to be tested. If female odor varies in relation to sex hormone concentrations, then does odor vary with reproductive status in the same individual: e.g., between mating and non-mating periods, from prepubescent to sexually mature, between cycling and non-cycling phases, and from pre- to post-conception phases? Do female age, genetic profile, and health status influence this relationship and how might this affect female reproductive capacity and success? And, finally, does female odor modulate socio-sexual interactions, especially in relation to the level of female-female competition, and if so, how? This last question is particularly interesting as, besides advertising their fertility, females would also benefit from receiving information about their rivals' fertility status to increase their own mating success. They may intensify competition with other fertile females through agonistic interactions (Baniel et al., 2018; A. C. Hurst et al., 2017) or by exaggerating their own attractivity (e.g., behavioral or vocal solicitations (Fallon et al., 2016)). However, whether and how female odor modulates the level of intra-sexual competition is little studied in mammals, including primates (M. L. Fisher & Burch, 2021; Stockley et al., 2013). Investigating these research questions will provide valuable information about primate olfactory communication. However, doing so will be challenging in both captive and wild populations as such studies require longitudinal observations and data collection, a considerable budget for data storage and analyses, and may conflict with population management plans (e.g., use of contraceptives in breeding programs preclude analyses of female cycles).

In conclusion, our study identifies 31 volatile urine compounds and 37 volatile vaginal compounds of possible importance in olfactory communication in Japanese macaques. We found evidence for inter-individual (urine and vaginal samples) and intra-cycle (urine samples) differences in female odors. While we cannot draw clear conclusions about the role of female odors in Japanese macaque sexual communication, our results contribute to studies investigating how olfaction mediates socio-sexual interactions in human and non-human primates. To assess whether female odor has an adaptive sexual signaling function, further work is needed to increase sampling effort and determine if males and females can perceive inter-individual and intra-cycle differences in female odors, and whether female odors modify the receiver's behavior.

## Acknowledgments

We thank the Center for Human Evolution Modeling Research of the Kyoto University Center for the Evolutionary Origins of Human Behavior (EHUB) (previously Primate Research Institute) for the use of the primate facilities. We also thank Keiko Mouri from EHUB for her help in the lab and Louise Ducroix (LD) for her help with animal training. Finally, we thank Prof. Luca Calamai and Dr Marco Michelozzi for their assistance with chemical analyses at the Laboratory for Analysis and Research in Environmental Chemistry – Italian National Research Council, Florence, Italy. We are thankful to two anonymous reviewers for their comments on earlier versions of this manuscript. This work was financially supported by the Japanese Society for the Promotion of Science (JSPS) to LR (Kakenhi 15J01067) and by the CNRS to CG (PICS N°7258 - Multimodal advertisement of reproductive status in Japanese macaques (*Macaca fuscata*)). The authors declare no conflict of interest.

## Figures

**Fig 1.** Nonmetric multidimensional scaling plots of similarity in urine sample chemical composition across cycles in relation to female identity and menstrual cycle phases.

**Fig 2.** Variation in urine (A, B, C) and vaginal (D, E, F) richness (A, D), Shannon H (B, E), and intensity (C, F) across females. Plots show the median (black horizontal line), first and third quartiles (top and bottom of the box), and the range (upper and lower whiskers) values.

**Fig 3.** Nonmetric multidimensional scaling plots of similarity in vaginal sample chemical composition in relation to female identity and menstrual cycle phases.

**Fig 4.** Variation in urine (A, B, C) and vaginal (D, E, F) richness (A, D), Shannon H (B, E), and intensity (C, F) across cycle phases. Plots show the median (black horizontal line), the first and third quartiles (top and bottom of the box), and the range (upper and lower whiskers) values.

**Fig S1.** Example of typical chromatographs from urine (A) and vaginal (B) samples during the fertile phase.

**Fig S2.** Individual variation in urine richness (A), Shannon H (B), and intensity (C) across cycle phases. Plots show the median (black horizontal line), first and third quartiles (top and bottom of the box), and the range (upper and lower whiskers).

**Fig S3.** Individual variation in vaginal richness (A), Shannon H (B), and intensity (C) across cycle phase. Plots show the median (black horizontal line), first and third quartiles (top and bottom of the box), and the range (upper and lower whiskers).

## References

Altmann, S. A. (1970). The pregnancy sign in savannah baboons. *The Journal of Zoo Animal Medicine*, 4(2), 8-12. <https://doi.org/10.2307/20094180>

- 515 Anderson, M. J. (2017). Permutational Multivariate Analysis of Variance (PERMANOVA). In  
 516 *Wiley StatsRef : Statistics Reference Online* (p. 1-15). John Wiley & Sons, Ltd.  
 517 <https://doi.org/10.1002/9781118445112.stat07841>
- 518 Baniel, A., Cowlshaw, G., & Huchard, E. (2018). Jealous females? Female competition and  
 519 reproductive suppression in a wild promiscuous primate. *Proc. R. Soc. B*, 285(1886),  
 520 20181332. <https://doi.org/10.1098/rspb.2018.1332>
- 521 Behboodi, E., Katz, D., Samuels, S., Tell, L., Hendrickx, A., & Lasley, B. (1991). The use of a  
 522 urinary estrone conjugates assay for detection of optimal mating time in the  
 523 cynomolgus macaque (*Macaca fascicularis*). *Journal of Medical Primatology*, 20(5),  
 524 229-234.
- 525 Beltran-Bech, S., & Richard, F.-J. (2014). Impact of infection on mate choice. *Animal*  
 526 *Behaviour*, 90, 159-170. <https://doi.org/10.1016/j.anbehav.2014.01.026>
- 527 Bercovitch, F. B., Bashaw, M. J., & del Castillo, S. M. (2006). Sociosexual behavior, male  
 528 mating tactics, and the reproductive cycle of giraffe *Giraffa camelopardalis*.  
 529 *Hormones and Behavior*, 50(2), 314-321.  
 530 <https://doi.org/10.1016/j.yhbeh.2006.04.004>
- 531 Boulet, M., Charpentier, M. J., & Drea, C. M. (2009). Decoding an olfactory mechanism of kin  
 532 recognition and inbreeding avoidance in a primate. *BMC Evolutionary Biology*, 9, 281.  
 533 <https://doi.org/10.1186/1471-2148-9-281>
- 534 Boulet, M., Crawford, J. C., Charpentier, M. J. E., & Drea, C. M. (2010). Honest olfactory  
 535 ornamentation in a female-dominant primate. *Journal of Evolutionary Biology*, 23(7),  
 536 1558-1563. <https://doi.org/10.1111/j.1420-9101.2010.02007.x>
- 537 Brauch, K., Pfefferle, D., Hodges, K., Möhle, U., Fischer, J., & Heistermann, M. (2007). Female  
 538 sexual behavior and sexual swelling size as potential cues for males to discern the

- 539 female fertile phase in free-ranging Barbary macaques (*Macaca sylvanus*) of  
 540 Gibraltar. *Hormones and Behavior*, 52(3), 375-383.  
 541 <https://doi.org/10.1016/j.yhbeh.2007.06.001>
- 542 Burriss, R. P., Troscianko, J., Lovell, P. G., Fulford, A. J. C., Stevens, M., Quigley, R., Payne, J.,  
 543 Saxton, T. K., & Rowland, H. M. (2015). Changes in women's facial skin color over the  
 544 ovulatory cycle are not detectable by the human visual system. *PLoS ONE*, 10(7),  
 545 e0130093. <https://doi.org/10.1371/journal.pone.0130093>
- 546 Candland, D. K., Blumer, E. S., & Mumford, M. D. (1980). Urine as a communicator in a New  
 547 World primate, *Saimiri sciureus*. *Animal Learning & Behavior*, 8(3), 468-480.  
 548 <https://doi.org/10.3758/BF03199636>
- 549 Candolin, U. (2003). The use of multiple cues in mate choice. *Biological Reviews*, 78(4),  
 550 575-595. <https://doi.org/10.1017/S1464793103006158>
- 551 Carosi, M., & Visalberghi, E. (2002). Analysis of tufted capuchin (*Cebus apella*) courtship and  
 552 sexual behavior repertoire : Changes throughout the female cycle and female  
 553 interindividual differences. *American Journal of Physical Anthropology*, 118(1), 11-24.  
 554 <https://doi.org/10.1002/ajpa.10083>
- 555 Cerda-Molina, A. L., Hernández-López, L., Rojas-Maya, S., Murcia-Mejía, C., & Mondragón-  
 556 Ceballos, R. (2006). Male-induced sociosexual behavior by vaginal secretions in  
 557 *Macaca arctoides*. *International Journal of Primatology*, 27(3), 791-807.  
 558 <https://doi.org/10.1007/s10764-006-9045-0>
- 559 Charpentier, M. J. E., Boulet, M., & Drea, C. M. (2008). Smelling right : The scent of male  
 560 lemurs advertises genetic quality and relatedness. *Molecular Ecology*, 17(14),  
 561 3225-3233. <https://doi.org/10.1111/j.1365-294X.2008.03831.x>

- 562 Charpentier, M. J. E., Crawford, J. C., Boulet, M., & Drea, C. M. (2010). Message 'scent' :  
 563 Lemurs detect the genetic relatedness and quality of conspecifics via olfactory cues.  
 564 *Animal Behaviour*, 80(1), 101-108. <https://doi.org/10.1016/j.anbehav.2010.04.005>
- 565 Clarke, K. R. (1993). Non-parametric multivariate analyses of changes in community  
 566 structure. *Australian Journal of Ecology*, 18(1), 117-143.  
 567 <https://doi.org/10.1111/j.1442-9993.1993.tb00438.x>
- 568 Clarke, P. M. R., Barrett, L., & Henzi, S. P. (2009). What role do olfactory cues play in chacma  
 569 baboon mating? *American Journal of Primatology*, 71(6), 493-502.  
 570 <https://doi.org/10.1002/ajp.20678>
- 571 Converse, L. J., Carlson, A. A., Ziegler, T. E., & Snowdon, C. T. (1995). Communication of  
 572 ovulatory state to mates by female pygmy marmosets, *Cebuella pygmaea*. *Animal*  
 573 *Behaviour*, 49(3), 615-621. [https://doi.org/10.1016/0003-3472\(95\)80194-4](https://doi.org/10.1016/0003-3472(95)80194-4)
- 574 Das, A., Lee, S.-H., Hyun, T. K., Kim, S.-W., & Kim, J.-Y. (2013). Plant volatiles as method of  
 575 communication. *Plant Biotechnology Reports*, 7(1), 9-26.  
 576 <https://doi.org/10.1007/s11816-012-0236-1>
- 577 Delbarco-Trillo, J., Burkert, B. A., Goodwin, T. E., & Drea, C. M. (2011). Night and day : The  
 578 comparative study of strepsirrhine primates reveals socioecological and phylogenetic  
 579 patterns in olfactory signals. *Journal of Evolutionary Biology*, 24(1), 82-98.  
 580 <https://doi.org/10.1111/j.1420-9101.2010.02145.x>
- 581 delBarco-Trillo, J., & Drea, C. M. (2014). Socioecological and phylogenetic patterns in the  
 582 chemical signals of strepsirrhine primates. *Animal Behaviour*, 97, 249-253.  
 583 <https://doi.org/10.1016/j.anbehav.2014.07.009>
- 584 Delbarco-Trillo, J., Harelimana, I. H., Goodwin, T. E., & Drea, C. M. (2013). Chemical  
 585 differences between voided and bladder urine in the aye-aye (*Daubentonia*



- 586 *madagascariensis*) : Implications for olfactory communication studies. *American*  
 587 *Journal of Primatology*, 75(7), 695-702. <https://doi.org/10.1002/ajp.22083>
- 588 Deschner, T., Heistermann, M., Hodges, K., & Boesch, C. (2003). Timing and probability of  
 589 ovulation in relation to sex skin swelling in wild West African chimpanzees, *Pan*  
 590 *trogodytes verus*. *Animal Behaviour*, 66(3), 551-560.  
 591 <https://doi.org/10.1006/anbe.2003.2210>
- 592 Deschner, T., Heistermann, M., Hodges, K., & Boesch, C. (2004). Female sexual swelling size,  
 593 timing of ovulation, and male behavior in wild West African chimpanzees. *Hormones*  
 594 *and Behavior*, 46(2), 204-215. <https://doi.org/10.1016/j.yhbeh.2004.03.013>
- 595 Douglas, P. H., Hohmann, G., Murtagh, R., Thiessen-Bock, R., & Deschner, T. (2016). Mixed  
 596 messages : Wild female bonobos show high variability in the timing of ovulation in  
 597 relation to sexual swelling patterns. *BMC Evolutionary Biology*, 16, 140.  
 598 <https://doi.org/10.1186/s12862-016-0691-3>
- 599 Drea, C. M., Boulet, M., Delbarco-Trillo, J., Greene, L. K., Sacha, C. R., Goodwin, T. E., &  
 600 Dubay, G. R. (2013). The « secret » in secretions : Methodological considerations in  
 601 deciphering primate olfactory communication. *American Journal of Primatology*,  
 602 75(7), 621-642. <https://doi.org/10.1002/ajp.22143>
- 603 Dubuc, C., Brent, L. J. N., Accamando, A. K., Gerald, M. S., MacLarnon, A., Semple, S.,  
 604 Heistermann, M., & Engelhardt, A. (2009). Sexual skin color contains information  
 605 about the timing of the fertile phase in free-ranging *Macaca mulatta*. *International*  
 606 *Journal of Primatology*, 30(6), 777-789. <https://doi.org/10.1007/s10764-009-9369-7>
- 607 Elwell, E. J., Walker, D., & Vaglio, S. (2021). Sexual dimorphism in crowned lemur scent-  
 608 marking. *Animals: An Open Access Journal from MDPI*, 11(7), 2091.  
 609 <https://doi.org/10.3390/ani11072091>

- 610 Emery, M. A., & Whitten, P. L. (2003). Size of sexual swellings reflects ovarian function in  
 611 chimpanzees (*Pan troglodytes*). *Behavioral Ecology and Sociobiology*, 54(4), 340-351.  
 612 <https://doi.org/10.1007/s00265-003-0648-6>
- 613 Engelhardt, A., Fischer, J., Neumann, C., Pfeifer, J.-B., & Heistermann, M. (2012). Information  
 614 content of female copulation calls in wild long-tailed macaques (*Macaca fascicularis*).  
 615 *Behavioral Ecology and Sociobiology*, 66(1), 121-134.  
 616 <https://doi.org/10.1007/s00265-011-1260-9>
- 617 Engelhardt, A., Hodges, J. K., & Heistermann, M. (2007). Post-conception mating in wild long-  
 618 tailed macaques (*Macaca fascicularis*) : Characterization, endocrine correlates and  
 619 functional significance. *Hormones and Behavior*, 51(1), 3-10.  
 620 <https://doi.org/10.1016/j.yhbeh.2006.06.009>
- 621 Engelhardt, A., Hodges, J. K., Niemitz, C., & Heistermann, M. (2005). Female sexual behavior,  
 622 but not sex skin swelling, reliably indicates the timing of the fertile phase in wild long-  
 623 tailed macaques (*Macaca fascicularis*). *Hormones and Behavior*, 47(2), 195-204.  
 624 <https://doi.org/10.1016/j.yhbeh.2004.09.007>
- 625 Enomoto, T., Seiki, K., & Haruki, Y. (1979). On the correlation between sexual behavior and  
 626 ovarian hormone level during the menstrual cycle in captive Japanese monkeys.  
 627 *Primates*, 20(4), 563-570. <https://doi.org/10.1007/BF02373438>
- 628 Espino-Díaz, M., Sepúlveda, D. R., González-Aguilar, G., & Olivas, G. I. (2016). Biochemistry of  
 629 apple aroma : A review. *Food Technology and Biotechnology*, 54(4), 375-397.  
 630 <https://doi.org/10.17113/ftb.54.04.16.4248>
- 631 Fallon, B. L., Neumann, C., Byrne, R. W., & Zuberbühler, K. (2016). Female chimpanzees  
 632 adjust copulation calls according to reproductive status and level of female

- 633 competition. *Animal Behaviour*, 113, 87-92.
- 634 <https://doi.org/10.1016/j.anbehav.2015.12.028>
- 635 Fernström, A.-L., Fredlund, H., Spångberg, M., & Westlund, K. (2009). Positive reinforcement  
636 training in rhesus macaques—Training progress as a result of training frequency.  
637 *American Journal of Primatology*, 71(5), 373-379. <https://doi.org/10.1002/ajp.20659>
- 638 Fisher, M. L., & Burch, R. L. (2021). Female-Female Competition. In T. K. Shackelford & V. A.  
639 Weekes-Shackelford (Éds.), *Encyclopedia of Evolutionary Psychological Science* (p.  
640 3043-3046). Springer International Publishing. [https://doi.org/10.1007/978-3-319-](https://doi.org/10.1007/978-3-319-19650-3_1408)  
641 [19650-3\\_1408](https://doi.org/10.1007/978-3-319-19650-3_1408)
- 642 Fisher, R. A. (1915). The evolution of sexual preference. *The Eugenics Review*, 7(3), 184-192.
- 643 Fooden, J., & Aimi, M. (2005). Systematic review of Japanese macaques, *Macaca fuscata*  
644 (Gray, 1870). *Fieldiana Zoology*, 104, 1-198.
- 645 Fox, G. J. (1982). Potentials for pheromones in chimpanzee vaginal fatty acids. *Folia*  
646 *Primatologica*, 37(3-4), 255-266. <https://doi.org/10.1159/000156036>
- 647 Fujita, S., Mitsunaga, F., Sugiura, H., & Shimizu, K. (2001). Measurement of urinary and fecal  
648 steroid metabolites during the ovarian cycle in captive and wild Japanese macaques,  
649 *Macaca fuscata*. *American Journal of Primatology*, 53(4), 167-176.  
650 <https://doi.org/10.1002/ajp.3>
- 651 Furuichi, T. (1987). Sexual swelling, receptivity, and grouping of wild pygmy chimpanzee  
652 females at Wamba, Zaïre. *Primates*, 28(3), 309-318.  
653 <https://doi.org/10.1007/BF02381014>
- 654 Gangestad, S. W., Thornhill, R., & Garver, C. E. (2002). Changes in women's sexual interests  
655 and their partners' mate-retention tactics across the menstrual cycle : Evidence for

- 656           shifting conflicts of interest. *Proceedings. Biological Sciences / The Royal Society*,  
657           269(1494), 975-982. <https://doi.org/10.1098/rspb.2001.1952>
- 658   Garcia, C., Shimizu, K., & Huffman, M. (2009). Relationship between sexual interactions and  
659           the timing of the fertile phase in captive female Japanese macaques (*Macaca*  
660           *fuscata*). *American Journal of Primatology*, 71(10), 868-879.  
661           <https://doi.org/10.1002/ajp.20717>
- 662   Gildersleeve, K. A., Haselton, M. G., Larson, C. M., & Pillsworth, E. G. (2012). Body odor  
663           attractiveness as a cue of impending ovulation in women : Evidence from a study  
664           using hormone-confirmed ovulation. *Hormones and Behavior*, 61(2), 157-166.  
665           <https://doi.org/10.1016/j.yhbeh.2011.11.005>
- 666   Gordon, T. P., Gust, D. A., Busse, C. D., & Wilson, M. E. (1991). Hormones and sexual  
667           behavior associated with postconception perineal swelling in the sooty mangabey  
668           (*Cercocebus torquatus atys*). *International Journal of Primatology*, 12(6), 585-597.  
669           <https://doi.org/10.1007/BF02547671>
- 670   Greene, L. K., & Drea, C. M. (2014). Love is in the air : Sociality and pair bondedness  
671           influence sifaka reproductive signalling. *Animal Behaviour*, 88, 147-156.  
672           <https://doi.org/10.1016/j.anbehav.2013.11.019>
- 673   Grogan, K. E., Harris, R. L., Boulet, M., & Drea, C. M. (2019). Genetic variation at MHC class II  
674           loci influences both olfactory signals and scent discrimination in ring-tailed lemurs.  
675           *BMC Evolutionary Biology*, 19(1), 171. <https://doi.org/10.1186/s12862-019-1486-0>
- 676   Hamilton, W. D., & Zuk, M. (1982). Heritable true fitness and bright birds : A role for  
677           parasites? *Science (New York, N.Y.)*, 218(4570), 384-387.

- 678 Harris, R. L., Boulet, M., Grogan, K. E., & Drea, C. M. (2018). Costs of injury for scent  
679 signalling in a strepsirrhine primate. *Scientific Reports*, 8(1), 9882.  
680 <https://doi.org/10.1038/s41598-018-27322-3>
- 681 Haselton, M. G., Mortezaie, M., Pillsworth, E. G., Bleske-Rechek, A., & Frederick, D. A. (2007).  
682 Ovulatory shifts in human female ornamentation : Near ovulation, women dress to  
683 impress. *Hormones and Behavior*, 51(1), 40-45.  
684 <https://doi.org/10.1016/j.yhbeh.2006.07.007>
- 685 Henkel, S., Lambides, A. R., Berger, A., Thomsen, R., & Widdig, A. (2015). Rhesus macaques  
686 (*Macaca mulatta*) recognize group membership via olfactory cues alone. *Behavioral*  
687 *Ecology and Sociobiology*, 69(12), 2019-2034. [https://doi.org/10.1007/s00265-015-](https://doi.org/10.1007/s00265-015-2013-y)  
688 2013-y
- 689 Henkel, S., & Setchell, J. M. (2018). Group and kin recognition via olfactory cues in  
690 chimpanzees (*Pan troglodytes*). *Proceedings of the Royal Society B: Biological*  
691 *Sciences*, 285(1889), 20181527. <https://doi.org/10.1098/rspb.2018.1527>
- 692 Heymann, E. W. (2006). The neglected sense-olfaction in primate behavior, ecology, and  
693 evolution. *American Journal of Primatology*, 68(6), 519-524.  
694 <https://doi.org/10.1002/ajp.20249>
- 695 Higham, J. P., Brent, L. J. N., Dubuc, C., Accamando, A. K., Engelhardt, A., Gerald, M. S.,  
696 Heistermann, M., & Stevens, M. (2010). Color signal information content and the eye  
697 of the beholder : A case study in the rhesus macaque. *Behavioral Ecology*, 21(4),  
698 739-746. <https://doi.org/10.1093/beheco/arq047>
- 699 Higham, J. P., Heistermann, M., Ross, C., Semple, S., & Maclarnon, A. (2008). The timing of  
700 ovulation with respect to sexual swelling detumescence in wild olive baboons.  
701 *Primates*, 49(4), 295-299. <https://doi.org/10.1007/s10329-008-0099-9>

- Higham, J. P., Heistermann, M., Saggau, C., Agil, M., Perwitasari-Farajallah, D., & Engelhardt, A. (2012). Sexual signalling in female crested macaques and the evolution of primate fertility signals. *BMC Evolutionary Biology*, 12, 89. <https://doi.org/10.1186/1471-2148-12-89>
- Higham, J. P., Hughes, K. D., Brent, L. J. N., Dubuc, C., Engelhardt, A., Heistermann, M., Maestriperi, D., Santos, L. R., & Stevens, M. (2011). Familiarity affects the assessment of female facial signals of fertility by free-ranging male rhesus macaques. *Proceedings of the Royal Society of London B: Biological Sciences*, 278(1723), 3452-3458. <https://doi.org/10.1098/rspb.2011.0052>
- Higham, J. P., MacLarnon, A. M., Ross, C., Heistermann, M., & Semple, S. (2008). Baboon sexual swellings : Information content of size and color. *Hormones and Behavior*, 53(3), 452-462. <https://doi.org/10.1016/j.yhbeh.2007.11.019>
- Higham, J. P., Semple, S., MacLarnon, A., Heistermann, M., & Ross, C. (2009). Female reproductive signaling, and male mating behavior, in the olive baboon. *Hormones and Behavior*, 55(1), 60-67. <https://doi.org/10.1016/j.yhbeh.2008.08.007>
- Huchard, E., Benavides, J. A., Setchell, J. M., Charpentier, M. J. E., Alvergne, A., King, A. J., Knapp, L. A., Cowlshaw, G., & Raymond, M. (2009). Studying shape in sexual signals : The case of primate sexual swellings. *Behavioral Ecology and Sociobiology*, 63(8), 1231-1242. <https://doi.org/10.1007/s00265-009-0748-z>
- Hurst, A. C., Alquist, J. L., & Puts, D. A. (2017). Women's fertility status alters other women's jealousy and mate guarding. *Personality and Social Psychology Bulletin*, 43(2), 191-203. <https://doi.org/10.1177/0146167216678859>
- Hurst, J. L., Payne, C. E., Nevison, C. M., Marie, A. D., Humphries, R. E., Robertson, D. H., Cavaggioni, A., & Beynon, R. J. (2001). Individual recognition in mice mediated by

- 726 major urinary proteins. *Nature*, 414(6864), 631-634.
- 727 <https://doi.org/10.1038/414631a>
- 728 Jacobs, G. H. (2008). Primate color vision : A comparative perspective. *Visual Neuroscience*,
- 729 25(5-6), 619-633. <https://doi.org/10.1017/S0952523808080760>
- 730 Jänig, S., Weiß, B. M., Birkemeyer, C., & Widdig, A. (2019). Comparative chemical analysis of
- 731 body odor in great apes. *American Journal of Primatology*, 81(6), e22976.
- 732 <https://doi.org/10.1002/ajp.22976>
- 733 Kelliher, K. R. (2007). The combined role of the main olfactory and vomeronasal systems in
- 734 social communication in mammals. *Hormones and Behavior*, 52(5), 561-570.
- 735 <https://doi.org/10.1016/j.yhbeh.2007.08.012>
- 736 Kücklich, M., Weiß, B. M., Birkemeyer, C., Einspanier, A., & Widdig, A. (2019). Chemical cues
- 737 of female fertility states in a non-human primate. *Scientific Reports*, 9(1), 1-12.
- 738 <https://doi.org/10.1038/s41598-019-50063-w>
- 739 Kuukasjärvi, S., Eriksson, C. J. P., Koskela, E., Mappes, T., Nissinen, K., & Rantala, M. J. (2004).
- 740 Attractiveness of women's body odors over the menstrual cycle : The role of oral
- 741 contraceptives and receiver sex. *Behavioral Ecology*, 15(4), 579-584.
- 742 <https://doi.org/10.1093/beheco/arh050>
- 743 Leonhardt, S. D., Menzel, F., Nehring, V., & Schmitt, T. (2016). Ecology and evolution of
- 744 communication in social insects. *Cell*, 164(6), 1277-1287.
- 745 <https://doi.org/10.1016/j.cell.2016.01.035>
- 746 Macdonald, E. A., Fernandez-Duque, E., Evans, S., & Hagey, L. R. (2008). Sex, age, and family
- 747 differences in the chemical composition of owl monkey (*Aotus nancymae*)
- 748 subcaudal scent secretions. *American Journal of Primatology*, 70(1), 12-18.
- 749 <https://doi.org/10.1002/ajp.20450>

- 750 Matsumoto-Oda, A., Oda, R., Hayashi, Y., Murakami, H., Maeda, N., Kumazaki, K., Shimizu, K.,  
 751 & Matsuzawa, T. (2003). Vaginal fatty acids produced by chimpanzees during  
 752 menstrual cycles. *Folia Primatologica*, 74(2), 75-79.
- 753 Mei, M., Grillot, R. L., Abbey, C. K., Emery Thompson, M., & Roney, J. R. (2022). Does scent  
 754 attractiveness reveal women's ovulatory timing? Evidence from signal detection  
 755 analyses and endocrine predictors of odour attractiveness. *Proceedings of the Royal*  
 756 *Society B: Biological Sciences*, 289(1970), 20220026.  
 757 <https://doi.org/10.1098/rspb.2022.0026>
- 758 Michael, R. P., Bonsall, R. W., & Warner, P. (1974). Human vaginal secretions : Volatile fatty  
 759 acid content. *Science (New York, N.Y.)*, 186(4170), 1217-1219.
- 760 Michael, R. P., & Keverne, E. B. (1970). Primate sex pheromones of vaginal origin. *Nature*,  
 761 225(5227), 84-85. <https://doi.org/10.1038/225084a0>
- 762 Milton, K. (1985). Urine washing behavior in the woolly spider monkey (*Brachyteles*  
 763 *arachnoides*). *Zeitschrift Für Tierpsychologie*, 67(1-4), 154-160.  
 764 <https://doi.org/10.1111/j.1439-0310.1985.tb01384.x>
- 765 Möhle, U., Heistermann, M., Dittami, J., Reinberg, V., Wallner, B., & Hodges, J. K. (2005).  
 766 Patterns of anogenital swelling size and their endocrine correlates during ovulatory  
 767 cycles and early pregnancy in free-ranging barbary macaques (*Macaca sylvanus*) of  
 768 Gibraltar. *American Journal of Primatology*, 66(4), 351-368.  
 769 <https://doi.org/10.1002/ajp.20161>
- 770 Møller, A. P., Christe, P., & Lux, E. (1999). Parasitism, host immune function, and sexual  
 771 selection. *The Quarterly Review of Biology*, 74(1), 3-20.



- 772 Morelli, T. L., Hayes, R. A., Nahrung, H. F., Goodwin, T. E., Harelimana, I. H., Macdonald, L. J.,  
 773 & Wright, P. C. (2013). Relatedness communicated in lemur scent. *Die*  
 774 *Naturwissenschaften*, 100(8), 769-777. <https://doi.org/10.1007/s00114-013-1074-x>
- 775 Nigi, H. (1975). Menstrual cycle and some other related aspects of Japanese monkeys  
 776 (*Macaca fuscata*). *Primates*, 16(2), 207-216. <https://doi.org/10.1007/BF02381418>
- 777 Oksanen, J., Blanchet, G. F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,  
 778 O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H.  
 779 (2020). *vegan : Community ecology package*. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)  
 780 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan)
- 781 Olsson, M. J., Lundström, J. N., Kimball, B. A., Gordon, A. R., Karshikoff, B., Hosseini, N.,  
 782 Sorjonen, K., Olgart Höglund, C., Solares, C., Soop, A., Axelsson, J., & Lekander, M.  
 783 (2014). The scent of disease : Human body odor contains an early chemosensory cue  
 784 of sickness. *Psychological Science*, 25(3), 817-823.  
 785 <https://doi.org/10.1177/0956797613515681>
- 786 O'Neill, A. C., Fedigan, L. M., & Ziegler, T. E. (2004). Relationship between ovarian cycle  
 787 phase and sexual behavior in female Japanese macaques (*Macaca fuscata*). *American*  
 788 *Journal of Physical Anthropology*, 125(4), 352-362.  
 789 <https://doi.org/10.1002/ajpa.20024>
- 790 Osorio, D., & Vorobyev, M. (2005). Photoreceptor spectral sensitivities in terrestrial animals :  
 791 Adaptations for luminance and colour vision. *Proceedings of the Royal Society of*  
 792 *London B: Biological Sciences*, 272(1574), 1745-1752.  
 793 <https://doi.org/10.1098/rspb.2005.3156>
- 794 Pfefferle, D., Brauch, K., Heistermann, M., Hodges, J. K., & Fischer, J. (2008). Female Barbary  
 795 macaque (*Macaca sylvanus*) copulation calls do not reveal the fertile phase but

- 796 influence mating outcome. *Proceedings. Biological Sciences / The Royal Society*,  
 797 275(1634), 571-578. <https://doi.org/10.1098/rspb.2007.1499>
- 798 Pfefferle, D., Heistermann, M., Pirow, R., Hodges, J. K., & Fischer, J. (2011). Estrogen and  
 799 progesterone correlates of the structure of female copulation calls in semi-free-  
 800 ranging Barbary macaques (*Macaca sylvanus*). *International Journal of Primatology*,  
 801 32(4), 992-1006. <https://doi.org/10.1007/s10764-011-9517-8>
- 802 Phillips, K. A., Buzzell, C. A., Holder, N., & Sherwood, C. C. (2011). Why do capuchin monkeys  
 803 urine wash? An experimental test of the sexual communication hypothesis using  
 804 fMRI. *American Journal of Primatology*, 73(6), 578-584.  
 805 <https://doi.org/10.1002/ajp.20931>
- 806 Pipitone, R. N., & Gallup, G. G. (2008). Women's voice attractiveness varies across the  
 807 menstrual cycle. *Evolution and Human Behavior*, 29(4), 268-274.  
 808 <https://doi.org/10.1016/j.evolhumbehav.2008.02.001>
- 809 Pipitone, R. N., & Gallup, G. G. (2012). The unique impact of menstruation on the female  
 810 voice : Implications for the evolution of menstrual cycle cues. *Ethology*, 118(3),  
 811 281-291. <https://doi.org/10.1111/j.1439-0310.2011.02010.x>
- 812 Pisanski, K., Oleszkiewicz, A., Plachetka, J., Gmiterek, M., & Reby, D. (2018). Voice pitch  
 813 modulation in human mate choice. *Proceedings of the Royal Society B: Biological*  
 814 *Sciences*, 285(1893), 20181634. <https://doi.org/10.1098/rspb.2018.1634>
- 815 Poirier, A. C., Waterhouse, J. S., Dunn, J. C., & Smith, A. C. (2021). Scent marks signal species,  
 816 sex and reproductive status in tamarins (*Saguinus spp.*, neotropical primates).  
 817 *Chemical Senses*, bjab008. <https://doi.org/10.1093/chemse/bjab008>
- 818 Poirier, A. C., Waterhouse, J. S., Watsa, M., Erkenwick, G. A., Moreira, L. A. A., Tang, J.,  
 819 Dunn, J. C., Melin, A. D., & Smith, A. C. (2021). On the trail of primate scent signals : A

- 820 field analysis of callitrichid scent-gland secretions by portable gas chromatography-  
 821 mass spectrometry. *American Journal of Primatology*, 83(3), e23236.  
 822 <https://doi.org/10.1002/ajp.23236>
- 823 Poirotte, C., Massol, F., Herbert, A., Willaume, E., Bomo, P. M., Kappeler, P. M., &  
 824 Charpentier, M. J. E. (2017). Mandrills use olfaction to socially avoid parasitized  
 825 conspecifics. *Science Advances*, 3(4), e1601721.  
 826 <https://doi.org/10.1126/sciadv.1601721>
- 827 Probst, F., Fischbacher, U., Lobmaier, J. S., Wirthmüller, U., & Knoch, D. (2017). Men's  
 828 preferences for women's body odours are not associated with human leucocyte  
 829 antigen. *Proceedings of the Royal Society B: Biological Sciences*, 284(1864),  
 830 20171830. <https://doi.org/10.1098/rspb.2017.1830>
- 831 Rasmussen, L. E., Schmidt, M. J., Henneous, R., Groves, D., & Daves, G. D. (1982). Asian bull  
 832 elephants : Flehmen-like responses to extractable components in female elephant  
 833 estrous urine. *Science*, 217(4555), 159-162. <https://doi.org/10.1126/science.7089549>
- 834 Rigaill, L. (2020). Fine-scale variation in lip and cheek colour according to the timing of  
 835 ovulation in women. *Behavioral Ecology and Sociobiology*, 74(6), 71.  
 836 <https://doi.org/10.1007/s00265-020-02851-y>
- 837 Rigaill, L., & Garcia, C. (2021). Does male mate choice select for female coloration in a  
 838 promiscuous primate species? *Animal Behaviour*, 177, 171-181.  
 839 <https://doi.org/10.1016/j.anbehav.2021.05.002>
- 840 Rigaill, L., Higham, J. P., Lee, P. C., Blin, A., & Garcia, C. (2013). Multimodal sexual signaling  
 841 and mating behavior in olive baboons (*Papio anubis*). *American Journal of*  
 842 *Primatology*, 75(7), 774-787. <https://doi.org/10.1002/ajp.22154>

- 843 Rigaill, L., Higham, J. P., Winters, S., & Garcia, C. (2019). The redder the better? Information  
 844 content of red skin coloration in female Japanese macaques. *Behavioral Ecology and*  
 845 *Sociobiology*, 73(8), 103. <https://doi.org/10.1007/s00265-019-2712-x>
- 846 Rigaill, L., MacIntosh, A. J. J., Higham, J. P., Winters, S., Shimizu, K., Mouri, K., Furuichi, T., &  
 847 Garcia, C. (2015). Multimodal advertisement of pregnancy in free-ranging female  
 848 Japanese macaques (*Macaca fuscata*). *PLoS ONE*, 10(8), e0135127.  
 849 <https://doi.org/10.1371/journal.pone.0135127>
- 850 Rigaill, L., MacIntosh, A. J. J., Higham, J. P., Winters, S., Shimizu, K., Mouri, K., Suzumura, T.,  
 851 Furuichi, T., & Garcia, C. (2017). Testing for links between face color and age,  
 852 dominance status, parity, weight, and intestinal nematode infection in a sample of  
 853 female Japanese macaques. *Primates*, 58(1), 83-91. [https://doi.org/10.1007/s10329-](https://doi.org/10.1007/s10329-016-0575-6)  
 854 [016-0575-6](https://doi.org/10.1007/s10329-016-0575-6)
- 855 Rigaill, L., Suda-Hashimoto, N., Ducroix, L., Mouri, K., Furuichi, T., & Garcia, C. (2017). Testing  
 856 for links between female urine odor and male sexual behaviors in Japanese  
 857 macaques (*Macaca fuscata*). *International Journal of Primatology*, 38(5), 823-837.  
 858 <https://doi.org/10.1007/s10764-017-9980-y>
- 859 Scordato, E. S., & Drea, C. M. (2007). Scents and sensibility : Information content of olfactory  
 860 signals in the ringtailed lemur, *Lemur catta*. *Animal Behaviour*, 73(2), 301-314.  
 861 <https://doi.org/10.1016/j.anbehav.2006.08.006>
- 862 Semple, S. (2001). Individuality and male discrimination of female copulation calls in the  
 863 yellow baboon. *Animal Behaviour*, 61(5), 1023-1028.  
 864 <https://doi.org/10.1006/anbe.2001.1692>

- 865 Semple, S., & McComb, K. (2000). Perception of female reproductive state from vocal cues in  
 866 a mammal species. *Proceedings of the Royal Society of London. Series B: Biological*  
 867 *Sciences*, 267(1444), 707-712. <https://doi.org/10.1098/rspb.2000.1060>
- 868 Semple, S., McComb, K., Alberts, S., & Altmann, J. (2002). Information content of female  
 869 copulation calls in yellow baboons. *American Journal of Primatology*, 56(1), 43-56.  
 870 <https://doi.org/10.1002/ajp.1062>
- 871 Setchell, J. M., Charpentier, M. J. E., Abbott, K. M., Wickings, E. J., & Knapp, L. A. (2009). Is  
 872 brightest best? Testing the Hamilton-Zuk hypothesis in mandrills. *International*  
 873 *Journal of Primatology*, 30(6), 825-844. <https://doi.org/10.1007/s10764-009-9371-0>
- 874 Setchell, J. M., Charpentier, M. J. E., Bedjabaga, I.-B., Reed, P., Wickings, E. J., & Knapp, L. A.  
 875 (2006). Secondary sexual characters and female quality in primates. *Behavioral*  
 876 *Ecology and Sociobiology*, 61(2), 305-315. [https://doi.org/10.1007/s00265-006-0260-](https://doi.org/10.1007/s00265-006-0260-7)  
 877 7
- 878 Setchell, J. M., Vaglio, S., Abbott, K. M., Moggi-Cecchi, J., Boscaro, F., Pieraccini, G., & Knapp,  
 879 L. A. (2011). Odour signals major histocompatibility complex genotype in an Old  
 880 World monkey. *Proceedings. Biological Sciences / The Royal Society*, 278(1703),  
 881 274-280. <https://doi.org/10.1098/rspb.2010.0571>
- 882 Setchell, J. M., Vaglio, S., Moggi-Cecchi, J., Boscaro, F., Calamai, L., & Knapp, L. A. (2010).  
 883 Chemical composition of scent-gland secretions in an old world monkey (*Mandrillus*  
 884 *sphinx*) : Influence of sex, male status, and individual identity. *Chemical Senses*, 35(3),  
 885 205-220. <https://doi.org/10.1093/chemse/bjp105>
- 886 Setchell, J. M., Wickings, E. J., & Knapp, L. A. (2006). Signal content of red facial coloration in  
 887 female mandrills (*Mandrillus sphinx*). *Proceedings of the Royal Society B: Biological*  
 888 *Sciences*, 273(1599), 2395-2400. <https://doi.org/10.1098/rspb.2006.3573>

- 889 Setchell, J. M., & Wickings, J. E. (2004a). Sexual swelling in mandrills (*Mandrillus sphinx*) : A  
 890 test of the reliable indicator hypothesis. *Behavioral Ecology*, 15(3), 438-445.  
 891 <https://doi.org/10.1093/beheco/arh027>
- 892 Setchell, J. M., & Wickings, J. E. (2004b). Social and seasonal influences on the reproductive  
 893 cycle in female mandrills (*Mandrillus sphinx*). *American Journal of Physical*  
 894 *Anthropology*, 125(1), 73-84. <https://doi.org/10.1002/ajpa.10375>
- 895 Shoup-Knox, M. L., Ostrander, G. M., Reimann, G. E., & Pipitone, R. N. (2019). Fertility-  
 896 dependent acoustic variation in women's voices previously shown to affect listener  
 897 physiology and perception. *Evolutionary Psychology*, 17(2), 1474704919843103.  
 898 <https://doi.org/10.1177/1474704919843103>
- 899 Singh, D., & Bronstad, P. M. (2001). Female body odour is a potential cue to ovulation.  
 900 *Proceedings of the Royal Society B: Biological Sciences*, 268(1469), 797-801.  
 901 <https://doi.org/10.1098/rspb.2001.1589>
- 902 Smith, T. D., & Bhatnagar, K. P. (2004). Microsmatic primates : Reconsidering how and when  
 903 size matters. *The Anatomical Record Part B: The New Anatomist*, 279B(1), 24-31.  
 904 <https://doi.org/10.1002/ar.b.20026>
- 905 Smith, T. E. (2006). Individual olfactory signatures in common marmosets (*Callithrix jacchus*).  
 906 *American Journal of Primatology*, 68(6), 585-604. <https://doi.org/10.1002/ajp.20254>
- 907 Smith, T. E., Abbott, D. H., Tomlinson, A. J., & Mlotkiewicz, J. A. (1997). Differential display of  
 908 investigative behavior permits discrimination of scent signatures from familiar and  
 909 unfamiliar socially dominant female marmoset monkeys (*Callithrix jacchus*). *Journal*  
 910 *of Chemical Ecology*, 23(11), 2523-2546.  
 911 <https://doi.org/10.1023/B:JOEC.00000006664.38169.e5>

- 912 Smith, T. E., Tomlinson, A. J., Mlotkiewicz, J. A., & Abbott, D. H. (2001). Female marmoset  
 913 monkeys (*Callithrix jacchus*) can be identified from the chemical composition of their  
 914 scent marks. *Chemical Senses*, 26(5), 449-458.  
 915 <https://doi.org/10.1093/chemse/26.5.449>
- 916 Snowdon, C. T. (2004). Sexual selection and communication. In C. P. van Schaik & P. M.  
 917 Kappeler (Éds.), *Sexual Selection in Primates : New and Comparative Perspectives* (p.  
 918 57-70). Cambridge University Press.  
 919 <https://doi.org/10.1017/CBO9780511542459.006>
- 920 Spence-Aizenberg, A., Kimball, B. A., Williams, L. E., & Fernandez-Duque, E. (2018). Chemical  
 921 composition of glandular secretions from a pair-living monogamous primate : Sex,  
 922 age, and gland differences in captive and wild owl monkeys (*Aotus spp.*). *American*  
 923 *Journal of Primatology*, 80(2), e22730. <https://doi.org/10.1002/ajp.22730>
- 924 Stockley, P., Bottell, L., & Hurst, J. L. (2013). Wake up and smell the conflict : Odour signals in  
 925 female competition. *Philosophical Transactions of the Royal Society B: Biological*  
 926 *Sciences*, 368(1631), 20130082. <https://doi.org/10.1098/rstb.2013.0082>
- 927 Swaisgood, R. R., Lindburg, D. G., & Zhang, H. (2002). Discrimination of oestrous status in  
 928 giant pandas (*Ailuropoda melanoleuca*) via chemical cues in urine. *Journal of Zoology*,  
 929 257(3), 381-386. <https://doi.org/10.1017/S0952836902000985>
- 930 Thrall, P. H., Antonovics, J., & Dobson, A. P. (2000). Sexually transmitted diseases in  
 931 polygynous mating systems : Prevalence and impact on reproductive success.  
 932 *Proceedings of the Royal Society B: Biological Sciences*, 267(1452), 1555-1563.
- 933 Townsend, S. W., Deschner, T., & Zuberbühler, K. (2011). Copulation calls in female  
 934 chimpanzees (*Pan troglodytes schweinfurthii*) convey identity but do not accurately

- 935 reflect fertility. *International Journal of Primatology*, 32(4), 914-923.
- 936 <https://doi.org/10.1007/s10764-011-9510-2>
- 937 Trivers, R. L. (1972). Parental investment and sexual selection. In *Sexual Selection and the*
- 938 *Descent of Man, 1871–1971*. B. Campbell.
- 939 Ueno, Y. (1994). Olfactory discrimination of urine odors from five species by tufted capuchin
- 940 (*Cebus apella*). *Primates*, 35(3), 311-323. <https://doi.org/10.1007/BF02382728>
- 941 Vaglio, S., Minicozzi, P., Kessler, S. E., Walker, D., & Setchell, J. M. (2021). Olfactory signals
- 942 and fertility in olive baboons. *Scientific Reports*, 11(1), 8506.
- 943 <https://doi.org/10.1038/s41598-021-87893-6>
- 944 Vaglio, S., Minicozzi, P., Romoli, R., Boscaro, F., Pieraccini, G., Moneti, G., & Moggi-Cecchi, J.
- 945 (2016). Sternal gland scent-marking signals sex, age, rank, and group identity in
- 946 captive mandrills. *Chemical Senses*, 41(2), 177-186.
- 947 <https://doi.org/10.1093/chemse/bjv077>
- 948 Van Belle, S., Estrada, A., Ziegler, T. E., & Strier, K. B. (2009). Sexual behavior across ovarian
- 949 cycles in wild black howler monkeys (*Alouatta pigra*) : Male mate guarding and
- 950 female mate choice. *American Journal of Primatology*, 71(2), 153-164.
- 951 <https://doi.org/10.1002/ajp.20635>
- 952 Walker, D., & Vaglio, S. (2021). Sampling and analysis of animal scent signals. *Journal of*
- 953 *Visualized Experiments: JoVE*, 168. <https://doi.org/10.3791/60902>
- 954 Wallis, J., & Lemmon, W. B. (1986). Social behavior and genital swelling in pregnant
- 955 chimpanzees (*Pan troglodytes*). *American Journal of Primatology*, 10(2), 171-183.
- 956 <https://doi.org/10.1002/ajp.1350100207>
- 957 Wickham, H. (2016). *ggplot2 : Elegant graphics for data analysis*. Springer-Verlag New York.
- 958 <https://ggplot2.tidyverse.org>



959 Young, C., Majolo, B., Heistermann, M., Schülke, O., & Ostner, J. (2013). Male mating  
960 behaviour in relation to female sexual swellings, socio-sexual behaviour and  
961 hormonal changes in wild Barbary macaques. *Hormones and Behavior*, 63(1), 32-39.  
962 <https://doi.org/10.1016/j.yhbeh.2012.11.004>  
963