


Supplementary data 1. PRISMA flow diagram about SAFJ. Textual research on pubmed with following keywords: (septic OR infect*) AND (osteomyelitis OR arthritis) AND (facet OR Zygapophyseal OR Zygapophysial)



From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed.1000097

For more information, visit www.prisma-statement.org.

References of the nine articles of 4 or more SAFJ [1–9] and references of SAFJ due to SAFJ due to *Mycobacterium tuberculosis* [4,10–12] :

- [1] Ross JJ, Ard KL, Carlile N. Septic Arthritis and the Opioid Epidemic: 1465 Cases of Culture-Positive Native Joint Septic Arthritis From 1990-2018. Open Forum Infect Dis 2020;7:ofaa089. <https://doi.org/10.1093/ofid/ofaa089>.
- [2] Muffoletto AJ, Ketonen LM, Mader JT, Crow WN, Hadjipavlou AG. Hematogenous pyogenic facet joint infection. Spine 2001;26:1570–6. <https://doi.org/10.1097/00007632-200107150-00014>.
- [3] Narváez J, Nolla JM, Narváez JA, Martínez-Carnicer L, De Lama E, Gómez-Vaquero C, et al. Spontaneous pyogenic facet joint infection. Semin Arthritis Rheum 2006;35:272–83. <https://doi.org/10.1016/j.semarthrit.2005.09.003>.
- [4] André V, Pot-Vaucel M, Cozic C, Visée E, Morrier M, Varin S, et al. Septic arthritis of the facet joint. Med Mal Infect 2015;45:215–21. <https://doi.org/10.1016/j.medmal.2015.04.001>.
- [5] Ergan M, Macro M, Benhamou CL, Vandermarcq P, Colin T, L'HironDEL JL, et al. Septic arthritis of lumbar facet joints. A review of six cases. Rev Rhum Engl Ed 1997;64:386–95.
- [6] Ahl T, Hedström M, von Heijne A, Hammers Stiernstedt S. Acute spinal epidural abscess without concurrent spondylodiscitis. Successful closed treatment in 10 cases. Acta Orthop Scand 1999;70:199–202. <https://doi.org/10.3109/17453679909011262>.
- [7] Doita M, Nabeshima Y, Nishida K, Fujioka H, Kurosaka M. Septic arthritis of lumbar facet joints without predisposing infection. J Spinal Disord Tech 2007;20:290–5. <https://doi.org/10.1097/01.bsd.0000211285.91271.b3>.
- [8] Michel-Batôt C, Dintinger H, Blum A, Olivier P, Laborde F, Bettembourg-Brault I, et al. A particular form of septic arthritis: septic arthritis of facet joint. Joint Bone Spine 2008;75:78–83. <https://doi.org/10.1016/j.jbspin.2007.02.006>.
- [9] Amadoru S, Lim K, Tacey M, Aboltins C. Spinal infections in older people: an analysis of demographics, presenting features, microbiology and outcomes. Intern Med J 2017;47:182–8. <https://doi.org/10.1111/imj.13300>.
- [10] Tiemtoré-Kambou BMA, Tiendrébéogo-Zabsonre J, Diallo I, Gnoumou A, Ouédraogo DD, Cissé R. Inflammatory lumbar radiculalgia in Ouagadougou: posterior facet arthritis due to tuberculosis. Med Sante Trop 2017;27:222–4. <https://doi.org/10.1684/mst.2017.0690>.
- [11] Kouki S, Landolsi M, Ben Lassoued M, Gharsallah I. Uncommon cause of cervicobrachial neuralgia: epidural abscess complicating tuberculous arthritis. BMJ Case Rep 2017;2017. <https://doi.org/10.1136/bcr-2017-219458>.
- [12] Ambulgekar RK, Iqbal Z, Masne P. A Rare Case of Thoracolumbar Epidural Abscess with Tuberculous Etiology in an Adolescent Male with Neurological Deficit. J Orthop Case Rep 2022;12:98–101. <https://doi.org/10.13107/jocr.2022.v12.i09.3336>.