



HAL
open science

First record of *Zaprionus tuberculatus* (Diptera: Drosophilidae) in mainland France

Romain Georges, Amir Yassin, Hervé Colinet

► To cite this version:

Romain Georges, Amir Yassin, Hervé Colinet. First record of *Zaprionus tuberculatus* (Diptera: Drosophilidae) in mainland France. *Entomological Research*, 2024, 54 (4), pp.e12725. 10.1111/1748-5967.12725 . hal-04541879

HAL Id: hal-04541879

<https://univ-rennes.hal.science/hal-04541879>

Submitted on 19 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



RESEARCH PAPER

First record of *Zaprionus tuberculatus* (Diptera: Drosophilidae) in mainland France

Romain GEORGES¹ , Amir YASSIN² and Hervé COLINET¹ ¹ CNRS, ECOBIO ((Ecosystèmes, Biodiversité, Évolution)) - UMR 6553, University of Rennes, Rennes, France² CNRS, IRD, Laboratoire Evolution, Génomes, Comportement, Ecologie (EGCE) - UMR 9191, Université Paris-Saclay, Gif-sur-Yvette, France

Correspondence

Romain Georges, CNRS, ECOBIO
[(Ecosystèmes, Biodiversité, Évolution)] - UMR
6553, University of Rennes, 263 Avenue du
Gal Leclerc, CS 74205, 35000 Rennes Cedex,
France.

Email: romain.georges@univ-rennes.fr

Received 20 November 2023;
accepted 18 March 2024.

doi: 10.1111/1748-5967.12725

Abstract

As most drosophilid species are not considered as primary pest, studies of their communities *in natura*, with standardized surveys, are relatively sparse. However, *Drosophila suzukii* and its rapid expansion through the world led to the implementation of many monitoring programs. As part of a research project on *D. suzukii*, we set up in 2022 an annual fly monitoring in 16 fruits plantations in France to understand population's dynamics of *D. suzukii* and to broadly survey drosophilid communities. The captures led to the first observation of *Zaprionus tuberculatus* in mainland France. Over the whole monitoring, we trapped a total of 111 specimens in a fig orchard located in southern France (Salses-le-Château), both in fig trees and nearby hedgerow. The first detection of *Zaprionus tuberculatus* occurred in July 2022 in the hedgerow, and captures continued until January 2023 with an interruption in November and December. In addition, in this orchard we collected overripe figs in September 2023 from which over 20 *Z. tuberculatus* have emerged in the following 2 weeks, confirming the establishment of a permanent population. The pest status of *Z. tuberculatus* and its potential risk for agriculture is not yet clear, but the pest behavior of the close-relative species *Zaprionus indianus*, especially on figs, should be a warning point for the entry of *Z. tuberculatus* into the EU and France, as they may have similar polyphagous habit. The pest status and range expansion of *Z. tuberculatus* should be assessed to estimate risk to fruit's production.

Key words: France, fruit fly, invasion, pest risk, *Zaprionus tuberculatus*

Introduction

Drosophilidae Rondani is a very diverse family with almost 4,700 described species distributed in 77 genera (Bächli 2023). The largest genus of Drosophilidae is *Drosophila* Fallén which comprises 1,676 described species. There is a strong heterogeneity in the existing knowledge on Drosophilidae, with a huge literature on genetics as well as cellular and developmental biology of *Drosophila* species, mainly *Drosophila melanogaster*, but little knowledge on other genera. Studies of the whole drosophilid natural communities, including other genera than *Drosophila*, with standardized surveys are relatively sparse (de Toni et al. 2007; Delbac et al. 2020; Mateus et al. 2006). A possible reason for this lack

of interest is likely the ecology and lifecycle of Drosophilidae. For most drosophilid species, larvae grow on decaying plant material and forage on associated microbial communities (Carson 1971; Markow & O'Grady 2008) and are therefore not considered as primary pest. In addition, these flies are small and cause no trouble to humans or animals, so they received relatively less attention by scientific community. However, the spotted-wing drosophila *Drosophila suzukii* (Matsumura, 1931) and its rapid expansion through the world changed the situation. The damage caused to cultivated fruits by this invasive species and the huge economic loss have led to a large number of studies in the last decade about its ecology and control measures (Tait et al. 2021). Many fly monitoring

programs have been set up in various countries to better understand and predict populations dynamics of *D. suzukii* (Briem *et al.* 2018; Hamby *et al.* 2014; Harris *et al.* 2014; Kremmer *et al.* 2017). As part of a research project on *D. suzukii* (DroFramb action within the framework of ANR Drothermal project, <https://www.drothermal.cnrs.fr/>), in 2022, we set up an annual fly monitoring in various regions of France. The main objective was to conduct a vast monitoring program of *D. suzukii* during a whole year, including during winter months, to get data on *D. suzukii* occurrence and abundance in various areas and climates. This monitoring effort also provided the opportunity to survey Drosophilidae communities in fruit plantations. Regular field monitoring is crucial to detect novel invaders and potential novel pests. Indeed, early detection is key to successful preventative strategies. Here, we will not present the whole capture data of *D. suzukii* across all the various sites of the national monitoring program. Instead, we present the data of a single site in the south of France where we document the first occurrence, and likely establishment, of a new invasive and potential pest fly, *Zaprionus tuberculatus* Malloch, 1932 (Diptera: Drosophilidae) in mainland France.

Materials and Methods

From March 2022 to February 2023, we conducted a national participative monitoring program of *D. suzukii* (Matsumura, 1931) within 14 raspberry plantations. Because there were

no raspberry growers willing to participate in the south of France, where raspberry cultivation is much less developed, we decided to expand our sampling effort to cherry and fig orchards. As a result, we monitored a total of 16 fruit plantations along a latitudinal gradient in France (Fig. 1). In each orchard, three traps were installed at least 10 m apart within the plantation at fruiting height and three others in a hedgerow or a grove nearby at approximately 1 m height which corresponds to the height at which most potential wild fruits are available for *D. suzukii* (*Rubus* spp., *Sambucus* spp., *Eleagnus* spp.). Traps were opened for 10 days at the end of each month resulting in twelve 10-days trapping sessions across seasons. Flies were captured using home-made upside down red-colored water bottle traps (320 mm height— \varnothing 79.5 mm, Badoit[®], France) pierced with six holes (\varnothing 5 mm each) in the larger part of the bottle (see Figure S1). The trap contained a liquid bait (80 mL) in an open but inaccessible plastic container positioned inside the bottle. Additionally, a drowning solution (30 mL) was present in a collection tube screwed to the neck of the bottle (see Figure - S1). For the bait, we used a mixture of 80% apple cider vinegar (5% acidity, Bouton d'Or[®], France) and 20% of cane sugar syrup (Canadou[®], France). Flies caught in the bottles fell in the drowning solution that was made of salt water (40 g.L^{-1}) with a drop of odorless soap (L'Arbre Vert[®]—Peaux Sensibles sans parfum, France). Bait and drowning solution were renewed at each trapping session. Local temperatures at each site were monitored during all the experiment in plantation and hedgerow using two TMS-4 Standard




FIGURE 1 French locality (red dot) where specimens of *Zaprionus tuberculatus* were collected. Black dots represent the other monitored localities where no *Z. tuberculatus* specimens were collected during the same period (© EuroGeographics for the administrative boundaries).

dataloggers (TOMST[®], Czech Republic) which collected data every 15 min in three different levels: 6 cm into the soil, at the surface and 15 cm above the soil surface. These data allowed gathering microclimate thermal measurements at a biologically relevant scale and allowed exploring whether the level of *D. sukuzii* captures responded to soil or surface temperature as well as air temperature, especially during winter months.

Results and Discussion

While identifying *D. sukuzii* specimens collected in different regions of France, we came across flies with white longitudinal stripes at a specific location within a fig orchard in southern France (Salses-le-Château; 42°50′0.169″N 2°55′5.447″E; see Figs. 1 and 2). Flies with these morphological features typically belong to the *Zaprionus* Coquillett, 1902 genus. According to the Köppen-Geiger classification (Geiger 1954; Kottek et al. 2006), the climate of this region is type Csa, corresponding to hot-summer Mediterranean climate with mild winters averaging above 0°C and hot summers averaging above 25°C. In this location, over the whole monitoring period in 2022, we trapped 111 specimens of *Zaprionus* sp. out of a total of 3766 drosophilids (Table 1).

Species identification was first done by R. Georges and then sent to A. Yassin for confirmation based on external morphological criteria. The genus *Zaprionus* Coquillett,

1902 exhibits characteristic white longitudinal stripes on the frons and the mesonotum (Fig. 2A,B) and we could also easily distinguish a protruding tubercle bearing a bristle on the forefemur of all our specimens (Fig. 2C). This feature is characteristic of a clade of seven Afrotropical species, known as the *Zaprionus tuberculatus* subgroup (Yassin & David 2010), in reference to the first species that has been discovered in the clade, that is, *Zaprionus tuberculatus* Malloch, 1932. Only one of the seven species, the species *Zaprionus tuberculatus*, has invasive capacities and has expanded its geographical range to the palaearctic region during the last four decades (Yassin 2008). The seven species can be distinguished on the basis of morphological characters of male and female genitalia and internal reproductive organs as well as immature stages (Tsacas et al. 1977; Yassin & David 2010). Of these characters, male and female genitalia provide the best diagnoses on non-living specimens, with DNA barcoding being of limited utility due to shared mitochondrial sequences among closely-related species (Yassin 2008). Dissection of a sample of captured specimens from Salses-le-Château supported the morphological identification of the introduced flies as *Zaprionus tuberculatus*. Female spermathecae (Fig. 3A) have predominantly smooth surface except on the tip wherein the surface becomes slightly rigged, whereas the phallosome of the male aedeagi (Fig. 3B) has a conspicuous ventral process and a densely toothed flap surrounding its border. Individuals are now kept in 70% alcohol and are available upon request from R.G. and H.C. (see authors affiliation for address).




FIGURE 2 *Zaprionus tuberculatus*: dorsal view (A), lateral view (B), detail of the protruding tubercle on the forefemur of a male (C) and detail of the ovipositor of a female (D).

Table 1 Occurrence of *Zaprionus tuberculatus* collected in France. Number of male and female *Z. tuberculatus* and all other Drosophilidae trapped in a fig orchard and a hedgerow nearby during 1-year monitoring in 2022–23. Temperatures indicated are monthly air average, minimum (min) and maximum (max) of average daily temperature recorded every 15 min by TMS-4 dataloggers (TOMST, Czech Republic) at 15 cm above the ground surface

Date collected	Fig orchard			Daily temperature (°C) average; min—max	Hedgerow			
	<i>Zaprionus tuberculatus</i>		Other Drosophilidae		<i>Zaprionus tuberculatus</i>		Other Drosophilidae	
	Males	Females			Males	Females		
March 2022	0	0	36	11.8; 8.9–17.6	0	0	67	11.9; 8.9–17.6
April 2022	0	0	58	15.1; 6.5–22.2	0	0	33	15.1; 5.9–22.3
May 2022	0	0	1	20.9; 17.1–26.2	0	0	32	20.2; 17.3–25.6
June 2022	0	0	131	25.5; 21.3–30.3	0	0	172	24.4; 20.2–29.3
July 2022	0	0	650	28.7; 24.7–32.7	11	19	1,356	27.5; 23.2–32.1
August 2022	12	13	390	28.3; 24.0–33.4	13	15	479	27.7; 23.7–32.6
September 2022	0	0	5	22.4; 16.2–27.7	4	4	168	22.4; 16.3; 27.7
October 2022	4	2	227	19.5; 17.2–21.8	6	6	330	19.6; 17.3–21.9
November 2022	0	0	2	13.0; 7.6–19.0	0	0	81	13.6; 8.6–19.9
December 2022	0	0	0	9.2; 2.7–14.1	0	0	2	10.0; 4–15.3
January 2023	0	0	27	7.6; 3.0–12.8	2	1	24	8.1; 3–13.8
February 2023	0	0	71	8.9; 3.6–14.4	0	0	50	9.5; 3.5–15.2




FIGURE 3 Detail of female spermatheca (A) and male aedeagus (B) of trapped individuals of *Zaprionus tuberculatus* (afb: aedeagal flap border; avp: aedeagal ventral process).

Zaprionus tuberculatus is widespread in Africa and the islands of the Indian Ocean (Madagascar, Comoro, Seychelles, Réunion and Mauritius) and Atlantic Ocean (St Helena and Cap Verde) (Chassagnard 1988; Chassagnard & Kraaijeveld 1991; Tsacas 1980; Tsacas *et al.* 1977; Yassin & David 2010). At the end of 20th century, *Z. tuberculatus* was recorded in Palearctic region from Canary Islands, Egypt, Cyprus, Malta and Israel (Chassagnard & Kraaijeveld 1991; Ebejer 2001). More recently, *Z. tuberculatus* occurred in Turkey (Patlar *et al.* 2012), Italy (Raspi *et al.* 2014), Romania (Chireceanu *et al.* 2015), Tunisia (Ben Halima Kamel *et al.* 2020) and Brazil (Cavalcanti *et al.* 2022).

The first detection of *Z. tuberculatus* occurred in July 2022 in the hedgerow and captures continued until January 2023

with an interruption in November and December (Table 1). In the fig orchard, we trapped *Z. tuberculatus* in August 2022 and then in October 2022, corresponding respectively to the beginning and the end of figs harvest. These results suggest that *Z. tuberculatus* could remain in natural woody habitats and colonize orchards when fruits come to maturity. To verify whether this observation was the result of a temporary accidental introduction or an established population, we revisited the plantation the following year, in September 2023, to harvest 100 ripe and 100 overripe figs directly from the trees at different locations within the fig orchard. The collected fruits were placed on vermiculite layer in emergence cages (four cages with ripe figs and four cages with overripe figs). Cages were then placed in a quarantine

chamber regulated at 20°C and 70% RH to assess emergence for 1 month. Subsequently, we identified and counted the *Z. tuberculatus* individuals that emerged. Among the different fly species emerging from ripe and overripe figs (mainly *D. suzukii* and *Ceratitidis capitata*, not counted), we found 13 females and 10 males of *Z. tuberculatus*, but only from the overripe figs. The emergence of *Z. tuberculatus* occurred during the 2 weeks following the harvest. The presence of *Z. tuberculatus* for two consecutive years led us to posit that this invasive fly may have permanently established itself, at least at this specific location.

Compared to other first reports, our trapping results show a longer period of occurrence of *Z. tuberculatus* in newly colonized countries. In Turkey, the first records of *Z. tuberculatus* were reported to be in August (Patlar et al. 2012; Zengin 2020), whereas Raspi et al. (2014) in Italy and Constantina et al. (2015) in Romania reported that *Z. tuberculatus* flies were found in the traps only during autumn from September to October. To our knowledge, until now the latest captures were reported in mid-December in Turkey and Italy (Amiresmaeili et al. 2019; Zengin 2020), so this is the first time that adults of *Z. tuberculatus* are trapped in winter in Europe. During January, mean daily temperature recorded in the surveyed hedgerow was 8.1°C with a minimum of 3°C (Table 1). All daily temperature data for the air, ground surface or ground is available in Table S1. Several studies showed evidence of climatic niche shift in exotic species during their expansion (Hill et al. 2017; Medley 2010; Petersen 2013; Wei et al. 2017). Such niche shift was also demonstrated in other invasive Drosophilidae, *D. suzukii* (Ørsted & Ørsted 2019) and *Zaprionus indianus* (da Mata et al. 2010). Our results suggest that *Z. tuberculatus* may experience a climatic niche shift to accommodate to climate colder than in its tropical origin, as also suggested by Cavalcanti et al. (2022). Yet, additional monitoring of *Z. tuberculatus* in others areas and for several consecutive seasons are highly needed to substantiate this “niche shift” assumption. With climate projection indicating that global warming will result in a progression of Mediterranean climate to the north of France during the next decades (Dubreuil 2022), climatic conditions might become more suitable to a possible northward expansion of *Z. tuberculatus*.

The pest status of *Z. tuberculatus* and its potential risk for agriculture is not yet clear. In its native range, *Z. tuberculatus* develops on decaying fruits, particularly on fallen rotting figs (Lachaise et al. 1982) and was thus not considered as a pest. However, the spread of this species and the fact that it has successfully been reared on 49 different species of tropical fruits (Lachaise & Tsacas 1984) should be a point of vigilance. This is particularly important in view of the first emergences we have observed from overripe figs. Moreover, Balmès and Mouttet (2019) reported emergences of *Z. tuberculatus* from imported fruits of *Citrus sinensis* from South Africa and *Litchi sinensis* from Reunion.

Zaprionus indianus is an invasive species that is closely related to *Z. tuberculatus* (Yassin & David 2010). The pest behavior of *Z. indianus* should also be a warning point for the entry of *Z. tuberculatus* into the EU and France, as they may have similar nutritional ecology and invasion routes. In Brazil, *Z. indianus* was reported attacking figs and was responsible for 50% of fig losses of fruits that were still on trees (Dettler et al. 2021; Vilela et al. 1999). The entry in Central and South America and the damages to fig orchards (van der Linde et al. 2006; Vilela 1999) led to its inscription on the alert list of the European and Mediterranean Plant Protection Organization (EPPO) in 2016. Under laboratory conditions, Bernardi et al. (2017) observed that *Z. indianus* could oviposit in ripe strawberries and the larvae were then able to develop in the berries. Yet, the ability of *Z. indianus* to oviposit and generate offspring in healthy strawberry fruit was clearly facilitated by injuries caused by *D. suzukii* or by mechanical injuries (Bernardi et al. 2017). Although no specific report of damage by *Z. indianus* in the EU has been reported so far, considering the damage caused to figs in South America, *Z. indianus* establishment and spread was considered as a threat with possible large economic consequences in Europe (EFSA P on PH (PLH) et al. 2022). Consequently, in 2022, *Z. indianus* was considered by European Food Safety Authority as a potential Union quarantine pest (EFSA P on PH (PLH) et al. 2022). Our results suggest that a particular attention should also be paid on *Z. tuberculatus*. Indeed, we detected males and females *Z. tuberculatus* in a fig orchard during the harvest period where fruits are systematically removed before rotting as a prophylactic measure against *D. suzukii* (personal communication). This may suggest a potential capacity to develop in healthy fruits (to be confirmed). Indeed, *Zaprionus* flies can infest figs on trees via oviposition on ostiole base (EFSA P on PH (PLH) et al. 2022). Moreover, climatic conditions in many EU member states and host plant availability in those areas are conducive for establishment of both *Z. indianus* and *Z. tuberculatus*.

Our monitoring of *D. suzukii* enabled the first detection of *Z. tuberculatus* in southern France in summer 2022, at a singular site (a fig orchard and its adjacent hedgerow). This finding was reaffirmed the subsequent year, based on fruit samples collected in late summer 2023 in the same fig orchard. We will conduct additional sampling effort, over multiple years and at various locations to better characterize the phenology and formally confirm the establishment of this species in France. Our study also highlights a potential risk associated with the discovery of *Z. tuberculatus*, a new invasive drosophilid that may have potential economic consequences on fruit production. We are currently raising individuals collected on the invasive front in quarantine to carry out research into their ecology and biology in order to clarify their pest potential, particularly on figs, citrus fruit and strawberries.

Acknowledgments

We thank the Teaching Lab and Project Lab of the University of Rennes for their help in designing and manufacturing the traps. We thank the fig producer for the help in collecting the traps during the monitoring and Jeanne Begouen-Demeaux and Damien Gourmelon for their help in sorting the insects. We also thank the AVFF (Association pour la Valorisation de la Framboise Française) for setting up the network of producers involved in the monitoring. This work was supported by ANR Drothermal (ANR-20-CE02-011-01).

Conflict of interest statement

The authors declare that they have no conflict of interest.

References

- Amiresmaeli N, Jucker C, Savodelli S, Lupi D (2019) Can exotic drosophilids share the same niche of the invasive *Drosophila suzukii*? *Journal Of Entomological And Acarological Research* **51**: 7861. <https://doi.org/10.4081/jeur.2019.7861>.
- Bächli, G (2023) Taxodros: the database on taxonomy of Drosophilidae. <http://www.taxodros.unizh.ch/>
- Balmès V, Mouttet R (2019) The drosophilid risk on imports. *EPPO Bulletin* **49**: 122–126. <https://doi.org/10.1111/epp.12536>.
- Ben Halima Kamel M, Ben Cheik Z, Zouari S, Mdellel L, Balmès V (2020) First report of *Zaprionus indianus* and *Z. tuberculatus* (Diptera: Drosophilidae) in Tunisia. *EPPO Bulletin* **50**: 307–310. <https://doi.org/10.1111/epp.12662>.
- Bernardi D, Andreazza F, Botton M, Baronio CA, Nava DE (2017) Susceptibility and Interactions of *Drosophila suzukii* and *Zaprionus indianus* (Diptera: Drosophilidae) in Damaging Strawberry. *Neotropical Entomology* **46**: 1–7. <https://doi.org/10.1007/s13744-016-0423-9>.
- Briem F, Dominic AR, Golla B *et al.* (2018) Explorative data analysis of *Drosophila suzukii* trap catches from a seven-year monitoring program in southwest Germany. *Insects* **9**: 125. <https://doi.org/10.3390/INSECTS9040125>.
- Carson HL (1971) *The ecology of drosophila breeding sites*. University of Hawaii Foundation Lyon Arboretum Fund.
- Cavalcanti FAGS, Ribeiro LB, Marins G *et al.* (2022) Geographic expansion of an invasive fly: first record of *Zaprionus tuberculatus* (Diptera: Drosophilidae) in the Americas. *Annals of the Entomological Society of America* **115**: 267–274. <https://doi.org/10.1093/aesa/saab052>.
- Chassagnard M & Kraaijeveld AR (1991) The occurrence of *Zaprionus sensu stricto* in the palearctic region (Diptera: Drosophilidae), *Annales de la Société entomologique de France* (N.S.), **27**(4): 495–496. <https://doi.org/10.1080/21686351.1991.12277851>
- Chassagnard MT (1988) Esquisse phylogénétique du genre *Zaprionus* Coq. (Diptera: Drosophilidae) et description de trois nouvelles espèces afrotropicales. *Naturaliste Canadien (Revue d'Ecologie et de Systématique)* **115**: 305–322.
- Chireceanu C, Chiriloaie A, Teodoru A (2015) First record of the spotted wing drosophila *Drosophila suzukii* (Diptera: Drosophilidae) in Romania. *Romanian Journal for Plant Protection* **8**: 86–95.
- Constantina C, Teodoru A, Chiriloaie A (2015) The first detection of fruit fly *Zaprionus tuberculatus* Malloch (Diptera: Drosophilidae) in the eastern part of Europe (Romania). *Annals of the University of Craiova* **20**: 377–382.
- da Mata RA, Tidon R, Côrtes LG, De Marco P, Diniz-Filho JAF (2010) Invasive and flexible: niche shift in the drosophilid *Zaprionus indianus* (Insecta, Diptera). *Biological Invasions* **12**: 1231–1241. <https://doi.org/10.1007/s10530-009-9542-0>.
- de Toni DC, Gottschalk MS, Cordeiro J, Hofmann PPR, Valente VLS (2007) Study of the Drosophilidae (Diptera) communities on Atlantic Forest islands of Santa Catarina State, Brazil. *Neotropical Entomology* **36**: 356–375. <https://doi.org/10.1590/S1519-566X2007000300004>.
- Delbac L, Rusch A, Thiery D (2020) Temporal dynamics of *Drosophila suzukii* in vineyard landscapes. *Entomologia Generalis* **40**: 285–295. <https://doi.org/10.1127/entomologia/2020/0858>.
- Dettler MA, Barrientos GN, Martinez E *et al.* (2021) Field infestation level of *Zaprionus indianus* and *Drosophila suzukii* (Diptera: Drosophilidae) in *Ficus carica* (Rosales: Moraceae) and *Rubus idaeus* (Rosales: Rosaceae) in the Northeast of Buenos Aires province. *Revista De La Sociedad Entomologica Argentina* **80**: 43–47. <https://doi.org/10.25085/rsea.800307>.
- Dubreuil V (2022) Le changement climatique en France illustré par la classification de Köppen. *La Météorologie* **116**: 37–47. <https://doi.org/10.37053/lameteorologie-2022-0012>.
- Ebejer MJ (2001) A contribution to the knowledge of the Ephydroidea (Diptera : Camillidae, Campichoetidae and Drosophilidae) of Malta. <https://www.um.edu.mt/library/oar/handle/123456789/14943> (Accessed: 11 July 2023)
- EFSA P on PH (PLH), Bragard C, Baptista P *et al.* (2022) Pest categorisation of *Zaprionus indianus*. *EFSA Journal* **20**: e07144. <https://doi.org/10.2903/j.efsa.2022.7144>.
- Geiger, R (1954) Klassifikation der Klimate nach W. Köppen, in Springer (eds) *Landolt-Börnstein—Zahlenwerte und Funktionen aus Physik, Chemie, Astronomie, Geophysik und Technik, alte Serie*, pp. 603–607. Springer, Berlin.
- Hamby KA, Bolda MP, Sheehan ME, Zalom FG (2014) Seasonal Monitoring for *Drosophila suzukii* (Diptera: Drosophilidae) in California Commercial Raspberries. *Environmental Entomology* **43**: 1008–1018. <https://doi.org/10.1603/EN13245>.
- Harris DW, Hamby KA, Wilson HE, Zalom FG (2014) Seasonal monitoring of *Drosophila suzukii* (Diptera: Drosophilidae) in a mixed fruit production system. *Journal of Asia-Pacific Entomology* **17**: 857–864. <https://doi.org/10.1016/j.aspen.2014.08.006>.
- Hill MP, Gallardo B, Terblanche JS (2017) A global assessment of climatic niche shifts and human influence in insect invasions.

- Global Ecology and Biogeography* **26**: 679–689. <https://doi.org/10.1111/geb.12578>.
- Kottek M, Grieser J, Beck C, Rudolf B, Rubel F (2006) World Map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift* **15**: 259–263. <https://doi.org/10.1127/0941-2948/2006/0130>.
- Kremmer L, Thaon M, Borowiec N *et al.* (2017) Field monitoring of *Drosophila suzukii* and associated communities in South Eastern France as a pre-requisite for classical biological control. *Insects* **8**: 124. <https://doi.org/10.3390/INSECTS8040124>.
- Lachaise D, Tsacas L (1984) Breeding sites in tropical African Drosophilids. In: Ashburner M, Carson HL, Thompson JN (eds) *The genetics and biology of Drosophila*, pp 221–332. Academic Press.
- Lachaise D, Tsacas L, Couturier G (1982) The Drosophilidae associated with tropical African figs. *Evolution* **36**: 141–151. <https://doi.org/10.2307/2407976>.
- Markow TA, O'Grady P (2008) Reproductive ecology of *Drosophila*. *Functional Ecology* **22**: 747–759. <https://doi.org/10.1111/j.1365-2435.2008.01457.x>.
- Mateus RP, Buschini MLT, Sene FM (2006) The *Drosophila* community in xerophytic vegetations of the upper Parana-Paraguay river basin. *Brazilian Journal of Biology* **66**: 719–729. <https://doi.org/10.1590/S1519-69842006000400016>.
- Medley KA (2010) Niche shifts during the global invasion of the Asian tiger mosquito, *Aedes albopictus* Skuse (Culicidae), revealed by reciprocal distribution models: niche shifts and global invasion. *Global Ecology and Biogeography* **19**: 122–133. <https://doi.org/10.1111/j.1466-8238.2009.00497.x>.
- Ørsted IV, Ørsted M (2019) Species distribution models of the spotted wing *Drosophila* (*Drosophila suzukii*, Diptera: Drosophilidae) in its native and invasive range reveal an ecological niche shift. *Journal of Applied Ecology* **56**: 423–435. <https://doi.org/10.1111/1365-2664.13285>.
- Patlar B, Koc B, Yilmaz M, Ozsoy ED (2012) First records of *Zaprionus tuberculatus* (Diptera: Drosophilidae) from the Mediterranean Region, Turkey. *Drosophila Information Service* **95**: 94–96.
- Petersen MJ (2013) Evidence of a climatic niche shift following North American introductions of two crane flies (Diptera; genus *Tipula*). *Biological Invasions* **15**: 885–897. <https://doi.org/10.1007/s10530-012-0337-3>.
- Raspi A, Grassi A, Benelli G (2014) *Zaprionus tuberculatus* (Diptera Drosophilidae): first records from the European mainland. *Bulletin of Insectology* **67**: 157–160.
- Tait G, Mermer S, Stockton D *et al.* (2021) *Drosophila suzukii* (Diptera: Drosophilidae): a decade of research towards a sustainable integrated pest management program. *Journal of Economic Entomology* **114**: 1950–1974. <https://doi.org/10.1093/JEE/TOAB158>.
- Tsacas L (1980) Catalogue of the Diptera of the Afrotropical region. In: Crosskey RW (ed) *Family: Drosophilidae*, pp 673–685. British Museum (Natural History), London.
- Tsacas L, David J, Allemand R, Pasteur G, Chassagnard MT, & Derridj S (1977) Biologie Évolutive du Genre *Zaprionus* Recherches sur le Complexe Spécifique de *Z. Tuberculatus* [Dipt. Drosophilidae], Annales de la Société entomologique de France (N.S.), **13**(2): 391–415. <https://doi.org/10.1080/21686351.1977.12278641>
- van der Linde K, Steck GJ, Hibbard K, Birdsley JS, Alonso LM, Houle D (2006) First records of *Zaprionus indianus* (Diptera: Drosophilidae), a pest species on commercial fruits from Panama and the United States of America. *Florida Entomologist* **89**: 402–404.
- Vilela C (1999) Is *Zaprionus indianus* Gupta, 1970 (Diptera, Drosophilidae) currently colonizing the Neotropical region? *Drosophila Information Service* **82**: 37–39.
- Vilela C, Teixeira E, Stein C (1999) Nova praga nos figos: *Zaprionus indianus* Gupta, 1970. *Informativo da Sociedade Entomológica do Brasil* **21**: 2.
- Wei J, Zhang H, Zhao W, Zhao Q (2017) Niche shifts and the potential distribution of *Phenacoccus solenopsis* (Hemiptera: Pseudococcidae) under climate change. *PLoS ONE* **12**: e0180913. <https://doi.org/10.1371/journal.pone.0180913>.
- Yassin A (2008) Molecular and morphometrical revision of the *Zaprionus tuberculatus* species subgroup (Diptera: Drosophilidae), with descriptions of two cryptic species. *Annals of the Entomological Society of America* **101**: 978–988. <https://doi.org/10.1603/0013-8746-101.6.978>.
- Yassin A, David J (2010) Revision of the Afrotropical species of *Zaprionus* (Diptera, Drosophilidae), with descriptions of two new species and notes on internal reproductive structures and immature stages. *ZooKeys* **51**: 33–72. <https://doi.org/10.3897/zookeys.51.380>.
- Zengin E (2020) Occurrence of invasive species and seasonal dynamics of fruit flies (Diptera: Drosophilidae) species in Uşak province, Turkey. *Revista de la Sociedad Entomológica Argentina* **79**: 21–30. <https://doi.org/10.25085/rsea.790104>.

Supporting information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Figure S1. Homemade trap used for the *Drosophila suzukii* monitoring (DroFramb action within the framework of ANR Drothermal project, <https://www.drothermal.cnrs.fr/>).

Table S1. Temperature data collected by TMS-4 Standard dataloggers (TOMST®, Czech Republic) in the fig orchard and the hedgerow nearby. Indicated values are average \pm standard deviation; minimum - maximum of daily temperature recorded every 15 minutes at 6 cm into the ground (T_ground), at the surface (T_surface) and 15 cm above the ground surface (T_air).